

$$\begin{aligned}
& -k_{10} (\zeta_1(t) - \zeta_2(t)) + k_{10} (\zeta_2(t) - \zeta_3(t)) + \\
& 2 m_0 \left( \cos(\theta(t)) R(t) + \cos(\alpha(t)) \cos(\theta(t) + \psi(t)) \left( \frac{7 L_0}{10} + \eta_2(t) + \eta_3(t) + \eta_4(t) + \eta_5(t) \right) \right) \zeta_2'(t) \\
& \left( \cos(\theta(t)) R'(t) - \cos(\theta(t) + \psi(t)) \sin(\alpha(t)) \left( \frac{7 L_0}{10} + \eta_2(t) + \eta_3(t) + \eta_4(t) + \eta_5(t) \right) \alpha'(t) - \right. \\
& \quad \left. R(t) \sin(\theta(t)) \theta'(t) - \cos(\alpha(t)) \sin(\theta(t) + \psi(t)) \left( \frac{7 L_0}{10} + \eta_2(t) + \eta_3(t) + \eta_4(t) + \eta_5(t) \right) \right. \\
& \quad \left. (\theta'(t) + \psi'(t)) + \cos(\alpha(t)) \cos(\theta(t) + \psi(t)) (\eta_2'(t) + \eta_3'(t) + \eta_4'(t) + \eta_5'(t)) \right) + \\
& \left( m_0 \left( \cos(\theta(t)) R(t) + \cos(\alpha(t)) \cos(\theta(t) + \psi(t)) \left( \frac{7 L_0}{10} + \eta_2(t) + \eta_3(t) + \eta_4(t) + \eta_5(t) \right) \right) \right)^2 + i_{\text{xm2}} \Big) \\
& \zeta_2''(t) = c_{10} (\zeta_2'(t) - \zeta_3'(t)) - c_{10} (\zeta_3'(t) - \zeta_4'(t))
\end{aligned}$$