

$$\begin{aligned}
& -k_{10} (\zeta_7(t) - \zeta_8(t)) + k_{10} (\zeta_8(t) - \zeta_9(t)) + \\
& 2 m_0 \left(\cos(\theta(t)) R(t) - \cos(\alpha(t)) \cos(\theta(t) + \psi(t)) \left(\frac{L_0}{2} + \eta_6(t) + \eta_7(t) + \eta_8(t) \right) \right) \zeta_8'(t) \\
& \left(\cos(\theta(t)) R'(t) + \cos(\theta(t) + \psi(t)) \sin(\alpha(t)) \left(\frac{L_0}{2} + \eta_6(t) + \eta_7(t) + \eta_8(t) \right) \alpha'(t) - \right. \\
& \quad \left. R(t) \sin(\theta(t)) \theta'(t) + \cos(\alpha(t)) \sin(\theta(t) + \psi(t)) \left(\frac{L_0}{2} + \eta_6(t) + \eta_7(t) + \eta_8(t) \right) (\theta'(t) + \psi'(t)) - \right. \\
& \quad \left. \cos(\alpha(t)) \cos(\theta(t) + \psi(t)) (\eta_6'(t) + \eta_7'(t) + \eta_8'(t)) \right) + \\
& \left(m_0 \left(\cos(\theta(t)) R(t) - \cos(\alpha(t)) \cos(\theta(t) + \psi(t)) \left(\frac{L_0}{2} + \eta_6(t) + \eta_7(t) + \eta_8(t) \right) \right)^2 + i_{\text{xm8}} \right) \zeta_8''(t) = \\
& c_{10} (\zeta_7'(t) - \zeta_8'(t)) - c_{10} (\zeta_8'(t) - \zeta_9'(t))
\end{aligned}$$