

https://theses.gla.ac.uk/

Theses Digitisation:

https://www.gla.ac.uk/myglasgow/research/enlighten/theses/digitisation/

This is a digitised version of the original print thesis.

Copyright and moral rights for this work are retained by the author

A copy can be downloaded for personal non-commercial research or study, without prior permission or charge

This work cannot be reproduced or quoted extensively from without first obtaining permission in writing from the author

The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the author

When referring to this work, full bibliographic details including the author, title, awarding institution and date of the thesis must be given

Enlighten: Theses <u>https://theses.gla.ac.uk/</u> research-enlighten@glasgow.ac.uk

STUDIES IN BACTERIAL ADHERENCE TO CANINE AND FELINE CORNEOCYTES

Bу

YI-FANG LU

Thesis submitted for the degree of Master of Veterinary Medicine in the Faculty of Veterinary Medicine, University of Glasgow

©Yi-Fang Lu

ProQuest Number: 10390719

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



ProQuest 10390719

Published by ProQuest LLC (2017). Copyright of the Dissertation is held by the Author.

All rights reserved. This work is protected against unauthorized copying under Title 17, United States Code Microform Edition © ProQuest LLC.

> ProQuest LLC. 789 East Eisenhower Parkway P.O. Box 1346 Ann Arbor, MI 48106 – 1346



TABLE OF CONTENTS

Page

Table of contents		i
Abstract		li
Publications from Thesis		iii
Contents		iv
List of Tables	•••••	viii
List of Figures		ix
List of Appendices		xi
Acknowledgements	•••••••	xii
Author's Declaration		xiii
Chapter 1		1
Chapter 2	•••••	31
Chapter 3	•••••	66
Appendices		111
References		136

1. A

ABSTRACT

In the first chapter of this thesis a three-year retrospective survey showed 41% of dogs referred to the University of Glasgow Veterinary School for investigation of skin disease were diagnosed as suffering from pyoderma with 62% of these cases being associated with atopic dermatitis. In contrast, pyoderma was rarely diagnosed in the cat. The study confirmed the importance of canine pyoderma and its common association with atopic dermatitis. Although canine pyoderma due to *Staphylococcus intermedius* is common, it is poorly understood. Adhesion by staphylococci has been suggested as a potentially important factor in the establishment of infection. As many assays to quantify adhesion by bacteria are both tedious and time consuming this has hampered studies in this interesting area.

Chapters two and three detail the development of a simple adhesion assay to quickly quantify bacterial adhesion to canine and feline corneocytes. As far as the author can determine there is no published data on the size and surface area of canine and feline corneocytes therefore morphometric studies were undertaken in chapter 2. The mean canine and feline corneocyte diameter and surface area were found to be approximately 38-48 μ m and 1100-1800 μ m² respectively. Of three body sites examined the inner aspect of the canine and feline pinna proved to be most suitable for the collection of corneocytes for an adhesion assay.

In chapter 3 a simple adhesion assay is described. Corneocytes were collected from the inner aspect of the pinna using adhesive discs (D-squame® discs), and incubated in a moist chamber with bacteria under test. Image analysis software employing a script (macro) written by the author was used to count adherent bacteria. Adhesion by bacteria was found to be dependent on time, temperature, and bacterial concentration. The assay was both repeatable and reproducible. Using the adhesion assay so developed the pathogenic staphylococci (*S. aureus*, *S. hyicus* and *S. intermedius*) all adhered well to canine and feline corneocytes. *S. hominus* and a Micrococus species adhered poorly.

ii

PUBLICATIONS AND PRESENTATIONS ARISING FROM THIS THESIS

PROCEEDINGS AND PRESENTATIONS

Lu, Y. and McEwan, N.A. (2003). Morphometric analysis of canine and feline corneocytes. British Veterinary Dermatology Study Group Proceedings. Spring meeting. 2nd April 2003. 49-50.

<u>Lu</u>, Y. and McEwan, N.A. (2003). An assay for the rapid quantification of bacterial adherence to corneocytes from the dog and cat. Proceedings of the 19th Annual Congress of the ESVD-ECVD. Tenerife, Spain $4^{th} - 5^{th}$ September 2003. p. 141.

INDEX OF CHAPTER CONTENTS

CHAPTER 1	Page
A Three Year Retrospective Study Of The	1
Prevalence Of Canine And Feline Skin	
Diseases With Special Reference To	
Pyoderma In A University Referral Hospital	
Population	
LITERATURE REVIEW	2
Introduction	2
Canine and feline skin flora	2
Pathogenic skin microorganisms	4
Canine and feline pyoderma	5
The stratum corneum as a barrier	6
Predisposing factors for infection	7
Conclusions	8
INTRODUCTION	9
AIMS	9
MATERIAL AND METHODS	9
Dermatology cases	9
General hospital population	10
Statistical analysis	10
RESULTS	11
Retrospective study of canine and feline	11
skin disease referred to the University of	
Glasgow Veterinary School (2000-2002)	
Canine skin disease	11
Signalment	11
Canine skin diseases diagnosed	13
General canine hospital population at	17
UGVS (2000-2002)	
	19
Feline skin disease population	
Signalment	19
Feline skin diseases diagnosed	21
General feline hospital population at UGVS	23
(2000-2002)	

iv

DISCUSSION	26
Canine skin disease	26
Feline skin disease	28
Conclusions	29
CHAPTER 2	
A Study Of Canine And Feline Corneocyte	31
Morphometrics	
LITERATURE REVIEW	32
Introduction	32
Epidermal function	32
Keratinocyte differentiation	34
Corneocyte morphometrics	35
Collection of corneocytes and their study	37
INTRODUCTION	39
AIMS	39
MATERIAL AND METHODS	39
STUDY 1. CORNEOCYTE	39
MORPHOMETRICS	
Populations studied	39
Body sites sampled	40
Corneocyte collection	42
Corneocyte staining and image acquisition	43
Calibration and measurements	43
STUDY 2. COMPARISON OF	45
CORNEOCYTES FROM HAIRED AND	
NON-HAIRED SKIN	
Corneocyte collection	45
Counting method	46
STATISTICAL ANALYSIS	47
RESULTS	48
Corneocyte morphometrics	48
Box plots of corneocyte morphometrics	51
Comparison of corneocytes from haired	60
and nonhaired sites	
DISCUSSION	64

CHAPTER 3

An Adhesion Assay To Quantify Bacterial	66
Adherence To Canine And Feline	
Corneocytes	
LITERATURE REVIEW	67
Introduction	67
Mechanisms of bacterial adherence and	68
infection	
Methods of assessing bacterial adherence	70
Staphylococcal adherence in human and	71
canine atopic dermatitis	
AIMS	74
MATERIAL AND METHODS	74
Populations studied	74
Corneocyte collection	74
Bacteria	75
Adhesion assay	76
Image acquisition and bacteria counting	78
Study 1. The effects of time, temperature	84
and bacterial concentration on adhesion	
Time study	84
Temperature study	84
Concentration study	85
Study 2. Comparison of visual counting	85
and counting by image analysis	
Study 3. Repeatability study	86
Study 4. A comparison of corneocytes	86
collected from the ear and groin	
Study 5. Comparison of adherence by 5	87
different staphylococcal species and a	
Micrococcus to canine and feline	
corneocytes	
STATISTICAL ANALYSIS	88
RESULTS	
Effect on time on adhesion to corneocytes	89
(table 18 and figure 25)	
Comparison of adherence at 38°C and 4°C	91
(table 19 and figure 27)	
Effects of bacterial concentration on	93

vi

adhesion to corneocytes (table 20, figures	
28 and 29)	
Effects of bacterial concentration on	95
adhesion to corneocytes (table 20, figures	
28 and 29)	
Repeatability of counting method (table 22,	97
figure 33)	
Comparison of ear and groin corneocytes	9 8
(table 23, figure 34)	
Comparison of bacterial adhesion to	101
consecutive sampling from ear (table 24)	
Adhesion by different bacteria species to	102
canine and feline corneocytes (table 25,	
figure 35)	
DISCUSSION	105
SUMMARY	1 10

List of Tables

.

Table		Page
1	Most common breeds referred for skin disease (2000-2002)	12
2	Summary of the skin diseases diagnosed over a 3-year period at the UGVS (2000-2002)	14
3	Secondary bacterial and <i>Malassezia</i> infections associated with primary diagnoses	16
4	Most common breeds of general hospital (GH) and skin disease (SD) populations	19
5	Cat breeds referred for skin disease (2000-2002)	20
6	Summary of the skin diseases diagnosed over a 3-year period at the UGVS (2000-2002)	22
7	Most common cat breeds of general hospital (GH) and skin disease (SD) populations	25
8	Summary results of corneocyte diameter	49
9	Summary results of corneocyte surface area	50
10a	Corneocyte diameter ears	52
10b	Corneocyte diameter ears	52
10c	Corneocyte diameter ears	53
1 1 a	Corneocyte diameter thorax	53
11b	Corneocyte diameter thorax	54
12a	Corneocyte diameter groin	54
12b	Corneocyte diameter groin	55
12c	Corneocyte diameter groin	55
13a	Corneocyte surface area ears	56
13b	Corneocyte surface area ears	5 6
13c	Corneocyte surface area ears	57
14a	Corneocyte surface area thorax	57
14b	Corneocyte surface area thorax	58
15a	Corneocyte surface area groin	58
15b	Corneocyte surface area groin	59
15c	Corneocyte surface area groin	59
16	Canine corneocytes from haired area to unhaired area	60
17	Feline corneocytes from haired area to unhaired	62

area		
Effect on	time	on

.

18	Effect on time on adhesion to corneocytes	89
19	Comparison of adherence at 38°C and 4°C	91
20	Effects of bacteria concentration on adhesion to corneocytes	93
21	Comparison of bacteria counts by image analysis and visual counting	95
22	Repeatability of counting method	97
23	Comparison of ear and groin corneocytes	100
24	Comparison of bacterial adhesion to consecutive sampling from ear	101
25	Adhesion by different bacteria species to canine and feline corneocytes	102

List of Figures

Figure		Page
1	Most common breeds referred for skin	12
	disease (2000-2002)	
2	The most common skin diseases diagnosed	15
	at the UGVS (2000 - 2002)	
3	Most common dog breeds in general hospital	18
	population (2000-2002)	
4	Breeds referred for skin disease (2000-2002)	20
5	The most common feline skin diseases	23
	diagnosed at the UGVS (2000 - 2002)	
6	Most common cat breeds in general hospital	24
	population (2000-2002)	
7	Ear sampling area	41
8	Groin sampling area	41
9	Thoracic sampling area	42
10	Type I corneocytes from canine thorax	44
11	Type II corneocytes from canine ear	44
12	Comparison of haired and non-haired skin	46
13a	Ear diameter	51
13b	Ear surface area	51

ix

14a	Thoracic diameter	51
14b	Thoracic surface area	51
15a	Groin diameter	51
15b	Groin surface area	51
16	Histogram of numbers of Type I and type II	61
	corneocytes collected from canine ear	
17	Histogram of numbers of Type I and type II	63
	corneocytes collected from feline ear	
18	Moist chamber within water bath	77
19	D-Squame® discs and bacteria suspension	77
	prepared for incubation	
20	Slides after washing and staining	78
21	Selection of the counting area	80
22	Image thresholding	81
23	Application of a round filter	82
24	The final count	83
25	Effect on time on bacterial adhesion	90
26	Effect of time on bacterial adhesion –	90
	individual animals	
27	Histogram of temperature effects	92
28	Concentration of bacteria and adherence	94
29	Concentration curves showing clumping	94
30	Comparison of image analysis counting with manual count	96
31	Bland-Altman Image analysis and Operator	96
	1:Difference vs average	
32	Bland-Altman Image analysis and Operator	96
	2:Difference vs average	
33	Comparison of count 1 and 2	98
34	Linear regression Ear vs Groin R=0.976	99
35	Histogram of adherence by different bacteria	103
	species to canine and feline corneocytes	
36	Adherence shown by different staphylococcal species	104

List of Appendices

Appendix		Page
A	Dog breeds examined for investigation of skin disease at the UGVS 2000-2002	111
В	Main diagnosis of skin disease seen in dogs referred to the UGVS 2000-2002	112
С	Cats referred for investigation of skin disease at the UGVS 2000-2002	121
D	Normal dogs corneocyte diameter (µm)	122
E	Normal dogs corneocyte surface area (μm²)	124
F	Atopic dogs corneocyte diameter (µm)	126
G	Atopic dogs corneocyte surface area (μm^2)	128
Н	Normal cats corneocyte diameter (µm)	130
I	Normal cats corneocyte surface area (μm^2)	132
J	Scripts used in image analysis studies	134

xi

à

ACKNOWLEDGEMENTS

I would like to thank Professor David Taylor for his help and the use of the facilities of the Bacteriology section of the Department of Veterinary Pathology. For Miss Kathleen Reynolds who guided me and for Dr. Dominic Mellor who advised on the statistical methods. I am indebted to Dr. Neil McEwan for his great help, patience, and kindness in the preparation of this thesis.

Thanks go to Professor Martin Sullivan and the department of veterinary clinical studies for support of using facilities. Thanks to Leo Animal Health who kindly provided sponsorship to Spain to present my studies.

Also thank my friends, Aik-Choon Tan, Bruce Huang, Chun-Mei Lin, Diana Lin, En-I Chiu, Kelly Chang, Ken Liu, Mariana Fontan, Mrs. Hseuh, and Shi Wui Lou for giving support during this period. For the help of Dr. Wu and his wife. Especially for Chun-Kit Chan, for his patience and encouragement.

Lastly I would like to thank my parents for supported me through financial aid in my studies and their advice during the preparation and study for this thesis. For the kind support from my two brothers, Chi-Han and Chi-Yan. The work presented in this thesis was carried out by the author in the Department of Veterinary Clinical Studies, University of Glasgow.

Signed:

YI-FANG LU BVM BSc

Date: 18, JUNE, 2004

CHAPTER 1

A Three Year Retrospective Study Of The Prevalence Of Canine And Feline Skin Diseases With Special Reference To Pyoderma

Ś

LITERATURE REVIEW

Introduction

The epidermis of mammalian skin forms a protective barrier which provides physical, chemical, and microbial defence (Lloyd, 1976; Roth and James, 1989; Hill and Moriello, 1994). The outer most layer of the epidermis, the stratum corneum has been shown to be an important barrier to the penetration of micro-organisms and their metabolites from the skin surface (Sulzburger, 1960; Lai a Fat and Van Furth, 1974; Lloyd and Jenkinson, 1980). Hair forms the first line of physical defence which protects against the contact of pathogens with the skin, but it may also harbour bacteria, especially staphylococci (White, Ihrke, Stannard, Sousa, Reinke, Rosser, Jr. and Jang, 1983; Cox, Hoskins, Newman, Turnwald, Foil, Roy and Kearney, 1985; Allaker, Lloyd and Simpson, 1992; Harvey, 1994; Harvey and Lloyd, 1995).

Canine and feline skin flora

The normal skin microflora contributes to skin defence mechanisms. Bacteria are located in the superficial epidermis and in the infundibulum of the hair follicles, where sweat and sebum provide nutrients (Lloyd, 1980a; Scott, 1992). The normal flora is a mixture of bacteria that live in symbiosis, probably exchanging growth factors. The flora may change with different cutaneous environments. These are affected by factors such as heat, pH, salinity, moisture, albumin level, and fatty acid level (Mason, 1996; Chesney, 1997). The close relationship between the host and microorganisms enables bacteria to occupy microbial niches and to inhibit colonization by invading organisms. In addition, many bacteria (*Bacillus* spp., *Streptococcus* spp., and *Staphylococcus* spp.) are capable of producing antibiotic substances, and

3

some bacteria can produce enzymes (e.g. β -lactamase) that inhibit antibiotics (Saijonmaa-Koulumies and Lloyd, 1995). Bacteria that are isolated from the skin are commonly divided into three categories: resident organisms, transient organisms, and common pathogenic organisms (nomads) (Price, 1938; Somerville-Millar and Noble, 1974; Scott, Miller and Griffin, 2001b). Resident bacterial organisms can be routinely and repeatedly isolated and cultured from the surface of the skin or haircoat without causing clinical disease. Transient bacteria may on occasion be cultured from the skin but are of no significance unless they become involved in a pathologic process as secondary invaders. However, there is still controversy as to which microorganisms are truly residents on the canine skin. Studies describing the canine 'resident bacterial flora' have been based on short-term studies that fail to describe the dynamic situation of the microflora on the skin and thus do not provide evidence of the persistence of the individual bacterial species (Saijonmaa-Koulumies and Lloyd, 1996). In addition, both methods and anatomical sites employed for sampling have varied considerably in different studies (Krogh and Kristensen, 1976; Ihrke, Schwartzman, McGinley, Horwitz and Marples, 1978; Allaker, Lloyd and Simpson, 1992; Harvey and Lloyd, 1994; Harvey and Lloyd, 1995). Transient bacteria (e.g. Escherichia coli, Proteus mirabilis, Pseudomonas spp., Corynebacterium spp., and Bacillus sp.) do not multiply on the skin of healthy dogs (Krogh and Kristensen, 1976; Kristensen and Krogh, 1978; Krogh and Kristensen, 1981). The resident bacteria of canine skin include coagulase-negative staphylococci (Staphylococcus epidermidis, S. cohnii, S. saprophyticus, S. hominis, S. haemolyticus, S. capitis, S. warneri, S. xylosus, S. simulans, and S. sciuri); coagulase-positive staphylococci (S. intermedius); Micrococcus spp.; alpha-hemolytic streptococci; Actinobacter sp., Clostridium spp. and Propionibacterium acnes (Cox, Hoskins, Roy, Newman and Luther, 1984; Cox, Hoskins, Newman, Foil, Turnwald and Roy, 1988; Woldehiwet and Jones, 1990; Lloyd and Allaker, 1991;

Harvey, 1993; Greene and Lammler, 1993; Harvey and Lloyd, 1995; Saijonmaa-Koulumies and Llovd, 1996; Scott, Miller and Griffin, 2001b). In cats, the resident flora includes *Micrococcus* spp.; coagulase-negative staphylococci, especially Staphylococcus simulans; α-hemolytic streptococci; and Acinetobacter spp. (Scott, 1980; Medleau, Long, Brown and Miller, 1986). Coagulase-positive staphylococci, including both Staphylococcus aureus and S. intermedius, are also commonly isolated from normal cat skin and should be considered residents (Devriese, Nzuambe and Godard, 1984; Cox et al, 1985; Medleau and Blue, 1988). In Krogh and Kristensen's study (1976), 50% of the cultures from cats were negative. Household cats have a higher frequency of isolation of coagulase-negative staphylococci (S. capitis, S. epidermidis, S. haemolyticus, S. hominis, S. sciuri, and S. warneri) and coagulasepositive staphylococci (S. aureus and S. intermedius) compared with the case in cattery cats, suggesting that these organisms may be transferred from humans (Scott, Miller and Griffin, 2001b).

4

Pathogenic skin microorganisms

Nomads are able to colonize and reproduce only for short periods. They are able to take advantage of changing conditions at the skin surface and, if such changes favour proliferation on a long-term basis, clinical infection may result (Lloyd, 1995). Primary pathogenic organisms are capable of tissue invasion and creating disease. They are usually coagulase-positive staphylococci (*S. intermedius*, *S. aureus*, *S. hyicus*), and in the dog *S. intermedius* is the single most common isolate from canine skin infections (Phillips, Jr. and Kloos, 1981; Berg, Wendell, Vogelweid and Fales, 1984; Biberstein, Jang and Hirsh, 1984; Cox, Newman, Roy and Hoskins, 1984; Ihrke, 1987). The pathogenesis of canine pyoderma is poorly understood. Various virulence factors can be produced by *S. intermedius* but studies have failed to correlate this with

pathogenicity (Greene and Lammler, 1993; Burkett and Frank, 1998). In the light of this, it has been suggested that host factors, such as poorly developed epidermal defences, are very important (Mason, 1999). Cats infrequently experience pyoderma but commonly have subcutaneous abscesses from biting wounds. Therefore, the mouth flora of the cat is an important factor, which includes Pasteurella multocida, β-hemolytic streptococci, Corvnebacterium spp., Actinomyces spp., Bacteroides spp., and Fusobacterium spp. (Scott, Miller and Griffin. 2001b). Staphylococcus felis is increasingly isolated from cats with otitis externa, subcutaneous abscesses, paronychia, and other skin infections (Igimi, Kawamura, Takahashi and Mmitsuoka, 1989; Aarestrup and Jacobeson, 1993; Patel, 2003).

Canine and feline pyoderma

Skin infections are traditionally classified as primary or secondary infections to reflect the absence or the presence of an underlying cause (Scott, Miller and Griffin, 2001b). It is generally accepted that most pyodermas occur as a sequel to other cutaneous, metabolic, or immunologic abnormalities (Scott, Miller and Griffin, 2001b), such as hypersensitivity, endocrinopathies, or defects in keratinisation which allow the pathogen to multiply and result in pyoderma (Ihrke, 1987; Hill and Moriello, 1994).

The importance of staphylococcal carriage on the skin, hair and mucous membranes of dogs and the role of staphylococci in canine pyoderma is well documented (Mason and Lloyd, 1989; Mason and Lloyd, 1990; Allaker, Lloyd and Simpson, 1992; Allaker, Lloyd and Bailey, 1992; Harvey and Lloyd, 1994; Saijonmaa-Koulumies and Lloyd, 1996). In dogs, pyoderma is commonly diagnosed (Scott and Paradis, 1990), while in cats, there are few reports on staphylococcal skin disease (Cox *et al*, 1985; Medleau and Blue, 1988; White, 1991; Espinola and

Lilenbaum, 1996; Patel, 2002). The few published clinical reports (Raychaudhuri and Raychaudhuri, 1993; Scott, Miller and Griffin, 2001b) have in the main identified underlying immunosuppressive conditions such as feline leukaemia virus (FeLv), feline immunodeficiency virus (FIV), and toxoplasmosis as predisposing factors. S. aureus and S. intermedius have been cultured from feline skin lesions (Medleau and Blue, 1988; White, 1991) and have been implicated as pathogens, however, their exact role in the pathogenesis of feline skin disease remains unknown. S. felis is a coagulase-negative species which was first identified from clinical specimens (Igimi et al, 1989) and has subsequently been implicated in the pathogenesis of otitis externa (Higgins and Gottchalk, 1991), paronychia (Aarestrup and Jacobeson, 1993) and skin disease (Patel, Lloyd, Howell and Noble, 2002). The most recent study of the prevalence of cutaneous staphylococci in feral cats, healthy cats and pet cats with skin lesions suggested that S. felis may be a resident on cat skin and S. intermedius may play a role in perpetuating skin disease in cats (Patel, 2002).

The stratum corneum as a barrier

The stratum corneum consists of tightly packed keratinised cells, which are permeated by an emulsion of sebum, sweat, and intercellular cement substance. The cells and emulsion function as an effective physical and chemical barrier to potential pathogens (Lloyd, 1980a; Berg *et al*, 1984; Mason and Lloyd, 1993; Saijonmaa-Koulumies and Lloyd, 1996). The intercellular material of the stratum corneum is likely to be an important component of this barrier. However, it would appear from studies that less is present in dog skin than has been reported in cattle and man. This may partly explain the high incidence of pyoderma in comparison with other species of domestic animal (Scott, 1988; Scott, Miller and Griffin, 2001b). In cattle and sheep, scanning electron

microscope studies have shown that the hair follicle infundibula are protected by layers of squames permeated with sebaceous lipid; this material fills the infundibula forming a raised conical mass (Lloyd, Dick and Jenkinson, 1979). In dogs, similar techniques have shown that smaller amounts of this material are present and the infundibula are seen as pits in the epidermal surface (Lloyd and Garthwaite, 1982). Examination of the skin surface topography in freeze-fracture specimens produced similar results (Mason and Lloyd, 1993). The effects of these differences are unknown but it is possible that the relatively unprotected hair follicles and thinner, more compact corneum of dogs are more permeable and therefore make them more susceptible to pyoderma. This hypothesis is supported by the results of previous studies (Mason and Lloyd, 1990) in which tritiated staphylococcal protein A was shown to pass through the canine epidermis and could be detected within the superficial dermis of normal dog skin 5 hours after topical application. In man, percutaneous absorption of protein A is impeded by the presence of an intact stratum corneum (White and Noble, 1980).

Predisposing factors for infection

The factors that may affect bacterial colonization on skin are environment (humidity and temperature) (Marples, 1965; Bibel and Lebrun, 1975; McBride, Duncan and Knox, 1977; Lloyd, 1980b; Ihrke, 1987; McBride, 1993), host factors (the hair coat and skin secretions) (McGinley, Leyden, Marples and Kligman, 1975; Ihrke *et al*, 1978; Lloyd and Garthwaite, 1982; Garthwaite, Lloyd and Thomsett, 1983; White *et al*, 1983; Allaker, Lloyd and Simpson, 1992; Mason and Lloyd, 1993; McEwan Jenkinson D., 1993; Harvey and Lloyd, 1994; Harvey and Lloyd, 1995; Scott, Miller and Griffin, 2001a), and bacterial factors (adherence and interaction between bacteria) (Murphy, 1975; Aly, Shinefield, Strauss and Maibach, 1977; Bibel, Aly, Shinefield, Maibach and Strauss, 1982;

Bibel, 1982; Feingold, 1986; Mason and Lloyd, 1989; McEwan, 1990; Romero-Steiner, Witek and Balish, 1990; Allaker and Noble, 1993; Aly and Bibel, 1993; Mason and Stewart, 1993; Harvey and Noble, 1994; Saijonmaa-Koulumies and Lloyd, 1995).

8

Bacterial adherence to host tissue is one of the prerequisite steps in the colonization and infection of the skin (Beachey, 1981; Aly and Bibel, 1993). It is a multifactorial process, and is influenced by host susceptibility and tissue tropism (Gibbons and van Houte, 1971; Aly *et al*, 1977). Both *S. aureus* and *S. intermedius* are known to adhere avidly to human (Ofek and Doyle, 1994; Lowy, 1998) and canine (McEwan, 2000) corneocytes respectively and this is likely to be a factor in the development of staphylococcal skin disease.

Conclusions

Canine bacterial pyoderma is one of the most common infections seen in small animal practice and *S. intermedius* appears to be involved in the majority cases (Berg *et al*, 1984). It is clear that bacterial skin disease is very common in the dog but apparently rare in the cat. In general cutaneous infections with *S. intermedius* are secondary to changes in the skin or are a complication of some other dermatosis. One factor in the development of infection may be the ability of *S. intermedius* to adhere to corneocytes. Recent studies have shown that *S. intermedius* adheres to canine corneocytes and that this may be important in the development of bacterial skin disease in canine atopic dermatitis (McEwan, 2000; Forsythe, Hill, Thoday and Brown, 2002; Saijonmaa-Koulumies and Lloyd, 2002).

INTRODUCTION

The literature review has shown bacterial skin disease to be common in the dog but rare in the cat. As a prelude to studies on bacterial adherence to canine and feline corneocytes a study to determine the frequency and types of skin disease within the referral clinic population at the University of Glasgow Veterinary School Hospital (UGVS) was conducted.

AIMS

- 1. To identify the prevalence of canine and feline skin disease over a three-year period by retrospective study of case records held by the UGVS Small Animal Hospital.
- 2. To identify cases of canine and feline pyoderma and any associated skin diseases referred to the UGVS over the three-year period.
- 3. To determine the breed, age and sex distributions (population denominators) of all dogs and cats referred to the UGVS Hospital over the same three-year period.
- 4. To identify any breed predisposition and disease trends by comparison of the study populations.

MATERIAL AND METHODS

Dermatology cases

Over a three-year period (2000 to 2002), all dogs and cats referred to the UGVS for investigation of skin disease were included in the study. Cases were referred mostly by veterinary surgeons in Glasgow and the surrounding areas but also from further afield, in Scotland and northern England. All cases were investigated by veterinary surgeons that hold the

Royal College of Veterinary Surgeons Diploma in Veterinary Dermatology. Cases underwent standard dermatological approaches to investigation and diagnosis as described by McEwan (2002a). Bacterial skin disease was diagnosed using the following criteria (McEwan, 2002b).

- 1. The presence of clinical findings characteristic of bacterial pyoderma
- II. Culture of a significant micro-organism
- III. A clinical response to antibiotic treatment

General hospital population

The computerised data base system (DataFlex 3.05b. Server edition. Data Access Corporation, Miami, Florida, USA) is used to record patient information at the UGVS. Information for the various dog breeds referred for the period 2000 to 2002 was obtained by interrogation of this data base system. This information so obtained was used to compare the breed frequency of skin disease during this period with that of the general referral hospital population.

Statistical analysis

Statistical analysis was carried out using computerised software GraphPad Prism, version 4 (GraphPad Software Inc. San Diego CA). A level of p < 0.05 was considered to indicate statistical significance. The Chi-Square test was used to compare sex and breed by analysing data in a 2 x 2 contingency table. The unpaired t-test was used to compare ages when two groups were compared. The Mann-Whitney rank sum test was employed when the data was drawn from a non-normal population.

RESULTS

Retrospective study of canine and feline skin disease referred to the University of Glasgow Veterinary School (2000-2002)

Canine skin disease

Signalment

During the 3 year period studied a total of 330 dogs were examined for skin disease. Of the 330 cases, 132 (40.00%) were entire male, 60 (18.18%) neutered male, 43 (13.03%) entire female and 94 (28.48%) neutered female. In 1 case the sex was not recorded. This gave a male to female ratio of 1.4: 1.

The mean age (\pm standard deviation) at examination was 4.01 \pm 2.81 years (minimum and maximum age: 0.33 and 14.00 years; median age 3.21 years). In 6 cases the age was not recorded.

The most common breeds examined were Cross breed (38), German Shepherd dog (34), West Highland White terrier (22), Labrador retriever (34), Boxer (17), Cocker spaniel (11), Golden retriever (9), Border collie (9) and Staffordshire bull terrier (8). Overall there were 73 different pure breed dogs but 9 breeds accounted for approximately half (42.47%) of the cases examined (Table 1; Figure 1; Appendix A).

Table 1. Most common breeds referred for skin disease (2000-2002)

Breed	Number	Percentage
Cross breed	38	12
German Shepherd dog	34	10
Labrador retriever	34	10
West Highland White terrier	22	7
Boxer	17	5
Cocker spaniel	11	3
Golden retriever	9	3
Border collie	9	3
Staffordshire bull terrier	8	2
Others	148	45

Figure 1. Most common breeds referred for skin disease (2000-2002)



Chapter I

Canine skin diseases dlagnosed

There were 340 primary skin diseases diagnosed in the 330 dogs examined. The most common skin condition diagnosed was atopic dermatitis (192 cases; 58.18 %). Other common skin disorders were: allergy not fully diagnosed, 24 cases (7.27%); primary bacterial pyoderma, 23 cases (6.97%); flea allergic dermatitis, 21 cases (6.36%), and parasitic dermatoses, 18 cases (5.45%). (Table 2, Figure 2, Appendix B).

Secondary infections were common. Superficial pyoderma was presented in 94 (28.5%) cases. Deep pyoderma was recorded in 18 (5.5%) cases. Deep and superficial pyoderma occurred together in 6 (0.2%) cases. There was a single primary *Malassezia* dermatitis case and secondary *Malassezia* dermatitis was identified in 44 (13.3%) cases.

Table 2. Summary of the skin disea	ises diagnosed over a 3-year
period at the UGVS	\$ (2000-2002)

D	Totals	
Allergy	Atopic dermatitis Flea allergic dermatitis Food sensitivity Allergy not diagnosed *	192 21 4 24
Immune mediated	Pemphigus foliaceus Pemphigus vulgaris Cutaneous lupus erythematosus Erythema multiforme	5 1 3 1
Parasitic	Cheyletiellosis Demodicosis Sarcoptic mange	6 8 4
Endocrine	Hyperadrenocorticism Hypothyroidism	1 2
Acquired alopecia	Follicular dysplasia Pattern baldness Flank alopecia	1 1 1
Infections	<i>Malassezia</i> dermatitis (primary) Primary pyoderma	1 23
Neoplasia	Cutaneous lymphoma	1
Keratinisation defect	Primary seborrhoea Schnauzer comedone syndrome	1 1
Miscellaneous	Others	38

*History and clinical investigations suggested an allergic dermatosis but specific allergy testing and / or food trials not conducted due to cost and / or owner wishes.

Full list of diagnoses in Appendix B

Figure 2. The most common skin diseases diagnosed at the UGVS (2000 - 2002)



Secondary Infections		Associated primary diagnosis		
Туре		Total cases	Туре	Total cases
Pyoderma	Superficial	94	Atopic dermatitis	60
			Flea allergic dermatitis	5
			Cheyletiellosis	3
			Others	26
	Deep	18	Atopic dermatitis	9
			Demodicosis	3
	No. Contraction		Others	6
Superficial and deep	Superficial	12	Atopic dermatitis	3
	and deep		Others	9
Malassezia		44	Atopic dermatitis	34
			Others	10

Table 3. Secondary bacterial and Malassezia infectionsassociated with primary diagnoses

General canine hospital population at UGVS (2000-2002)

Excluding dogs referred for skin disease, a total of 4505 dogs were examined over the three-year period, 2000 to 2002. There were 991 (21.99%) entire female, 1064 (23.62%) neutered females, 1724 (38.27%) entire males and 664 (14.74%) neutered males. In 62 cases the sex was not recorded. This gave a male female ratio of 1.2: 1. There was no significant statistical difference in the overall sex ratios between the skin disease population and the general hospital population but there were significantly fewer intact females in the skin disease group compared to general hospital population (P<0.001). The mean (± standard deviation) age at examination was $5.17 (\pm 3.82 \text{ years})$ with a median age of 4.80 years. The minimum age was 0.01 years and maximum age was 16.99 years. In 66 cases the age was not recorded. The age of the canine skin disease population was statistically significantly lower (P<0.001) when compared to the general canine hospital population by using Mann-Whitney Rank Sum Test.

The most common breeds seen in the general canine hospital population were Cross breed (483 cases; 10.72%), Labrador retriever (338 cases, 7.50%), German Shepherd dog (308 cases; 6.84%), Retriever (317 cases, 7.04%), Border collie (204 cases, 4.53%), Boxer (178 cases; 3.95%), Cocker spaniel (178 cases, 3.95%), Cavalier King Charles spaniel (174 cases, 3.86%), Golden retriever (158 cases, 3.51%), West Highland White terrier (116 cases, 2.57%), English Springer spaniel (134 cases, 2.97%), and Yorkshire terrier (96 cases, 2.13%). (Figure 3; Table 4)

When the breeds with skin disease were compared to the general hospital population using the Chi squared test, the West Highland White terrier was significantly over-represented in the dermatological population

while there was no statistical difference between dermatological population and general hospital population for the other breeds. (Table 4)

Figure 3. Most common dog breeds in general hospital population (2000-2002)



Breed	SD*	GH*	P value	Relative Risk
Cross breed	1	1	0.8703	0.9563
German Shepherd dog	2	3	0.0171	1.543
Labrador retriever	2	2	0.1104	1.344
West Highland White terrier	4	6	< 0.0001	2.492
Boxer	5	5	0.3741	1.281
Cocker spaniel	6	5	0.6655	0.8427
Golden retriever	7	5	0.5410	0.7792
Border collie	7	4	0.2174	0.6401
Staffordshire bull terrier	8	9	0.3722	1.467
Cavalier King Charles spaniel	9	5	0.0786	0.4762
English Springer spaniel	10	7	0.1690	0.5168
Yorkshire terrier	11	8	0.3448	0.5777

Table 4. Most common breeds of general hospital (GH) and skin disease (SD) populations

*The order of the breeds in total population (e.g. 1: the most common breed in the population.)

Feline skin disease population

Signalment

During the 3 year period studied a total of 40 cats were examined for skin disease. Of the 40 cases, there were no entire males, 19 (47.50%) neutered male, 2 (5.00%) entire female and 17 (42.50%) neutered female. In 2 cases the sex was not recorded. This gave a male to female ratio of 1:1.

The mean age (\pm standard deviation) at presentation was 5.00 \pm 3.21 years (minimum and maximum age: 0.58 and 13.00 years; median age 4.25 years).
There were in total 5 different breeds. They were Domestic short hair (30), Persian (long hair) (4), Siamese (2), Burmese (1), Domestic long hair (1); Ragdoll (1) and Turkish Van (1). (Table 5; Figure 4)

Breed	Number	Percentage
Domestic short hair	30	75
Persian (long hair)	4	10
Siamese	2	5
Ragdoll	1	2.5
Domestic long hair	1	2.5
Burmese	1	2.5
Turkish Van	1	2.5

Table 5. Cat breeds referred for skin disease (2000-2002)





Feline skin diseases diagnosed

There were 42 primary skin diseases diagnosed in the 40 cats examined. The most common skin condition diagnosed was atopic dermatitis (12 cases; 28.6 %). Other common skin disorders, in descending order were: 9 cases (21.4%) of flea allergic dermatitis, 5 cases (11.9%) allergy not fully diagnosed, 4 cases (9.5%) of facial dermatitis and 2 cases (4.8%) of pemphigus foliaceus. (Table 6; Figure 5; Appendix C)

Secondary infection in cats was rarely diagnosed. There was only one case of superficial pyoderma in a cat with food sensitivity and a cat with feline acne had a deep pyoderma.

Table 6. S	Summary	of the	feline	skin	diseases	diagnosed	over	a	3-
	year	perio	d at the	e UG	VS (2000-	-2002)			

	Diagnosis	Totals
Allergy	Atopic dermatitis	12
	Flea allergic dermatitis	9
	Food sensitivity	2
	Allergy not diagnosed*	5
Immune mediated	Pemphigus foliaceus	2
Neoplasia	Ceruminous gland adenoma	1
Miscellaneous	Symmetrical alopecia	1
	Eosinophilic granuloma	1
	Idiopathic facial dermatitis	4
State of the second	Feline acne	1
	Plasma cell pododermatitis	1
	Psychogenic alopecia	1
	Cutaneous asthenia	1
	Chin oedema	1

*History and clinical investigations suggested an allergic dermatosis but specific allergy testing and / or food trials not conducted due to cost and / or owner wishes.

Full list of diagnoses in Appendix C



Figure 5. The most common feline skin diseases diagnosed at the UGVS (2000 - 2002)

General feline hospital population at UGVS (2000-2002)

Excluding cats referred for skin disease, a total of 912 cats were examined over the three-year period, 2000 to 2002. There were 105 (11.51%) entire female, 328 (35.96%) neutered females, 60 (6.58%) entire males and 405 (44.41%) neutered males. In 14 cases the sex was not recorded. This gave a male female ratio of 1.07: 1. There was no statistical difference between the sex of skin disease population and general hospital population. The mean (\pm standard deviation) age at

Chapter 1

examination was 7.20 (\pm 4.92 years) with a median age of 7.00 years. The minimum age was 0.06 years and maximum age was 18.00 years. In 31 cases the age was not recorded. The age of the skin disease population was significantly lower than the general hospital population (P=0.024). The most common breeds seen in the general hospital population were Domestic short hair (664 cases; 72.81%), Persian (Long hair) (98 cases; 10.75%), Domestic long hair (43 cases; 4.71%), Siamese (22 cases; 2.41%), Exotic (13 cases; 1.43%), and Burmese (12 cases, 1.32%). (Figure 6; Table7)

Figure 6. Most common cat breeds in general hospital population (2000-2002)



Table.7 Most common cat breeds of general hospital (GH) and skin disease (SD) populations

Breed	SD*	GH*	P value	Relative Risk
Domestic short hair	1	1	0.5888	1.274
Persian (long hair)	2	2	1.0000	0.9259
Siamese	3	4	0.2672	2.035
Turkish Van	4		0.0823	12.18
Domestic long hair	4	3	1.0000	0.5291
Burmese	4	5	0.4297	1.852
Exotic		5	1.0000	0

*The order of the breeds in total population (e.g. 1: the most common breed in the population.)

Chapter I

DISCUSSION

Canine skin disease

The overall prevalence of skin disease compared to the general hospital population of dogs at UGVS was 7.37% for the three-year period of 2000-2002. This figure is similar to a previous study of prevalence rates for canine skin disease in a ten-year (1990 – 1999) retrospective study undertaken in the same hospital conducted by McEwan (2002) where 7.69% prevalence for canine skin disease was found. There are few detailed published studies of the prevalence of canine skin disease but several studies and text books have estimated that between 20% and 75% of small animals examined in the average practice have skin problems as a chief or concurrent owner complaint (Schwartzman and Orkin, 1962; Nesbitt, 1983; Wilkinson, 1985; Ihrke, Stannard, Ardans and Griffin, 1985; Grant, 1991). In a comprehensive study conducted in a 50% referral and 50% first opinion practice Scott and Paradis (1990) found a prevalence of 18.8% for canine skin diseases. The variation in prevalence rates between various studies may reflect different case origins and in the resources available at the study hospitals. The study presented here was conducted in an entirely referral population while other studies have been in first opinion practice or a combination of first opinion and referral practice. Referral populations are unlikely to reflect the prevalence of skin disease in the general dog population but may reflect the number and types of chronic skin complaints in the general dog population.

The Mean age of dogs with skin disease in this study (4.01 years) was statistically significantly younger when compared to the general hospital population. The Mean age of general hospital population (5.17 years) was approximately one year older than the skin

population. This may be due to the high prevalence of atopic dermatitis in the skin disease group. There was no overall sex predilection for skin disease found in this study, which is consistent with the findings of McEwan (2002) and Scott and Paradis (1990). Entire female dogs appeared to be at increased risk for skin disease in this study but the reasons for this are not clear.

27

When the most common breeds referred for skin disease in the study were examined, it was found that the West Highland White terrier was over-represented in the study presented here. In the ten-year study conducted in the same hospital by McEwan (2002) both West Highland White terrier and the Boxer were found to be predisposed to skin disease. There is considerable information available on apparent breed predilections for specific canine dermatoses (Halliwell and Schwartzman, 1971; Scott, 1981; McDonald and Kunkle, 1983; Scott, Walton, Slater, Smith and Lewis, 1987a; Scott, Walton, Slater, Smith and Lewis, 1987b; Grant, 1991; Scott, Miller and Griffin, 2001b). Ihrke and Franti (1985) reported that 31 breeds of dogs showed an elevated risk in northern California whereas only three breeds appeared to be at increased risk in Scott and Paradis' study, which were Boxer, Dachshund, and Bichon Frise. Of these three breeds only the Dachshund was also at increased risk in the northern California study. There appears to be variation in breed prevalence for skin disease reported. This may be due to differences in the genetic pools and possibly due to geographic variations in certain dermatoses. In referral centres particular interest in a specific breed or dermatosis may influence both the breed prevalence and prevalence of specific dermatoses.

The most common main diagnosis was atopic dermatitis (58%). As the condition tends to be chronic and so very likely to be referred, this finding is not surprising. This finding is consistent with a recent study (McEwan, 2002a) while Scott and Paradis (1990) found atopic dermatitis was their second most common diagnosis.

Secondary bacterial skin disease was found to be very common with 28.5 % of dogs in the study being diagnosed with a superficial pyoderma and 5.5% having a deep pyoderma. These figures are very close to those found by McEwan (2002) 23.6% (superficial pyoderma), 4.2% (deep pyoderma) and Scott and Paradis (1990) who recorded a prevalence of 25.3% for bacterial skin disease (folliculitis/ furunculosis). Slightly over 50% of the cases of secondary pyoderma in this study were associated with atopic dermatitis, which is again consistent with McEwan (2002) and Scott and Paradis (1990).

Feline skin disease

The prevalence of feline skin disease was 4.39% during 2000-2002 compared to the general feline hospital population. There is little information available concerning the demographics of feline skin disorders compared to canine. The figure for this study is much lower than the study of Scott and Paradis (1990) where 15.2% of cases were feline skin diseases. Again this may reflect the different case resources between these two studies.

The two most common feline dermatologic disorders in this study were atopic dermatitis and flea allergic dermatitis. A survey conducted in 1981 by the American Academy of Veterinary Dermatology revealed parasitic dermatoses and miliary dermatitis to be common. Miliary dermatitis is not a specific diagnosis but rather a skin reaction pattern that may result from several potential causes. Scott & Paradis 1990 found abscess (18.5%), otodectic mange (12.9%), atopic dermatitis (5.6%) to be common skin disorders. The differences between studies probably

reflect that different populations, geographical location, type of veterinary practice, and clients' economic consideration may influence the recorded prevalence of skin diseases. As in the dog the Mean age of cases of feline skin disorders (5 years) was significantly lower than the general hospital population of cats (7.20 years). Again this would be consistent with the high prevalence of atopic dermatitis found.

29

There was no sex predilection for skin disease found and the most common breeds were the Domestic short hair, Persian, Domestic long hair, Siamese and Burmese.

Although atopic dermatitis was a relatively common diagnosis compared to dogs, there were very few cases of feline bacterial skin disease. In general bacterial skin disease appears to be rare in the cat.

Conclusions

The result of this study and the findings of the literature reviewed emphasize the importance of dermatology in small animal practice. This study and others show that allergic skin disease is common in both the cat and dog (Ihrke and Franti, 1985; Sischo, Ihrke and Franti, 1989; Lund, Armstrong, Kirk, Kolar and Klausner, 1999; Nagata and Sakai, 1999). In the dog, and particularly in referral practices, atopic dermatitis is very common (Ihrke and Franti, 1985; Sischo, Ihrke and Franti, 1989; Scott and Paradis, 1990; Lund et al, 1999; Nagata and Sakai, 1999; McEwan, 2002a). In dogs, bacterial skin disease due to S. intermedius is common (Ihrke and Franti, 1985; Sischo, Ihrke and Franti, 1989; Scott and Paradis, 1990; Lund et al, 1999; Nagata and Sakai, 1999; McEwan, 2002a). Much of this is secondary to other skin diseases, especially atopic dermatitis (Ihrke, Halliwell and Deubler, 1977; White and Ihrke, 1987; Scott and Paradis. 1990: McEwan. 2002a). In man.

Staphylococcus aureus skin infections are common in atopic dermatitis patients (Jones, Schmitz, Fluit, Acar, Gupta and Verhoef, 1999; Sader, Pfaller, Jones, Doern, Gales, Winokur and Kugler, 1999; Jones, Karlowsky, Draghi, Thornsberry, Sahm and Nathwani, 2003). Several theories have been advocated as to why both canine and human atopic dermatitis patients suffer from staphylococcal skin infections. One theory suggests that increased adhesion by staphylococci to skin could be a factor (Feingold, 1986; Roth and James, 1989; McEwan, 1990; Scott, Miller and Griffin, 2001b; Saijonmaa-Koulumies and Lloyd, 2002). A major problem when assessing bacterial adhesion to corneocytes is that many assays are tedious and time consuming to perform. A simple test to quantify adhesion by bacteria to corneocytes would be a useful tool. In the next two chapters the development of a simple assay to determine bacterial adhesion is described.

CHAPTER 2

A study of Canine and Feline Corneocyte Morphometrics

LITERATURE REVIEW

Introduction

The microscopic anatomy and physiology of the skin of the dog and the cat has been the subject of several studies (Lovell and Getty, 1957; Strickland and Calhoun, 1963; Kral and Schwartzman, 1964; Lyne and Short, 1965; Montagna, 1967; Schwarz, 1979; Schummer, 1981; Dyce, 1987; Suter, 1987; Evans and Christensen, 1993; Yager and Scott, 1993; Scott, Miller and Griffin, 2001b). The outer part of the skin is the epidermis, which is divided into stratum corneum (horny layer or cornified cell layer), stratum granulosum (granular layer), stratum spinosum (prickle cell layer or spinous cell layer), and stratum basale (basal layer). The epidermis rests on and is attached to the dermis, which in turn lies on the subcutis or hypodermis. The stratum corneum is the outer layer of terminally differentiated keratinocytes that is constantly being shed (Webb and Calhoun, 1954; Strickland and Calhoun, 1963; Schwarz, 1979; Scott, 1980; Greene and Lammler, 1993). It is a multilayered zone of corneocytes suspended in an extracellular lipid matrix, often likened to a series of bricks (corneocytes) with mortar (lipids) (Leigh, 1994). Corneocytes, which contain a variety of humectants and natural sunscreens synthesized from proteins, are the flattened, anuclear eosinophilic cells in stratum corneum which have no true cell membrane (Priestley, 1993). The stratum corneum varies in thickness from 3 to 35 μm in cats and from 5 to 1500 μm in dogs (Scott, Miller and Griffin, 2001b).

Epidermal function

The general functions of the epidermis are as follows (Moriello and Mason, 1995): (1) Barrier. In order to prevent loss of water and

electrolytes and limit the penetration of bacterial, viruses, and fungal organisms and their metabolites. (2) Sensory perception. Free nerve endings and sensory end organs (Schwarz, 1979; Scott, 1980; Scott, 1984; Lever and Schaumburg-Lever, 1990). (3) Pigmentation, which can protect against UV radiation. (4) Immuno-regulation and defence. Protection against infection. Langerhans' cells, keratinocytes, epidermotropic T lymphocytes, and draining peripheral lymph nodes are thought to form collectively an integrated system of skin-associated lymphoid tissue (SALT) that mediates cutaneous immuno-regulation (Thoday and Friedmann, 1986). Langerhans' cells, which are mononuclear dendritic cells located suprabasally (Lever and Schaumburg-Lever, 1990), stimulate the proliferation of relevant helper T lymphocytes by the presentation of antigen; they also induce cytotoxic T lymphocytes directed to allergens and modified self-determinants, produce interleukin (IL) 1 and other cytokines, contain numerous enzymes, and are phagocytic (Thoday and Friedmann, 1986).

The stratum basale, spinosum and granulosum are known as the living epidermis while the stratum corneum constitutes the terminally differentiated non-living epidermis. The normal mammalian epidermis is a primary barrier against microbial invasion. Anatomical features of this region impede the penetration of microorganisms or their potentially toxic secretions and metabolites into the host. The arrangement of inert squames in a lipid-rich matrix prevents the penetration of aqueous substances. The keratinocyte also plays an important role in the immune function of epidermis (Thoday and Friedmann, 1986). Keratinocytes can no longer be regarded as simply the 'building blocks' of the tissue. They are phagocytic and can process antigens (Thoday and Friedmann, 1986). They produce IL-1 constitutively and, following antigenic stimulation, can produce a wide range of cytokines (e.g., IL-1, IL-3, IL-6, iL-7, IL-8, IL-10, IL-12, IL-15, IL-18, IL-20, prostaglandins, leukotrienes, interferon, and

33

tumor necrosis factor α (TNF- α)), which influence the immune responses (Grone, 2002). In addition, keratinocyte-derived IL-7 and 15 are considered to be significant in T-cell trafficking, possibly even in the pathogenesis of cutaneous T-cell lymphoma (Grone, 2002).

Keratinocyte differentiation

Normal epidermal homeostasis requires a finely tuned balance between growth and differentiation of keratinocytes (Suter, 1987; Suter, Crameri, Olivry, Mueller, von Tscharner and Jensen, 1997). This balance must be greatly shifted in the direction of proliferation in response to injury and then must return to a state of homeostasis with healing. In addition, epidermal keratinocytes have important functions as regulators of cutaneous immunity and inflammation. The epidermis is ectodermal in origin and normally undergoes an orderly pattern of proliferation, differentiation, and keratinisation. As the primary cell type, keratinocytes define the structure and organization of the epidermis through their program of terminal differentiation. Keratinocytes are arranged in overlapping layers, which are indicative of their states of differentiation. The basal layer of keratinocytes, which abuts the basement membrane, is the least differentiated, proliferative layer. The next layer, stratum spinosum, is composed of keratinocytes derived from its basal layer. With increasing differentiation, the keratinocyte continues to migrate upwards. When it reaches the upper zone of the granular layer, the keratinocyte undergoes a type of programmed cell death, leading to dissolution of the nucleus and other organelles and formation of the cross-linked envelope characteristic of squames. Finally, the squames are lost from the surface of the skin, and the end product is the fully keratinised corneocyte. In healthy Beagle dogs, similar to human keratinocytes, the time required for a keratinocyte to migrate from the basal layer to the surface of the skin is 22 days (Baker, Maibach, Park, McFarland and O'Brien, 1973; Kwochka

and Rademakers, 1989). In normal epidermis there is a tight coordination between vertical migration and differentiation state, with characteristic molecules expressed and morphology manifested at each level. Keratinisation is the term used to describe the maturation of a keratinocyte from the stratum basale to the stratum corneum (Moriello and Mason, 1995).

35

Corneocyte morphometrics

The human stratum corneum constitutes less than 10% of the cells of the intact epidermis and is made of about 15 layers of a tightly packed syncytium of flattened non-nucleated keratinised cells known as corneocytes (Christophers and Kligman, 1964; Amer, Mostafa, Tosson and Nasr, 1996). The canine stratum corneum consists of approximately 50 cell layers (Lloyd and Garthwaite, 1982). The studies that have been carried out to investigate the morphology of corneocytes show that human corneocytes are extremely flat, but very large cells lacking nuclei (Goldschmidt and Kligman, 1967a; Blair, 1968) with their largest diameter measuring between 30-50µm (Amer et al, 1996; Mihara, 1997). Usually, corneocytes have a regular pentagonal or hexagonal shape with a surface area of 900-1200 μm^2 (Plewig and Marples, 1970). As a result of keratinization of cells, and transepidermal water loss, corneocytes become progressively flatter and are finally desquamated at the skin surface. The rate of desquamation in normal skin matches the rate of cell production so epidermal thickness is maintained. This physiolological desquamation goes unnoticed, but it results in visible scales when the rate of keratinisation is pathologically raised, such as in parakeratosis (Amer et al, 1996). Morphological variations of corneocytes depend on biologic variables, such as body region, age, season, race and sex. (1) Regional difference. Corneocytes vary in cell size and thickness with topographical site (Plewig, 1970). In man large cells are found in the

axilla, on the trunk, thigh, and upper arm. These cells also tend to be thin. Experimental cytologic examinations of the trunk and arms revealed statistically significant variations of cell sizes even when corneocytes were removed from regions only 10 cm apart (Holzle and Plewig, 1977; Haidl and Plewig, 1988). (2) Effects of age. Cell surface area of corneocytes increases with age (Plewig and Marples, 1970; Marks, 1980; Toyoshima, 1989). The mean surface area of corneocytes taken from the chest area in infancy is 900 μ m². In adults the surface area increases with the largest values occurring in old age (1200 μ m²). An age-dependent increase in size, often significant, is also seen in most other regions, with the exception of the forehead and mucous membrane of the tongue. (3) Seasonal changes. Morphological variations of corneocytes are seen as a consequence of climatic fluctuations, such as temperature and humidity. It is known that increased epidermopolesis leads to a decrease in corneocyte cell surface areas. Stratum corneum turnover is faster in summer than in winter. In one study, the corneocyte surface area and cell counts were measured in four groups of ten subjects monthly throughout the year (Herrmann, Scheuber and Plewig, 1983). The paraumbilical region and the lower lateral portion of the right leg were used as sampling sites. In the warm season during the months from March to September, the corneocytes were smaller than in the cold season. The cell counts increased correspondingly. The seasonal changes were most pronounced on lower legs in both older and younger women. (4) Sex and racial differences. It was found that the size of corneocytes in females was larger than those of males in 8 out of 10 different collection sites. The exceptions being the popliteal fossa and the heel (Lee and Lee, 1986). However, there was no significant difference between black, white and oriental skin (Corcuff, Lotte, Rougier and Maibach, 1991).

The cause of the difference in size according to the site of the body might be due to the difference in: (1) the turnover time of the stratum

corneum, (2) intercorneal cohesion, and (3) the thickness and shape of the corneocytes (Marks and Barton, 1983). In a study of mammalian corneocytes using hairless guinea pigs (HL-GP) and normal-haired guinea pigs (HD-GPs), it was concluded that HL-GP skin has a greater similarity to human skin than that of HD-GPs and other rodents (Sueki, Gammal, Kudoh and Kligman, 2000). In this same study, the measurements of the mean projected area showed that guinea pig corneocytes are slightly larger than human corneocytes. The mean corneocyte surface area of the guinea pig varied between 1034 and 1415 μ m². However, since corneocyte size varies depending on the body site and age this should be taken into consideration.

Collection of corneocytes and their study

Most of the literature concerning skin structure pertains to the human species and rodents (mice and rats) (Mackenzie, 1969; Christophers, 1971; Menton and Eisen, 1971a; Menton and Eisen, 1971b; Mackenzie, 1972; Goerttler, Reuter and Stahmer, 1973; Mackenzie and Linder, 1973; Allen and Potten, 1974). Several techniques have been developed over the past decades for the study of the human stratum corneum, including potassium hydroxide mount (Lookingbill, 1985), Tzanck smear (Tzanck, 1947), biopsy (Lookingbill, 1985), skin replica methods (Thomson, 1953; Sarkany, 1962; Makki, Barbenel and Agache, 1979), tape stripping (Pincus, 1951; Keddie and Sakai, 1965), liquid non-drying adhesive stripping (Goldschmidt and Kligman, 1967b), cryanoacrylate adhesive specimens (Keddie, Shadomy and Shadomy, 1963; Goldschmidt and Kligman, 1967b; Marks and Dawber, 1971; Marks and Dawber, 1972; Marks, 1972; Holland and Roberts, 1974; Tring and Murgatroyd, 1974; Tring and Murgatroyd, 1975; Marks, 1980; Mills, Jr. and Kligman, 1983; Katz, Prawer, Hien and Mooney, 1985), scanning electron microscope (SEM) (Hashimoto and Kanzaki, 1975), transmission electron microscope

37

(TEM) (Hashimoto and Kanzaki, 1975), fluorescent microscope (Christophers, 1971), confocal laser scanning microscope (Schatzlein and Cevc, 1998), and atomic force microscope (Kashibuchi, Hirai, O'Goshi and Tagami, 2002). The adhesive D-Sqame[®] disc (CuDerm Corp., Dallas, TX, USA), is a circular crystal clear foil 25 mm in diameter coated with a special adhesive substance that has been used as a convenient harvesting method for the superficial desquamating layer of the stratum corneum in several research studies (Serup, Winther and Blichmann, 1989; Jemec and Serup, 1992; Serup, 1992a; Serup, 1992b; Pierard and Pierard-Franchimont, 1993; El Gammal, Pagnoni, Kligman and el Gammal, 1996; Kawakami, Callicott and Zhang, 1998; De Paepe, Janssens, Hachem, Roseeuw and Rogiers, 2001; Yoon, Baik and Oh, 2002; Hill and Edwards, 2002).

In conclusion while there are occasional reports on the microanatomy of the skin of domestic large animals (Lloyd, Sasiak, Kitson, McEwan and Elder, 1993), and that of the dog and cat (Webb and Calhoun, 1954; Baker *et al*, 1973) there is little information on corneocyte morphometrics in these species. In man and rodents several different methods have been employed to collect corneocytes for study. The D-Squame® disc has been shown to be a valuable tool for the collection of corneocytes and would be potentially suitable for use in studies in the dog and cat.

38

INTRODUCTION

The literature reviewed shows that there is little published information on the size and shape of canine and feline corneocytes. Prior to developing an adhesion assay to study bacterial adhesion to corneocytes, studies were undertaken to determine size and surface area of canine and feline corneocytes collected from different body sites.

AIMS

The aims of the study were:

- 1. To determine the size and shape of canine and feline corneocytes
- 2. To assess three body sites for the collection of corneocytes that would be suitable for use in an assay for bacterial adherence using image analysis

MATERIAL AND METHODS

STUDY 1. CORNEOCYTE MORPHOMETRICS

Populations studied

Three groups of animals were studied:

1. Normal dogs. Normal dogs were obtained from the following sources: blood donor dogs held in the University of Glasgow Veterinary School, pet dogs belonging to Veterinary School staff and dogs referred with minor medical complaints. All dogs in this group were given a full clinical examination and were free from skin disease.

- 2. Atopic dermatitis dogs. These dogs fulfilled the critera for diagnosis as described by Willemse (1986) and were cases referred to the UGVS dermatology clinic.
- 3. Normal cats. Normal cats were obtained from the following sources: pet cats belonging to Veterinary School staff, cats undergoing screening studies for polycystic kidney disease or arthritis and cats referred with minor medical complaints. All cats in this group were given a full clinical examination and were free from skin disease.

None of the animals used in the studies had received either systemic or topical treatments for a period of at least three weeks prior to corneocyte sampling.

The breed, sex and age of individual animals used in the study are documented in Appendices D-I.

Body sites sampled

Three body sites were sampled from each of 5 animals from each group studied (atopic dogs, normal dogs and cats). The body sites sampled were:

- 1. Inner aspect of the base of the pinna (Figure 7)
- 2. Inner thigh in the femoral triangle area (Figure 8)
- 3. The left lateral thorax at the 10/11th intercostal space (Figure 9)

Figure 7. Ear sampling area



Figure 8. Groin sampling area







Felt pen marks can be seen marking sites for intradermal skin testing

Hair was removed by gently clipping (electric clippers; size 40 blade) from the thorax and groin area. In the atopic dog group a total of five dogs were studied and for each dog all three sites sampled. For the normal dogs and cats due to owner constraint on clipping of some sample sites it was not possible to obtain samples from all three sites in each individual animal. Seven normal dogs and ten cats were used in the study.

Corneocyte collection

Prior to sampling, surface debris was removed from the skin by applying five successive adhesive tape strips (Sellotape® Original). Corneocytes were then collected using 25 mm diameter adhesive discs (D-squame®, CuDerm Corporation, Dallas, Texas USA). From each collection site five consecutive samples were obtained using five D-squame® discs. Each disc was applied to the sample site once.

Corneocyte staining and Image acquisition

Corneocytes were stained using 2.5% rhodamine B and 0.75% methylene blue (Sueki *et al*, 2000) for 30 minutes, rinsed in tap water then air dried. A Leica DMLS microscope with a Leica ICCA digital video camera was used to acquire images. Images were obtained from each disc at a microscope magnification of x400 and saved as TIFF files onto the hard drive of a Viglen computer (Viglen Genie Pentium® 4). From each body site five D-Squame® discs were collected and from each disc ten corneocytes were measured giving a total of 50 corneocytes examined from each collection site per animal.

Calibration and measurements

Measurements were carried out using an image processing and analysis program (UTHSCSA ImageTool Version 3.0). Prior to measurement the image analysis program measuring tools were calibrated by use of a stage micrometer (Agar Scientific Ltd., Essex, UK).

Preliminary studies showed two distinct populations of corneocytes obtained from dog and cat skin. For the purposes of study these were designated type I and type II corneocytes. Type I corneocytes (Figure 10) were small, elongated and deeply staining. Type II corneocytes (Figure 11) were poorly staining and polygonal. The diameter and surface area of polygonal type II corneocytes were calculated and analysed. The diameter was calculated as an Mean of two measurements (longest and shortest breadth) for each corneocyte. The programme software automatically calculated the surface area after outlining the perimeter of each corneocyte using the software draw tool.



Figure 10. Type I corneocytes from canine thorax

Type I corneocytes (*) predominate. Rhodamine B and methylene blue staining. Original magnification x400.



Figure 11. Type II corneocytes from canine ear

Rhodamine B and methylene blue staining. Original magnification x400.

STUDY 2. COMPARISON OF CORNEOCYTES FROM HAIRED AND NON-HAIRED SKIN

It was observed that type I corneocytes were plentiful in samples from densely haired regions and type II corneocytes more readily obtained from less haired regions. In order to investigate the origin of the two populations of corneocytes, a study of haired and non-haired skin was undertaken.

Corneocyte collection

Five normal dogs and five normal cats were used in this study. D-Squame® discs were used to collect corneocytes from three sites on the inner aspect of the right pinna from the base of pinna to the more haired apex (Figure12). Prior to corneocyte collection surface debris was removed using successive adhesive tape strips as described above. Corneocytes were stained using the method detailed above.



Figure 12. Comparison of haired and non-haired skin

Counting method

From each D-Squame® disc, five images were collected at x400 magnification and saved as TIFF files. Using the image analysis software (ImageTool) a script (macro program) was written that produced a box out line 400 x 400 pixels (approximately $13000\mu m^2$) in size centrally positioned over the image to be studied (Appendix J). The numbers of the two types of corneocytes were counted manually within the centrally positioned counting area. Any corneocytes on the edge of counting box were counted if they were at least 50% within the square counting area.

STATISTICAL ANALYSIS

Statistical analysis and graphical presentation of data was conducted using computerised statistical packages, GraphPad Prism, version 4 (Graphpad Software Inc. San Diego CA). Comparisons between groups were conducted using simple one-way analysis of variance (ANOVA), where data was not normally distributed; Kruskal-Wallis ANOVA on ranks was used. To isolate the group or groups that differed from the others a multiple comparison procedure (Dunn's method) was employed. A value of p< 0.05 was considered to be significant.

RESULTS

Corneocyte morphometrics

Two broad populations of corneocytes could be identified. One consisted of small, single, irregular and often fragmented cells that stained darkly (type I corneocytes Figure 10) and a second population of lightly pink-staining, transparent and polygonal cells that formed a honeycomb pattern with slightly overlapping edges (type II corneocytes Figure 11). Type I corneocytes appear to arise from hair follicles (tables 16 and 17; figures 16 and 17).

A summary of the results for the mean diameter and surface area of type II corneccytes collected from normal dogs, atopic dermatitis dogs and normal cats are shown in tables 8 and 9, below together with a statistical analysis comparing data between animals and between discs for each site sampled. The mean diameter of corneccytes from the ear, thorax and groin of normal dogs was 37.92 μ m, 42.82 μ m and 43.47 μ m. For atopic dermatitis dogs equivalent figures were 38.38 μ m, 43.24 μ m and 40.19 μ m. For cats the figures were 40.41 μ m, not done and 49.92 μ m. The mean surface area of corneccytes from the ear, thorax and groin of normal dogs was 1091.96 μ m², 1399.40 μ m² and 1435.72 μ m². For atopic dermatitis dogs equivalent figures were 1160.86 μ m², 1439.67 μ m² and 1249.75 μ m². For cats the figures were 1222.17 μ m², not done and 1870.71 μ m². In general cat corneccytes were larger than dogs (both normal and atopic) and corneccytes from the groin appear to be larger than form the ear or thorax.

When corneocytes were from each set of five discs were compared from an individual animal there was generally little within animal difference between each of the five discs (tables 10-15). Differences were noted when corneocytes were compared between individual animals of the same group (tables 10-15).

Studies were not conducted on samples obtained from cat thorax as there were insufficient type II corneocytes obtained for this site to count. The results of the mean findings for each body site with statistical summaries are showed in tables 10a – 15c. Figures 13a – 15b depict box plots of the corneocyte diameters and surface areas from each body site studied. Appendices D-I contain the raw data for these studies.

	Ear		Tho	rax	Groin	
	Mean (µm) ± SD	Median (µm)	Mean (µm) ± SD	Median (µm)	Mean (µm) ± SD	Median (µm)
Normal dogs	37.92 ± 4.27	38.30	42.82 ± 4.36~~	42.61	43.47 ± 4.32~	43.19
Atopic dogs	38.38 ± 5.44	38.40	43.24 ± 4.61##	43.40	40.19 ± 6.42**	40.80
Cats	40.41 ± 4.74*	40.19	Not Done	Not Done	48.48 ± 7.32#	49.92

Table 8. Summary results of corneocyte diameter

* Significantly larger than normal and atopic ear corneocytes

Significantly larger than normal and atopic groin corneocytes

~ Significantly larger than atopic groin corneocytes

Groin corneocytes from normal and cat significantly bigger than ear for equivalent groups

**Atopic groin corneocytes significantly bigger than atopic ear corneocytes

##Atopic thorax corneocytes significantly bigger than atopic groin and ear corneocytes

~~Normal thorax corneocytes significantly bigger than normal ear corneocytes

The second	Ear		Tho	ax	Groin	
	Mean (µm ²) ± SD	Median (µm²)	Mean (µm²) ± SD	Median (µm²)	Mean (μm²) ± SD	Median (µm²)
Normal dogs	1091.96 ± 208	1109.44	1399.40 ± 29.60##	1411.58	1435.72 ±236.87‡	1415.36
Atopic dogs	1160.86 ±72.28#	1142.51	1439.67 ± 74.80**	1450.63	1249.75 ±368.55	1236.29
Cats	1222.17 ± 39.02*	1197.70	Not Done	Not Done	1772.11 ± 461.11~	1870.71

Table 9. Summary results of corneocyte surface area

* Significantly larger than normal and atopic ear corneocytes

Significantly larger than normal ear corneocytes

~ Significantly larger than normal and atopic groin corneocytes

‡ Significantly larger than atopic groin corneocytes

All groin corneocytes significantly bigger than ear corneocytes for equivalent groups

**Atopic thorax corneocytes significantly bigger than atopic ear and groin corneocytes

##Normal thorax corneocytes significantly bigger than normal ear corneocytes



Box plots of corneocyte morphometrics

Box and whisker plots. The horizontal line in the middle shows the median of the sample. The top and bottom of the box show the 75th and 25th percentiles and the top and bottom of the whiskers show the maximum and minimum values.

				-					
Normal dogs ears (Mean±SD) (μm)									
	Disc 1	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5			
N1	40.00± 2.72	35.85± 5.34	39.55± 5.20	37.84± 2.62	41.09± 2.75	38.99± 4.03			
N2	40.00± 3.93	39.90± 2.18	38.01± 3.50	36.95± 1.96	42.30± 4.04	39.16± 3.86			
N3	39.70± 4.75	38.41± 2.14	39.07± 3.78	39.63± 4.69	40.31± 4.60	39.41± 3.86			
N4	35.36± 2.92	38.72± 3.03	37.42± 2.64	40.45± 2.84	39.46± 4.47	38.47± 3.55			
N5*	34.91± 2.73	32.41± 2.72	32.64± 2.55	35.66± 3.52	32.89± 2.70	33.57± 3.06			
Ali	37.99± 4.11	37.24± 4.08	37.18± 4.44	38.03± 3.67	39.16± 4.84	37 .92 ± 4.27			

Table 10a. Corneocyte diameter ears

No significant difference between discs 1-5

* N5 significantly smaller than N1, 2, 3 and 4

Atopic dogs ears (Mean±SD) (μm)									
	Disc 1*	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5			
AD1	36.76± 2.3	37.21± 2.44	40.03± 4.49	38.19± 2.88	40.32± 2.04	38.68± 3.54			
AD2**	32.31± 3.46	35.44± 3.74	31.41± 3.98	32.75± 4.68	35.11± 2.64	33.37± 4.01			
AD3	35.47± 2.42	39.40± 3.73	43.59± 3.19	37.58± 3.61	41.68± 3.24	39.58± 4.17			
AD4	29.50± 4.40	39.55± 2.89	38.42 <u>+</u> 4.68	37.76± 2.98	40.57± 2.54	37.59± 5.72			
AD5#	41.56± 2.88	45.16± 3.60	43.99± 4.29	45.11± 4.45	37.30± 4.86	42.65± 5.00			
All	35.12± 5.14	39.44± 4.65	39.50± 6.16	38.53± 5.64	39.30± 4.27	38.38± 5.44			

Table 10b. Corneocyte diameter ears

* Disc 1 significantly smaller that discs 2, 3 and 4 ** AD2 significantly smaller than AD1, 3, 4 and 5 # AD5 significantly bigger than AD1, 2, 3 and 4

Cat ears (Mean±SD) (μm)									
	Disc 1	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5			
C1#	40.93± 2.89	40.69± 2.94	43.06± 3.87	41.88± 2.46	42.47± 3.02	41.81± 3.08			
C2	39.62± 4.95	37.40± 4.18	36.53± 2.59	38.12± 4.23	38.82± 3.33	38.10± 3.93			
C3*	40.14± 4.39	43.25± 4.24	41.70± 4.08	45.05± 3.99	45.06± 5.49	43.04± 4.70			
C4	37.01± 6.77	37.09± 7.23	36.47± 3.74	36.81± 3.72	37.86± 2.98	37.05± 5.00			
C5	40.82± 3.10	41.49± 4.22	43.18± 4.22	42.79± 3.60	42.09± 3.11	42.09± 3.63			
All	39.70± 4.67	39.99± 5.17	40.19± 4.73	40.93± 4.65	41.28± 4.43	40.41± 4.74			

Table 10c. Corneocyte diameter ears

No significant differences between discs 1-5

* C3 significantly bigger than C2, and 4 # C1 significantly bigger than C2, and 4

Table 11a. Corne	ocyte diameter thorax
------------------	-----------------------

Normal dogs thorax (Mean±SD) (μm)									
	Disc 1	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5			
N1	40.95± 2,34	41.24± 2.23	41.14± 2.63	40.20± 3.88	41.48± 2.89	40.83± 3.05			
N2	37.41± 6.31	41.08± 4.91	40.28± 3.51	42.68± 5.96	42.46± 4.98	40.82± 5.18			
N3	48.41± 2.86	46.68± 4.38	48.58± 2.83	48.63± 2.56	45.84± 2.38	47.53± 3.13			
N4*	45.501± 5.94	47.39± 4.26	43.02± 3.29	43.24± 4.52	46.10± 3.61	45.05± 4.57			
N5#	46.60± 3.71	43.81± 2.88	43.25± 2.16	43.70± 2.84	44.79± - 3.53	44.63± 3.18			
All	42.91± 5.56	42.84± 4.41	42.20± 3.36	42.58± 4.18	43.58± 4.04	42.82± 4.36			

No significant differences between discs 1-5

* N4 significantly bigger than N1, and 2

N5 significantly bigger than N1 and 2

	Disc 1	Disc 2*	Disc3	Disc 4	Disc 5	Disc1-5
AD1	41.20± 3.75	43.79± 2.80	41.29± 3.97	38.83± 5.08	37.62± 8.18	40.69± 5.14
AD2	40.71± 2.74	43.10± 1.90	44.07± 1.78	41.66± 3.10	41.20± 1.92	42.27± 2.75
AD3	42.23± 3.08	41.57± 4.68	43.58± 4.26	41.11± 3.27	40.72 ± 3.05	41.79± 3.96
AD4**	44.42± 4.24	47.04± 3.41	44.83± 3.64	49.74± 3.47	43.10± 3.25	46.11 ± 4.2 8
AD5#	45.36± 5.25	43.98± 3.89	46.60± 2.76	46.04± 5.86	45.21± 3.67	45.33± 4.18
All	42.78± 4.17	44.14± 3.74	44.00± 4.14	43.57± 5.63	41.71± 4.87	43.24 ± 4.6 1

Table 11b Corneocyte diameter thorax

* Disc 2 significant bigger than disc 5

** AD4 significantly bigger than AD1, 2 and 3 # AD5 significantly bigger than AD1, 2 and 3

Normal dogs groin (Mean±SD) (μm)						
	Disc 1	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5
N1	40.54± 4.46	39.38± 3.85	42.74± 4.11	43.63± 2.67	39.93± 2.67	41.21± 3.88
N2#	43.11± 3.32	46.43± 2.54	46.24± 5.10	42.73± 2.60	42.79± 2.72	44.40± 3.68
N3*	48.41± 2.86	46.62± 4.13	48.08± 3.11	48.28± 2.65	46.25± 2,60	47.53± 3.13
N4~	44.89 <u>-</u>) 4.30	41.51± 2.19	42.30± 1.83	44.49± 3.89	44.98± 3.97	43.73± 3.46
N5	39.36± 2.96	41.16± 4.82	40.72± 3.47	42.17± 2.52	39.37± 3.74	40.46± 3.48
All	43.26± 4.77	43.08± 4.64	44.09± 4.45	44.26± 3.47	42.64± 4.09	43.47± 4.32

Table 12a. Corneocy	rte diameter groin
---------------------	--------------------

No significant differences between discs 1-5 * N3 significantly bigger than N1, 2, 4 and 5

N2 and ~N4 significantly bigger than N1 and 5

Atopic dogs groin (Mean±SD) (μm)						
	Disc 1	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5
AD1~	33.79± 4.95	39.33± 3.55	39.17± 4.61	39.39± 11.08	41.03± 2.67	38.56± 6.17
AD2	32.64± 4.33	31.67± 3.57	30.97± 2.61	30.57± 4.40	36.62± 2.48	32.43± 4.11
AD3#	45.10± 3.71	41.74± 5.70	42.82± 4.48	44.08± 3.76	45.42± 3.00	43.84± 4.13
AD4*	46.63± 4.23	45.59± 5.35	46.23± 4.79	45.32± 3.18	44.97± 2.27	45.86± 3.84
AD5	41.07± 3.10	37.42 <u>+</u> 4.52	38.32± 1.67	41.18± 3.00	36.62± 2.48	38.93± 3.39
All	39.87± 7.00	39.28± 6.63	40.78± 6.24	40.00± 7.53	40.42± 3.67	40.19± 6.42

Table 12b. Corneocyte diameter groi	Table 1	12b.	Corneocyte	diameter	groin
-------------------------------------	---------	------	------------	----------	-------

No significant differences between discs 1-5

* AD4 and # AD3 significantly bigger than AD1, 2 and 5

~ AD1 significantly bigger than AD2

Cats groin (Mean±SD) (μm)						
	Disc 1	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5
C1~	52.03± 3.96	52.81± 4.55	55.83± 2.56	53.96± 4.52	48.47± 0.99	5 2.4 1± 4.4 2
C2‡	37.49± 3.98	41.25± 4.86	47.00± 4.75	43.07± 3.43	47.25± 5.92	43.16± 5.75
C3	37,78± 3.21	39.19± 3.67	43.10± 3.16	42.41± 3.70	38.19± 5.10	40.13± 4.29
C4#	54.03± 2.98	53.03± 5.87	51.18± 3.14	52.66± 4.54	53.51± 3.31	52.88± 4.07
C5*	54.39± 5.68	51.85± 2.45	55.07± 3.36	51.75± 3.56	55.94± 5.76	$\textbf{53.80}{\pm}\textbf{ 4.53}$
All	47.14± 8.80	47.49± 7.48	50.44± 5.98	48.69± 6.20	48.62± 7.62	48.48± 7.32

Table 12c. Corneocyte diameter groin

No significant differences between discs 1-5

* C5, #C4 and ~C1 significantly bigger than C2 and 3

‡ C2 significantly bigger than C3
		a hit on a page of the second second second			المروم الم المراجع الم				
Normal dogs ears (Mean±SD) (μm²)									
	Disc 1	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5			
N1	1,230,14±	1,085.03±	1,166.60±	1,128.03±	1,161.00±	1,1 54.1 6±			
	191.36	237.09	146.13	170.36	122.19	177.14			
N2	1,194.79±	1,234.44±	1,018.23±	1,082.06±	1,260.86±	1,158.07±			
	116.14	145.99	178.31	200.16	160.13	181.91			
N3	1,356.17±	1,152.23±	1,121,83±	1114.94±	971.47±	1143.33±			
	130.4	145.34	172,71	126.38	165.26	189.68			
N4	967.14±	1,191.92±	1,221.67±	1,287.91±	1,110.81±	1,155.89±			
	107.10	193.93	161.28	150.24	101.54	179.50			
N5*	908.17±	786.00±	786.46±	888.87±	872.35±	848.37±			
	99.59	130.78	112.97	96.21	93.82	115.68			
All	1,131.28±	1089.93±	1,062.96±	1100.36±	1,075.30±	1091.96±			
	212.25	232.81	215.44	195.05	188.06	208.95			

Table 13a. Corneocyte surface area ear
--

No significant differences between discs 1-5 * N5 significantly smaller than N1, 2, 3 and 4

	Atopic dogs ears (Mean±SD) (μm ²)									
	Disc 1	Disc 2#	Disc3*	Disc 4	Disc 5	Disc1-5				
AD1	1,063.87±	1,032.97±	1,260.56±	1,080.58±	1,179.29±	1,133.56±				
	158.63	108.77	169.43	165.67	142.8	169.71				
AD2	856.95±	1,117.51±	737.28±	906.41±	936.84 .:	910.17±				
	82.07	150.66	99.18	140.43	125.39	164.11				
AD3	1,23.54±	1,182.24±	1,518.94±	1,159.32±	1,276.74±	1,252.17				
	157.11	161.83	98.99	177.09	254.22	±222.61				
AD4	699.51±	1,115.27±	1,219.25±	1,154.43±	1,274.17±	1,112.18±				
	11 2.9 3	173.01	167.2	117.14	129.27	267.84				
AD5~	1,429.91±	1,505.46±	1,456.65±	1,482.13±	1,064.95±	1,396.22±				
	149.66	214.2	243.24	264.44	104.84	259.56				
All	1,034,76±	1192.65±	1243.64±	1,166.64±	1,166.59±	1,160.86±				
	282.64	223.93	318.73	273.91	213.93	272.28				

* Disc3 and #Disc 2 significantly bigger than Disc 1 ~ AD5 significantly bigger than AD1, 2 and 4 ‡ AD2 significantly smaller than AD1, 3, 4 and AD5

	Cats ears (Mean±SD) (μm²)									
	Disc 1*	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5				
C1	1,017.67±	1,209.12±	1,183.39±	1,311.74±	1,281.46±	1,200.68±				
	196.74	171.39	129.59	142.05	218.66	197.07				
C2	1,069.80±	1,184.45±	1,043.28±	1,113.28±	1,248.98±	1,121.13±				
	129.1	224.72	296.05	210.38	122.63	211.12				
C3#	1,229.81±	1,418.24±	1,309.68±	1,446.31±	1,553.25±	1,379.85±				
	188.69	332.37	192.65	207.14	305.95	257.57				
C4	1,121.60±	1,128.86±	1,065.57±	1,072.90±	1,011.76±	1,093.85±				
	117.89	100.79	136.1	153.49	230.32	158.5				
C5~	1,161.38±	1,357.03±	1,391.12±	1,430.52±	1,236.57±	1,315.32±				
	224.34	226.80	281.11	147.75	180.27	231.35				
All	1,120.59 \pm	1,258.35±	1,202.17 ±	1 ,271.43 ±	1 ,258.21 ±	1,183.33±				
	188.16	237.13	242.46	231.92	264.62	232.73				

Table 13c. Corneocyte surface area ears

* Disc 1 significantly smaller than Discs 4 and 5
C3 significantly bigger than C1, 2 and 4
~ C5 significantly bigger than C2 and 4

Table	14a.	Corneoc	vte	surface	area	thorax
						•

	Normal dogs thorax (Mean±SD) (μm²)								
_	Disc 1	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5			
N1	1,251.51±	1,126.04±	1,246.77±	1,336.07±	1,251.93±	1,242.46±			
	217.31	239.45	150.78	131.35	220.61	200.2			
N2	1,051.62±	1,251.35±	1,391.06±	1,377.73±	1,426.28±	1,299.61±			
	160.85	182.28	128.68	243.49	172.15	222.44			
N3~	1,413.21±	1,418.87±	1,467.52±	1,286.50±	1,486.63±	1,414.55±			
	171.24	228.79	165.32	170.9	202.44	194.67			
N4*	1,583.76±	1,464.43±	1,503.64±	1,540.98±	1,531.76±	1,524.91±			
	219.17	299.48	16 2.6 5	298.04	131.93	226.44			
N5#	1,617.90±	1,531.92±	1,490. 79 ±	1,431.00±	1,505.71±	1,515.47±			
	91.06	196.43	142.41	161.48	125.42	154.3			
All	1383.60±	1,358.52±	1,419.96±	1,394.45±	1,440.46±	1,399.40±			
	273.44	268.98	173.28	219.8	195.69	229.6			

No significant differences between discs 1-5 * N4 significantly bigger than N1, 2 and 3 #N5 and ~N3 significantly bigger than N1 and 2

					COMPANY OF A DESCRIPTION OF A DESCRIPTIO				
Atopic dogs thorax (Mean±SD) (μm²)									
	Disc 1	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5			
AD1	1,247.35±	1,437.98±	1,101.52±	1,110.08±	1,100.57±	1,210.99±			
	323.74	255.95	237.27	335.17	317.79	303.05			
AD2	1,307.85±	1,391.32±	1,458.90±	1,390.47±	1,329.60±	1,387.97±			
	191.7	222.68	118.92	129.96	148.58	180.37			
AD3~	1,399.20±	1,333.94±	1,543.14±	1,466.59±	1,313.97±	1,398.96±			
	195.93	261.45	179.08	334.75	134.3	248.13			
AD4#	1,597.22±	1,723.51±	1,577.18±	1,772.55±	1,502.05±	1,636.83±			
	204.3	178.12	190.2	239.01	157.39	207.55			
AD5*	1,708.66±	1,466.21±	1,456.38±	1,558.77±	1,592.68±	1,563.59±			
	197.94	165.7	254.58	207.07	135.84	203. 9 8			
All	1,452.06±	1,485.15±	1,428.74±	1,456.91±	1,375.49±	1,439.67±			
	281.1	245.14	274.53	318.97	246.21	274.8			

Table 14b. Corneocyte surface area thorax

No significant differences between discs 1-5

* AD5 and #AD4 significantly bigger than AD1, 2 and 3

~AD3 significantly bigger than AD1

Table 15a. Corneocyte surface area groin

Normal dogs groin (Mean±SD) (μm²)									
	Disc 1	Disc 2	Disc3	Disc 4	Disc 5	Disc1-5			
N1	1,364.14±	1,308.43±	1 ,3 16.14±	1,300.40±	1 ,27 1.71±	1,312.17±			
	183.34	202.53	132.84	184.94	123.6	164.19			
N2*	1,721.94±	1,641.21±	1,630.60±	1,794.27±	1,602.62±	1,678.13±			
	254.8	259.83	179.65	116.99	176.6	208.67			
N3#	1,457.36±	1,522.01±	1,498.72±	1,430.90±	1,433.21±	1,468.44±			
	173.94	114.98	249.31	242.92	150.98	189.12			
N4~	1,405.62±	1,344.22±	1,299.91±	1,608.70±	1,509.36±	1,433.56±			
	126.13	144.26	136.6	302.87	171.02	212.74			
N5	1,296.18±	1,309.61±	1,226.31±	1,297.67±	1,301.76	1,286.30±			
	161.04	210.15	176.47	175.70	221.11	184.89			
All	1,449.05±	1,425.10±	1,394.34±	1,486.39±	1,423.73±	1,435.72±			
	230.61	229.3	228.42	281.97	207.05	236.87			

No significant differences between discs 1-5

* N2 significantly bigger than N1, 3, 4, and 5

N3 and ~ N4 significantly bigger than N1 and 5

MBA 448 (28 YEA) 467 HEL	Atopic dogs groin (Mean±SD) (μm²)									
AD1 ‡	965.22±	1,063.21±	1,138.99±	1 ,214 .99±	$1,085.60\pm$	1,099.52±				
	254.25	413.20	232.39	272.78	188.08	273.18				
AD2	780.05±	812.26±	796.41±	$661.57\pm$	$1,055.70 \pm$	825.10±				
	87.72	84.51	103.83	88.41	168.28	165.97				
AD3#	1,656.34±	1,333.17±	1,414.81±	1,460.66±	$1,474.47\pm$	$1,\!476.10\pm$				
	130.09	2 2 7.49	290.97	221.23	188.27	227.48				
AD4*	1,688.51±	1,667.44±	1,569.39±	$1,518.54\pm$	$1,\!585.86\pm$	$1,\!631.91 \pm$				
	224.54	408.53	245.69	286.16	142.48	276.75				
AD5~	1,181.22±	1,130.46±	1,098.88	1,181.74±	$1,\!058.43\pm$	$\textbf{1,216.13} \pm$				
	233.71	193.17	213.65	181.51	158.89	268.10				
All	1,254.27±	1,223.05±	1,212.79±	1,213.02±	1,259.67 ±	1,249.74±				
When a full charge of the	413.76	421.48	348.01	367.94	287.72	368.55				

Table 15b. Corneocyte surface area groin

No significant differences between discs 1-5 * AD4 significantly bigger than AD1, 2, 3, and 5 # AD3 significantly bigger that AD1, 2 and 5 ~AD5 and ‡AD1 significantly bigger than AD2

Table 15c. Corneocyte surface area groin

	Cats groin (Mean±SD) (μm²)									
C1~	2,044.77±	2,103.23±	2,080.69±	2,022.30±	1,680.96±	1,986.75±				
	208.86	130.24	423.21	181.32	177.64	276.04				
C2 ‡	1,103.03±	1,222.70±	1,513.67±	1,338.72±	1,753.24±	1,404.73±				
	111.22	140.85	206.2	176.3	229.82	295.85				
C3	1,148.41±	1,170.70±	1,199.26±	1,327.68±	1,257.95±	1,227.36±				
	215.11	153.42	234.6	176.57	196.67	195.21				
C4#	2,238.97±	2,003.97	2,021.59±	2,130.68±	2,053.71±	2,077.76±				
	273.52	387.64	.181.27	285.27	172.52	269.7				
C5*	2,205.50±	2,098.67±	2,175.93±	2,139.09±	2,177.81±	2,163.95±				
	109.27	206.3	153.73	173.76	236.42	257.58				
All	1,755.41±	1,719.31±	1,796.40±	1,778.60±	1,810.82±	1,772,11±				
	540.77	464.64	452.1	416.23	434.98	461.11				

No significant differences between discs 1-5

* C5 significantly bigger than C1, 2 and 3

C4 and ~C1 significantly bigger than C2 and 3

‡ C2 significantly bigger than C3

Comparison of corneocytes from haired and nonhaired sites

The numbers of polygonal type II corneocytes were highest in the sparsely haired sampling site 1 at the base of the pinna becoming progressively less common in discs 2 to 3. The converse finding was obtained for type I corneocytes.

Dog	 Corneocyte 						
1	Туре	Image1	Image2	lmage3	Image4	lmage5	Total
Disc1	1	10	2	1	7	1	21
	11	13	7	12	8	13	53
Disc2	2 1	12	16	13	14	7	62
	11	4	5	5	7	2	23
Disc3	5 I	26	40	12	27	31	136
	11	5	1	2	2	2	12
2	Туре	Image1	lmage2	lmage3	lmage4	lmage5	Total
Disc1	Î	1	0	Ō	1	2	4
	I I	14	1 1	17	22	23	87
Disc2	2	5	3	2	0	1	11
	11	24	17	17	22	7	87
Disc3	3 L	1	4	9	6	1	21
	11	4	2	5	2	5	18
3	Туре	Image1	Image2	lmage3	lmage4	lmage5	Total
Disc1	1	2	2	2	3	5	14
	11	9	9	12	5	9	44
Disc2	2	2	6	9	4	4	25
	11	9	9	5	5	15	43
Disc3	3 1	4	8	4	3	4	23
	11	2	7	1	3	2	15
4	Туре	lmage1	Image2	lmage3	lmage4	lmage5	Total
Disc1	1 1	1	4	0	1	3	9
	II	13	3	13	14	8	51
Disc2	2 1	1	4	6	0	3	14
	11	10	19	7	22	9	67
Disc3	31	12	5	5	7	5	34
	11	8	11	2	1	1	23
5	Туре	Image1	lmage2	Image3	lmage4	lmage5	Total
Disc1	1	0	1	3	1	2	7
	11	24	8	11	7	3	53
Disc2	21	7	7	5	3	3	25
	11	3	3	2	5	12	25
Disc3	3 I	9	4	4	5	8	30
	H	3	2	2	2	9	18

Table 16. Canine corneocytes from haired area to unhaired area





collected from canine ear

Columns represent the total counts for 5 dogs from each of three sites.

Cat	Corneocyte	The Prophyle of		A PARA			
1	Туре	Image1	Image2	Image3	Image4	Image5	Total
Disc1	1	1	1	0	2	0	4
		7	10	5	14	17	53
Disc2	I	0	1	3	4	3	11
	Ш	5	22	15	8	8	58
Disc3	1	4	3	0	1	2	10
	Ш	11	3	7	14	3	38
2	Туре	Image1	Image2	Image3	Image4	Image5	Total
Disc1	1	1	3	1	0	2	7
	11	19	18	11	18	7	73
Disc2	I	0	1	3	1	5	10
	11	20	10	5	5	1	41
Disc3	I.	1	2	3	4	3	13
	П	2	0	2	4	1	9
3	Туре	Image1	Image2	Image3	Image4	Image5	Total
Disc1	I	0	0	0	0	0	0
	П	14	14	25	12	21	86
Disc2	1	0	0	4	0	0	4
	П	14	20	11	19	27	91
Disc3	. I	0	5	2	4	6	17
	П	26	1	4	1	3	35
4	Туре	Image1	Image2	Image3	Image4	Image5	Total
Disc1	1	0	0	0	0	0	0
	11	24	4	28	15	18	89
Disc2	: I	0	0	0	1	0	1
	11	6	9	3	14	12	44
Disc3	i I	4	0	1	1	2	8
	11	4	1	1	6	0	12
5	Туре	Image1	Image2	Image3	Image4	Image5	Total
Disc1	I	0	1	0	0	0	1
	11	12	6	9	24	15	66
Disc2	: I	1	1	0	1	0	3
	Ш	21	17	17	14	12	81
Disc3	; I	2	1	1	1	2	7
	11	1	3	11	2	0	17

Table 17. Feline corneocytes from haired area to unhaired area



Figure 17. Histogram of numbers of type I and type II corneocytes



collected from feline ear

Columns represent the total counts for 5 cats from each of three sites.

DISCUSSION

The average diameter of canine and feline corneocytes in this study was approximately 38-48 µm. The equivalent figures for surface area were found to be 1100-1800 μ m². This is similar to that reported for man (El Gammal, Kligman and el Gammal, 1997)(30-40µm) and larger than that reported for the mouse (Schatzlein and Cevc 1998) (25-30µm). This study appears to be the first to define the size and shape of canine and feline corneocytes. Because not all sites were sampled from each individual animal and some sites required clipping before sampling it was not possible to directly compare the various sampling sites of the different animal groups studied. This was not the major purpose of the study but in general, cat corneocytes were larger than those from dogs. Corneocytes from the groin tended to be largest and corneocytes from the ears tended to be smaller than those from the other two sites. There was therefore some species and site differences appreciated which may justify further larger studies. There was little variation in the size of corneocytes between each of the five discs collected from the same site suggesting that samples from the same site would be suitable for comparative studies in an adhesion assay and that the corneocytes do not vary in size between the layers sampled.

In man corneocyte size has been shown to be determined by the epidermal turn over rate which in turn may be influenced by body site, sex and inflammatory skin disease (Fluhr, Kao, Jain, Ahn, Feingold and Elias, 2001; Kashibuchi *et al*, 2002). Because inflamed skin is likely to have an increased epidermal turnover time it could be anticipated that corneocytes from atopic dogs would be smaller than those from normal dogs. There were individual atopic dogs that did appear to have small corneocytes but in general there was no overall tendency for atopic corneocytes to be smaller than normal ones. In general there were significant variations in

size between individual animals, which may suggest possible breed, sex or age influences. Much larger studies would be required to identify any such differences positively.

The two types of corneocytes identified based on shape are of interest. The elongated type I corneocytes appeared to be more common in haired skin areas suggesting that their origin is the hair follicle. The lighter staining and polygonal type II corneocytes are more typical of those identified in human skin and are likely to originate from the interfollicular skin area.

Type II corneocytes being flat and lightly staining are better suited for use in adhesion assays and particularly those that employ image analysis as a counting tool. When counting bacteria using image analysis good distinction between the bacteria to be counted and the background is necessary. Sheets of type II corneocytes could be most readily obtained from the inner pinnae of both the dog and cat and this site was considered to be most suited for the collection of corneocytes to be used in an adherence assay.

2

strains a failer.

CHAPTER 3

An Adhesion Assay To Quantify Bacterial Adherence To Canine And Feline Corneocytes

LITERATURE REVIEW

Introduction

Although bacteria were observed to adhere to red blood cells as far back as 1908, it was not until the 1970's that bacterial adherence was studied in any detail (Gibbons and van Houte, 1971; Gibbons and Houte, 1975). Adhesion is known to be essential for maintaining the normal microflora in or on their host, and it is also the crucial first stage in any infectious disease. It is now recognized that bacterial adhesion is a prerequisite to colonization and infection (Feingold, 1986; Roth and James, 1989). Over the last twenty years, research has revealed the enormous complexity underlying the phenomenon of bacterial adhesion. The initial research goal was to understand the mechanism of attachment and its effects on bacteria as well as the host. As research progressed, however, it became evident that many different attachment mechanisms exist. Various studies have shown that certain species of bacteria will adhere preferentially to particular tissues whilst other bacteria fail to adhere. For example, Staphylococcus mitis adheres well to the surface of teeth and to cheek epithelium while Escherichia coli adheres well to intestinal epithelium but not to oral tissues (Gibbons and van Houte, 1971; Gibbons and Houte, 1975; Aly and Bibel, 1993). Similarly, host species specificity can be shown (Gibbons, Spinell and Skobe, 1976; Valentin-Weigand, Chhatwal and Blobel, 1988). These diverse forms of adhesion are the results of numerous evolutionary pressures, and each may be part of a larger survival strategy. This suggests that bacteria have developed specific adhesion molecules to allow them to colonise and flourish in specific niches on specific host animal species. The ability of bacteria to adhere has obviously evolved as an important factor and is one of the significant initial steps in the process of colonization (Feingold, 1986).

le inte del constitue

i. K

a ser a la ser a la ser a la ser a la ser a s

14-10-11

Mechanisms of bacterial adherence and infection

Bacterial adhesion is a complex process influenced by both the host and the organism. Bacteria adhere only to complementary substrata. They adhere by ionic or Coulombic interactions, by hydrogen bonding (Pimentel and McClellan, 1960), by the hydrophobic effect (Duncan-Hewitt, 1990), and by coordination complexes involving multivalent metal ions. Adhesion is highly stereospecific in that a bacterium will bind to a substratum only if the substratum possesses a certain kind of receptor. Bacteria possess surface adhesion molecules, which influence their ability to bind to cells. For those bacteria that possess pilli and fimbriae such as the *Enterobacteriaceae* those structures have adhesins at their tips that enables specific binding to receptors. Of the staphylococci *S. aureus* has been the most studied and here teichoic acids, fibronectin binding proteins, protein A, collagen binding proteins and capsular polysaccharide have all been shown to be involved in adhesion (Van Belkum, Kools-Sijmons and Verbrugh, 2002).

Several studies have shown that adhesion is dependent on time, temperature, and concentration of bacteria as well the influence of certain disease states (Berg *et al*, 1984; Harvey and Noble, 1994; Forsythe *et al*, 2002; Saijonmaa-Koulumies and Lloyd, 2002). Although there is evidence that adhesion by staphylococci may play a part in bacterial infections in both human and canine atopic dermatitis (Cole and Silverberg, 1986; McEwan, 2000) there is currently little evidence that the ability to adhere by *S. intermedius* can be considered a virulence factor. Using *S. intermedius* strains isolated from normal and infected canine skin Cree and Noble (1995) found increased adherence by pyoderma strains while Saijonmaa-Koulumies and Lloyd (2002) found no such relationship. It may be better to think of adhesion as an indirect virulence factor in that

68

and the second of the second of the second second

that the difference of the second

¥**************

1. St. 1.

after attachment direct virulence factors can be delivered at close proximity to host cells (Van Belkum, Kools-Sijmons and Verbrugh, 2002).

A number of approaches have been used to examine the relationship between adhesion and infectivity. One method was to compare the infectivity phenotypes or genotypes expressing or not expressing adhesions. Another method to assess the role of adhesion in the infectious process is to compare the sensitivity or resistance to infection by hosts that either over express or do not express, respectively, adhesin receptors. In general, there is a positive correlation between the ability of the tissue cells to bind the bacterial pathogen and susceptibility of the host to develop infection caused by the pathogen. Ofek and Beachey (1980) reviewed the bacterial binding capacity of epithelial cells from infection-prone people finding it to be greater than that from similar cells of people not prone to infection. One piece of convincing evidence described the relationship between resistance to infection and the capacity of host cells to bind pathogenic bacteria (Sellwood, Gibbons, Jones and Rutter, 1975). Intestinal epithelial cell brush borders isolated from piglets that were resistant to the diarrhoeal disease failed to bind E. coli K88 bacteria in vitro, whereas those from susceptible animals were able to bind the organisms in high numbers. (Mouricout, Petit, Carias and Julien, 1990) were able to block the adhesion by E. coli K99 to calf intestine, and hence disease, by administration of glycoprotein glycans derived from bovine plasma. Thus demonstrating that adhesion is important in the infection process and its study can lead to potentially novel treatments.

There are a number of reasons to support the view that adhesion is important in infectivity. One of the most widely accepted is that bacterial adhesion endows the pathogen with the ability to withstand cleansing mechanisms operation on mucosal and endothelial surfaces 가슴을 알았다. 아들은 옷을 가는 것을 가지? 않는 것 같은 것을 알았다.

and a second second

No. of the other states of

(Ofek and Beachey, 1980; Beachey, 1981). It has been pointed out (Freter, 1978; Freter, O'Brien and Halstead, 1978; Freter, 1980) that successful association of the pathogen with mucosal surfaces such as intestinal cells is a complex process. Adhesion to animal cells seems to confer at least three main advantages in addition to the ability to withstand mechanical cleansing mechanisms, including growth advantages, toxin targeting, and escape from antimicrobial or antitoxin activities (Ofek and Doyle, 1994).

Methods of assessing bacterial adherence

Several methods have been used to quantify microbial adhesion to cells (Evans-Hurrell, Adler, Denver, Rogers and Williams, 1993; Ofek and Doyle, 1994), including direct cell counting using light microscopy (Tavendale, Jardine, Old and Duquid, 1983; Miyake, Kohada, Fujii, Sugai and Suginaka, 1989; Miyake, Sugai, Kohada, Minagi and Suginaka, 1990; Romero-Steiner, Witek and Balish, 1990), plate counting (Cooper, Jeffery-Wiseman, Bortner, Wheat and Wadstrom, 1990), and radiometric assays (Jordens, 1990; Verbrugh, 1990; Wyatt, Poston and Noble, 1990). Generally, cells are incubated with the microbe under test either in vivo or in vitro. Non-adherent bacteria are commonly removed by either filtration or by differential centrifugation. Adhesion is then quantified either by direct visualisation under light or electron microscopy or quantitatively by the use of markers such as fluorescent dyes or radioactivity (Bartelt and Duncan, 1978; Bibel, Aly, Lahti, Shinefield and Maibach, 1987; Valentin-Weigand, Chhatwal and Blobel, 1987; Kanzaki, Morishita, Akiyama and Arata, 1996). The major limitations of these methods include extended time for analysis, operator fatigue, subjectivity, and in the case of radiometric assays, an inability to discriminate between bacteria specifically adhering to the epithelial tissue and those attached to the inert support (Evans-Hurrell et al, 1993). Experimental design is also

70

and the star was a weather

11 12

2月27日、第七次、安安、季季を置ける

ι.

الكافية تتطليك فالمحاد

interesting to the second s

A week

important and several studies have shown that fluorescein used as a marker may significantly alter bacterial adherence, presumably by modification of bacterial adhesins (Evans-Hurrell *et al*, 1993; Barthelson, Hopkins and Mobasseri, 1999).

71

As expected, adherence is dependent on temperature, bacterial concentration and also pH (Valentin-Weigand, Chhatwal and Blobel, 1987; Saijonmaa-Koulumies and Lloyd, 1996). However, one study showed that too high a bacterial concentration may result in cytotoxic effects, which may increase or decrease adhesion by exposing, inducing or decreasing binding sites (Charboneau and Rubins, 1998).

Staphylococcal adherence in human and canine atopic dermatitis

There is a growing body of evidence to suggest that all of the pyogenic Gram-positive cocci rely on multiple adhesins in order to adhere avidly to substrata (Hasty, Ofek, Courtney and Doyle, 1992). The most successful pyogenic cocci have the ability to adhere to and grow on mucosal tissues without causing symptoms. This is commonly known as the carrier state.

In human and canine atopic dermatitis patients, bacterial skin disease is a very common and important complication. The major microbial pathogens involved are the pathogenic staphylococci; *Staphylococcus aureus* in man (Leyden, Marples and Kligman, 1974; Dahl, 1983; Hanifin and Homburger, 1986; Masenga, Garbe, Wagner and Orfanos, 1990) and *Staphylococcus intermedius* in dogs (Phillips, Jr. and Kloos, 1981; Berg *et al*, 1984; Biberstein, Jang and Hirsh, 1984; Cox *et al*, 1984; Ihrke, 1987). In human atopics, the normal microflora changes with an increase in density and carriage of pathogenic staphylococci on the skin (Aly *et al*, 1977). A similar situation is likely to exist in the dog

(Kristensen and Krogh, 1978). Mason and Lloyd (1989) investigated the hypothesise that changes in the skin surface defence mechanisms in atopic dogs led to their susceptibility to invasion by *S. intermedius* and suggested that the pathogenesis of pyoderma secondary to hypersensitivity may be via an effect on epidermal permeability, promoting penetration of staphylococcal antigens from the stratum corneum which then cause the lesions of pyoderma.

72

an the second of the second state and a second state of the second second second second second second second se

Cole and Silverberg (1986) were the first to hypothesise that human atopic dermatitis patients may be susceptible to S. aureus infections due to increased adherence to atopic corneocytes. Foster and Höök (1998) reviewed S. aureus adherence to host extracellular matrix. They concluded that adherence is mediated by protein adhesions of the MSCRAMM (microbial surface components recognizing adhesive matrix molecules) family, such as fibronectin-binding, fibrinogen-binding and collagen-binding proteins. For adhesion to the skin the closely related fibronectin-binding proteins (FnBPA and FnBPB) appear to be important. Most strains of S. aureus express both fibronectin-binding proteins which are encoded by two closely linked genes (Greene, McDevitt, Francois, Vaudaux, Lew and Foster, 1995; Johansson, Flock and Svensson, 2001). More recent work has identified that S. aureus does bind to fibrinogen and fibronectin in skin (Kanzaki et al, 1996; Cho, Strickland, Boguniewicz and Leung, 2001), who also provided further evidence of enhanced binding by S. aureus to human atopic skin compared to normal human skin.

In atopic dogs, there are few published studies on the adherence by *S. intermedius*. Adherence by *S. intermedius* to normal canine corneocytes was shown to depend on duration of incubation, temperature of incubation and the concentration of bacteria used (Saijonmaa-Koulumies and Lloyd, 2002). McEwan (2000, 2002c) found a statistically

significantly greater adherence by *S. intermedius* to corneocytes from atopic dogs and dogs suffering from pyoderma when compared with a normal group. Forsythe *et al.* (2002) suggested that *S. intermedius* adheres to canine corneocytes by a specific receptor-ligand interaction and there may be a breed predisposition for adherence and that different body sites also influence bacterial adherence.

73

From the available evidence of the literature the ability of staphylococci to adhere to corneocytes is likely to be important in the development of pyoderma. In both human and canine atopic dermatitis pathogenic staphylococci adhere well to corneocytes. A major limitation of adherence studies is the time consuming and tedious method of many adherence assays. A simple and quick method to assess bacterial adherence to corneocytes would be a useful tool for the investigation of this phenomenon, and this was the goal of the studies that follow.

AIMS

The aims of the following studies were to:

- 1. Develop a simple assay to quantify bacterial adherence to canine and feline corneocytes
- 2. To develop image analysis techniques to count bacteria adherent to canine and feline corneocytes

MATERIAL AND METHODS

Populations studied

Three groups of animals were used as sources of corneocytes:

- 1. Normal dogs
- 2. Dogs suffering from atopic dermatitis
- 3. Normal cats

These groups were defined as detailed in Chapter 2 of this thesis. None of the animals used in the studies had received either systemic or topical treatments for a period of at least three weeks prior to corneocyte sampling.

Corneocyte collection

Corneocytes were collected in a similar fashion to that described in Chapter 2 of this thesis. The skin area to be sampled was first prepared by removing surface debris by five successive applications of ordinary adhesive tape. Corneocyte samples for the adhesion assay were collected by applying a 25mm D-Sqaume® disc that was gently pressed onto the skin surface once. Preliminary studies (study 4) indicated that up to 5 consecutive D-Squame® discs from the same area would sample corneocytes with similar adhesive properties.

Bacteria

All staphylococci used in the study were collected from clinical cases. Specific species of staphylococci were identified by a microtray method (API Staph. System. BioMerieux, Lyons, France). The *Micrococcus* sp. used was isolated from normal human skin and identified by conventional microbiological techniques.

- 1. Staphylococcus intermedius. Isolated from purulent material collected from a case of canine deep pyoderma (muzzle furunculosis)
- 2. Staphylococcus hominis. Isolated from a bovine nasal swab
- 3. Staphylococcus aureus. Isolated from a case of bovine mastitis
- 4. Staphylococcus hyicus. Isolated from a case of bovine mastitis
- 5. Micrococcus sp. Isolated from human skin

All bacteria were cultured on sheep blood agar, subcultured into liquid medium (Oxoid Nutrient Broth. No. 2. Unipath Ltd., Basingstoke, Hampshire, England) and frozen at -70° C in one millilitre aliquots.

Suspensions of *S. intermedius* and *S. hominis* increasingly diluted in sterile phosphate buffered saline (PBS) were made at optical densities (OD) ranging from 0.2-0.8 at 570 nanometers (Cole and Silverberg, 1986) on a spectrophotometer (Cecil instruments, Cambridge, England) and plated on sheep blood agar at ten fold dilutions to determine the number of colony forming units (CFU) per ml. By extrapolation an OD₅₇₀ of 0.15

Anthe Antheory and

1

.

- and the set of the second of the second se

And the second s

was found to give approximately 1×10^{6} CFU per ml and this dilution was used for each bacterial suspension in the final assay.

76

مىيە بەيغۇرىرى بەيغۇر ، ئەيمىيە بەيدىيە بەيدىيە بەيدىيە بەيدە بەيدىيە بەيدىيە بەيدىيە بەيدىيە بەيدىيە بەيدىيە ب

When required, a frozen aliquot of staphylococci was thawed and plated on sheep blood agar and incubated at 38° C for 24 hours. Colonies were harvested and placed in a centrifuge tube with 10 millilitres of sterile PBS. The cocci were washed in twice in PBS by centrifugation and the resulting suspension adjusted to an OD of approximately 0.15 at 570 nanometers. Preliminary studies (study 1) indicated that an incubation time of 45 minutes and a bacterial concentration of 0.15 at OD₅₇₀ was suited to the adhesion assay used.

Adhesion assay

A 25mm D-Squame® disc with adherent corneocytes was placed corneocyte surface uppermost on a microscope slide. A 0.5 ml bacterial suspension in PBS was placed over the corneocyte layer and incubated at 38°C for 45 minutes in a moist chamber held within a water bath (figures 18 and 19). After incubation the disc was washed in tap water, stained for 10 seconds with crystal violet and washed to remove excessive stain (figure 20). Washing and staining procedures followed a standard protocol. After washing the disc was placed on a glass microscope slide and air-dried.



Figure 18. Moist chamber within water bath

Figure 19. D-Squame® discs and bacteria suspension prepared for incubation





Figure 20. Slides after washing and staining

Image acquisition and bacteria counting

Adherent bacteria were counted by an image processing and analysis program (UTHSCSA ImageTool Version 3.0 and The Image Processing Tool Kit v.5, Reindeer Graphics Inc., Asheville, NC). UTHSCSA ImageTool is an image processing and analysis program for Microsoft Windows available from the following web site http://ddsdx.uthscsa.edu/dig/itdesc.html.

From each D-Squame® disc ten images at x 630 magnification were collect and saved as black and white TIFF files. The process of image thresholding and counting was semi-automated by a macro or script

written by the author (Appendix J). This script was identified as "Bacteria Count".

79

「おいている」

And the strength of the state of the

The various steps in the image analysis bacteria counting are depicted in figures 21 - 24. Figure 21 shows the initial image collected from the disc and the creation of a 400 x 400 pixel box (an area of approximately 13060 μ m², approximately the surface area of 10 corneocytes), which could then be manually placed over a selected counting area. Figure 22 shows the thresholding process. Automatic bilevel thresholding using the plugin provided generally produced good separation of bacteria from corneocytes but the program also allowed for manual selection of the threshold value. Next (figure 23) a round filter was applied which allowed better selection of bacteria from "background noise" created by debris and corneocyte folds. Finally the total count for the area was displayed (figure 24). For each study a total of ten images were collected and counted from each disc this gave a final count, which equated to a surface area equivalent to 100 corneocytes.



Figure 21. Selection of the counting area

A 400 x 400 pixel box is generated which can be manually placed over the desired counting area.





A greyscale threshold value is selected that separates bacteria from background to produce a binary image.



Figure 23. Application of a round filter

A round filter was chosen to differentiate bacteria cocci from non-round background "noise"

Figure 24. The final count



After application of the round filter the final count is displayed.

For each study control discs were included that consisted of discs incubated with sterile PBS and processed as for the discs incubated with bacteria. Control discs were scanned for adherent bacteria and were subjected to image analysis and counting.

Study 1. The effects of time, temperature and bacterial concentration on adhesion

Time study

A set of 5 D-Squame® discs with adherent corneocytes were obtained from the right pinna of 3 dogs and 2 cats. All animals were normal and free of skin disease apart from Cat 2, which had been diagnosed as suffering from atopic dermatitis. Corneocytes were incubated with either *S. intermedius* or *S. hominis* for 15, 30, 45 and 60 minutes. From each disc 10 images were obtained and stored as TIFF files for counting. The number of adherent bacteria was determined using image analysis and the "Bacteria Count" script.

Dog 1. Three-year-old male Greyhound Dog 2. Ten-year-old female neutered Labrador retriever Dog 3. Four-year-old male Greyhound Cat 1. Eight-year-old female neutered domestic shorthair Cat 2. Ten-year-old male neutered domestic shorthair

Temperature study

Two dogs and one cat were used in this study. All animals were normal with no skin disease. Samples were collected from the right and left ears of all three animals and incubated at 38°C and 4°C as described above. Control samples were incubated with sterile PBS at 38°C and 4°C. Bacteria were counted using the method described above.

Dog 1. Ten-year-old female neutered Labrador retriever Dog 2. Four-year-old male Cross breed Cat 1. Three-year-old FN Domestic shorthair 84

Concentration study

Bacterial suspensions of *S. intermedius* and *S. hominis* were made in sterile PBS over a range of optical densities (570nm) from 0.045 to 1.156 and were incubated at 38°C with corneocytes using the assay technique as above. Two cats and three dogs were used in this study. All animals were normal with no skin disease.

Cat 1, A six-year-old female neutered domestic shorthair

Cat 2. A three-year-old male neutered domestic shorthair

Dog 1. A nine-year-old male Bull terrier

Dog 2. An eleven-year-old female neutered German Shepherd dog

Dog 3. A three-year-old male Greyhound

Study 2. Comparison of visual counting and counting by image analysis

Ten images were randomly selected from each of the following four groups:

- 1. Cat corneocytes incubated with S. intermedius
- 2. Cat corneocytes incubated with S. hominis
- 3. Dog corneocytes incubated with S. intermedius
- 4. Dog corneocytes incubated with S. hominis

Using the image analysis software a centrally placed rectangular counting area 200 x 200 pixels ($3265 \ \mu m^2$) was generated (Appendix J). The numbers of cocci in each counting area, including those at the edge of the counting area, were then calculated by firstly direct visual (manual) counting by two operators using the image analysis program count and

and an a faith an a wate a start of the activity of the start of the

يستكمأ أيراك ستقتيب بالكرام الالالال

tag tool and then by the image analysis program using the Image Tool script "Bacteria Count".

Study 3. Repeatability study

Twelve D-Squame® discs (six incubated with *S. intermedius* and six incubated with *S. hominis*) were randomly selected from a pool of discs that had been prepared for previous studies. For each D-Squame® disc the operator selected and stored 10 images as TIFF files. The number of bacteria in each image was then calculated using the image analysis program and the script "Bacteria Count". The same operator repeated this entire procedure (selected different images) a second time after a time interval of at least one week. The operator was blinded as to the identity of individual discs. The two sets of data were then compared to assess repeatability of the counting method.

Study 4. A comparison of corneocytes collected from the ear and groin

Corneocytes were collected from the pinna and groin area in each of four dogs for comparison. In two cats corneocytes were collected from the pinna only. From each site five consecutive D-Squame® discs were obtained. The corneocytes so collected were incubated with either *S. intermedius* or *S. hominis*. A sixth disc was collected from each site and used, as a control disc and incubated with sterile PBS. D-Squame® discs and adhesion assays were performed as detailed above. The operator was blinded as to the identity of individual discs.

Dog 1. Two years and 2 month old male neutered Bull terrier suffering from atopic dermatitis

an tit i militie and an

Ń

Dog 2. Three-year-old male neutered West Highland white terrier suffering from atopic dermatitis

Dog 3. Normal four years and 4 months old Labrador retriever

Dog 4. Normal five-year-old Grey hound

Cat 1. Normal three-year-old male neutered Domestic shorthair

Cat 2. Normal six-year-old female neutered Domestic shorthair

Study 5. Comparison of adherence by 5 different staphylococcal species and a *Micrococcus* to canine and feline corneocytes

Five species of staphylococcus and one *Micrococcus* were incubated with canine and feline corneocytes as described above. The species of bacteria used and their origin are detailed above. Five consecutive D-Squame® discs were obtained from the right pinna of five normal cats and dogs. A sixth disc was included as a control. The corneocytes collected from each animal were incubated with each of the five bacteria species and the adherent bacteria counted as described above. The operator was blinded as to the identity of individual discs.

Dog 1. Four-year-old male Greyhound

Dog 2.Three-year-old female neutered Greyhound

Dog 3. Ten-year-old female neutered Labrador retriever

Dog 4. Six-year-old female neutered Cocker spaniel

Dog 5. 4-year-old female neutered Cross breed

Cat 1. Five-year-old male neutered Domestic shorthair

Cat 2. One year and 2 months old female entire Exotic shorthair

Cat 3. Six-year-old female neutered Domestic shorthair

Cat 4. Two years and 3 month old male entire Exotic shorthair

Cat 5. Eleven-month-old female entire Exotic shorthair

and a second state of the second

STATISTICAL ANALYSIS

Statistical analysis and graphical presentation of data was conducted using computerised statistical packages, GraphPad Prism, version 4 (Graphpad Software Inc. San Diego CA). Comparisons between groups of more than two were conducted using simple one-way analysis of variance (ANOVA), where data was not normally distributed; Kruskal-Wallis ANOVA on ranks was used. To isolate the group or groups that differed from the others a multiple comparison procedure (Dunn's method) was employed. Where two groups were compared the paired t test was used (comparison of ear and groin, table 23). Comparison between two or more groups was also assessed by linear regression and calculation of the correlation coefficient. Bland-Altman plots were used to compare methods for assessing bacteria adherence. A value of p< 0.05 was considered to be significant.

and the second of the second of the second

Ì

RESULTS

Effect on time on adhesion to corneocytes (table 18 and figure 25)

For *S. intermedius* there was a clear increase in the number of adherent cocci with time. In contrast there appeared to be little increase in the numbers of adherent over time shown by *S. hominis*.

	15 minutes	30 minutes	45 minutes	60 minutes	Control
S. intermedius			a second seco		
	162.80±	185.20±	197.60±	194.20±	36.20±
Dog 1	36.11	24.01	32.15	37.46	22.88
A start the second start	108.30±	131.30±	248.90±	282.00±	24.80±
Cat 1	21.38	43.72	42.42	48.02	22.71
	186.40±	332.20±	385.80±	498.50±	38.70±
Dog 3	34.35	51.98	91.35	26.58	25.16
Mean	152.50	216.23	277.43	324.90	33.23
S. hominis					
A STATE OF A STATE OF A	47.30±	55.30±	53.80±	56.20±	28.50±
Dog 4	22.49	36.33	24.68	38.22	26.31
	45.00±	51.40±	50.90±	48.50±	8.80±
Dog 2	23.78	23.78	19.35	9.63	7.63
Mean	46.15	53.35	52.35	52.35	16.97

Table 18. Effect on time on adhesion to corneocytes (mean values ± standard deviation)



Figure 25. Effect of time on bacterial adhesion





Comparison of adherence at 38°C and 4°C (table 19 and figure 27) (mean values ± standard deviation)

For both *S. intermedius* and *S. hominis* there was a clearly increased adherence at 38°C compared to 4°C.

		Test s 38°C	amples Mean bacteria 4ºC	Conf a count ± SD 38°C	trol 4ºC
Dog 1	Right Ear 1	379.90± 71.90	157.50± 30.74	30.30± 13.50	29.10± 24.35
S. intermedius	Right Ear 2	371.00± 33.54	163.10± 26.30		
	Left Ear 1	315.50± 92.77	165.90± 30.37		
State of the second	Left Ear 2	376.00± 74.42	164.00± 26.06		
Dog 2	Right Ear 1	95.90± 31.48	41.20± 22.44	13.30± 8.97	18.50± 11.64
S. hominis	Right Ear 2	115.30± 42.35	22.80± 5.94		
	Left Ear 1	135.00± 31.76	34.60± 9.96		
A. C.	Left Ear 2	117.30± 30.97	54.20± 16.7		
Cat 1	Right Ear 1	384.00± 61.98	229.20± 52.97	30.80± 24.43	26.00± 23.40
S. intermedius	Right Ear 2	402.70± 53.34	187.60± 44.09		

Table 19. Comparison of adherence at 38°C and 4°C


Figure 27. Histogram of temperature effects

Effects of bacterial concentration on adhesion to corneocytes (table 20, figures 28 and 29)

In general bacteria adherence increased with increasing concentration. At higher bacteria concentrations (optical densities above 0.200) clumping of bacteria was commonly encountered. Clumping was rarely encountered at an OD of under 0.200.

		Mean bacteria count ± SD			
		OD at			
		570 nm	Count	Control	
Cat 1	S. intermedius	0.698	Clumped	28.70±	26.80
R ear		0.433	Clumped		
		0.321	Clumped		
College		0.133	239.30± 5	1.18	_
Cat 2	S. intermedius	0.476	Clumped	54.4±	26.72
R ear		0.303	Clumped		
		0.148	Clumped		
- Carlos - C		0.138	264.30± 30	0.43	
		0.119	158.90± 4	7.12	
Dog 1	S. intermedius	0.448	Clumped	21.3±	22.47
R ear		0.237	252.60± 42	2.16	
		0.095	206.40± 5	1.70	
- ANALIS -		0.045	170.50± 4	7.21	
Dog 2	S. intermedius	1.156	Clumped	24.3±	14.46
R ear		0.810	Clumped		
		0.413	424.00± 5	6.20	
Sec. Sec.		0.225	407.20± 4	7.06	
Contraction of the		0.143	303.40± 5	3.00	
Dog 2	S. intermedius	0.810	Clumped	36.6±	19.42
L ear		0.413	478.10± 5	1.61	
		0.225	340.60± 5	8.45	
A State State		0.143	249.40± 6	0.70	
Dog 3	S. hominis	0.443	121.00± 3	7.72 14.6±	9.97
R ear		0.243	60.60± 10	6.86	
ALC P		0.135	27.00± 9	.13	
		0.076	20.70± 5	.42	
Dog 3	S. hominis	0.443	47.90± 2	2.08 14.5±	10.12
L ear		0.243	34.70± 14	4.38	
The second		0.135	21.00± 7	.41	
NE EST		0.076	19.50± 4	.84	

Table 20.	Effects	of bacterial	concentration	on adhesion
-----------	---------	--------------	---------------	-------------

OD - optical density

600-S. intermedius Average bacteria count 500 S. hominis 400-300-200 100 0-0.0 0.2 0.3 0.4 0.5 0.1 Optical density at 570 nm

Comparison of bacterial counts by image analysis and visual counting

Image analysis correlated well with the visual counting by both operator 1 (R= 0.955) and operator 2 (R=0.835). Bland-Altman plots for both operators 1 and 2 when compared to image analysis showed similar results in that discrepancies between the two methods became larger when the bacterial counts were high.

Figure 28. Concentration of bacteria and adherence

Chapter 3

Table 21. Comparison	of bacterial	counts b	by image	analysis a	ind
	visual co	unting			

Dog		lmage analysis	Operator 1	Operator 2	Cat	Image analysis	Operator 1	Operator 2
S. hominis	1	6	9	8		32	32	40
	2	35	51	55		32	41	53
and see the set	3	13	15	16		7	6	10
In Allen States	4	29	35	32		6	6	6
	5	6	10	11		5	5	9
	6	48	49	60		7	5	6
	7	13	15	17		7	5	6
A States	8	12	13	13		8	7	8
La Maria Maria	9	5	3	3		24	21	20
	10	59	50	66		26	26	33
S. intermedius	1	85	60	63		109	129	204
	2	43	45	59		45	123	60
A State of the second s	3	90	97	111		79	81	101
	4	129	125	131		62	63	76
	5	97	98	88		111	137	180
	6	95	104	108		88	109	123
	7	108	120	112		70	84	119
All Marine and	8	121	122	115		105	138	197
Contraction of the second	9	86	90	89		51	65	73
A CARLE	10	83	88	102		44	57	77



Figure 30. Comparison of Image analysis counting with manual count

Figure 31. Bland-Altman Image analysis and Operator 1:Difference vs average



Figure 32. Bland-Altman Image analysis and operator 2:Difference vs average



Repeatability of counting method (table 22, figure 33)

There was no significant statistical difference between the results of repeated counts of adherence on two separate occasions (paired t-test p = 0.228). The two counts also showed a high degree of correlation (R= 0.934).

97

Case	Bacterium used	Count 1	Count 2
Dog 1	S. intermedius	290.8	249.3
Dog 2	S. hominis	56.7	58.3
Dog 3	S. intermedius	285.6	296.2
Dog 4	S. hominis	83.5	37.1
Dog 5	S. intermedius	126.5	110.2
Dog 6	S. hominis	10.9	24.8
Cat 1	S. intermedius	107.0	133.9
Cat 2	S. hominis	62.5	65.2
Cat 3	S. intermedius	76.7	79.4
Cat 4	S. hominis	28.7	18.7
Cat 5	S. intermedius	173.1	130.1
Cat 6	S. hominis	47.6	41.0

Table 22. Repeatability of counting method

0-

Ó



200

count 1

100

400

300

Comparison of ear and groin corneocytes (table 23, figure 34)

In the dog the degree of adherence by both *S. intermedius* and *S. hominis* to corneocytes from the ear and groin were comparable (p = 0.407). The adherence counts for ear and groin were highly correlated (R = 0.976).



			Mean	± SD	Passage
	Disc	Ea	1	Groin	
Dog 1	1	286.50±	65.83	305.80±	58.50
S. intermedius	2	282.90±	55.02	291.80±	30.10
Right side	3	254.90±	21.87	279.60±	50.90
	4	272.60±	45.70	285.20±	22.86
	5	291.90±	23.61	297.30±	65.90
and the second second	Control	20.70±	6.33	38.90±	12.93
Dog 2	1	238.50±	43.71	311.10±	64.15
S. intermedius	2	242.90±	33.68	335.60±	56.73
Right side	3	*304.90±	38.00	301.10±	55.05
	4	272.20±	36.06	312.90±	27.81
	5	303.30±	78.48	329.30±	59.26
All the second	Control	48.10±	11.59	72.90±	13.34
Desta de la compañía	-	004.001	05 07	045.001	15.05
Dog 3	1	381.90±	85.07	345.60±	45.25
S. Intermedius	2	382.00±	71.95	344.90±	25.19
Right side	3	359.80±	10.45	338.30±	20.19
	4	304.90±	82.19	344.70±	32.75
	Control	370.70±	41.77	342.10±	29.10
- Continue	Control	20.00±	22.53	39.00±	20.13
Dog 3	1	60.80±	22.52	57.90±	15.95
S. hominis	2	53.10±	16.49	54.90±	24.91
Left side	3	62.90±	26.54	47.00±	15.16
and the second s	4	55.00±	15.71	55.20±	28.69
and the same of the same of the	5	58.50±	19.30	55.50±	13.28
	Control	25.10±	18.27	25.60±	16.68
Contraction of the local data					
Dog 4	1	72.00±	22.72	67.50±	22.53
S. hominis	2	79.10±	26.00	70.20±	32.48
Right side	3	79.20±	37.11	56.50±	24.98
	4	62.70±	21.95	69.00±	33.47
Constant of the second	5	67.50±	17.03	69.60±	17.89
REPART	Control	25.10±	10.81	27.90±	17.19

Table 23. Comparison of ear and groin corneocytes

*Dog2. Statistically significantly larger than 1 and 4 (p< 0.05) (one way ANOVA)

Paired t test ear v groin p=0.407 (not significant)

Comparison of bacterial adhesion to consecutive sampling from ear (table 24)

There were no statistical differences (ANOVA) between the adherence by *S. intermedius* or *S. hominis* on consecutive samples from the same individual animal. That is there was no within animal differences in the adhesion from each of five consecutive samples.

Table 24. Comparison of bacterial adhesion to consecutive sampling from ear

		and the state of	A DATE AND A	
		Mean	± SD	
	Disc		ar	
Cat 1	1	148.70±	33.26	
S. intermedius	2	145.50±	43.62	
Right ear	3	140.30±	68.33	
	4	144.30±	38.21	
A State of the	5	146.50±	50.45	
	Control	34.20±	17.73	
Cat 1	1	56.50±	15.06	-
S. hominis	2	57.60±	20.06	
Left ear	3	50.70±	18.07	
and the second second	4	52.50±	18.40	
	5	52.00±	21.31	
	Control	18.00±	8.00	
Cat 2	1	184.20±	25.63	
S. intermedius	2	181.90±	31.14	
Right ear	3	183.70±	36.89	
Sector Sector	4	175.30±	30.88	
	5	187.80±	35.49	
	Control	23.10±	13.52	
Cat 2	1	48.00±	18.12	_
S. hominis	2	46.60±	19.71	
Left ear	3	46.30±	24.17	
	4	53.70±	17.19	
	5	49.20±	27.23	
	Control	23.00±	14.08	

Adhesion by different bacteria species to canine and feline corneocytes (table 25, figure 35)

S. intermedius, *S. hyicus* and *S. aureus* all adhered well to both canine and feline corneocytes but in general was higher to canine corneocytes. *S. hominis* and a *Microcuccus* sp. Adhered poorly to both canine and feline corneocytes.

Table 25. Adhesion by different bacteria species to canine and feline corneocytes

	S. aureus	S. hvicus	Micrococcus	S. hominis	S. intermedius	Control
Cat 1	101.10±	184.60±	15.90±	62.50±	107.00±	23.50±
	31.64	12.84	9.69	30.23	36.20	11.82
Cat 2	95.70±	116.70±	20.70±	32.20±	58.40±	24.60±
	38.86	27.18	8.39	14.99	30.21	15.26
Cat 3	27.00±	108.00±	14.00±	28.70±	76.70±	19.10±
	15.06	15.63	10.62	15.85	47.78	8.31
Cat 4	26.60±	130.70±	8.70±	7.50±	47.60±	20.40±
	9.88	26.84	7.12	7.63	15.89	10.27
Cat 5	162.50±	285.80±	73.40±	76.90±	115.20±	16.90±
	73.42	52.32	27.06	49.68	51.36	10.06
Mean	82.58	165.16	26.54	41.56	80.98	20.90
Dog 1	107.30 ±	251.60 ±	22.00 ±	35.60±	122.60±	21.80±
	57.02	34.62	21.57	28.07	45.24	12.40
Dog 2	116.00 ±	227.20±	28.80 ±	40.10±	113.80±	29.70±
	70.33	39.80	14.34	24.51	32.27	12.75
ST ALL MARKED	176.60±	257.00±	25.80±	10.90±	126.50±	19.20±
Dog 3	38.31	51.29	18.67	8.20	94.26	9.60
Dog 4	101.00±	198.00±	31.70±	32.70±	245.40±	25.10±
	48.26	79.09	17.78	16.74	45.24	13.75
Dog 5	35.30±	166.50±	5.70±	16.80±	105.80±	22.20±
The second	25.21	81.36	4.30	13.34	63.30	12.59
Mean	107.24	220.06	22.80	27.22	142.82	23.6



Figure 35. Histogram of adherence by different bacteria species to canine and feline corneocytes

Figure 36. Adherence shown by different staphylococcal species

S. intermedius dog corneocytes

S. hominis dog corneocytes

Ł

- S. hyicus dog corneocytes
- S. hyicus cat corneocytes







S. intermedius cat corneocytes







DISCUSSION

The findings from this study indicate that the assay developed is a reliable and relatively quick method for the assessment of adhesion by bacteria cocci to both canine and feline corneocytes. The assay was able to differentiate between a known high adhering bacteria strain of *S. intermedius* and known low adhering bacteria stain of *S. hominis*. This is the first study of bacterial adherence to feline corneocytes.

The study on effects of temperature produced results similar to those of previous studies (Forsythe et al, 2002; Saijonmaa-Koulumies and Lloyd, 2002) in that bacterial adherence was higher at a higher temperature. The adherence shown by S. intermedius was also directly correlated with time, which too is similar to previous studies. The adherence shown by S. hominis did not appear to differ greatly over the time interval studied. This may suggest that S. hominis does not actively adhere to canine corneocytes and that the adherence shown was passive in nature. The findings of the temperature study in which the degree of adherence was higher at the higher temperature for both does not, however, support a passive mode of adhesion. A larger study of S. hominis over a longer period of time would provide information that may help to resolve this contradiction. The concentration study provided valuable information. As expected the higher the concentration of bacteria used the greater was the adherence but problems were encountered with clumping of bacteria at the higher concentrations. In general S. intermedius was very likely to clump at concentrations equivalent to an OD_{570nm} of 0.3 to 0.4 or above. Clumping of both S. intermedius and S. hominis was rare under an OD_{570nm} of 0.2. Clumping of bacteria made counting difficult or impossible when there was severe clumping. Bacterial clumping has been a problem recognised in other studies (Saijonmaa-Koulumies and Lloyd, 2002). It has been noted that staphylococci tend to

计算机输入器 化胆油 医甲基甲基基 法常期 计学校 化化物学 化化物学 计分析

and the second second

form clumps in both a strain-dependent and growth-dependent manor (Van Wamel, Vandenbroucke-Grauls, Verhoef and Fluit, 1998; Van Belkum, Kools-Sijmons and Verbrugh, 2002). In the assay used by (Saijonmaa-Koulumies and Lloyd, 2002) visual enumeration of bacteria was used to assess adhesion. Clumping was encountered in this study but because the authors considered that it was not possible to count individual bacteria in a clump each clump was counted as a single unit. The problem of clumping was overcome by Forsythe *et al* (2002) by using image analysis to calculate the surface area of adherent bacteria rather than counting individual bacteria. It would appear, however, that it would be advisable to conduct preliminary studies to identify potential problems with bacterial clumping for any adherence assay being developed. An incubation time of 45 minutes at 38°C and a bacterial concentration equivalent to 0.15 at OD_{570nm} was chosen as being both suitable and practical for the adhesion assay developed.

There was a high degree of correlation between manual (direct visual) counting and counting by image analysis. This was shown for both observers. This does not, however, imply that the results of the two methods gave identical results. Bland-Altman plots are useful to compare two methods of measurement (Bland and Altman, 1986; Bland and Altman, 1999). The Bland-Altman plots obtained in this study indicate that both observers tended to produce higher counts than the results obtained by image analysis. There was a tendency for this discrepancy to become wider as the number of adherent bacteria increased. Both observers were able to recognise small bacterial clumps of 2-4 cocci and count these as individual bacteria while the image analysis program would either exclude these or count them as a single count. The two observers did not give identical results with again a tendency for greater discrepancy when counting large numbers of adherent bacteria. Observer 1 tended to count consistently lower than observer 2 which may suggest that there might be

106

- 第二 - ころう デーシー・モーチー つ

- いたいかいない ひゃんかか たいかいかい 男性的 あかくたい シー・ション・シーンシント

يفيقطكمقا عاديره

the estimated and direction of the Press

South States

some subjectivity in assessing the numbers of adherent bacteria. More constant results may be obtained between observers with practice.

Since the images collected for analysis were individually selected by the operator there was the opportunity for variation here. The study comparing repeat analysis of the image selection and counting technique using image analysis showed that the method was repeatable.

Each control disc was visually scanned for the presence of bacteria and in all discs so examined no bacteria were seen. Control discs when counted using image analysis tended to give a count of approximately 20-30. After processing for image analysis and thresholding portions of the image that represented folds and edges of the corneocyte were visible and some of these were counted by the software program. Corneocytes and adherent bacteria were examined under a x 63 dry objective and the discs were not mounted. Mounting the discs in various mountants would largely abolish visible corneocyte edges and folds but had two disadvantages over the images obtained from unmounted discs. Firstly all mountants used resulted in the adhesive on the D-Squame® discs dissolving so that corneocytes and adherent bacteria tended to float loose in the preparation. Secondly when a mountant was used the image was cleared of corneocyte folds and edges but this made it impossible to identify clearly individual corneocytes. Using no mountant ensured that the distribution of bacteria adherent to corneocytes could be assessed.

Forsythe *et al* (2002) did find significant differences in the adhesion by *S. intermedius* to corneocytes collected from different body sites but this was not correlated to predilections sites for canine pyoderma. In this study both *S intermedius* and *S. hominis* appeared to adhere equally as well or poorly to corneocytes from the ear and groin of

 $\Delta \sqrt{2} M_{\rm eff} = 1000 M_{\rm eff} + 10$

1

a-description in the

dogs. Superficial pyoderma is rarely encountered on the inner aspect of the canine ear but is common in the inner thigh and groin region. This study suggests that in the dogs studied adherence to the ear corneocyte samples reflects adherence to groin corneocytes. Apart from a single instance there were no statistical differences between the numbers of adherent bacteria from each of the five discs collected from a single site of either the ear (cat and dog) or the groin (dog). This suggests that up to five discs from the one site may be used in comparative studies into potential factors that may affect adhesion. In cats comparison between the ear and groin was not conducted, as it was difficult to obtain suitable corneocyte samples from haired body sites such as the groin.

The final study compared the adhesion by five different staphylococcal species and a *Micrococcus* species, Pathogenic staphylococci (S. intermedius, S. aureus and S. hyicus) adhered well to both canine and feline corneocytes. The non-pathogenic S, hominis and the Micrococcus species tested did not adhere well. The strain of S. intermedius and S. hominis had been shown to adhere well and poorly respectively to canine corneocytes in a previous study (McEwan, 2002a). The adhesion assay used by McEwan (2002c) employed incubating corneocytes and bacteria in a suspension of PBS. Both studies using the same strains of S. intermedius and S. hominis but different methods produced similar results suggesting that both assays accurately assessed adhesion. Saijonmaa-Koulumies and Lloyd (2002) found two strains of S. aureus to adhere poorly to canine corneocytes while Forsythe et al (2002) using a single strain of S. aureus also found poor adherence to canine corneocytes. The S. aureus strains used by both Forsythe et al (2002) and Saijonmaa-Koulumies and Lloyd (2002) were of human origin while the strain used in the study described here was of bovine origin. Strain variation for adherence is recognised for S. aureus (Wyatt, Poston and Noble, 1990; Peacock, Day, Thomas, Berendt and Foster, 2000). Strain

108

services and the service of the property of the service of the ser

 C. C. Serie and Y. Maridia, A. M. Michell, Marian Mathematical Sciences and American Sciences an American Sciences and American S American Sciences and American Sciences American Sciences and American Scienc

ŝ.

variation may be one explanation for the different findings from different studies. Other factors may be the adherence assay methods used including the concentrations of bacteria, growth stage of the bacteria and possibly the media on which test bacteria were grown. The high adherence shown by *S. hyicus* to both dog and cat corneocytes is interesting. Since *S. hyicus* is not commonly recognised as a pathogen in the dog or cat this finding may suggest that the ability to adhere does not play a role in the development of infection or that it is but one factor in many that contribute to infection. Again larger studies using several strains are needed to clarify this situation.

In its present form the assay and counting procedure is a simple and reliable method for the assessment adhesion by cocci to canine and feline corneocytes. The inner aspect of the ear is a convenient sampling site that is tolerated well by both dogs and cats. Multiple samples from the same site can be obtained and used in studies to compare variables. Image analysis proved a useful and time saving method for bacteria counting. Further improvements in counting may be achievable by the use of more sophisticated software and through additional automation of the image manipulation and processing.

SUMMARY

The main findings of the various studies in this thesis are as follows:

- The three year retrospective survey of skin disease at the UGVS showed 62% of canine patients suffered from pyoderma, which was associated with atopic dermatitis, however, pyoderma was rarely diagnosed in the cat.
- The corneocyte morphometric studies appear to be the first study of its kind in the dog and cat.
- Two morphological types of corneocytes were identified. One a deeply staining elongated cell than probably arises from the hair follicle and a second mostly hexagonal cell that stained poorly.
- The mean canine and feline corneocyte diameter and surface area were found to be 38-48 μm and 1100-1800 $\mu m^2.$
- According to studies, corneocytes collected by D-squame® discs from pinna area of dogs and cats proved to be suitable for the collection of corneocytes for use in bacterial adhesion assays.
- Image analysis simplified more traditional methods for the counting of bacteria adherent to cells.
- S. intermedius, S. hyicus and S. aureus all adhered well to both canine and feline corneocytes while S. hominis and a Micrococcus sp. adhered poorly.

The studies conducted here provide a good starting point for several future studies. Larger studies could be usefully conducted to determine if there are truly differences in corneocytes from different body areas and animal breeds. The adherence assay opens the way to process large numbers of samples. Of particular interest here are identifying differences in strains of bacteria and methods for the blocking of adhesion by *S. intermedius* to canine skin.

Appendix A. Dog breeds examined for investigation of skin disease at the UGVS 2000-2002

Breed	Number	Breed	Number
Airedale terrier	2	Leonberger	1
Akita	4	Lhasa Apso	1
Alaskan Malamute	1	Lurcher	1
Basset hound	2	Maltese terrier	3
Beagle	4	Mastiff Neopolitan	1
Bearded collie	2	Miniature Poodle	1
Bernese Mountain Dog	1	Miniature Schnauzer	4
Bichon Frise	3	Munsterlander	1
Border Collie	9	Newfoundland	1
Border terrier	3	Old English sheepdog	1
Boxer	17	Papillon	1
Bull terrier	4	Pekinese	2
Bulldog	4	Pomeranian	2
Bullmastiff	2	Rhodesian ridgeback	2
Cairn terrier	1	Rottweiler	6
Cavalier King Charles spaniel	6	Rough collie	4
Chow Chow	2	Saint Bernard	1
Clumber spaniel	1	Saint Bernard	2
Cocker spaniel	11	Samoyed	1
Collie	1	Schnauzer	1
Dachshund (Short Hair)	1	Scottish terrier	1
Dalmatian	1	Shar Pei	5
Deerhound	1	Shetland sheepdog	5
Dobermann	6	Shih Tzu	2
Elkhound	1	Spinone	1
English setter	5	Spitz	1
English Springer spaniel	5	Staffordshire bull terrier	8
Field spaniel	2	Tibetan spaniel	1
Flat coated retriever	2	Tibetan terrier	3
French bulldog	1	Weimeraner	3
German Shepherd	34	White Highland White terrier	22
German Short Haired pointer	2	Wire Haired Fox terrier	1
German Wired Haired pointer	1	Yorkshire terrier	6
Golden retriever	9	х	15
Greyhound	3	Collie X	11
Husky	1	German Shepherd X	2
Irish setter	1	Jack Russell terrier X	1
Jack Russell terrier	4	Labrador X	5
Labrador retriever	34	Rottweiler X	1
Lakeland terrier	2	Terrier X	3

ų

÷

يىيى بىرى يەرىكى بىرىيە بىرىغان يەرىكى بەر يەرىكى بىرىيە بىرىكى بىرىكى بىرىكى بىرىكى بىرىكى بىرىكى بىرىكى بىرىك بىرىكى بىرىكى

:

and the second second

ŧ.

Appendix B. Main diagnosis of skin disease seen in dogs referred to the UGVS 2000-2002

Case	Breed	Sex	Age in years	Main diagnosis
141811	Border collie	Μ	5.00	Acral lick dermatitis
141894	German Shepherd dog	М	4.42	Acral lick dermatitis
141068	Labrador retriever	FN	2.00	Acral lick dermatitis
140514	Labrador retriever	М	6.58	Acral lick dermatitis
123121	Saint Bernard	М	2.50	Acral lick dermatitis
142082	Beared collie	Μ	7.42	Allergic dermatosis
144692	Bichon Frise	MN	10.00	Allergic dermatosis
145727	Boxer	Μ	0.67	Allergic dermatosis
142695	Boxer	F	2.50	Allergic dermatosis
143376	Cavalier King Charles spaniel	Μ	2.00	Allergic dermatosis
140543	Cocker spaniel	FN	7.58	Allergic dermatosis
140680	Dobermann	M	2.00	Allergic dermatosis
141369	English setter	M	0.92	Allergic dermatosis
200352	German Shepherd dog	FN	1.67	Allergic dermatosis
142007	German Shepherd dog	Μ	5.00	Allergic dermatosis
144732	German Shepherd dog	Μ	5.00	Allergic dermatosis
139570	Golden retriever	M	8.00	Ailergic dermatosis
138744	Labrador retriever	FN	3.00	Allergic dermatosis
144010	Labrador retriever	М	5.67	Allergic dermatosis
144326	Labrador retriever	M	1.50	Allergic dermatosis
143415	Labrador retriever	Μ	8.00	Allergic dermatosis
139752	Labrador X	FN	5.00	Allergic dermatosis
139142	Schnauzer	F	4.00	Allergic dermatosis
145596	Shar pei	М	2.00	Allergic dermatosis
145663	Shar pei	MN	2.83	Allergic dermatosis
144161	White Highland White terrier	FN	1.83	Allergic dermatosis
141708	White Highland White terrier	M	1.00	Allergic dermatosis
140684	X	MN	6.00	Allergic dermatosis
200863	X	M	4.00	Allergic dermatosis
139000	Alaskan Malamute	М	3.00	Alopecia X
141097	Pomeranian	М	2.92	Alopecia X
139783	Pomeranian	MN	6.00	Alopecia X
145646	Lhasa Apso	MN	6.25	Arnold Chiari syndrome
142531	Airedale terrier	FN	3.17	Atopic dermatitis
145469	Airedale terrier	MN	2.42	Atopic dermatitis
144227	Beagle	FΝ	8.00	Atopic dermatitis
140972	Bernese Mountain Dog	М	4.50	Atopic dermatitis
200253	Bichon Frise	М	1.00	Atopic dermatitis
139314	Bichon Frise	FN	4.00	Atopic dermatitis
143973	Border collie	M	0.67	Atopic dermatitis
139424	Border collie	М	2.00	Atopic dermatitis

化过去式 化过去式 化化合金属 化合金属 化合金 化合金 化合金 化合金合金 化合金合金 化合金合金 化合金合金 化合金合金 化合金合金 化合金

And the second second

al and the second

i.

÷

a bill a mattach an an A an a sao

ŝ

Case Breed	Sex	Age in years	Main diagnosis
200350 Border collie	F	5.00	Atopic dermatitis
139841 Border terrier	FN	2.83	Atopic dermatitis
139674 Border terrier	MN	6.00	Atopic dermatitis
142207 Boxer	М	1.00	Atopic dermatitis
140336 Boxer	Μ	1.08	Atopic dermatitis
140790 Boxer	FN	1.42	Atopic dermatitis
139574 Boxer	М	1.67	Atopic dermatitis
141011 Boxer	Μ	2.00	Atopic dermatitis
138946 Boxer	М	2.00	Atopic dermatitis
200326 Boxer	MN	3.17	Atopic dermatitis
141004 Boxer	F	3.25	Atopic dermatitis
139987 Boxer	FΝ	3.58	Atopic dermatitis
144566 Boxer	Μ	4.50	Atopic dermatitis
141624 Boxer	FΝ	5.00	Atopic dermatitis
140879 Boxer	Μ	0.67	Atopic dermatitis
140122 Buil terrier	Μ	2.92	Atopic dermatitis
141634 Bull terrier	Μ	5.00	Atopic dermatitis
139837 Bull terrier	FN	6.17	Atopic dermatitis
144127 Bull terrier	М	7.58	Atopic dermatitis
145819 Bulldog	۴	1.00	Atopic dermatitis
140353 Bulldog	۴	1.00	Atopic dermatitis
143816 Bulldog	Μ	2.50	Atopic dermatitis
144520 Cairn terrier	MN	7.00	Atopic dermatitis
144282 Cavalier King Charles spaniel	Μ	3.00	Atopic dermatitis
145695 Cavalier King Charles spaniel	Μ	5.00	Atopic dermatitis
143746 Chow Chow	Μ	5.67	Atopic dermatitis
200250 Clumber spaniel	F	6.33	Atopic dermatitis
142191 Cocker spaniel	MN	3.00	Atopic dermatitis
143910 Cocker spaniel	F	3.00	Atopic dermatitis
140194 Cocker spaniel	М	3.00	Atopic dermatitis
141042 Cocker spaniel	F	4.00	Atopic dermatitis
143431 Cocker spaniel	RN	4.58	Atopic dermatitis
140073 Cocker spaniel	M	7.58	Atopic dermatitis
141977 collie X	ΕN	0.58	Atopic dermatitis
144909 collie X	Μ	4.08	Atopic dermatitis
139678 collie X	М	7.00	Atopic dermatitis
200473 collie X	FN	9.58	Atopic dermatitis
139006 Dobermann	М	1,50	Atopic dermatitis
140799 Dobermann	ËΝ	2.00	Atopic dermatitis
145169 Dobermann	MN	2.00	Atopic dermatitis
141769 English setter	M	1.58	Atopic dermatitis
140278 English setter	MN	2.50	Atopic dermatitis
144338 English setter	FN	4.00	Atopic dermatitis
138856 English setter	FN	4.00	Atopic dermatitis

Ì

Middle of a court of a court of a second second of a state of the state of the court of a court of the second of t

Appendices

Case Breed	Sex	Age in years	s Main diagnosis
	1-1 R.4	0.00	Alexia dama ditia
141369 English Springer span		2.00	Atopic dermatitie
138793 Field spanlel		7.00	Atopic dermatitie
141751 Mat coaled reulever	EVIIN	4.20	Atopic dermanus
140032 Flat coated retriever	IVI 1.4	7.00	Atopic dermatitie
145831 French Dulldog	- N	1.00	Atopic dermatitie
143167 German Shepherd do	g ivi ∾ ENI	1.75	Atopic dermatitie
141542 German Snepherd do	g FN - M	2.00	Atopic dermatitis
139872 German Shepherd dog	g IVI ~ M	2.00	Atopic dermatitis
145094 German Shepherd do	g w	2.50	Atopic dermatitie
139658 German Snepherd do	g ₁vi 	3.4Z	Atopic dermatus
142089 German Snepherd do	g FN	4.00	Atopic dermatitis
141268 German Shepherd do	g MN	4.42	Atopic dermatitis
143991 German Shepherd do	g M	5.67	Atopic dermatitis
140929 German Shepherd do	g FN	6.00	Atopic dermatitis
145613 German Shepherd do	g FN	6.00	Atopic dermatitis
140862 German Shepherd do	g FN	7.00	Atopic dermatitis
200528 German Shepherd do	g Mi	7.75	Atopic dermatitis
142744 German Shepherd do	g Mi	7.92	Atopic dermatitis
125279 German Shepherd do	g MN	8.00	Atopic dermatitis
143142 German Shepherd do	g FN	8.00	Atopic dermatitis
145117 German Shepherd do	g	8.00	Atopic dermatitis
143215 German Shepherd do	g M	0.75	Atopic dermatitis
200155 German Short Haired	Pointer M	1.00	Atopic dermatitis
143842 German Wired Haired	Pointer FN	2.75	Atopic dermatitis
139920 Golden retriever	M	2.00	Atopic dermatitis
143495 Golden retriever	MN	2.50	Atopic dermatitis
141924 Golden retriever	MN	2.75	Atopic dermatitis
138873 Jack Russell terrier	FN	4.00	Atopic dermatitis
144703 Jack Russell terrier	MN	4.92	Atopic dermatitis
141664 Jack Russell terrier X	MN	8.00	Atopic dermatitis
140711 Jack Russell terrier	MN	4.00	Atopic dermatitis
145450 Jack Russell terrier	М	6.00	Atopic dermatitis
139563 Labrador retriever	FN	2.25	Atopic dermatitis
144588 Labrador retriever	MN	4.25	Atopic dermatitis
145659 Labrador retriever	FN	5.17	Atopic dermatitis
141156 Labrador retriever	M	6.50	Atopic dermatitis
143287 Labrador retriever	M	7.83	Atopic dermatitis
145886 Labrador retriever	M	9.00	Atopic dermatitis
145740 Labrador retriever	M	5.00	Atopic dermatitis
144088 Labrador refriever	M	2.00	Atopic dermatitis
1/1728 Labrador retriever	M	4.00	Atopic dermatitis
138204 Labrador ratiovor	кл Кл	6 00	Atopic dermatitis
143218 Labrador rotriaver	1VI 7.4	5.50 5.58	Atopic dermatitie
142470 Labrador rotriover		1 50	Atonic dermatitis
140470 Lapiador reulever	FIN	1.00	Auple demidulis

114

άŝ

and the set of the set of the formation of the set of the set

1

· · · · ·

с. К

والمستقدين والمسترجين والمسترجين

and and a state of the state o

Case Breed Se	ex Age in years	Main diagnosis
143679 Labrador retriever M	N 1.00	Atopic dermatitis
141752 Labrador retriever F	N 2.00	Atopic dermatitis
140546 Labrador retriever N	1 2.00	Atopic dermatitis
142096 Labrador retriever F	N 4.00	Atopic dermatitis
140761 Labrador retriever M	N 4.00	Atopic dermatitis
141749 Labrador X	/ 7.00	Atopic dermatitis
200362 Lurcher F	N 4.00	Atopic dermatitis
142020 Maltese terrier M	N 3.00	Atopic dermatitls
143861 Maltese terrier	8.00	Atopic dermatitis
142900 Miniature Schnauzer F	N 2.17	Atopic dermatitis
144537 Miniature Schnauzer M	N 4.00	Atopic dermatitis
126905 Munsterlander	1 2.00	Atopic dermatitis
138788 Labrador retriever	A 1.83	Atopic dermatitis
140894 Labrador retriever	= 2.67	Atopic dermatitis
138912 Labrador retriever	A 4.00	Atopic dermatitis
139978 Labrador retriever	- 5.92	Atopic dermatitis
140503 Labrador X	1 5.00	Atopic dermatitis
140957 Rhodesian Ridgeback	= 1.17	Atopic dermatitis
142529 Rhodesian Ridgeback F	N 2.00	Atopic dermatitis
141731 Rottweiler	- 2.00	Atopic dermatitis
127740 Rottweiler	A 1.42	Atopic dermatitis
127740 Rottweiler F	N 7.00	Atopic dermatitis
143417 Rottweiler X M	N 2.00	Atopic dermatitis
145609 Rough collie M	N 5.50	Atopic dermatitis
142049 Rough collie	A 5.50	Atopic dermatitis
143465 Saint Bernard	- 2,75	Atopic dermatitis
140842 Shar Pei	= 0.58	Atopic dermatitis
139601 Shar pei	= 1.50	Atopic dermatitis
144763 Shetland Sheepdog F	N 3.17	Atopic dermatitis
144416 Shetland sheepdog	A 4.92	Atopic dermatitis
143661 Shetland Sheepdog	N 7.00	Atopic dermatitis
145259 Shih Tzu F	N 4.00	Atopic dermatitis
142759 Saint Bernard M	N 2.00	Atopic dermatitis
145494 Staffordshire Bull terrier	и 0.92	Atopic dermatitis
145417 Staffordshire Bull terrier	F 1.83	Atopic dermatitis
141416 Staffordshire Bull terrler	F 1.92	Atopic dermatitis
138733 Staffordshire Bull terrier F	N 2.00	Atopic dermatitis
138733 Staffordshire Bull terrier	M 3.50	Atopic dermatitis
141864 Staffordshire Bull terrier	4.50 V	Atopic dermatitis
144071 Terrier N	IN 4.00	Atopic dermatitis
200303 Terrier X	N 1.25	Atopic dermatitis
141235 Terrier X	vi 3.42	Atopic dermatitis
139598 Tibetan terrier F	N 2.00	Atopic dermatitis
142090 White Highland White terrier	vi 0.67	Atopic dermatitis

tertuken in dia serie turun turun turun turun antara serie etter antara serie etter

ļ

المفاريح أمطم والمراجعة المؤسسة فرجار والمتعاركين فالمراجع والمراجع فالمناطق والمعادية والمعارية والمعا

and a straight of the second

and and a

Case Breed	Sex /	Age in years	Main diagnosis
143942 White Highland White terrier	F	0.67	Atopic dermatitis
144284 White Highland White terrier	MN	1.00	Atopic dermatitis
141029 White Highland White terrier	FN	1.75	Atopic dermatitis
141134 White Highland White terrier	MN	1.83	Atopic dermatitis
142297 White Highland White terrier	MN	2.17	Atopic dermatitis
144638 White Highland White terrier	Μ	4.50	Atopic dermatitis
141825 White Highland White terrier	FN	5.00	Atopic dermatitis
200866 White Highland White terrier	MN	5.00	Atopic dermatitis
143606 White Highland White terrier	FN	8.08	Atopic dermatitis
128055 X	M	0.75	Atopic dermatitis
144407 X	MN	1.00	Atopic dermatitis
141908 X	MN	1.58	Atopic dermatitis
139921 X	FN	2.00	Atopic dermatitis
200313 X	FN	2.50	Atopic dermatitis
141414 X	F	4.00	Atopic dermatitis
139750 X	FN	4.00	Atopic dermatitis
140344 X	FN	4,25	Atopic dermatitis
144302 Yorkshire terrier	М	0.83	Atopic dermatitis
145951 Yorkshire terrier	M	1.58	Atopic dermatitis
145487 Yorkshire terrier	FN	4.00	Atopic dermatitis
141015 Yorkshire terrier	FN	4.00	Atopic dermatitis
144981 Yorkshire terrier	FN	2,80	Atopic dermatitis
140347 Basset hound	М	1.00	Atopic dermatitis
145367 Beagle	FN	1.00	Atopic dermatitis
145541 Beagle	FN	7.08	Atopic dermatitis
145869 Bullmastiff	М	1.00	Atopic dermatitis
142404 Bullmastiff	М	1.67	Atopic dermatitis
145477 collie	FN	1.00	Atopic dermatitis
140372 collie X	MN	3.83	Atopic dermatitis
140243 German Shepherd dog	MN	2.00	Atopic dermatitis
144341 Golden Retreiver	М	3.00	Atopic dermatitis
140033 Labrador retriever	М	1.33	Atopic dermatitis
141455 Labrador retriever	М	1.00	Atopic dermatitis
142293 Maltese terrier	MN	3.25	Atopic dermatitis
141132 Mastiff Neopolitan	F	4.00	Atopic dermatitis
139227 Tibetan terrier	FN	2.00	Atopic dermatitis
143938 Tibetan terrier	М	2.00	Atopic dermatitis
141439 White Highland White terrier	FN	1.00	Atopic dermatitis
141547 White Highland White terrier	F	1.00	Atopic dermatitis
139300 White Highland White terrier	MN	3.00	Atopic dermatitis
200203 White Highland White terrier	F	3.00	Atopic dermatitis
145545 White Highland White terrier	MN	4.00	Atopic dermatitis
142794 White Highland White terrier	F	6.00	Atopic dermatitis
140921 collie X	FN	6.00	Atopic dermatitis / FAD

** - **

-ţ

1000

وسيه كالمناط المناز أركركم

Appendices

Case Breed	Sex /	Age in years	Main diagnosis
	-	0.07	Ad
141326 White Highland White terrier	-1	0.67	Atopic dermatitis / demodectic
140389 German Shepherd dog	FN	2.00	Atopic dermatitis / FAD
142862 German Shepherd dog	M	5.00	Atopic dermatitis / FAD
200251 Labrador retriever	FN	4.00	Atopic dermatitis / FAD
200332 X	FN	9.00	Atopic dermatitis / FAD
142693 Yorkshire terrier	F	1.00	Atopic dermatitis / FAD
140542 Labrador retriever	F	1.00	Atopic dermatitis / food sensitivity
143067 White Highland White terrier	MN	3.00	Atopic dermatitis / food sensitivity
137559 Greyhound	FN	2.00	Bald thigh syndrome
200410 Greyhound	FN	3.00	Baid thigh syndrome
141031 Rottweiler	М	6.58	Calcinosis cutis iatrogenic
144340 Elkhound	MN	6.92	Cheyletiellosis
127238 Cocker spaniel	FN	2.00	Cheyleteillosis / hypothyroidism
139066 Cocker spaniel	М	9.00	Cheyletiellosis
143412 Labrador retriever	М	3.33	Cheyletiellosis
144788 Spitz	MN	5.00	Cheyletieliosis
142766 Old English Sheepdog	FN	3.00	Cheyletiellosis / FAD
144684 German Short Haired Pointer	r F	0.42	CLE (cutaneous lupus erythematosis)
143311 German Shepherd dog	FN	7.00	CLE (cutaneous lupus erythematosis)/ DLE
145111 Shetland sheepdog	М	11.00	Cutaneous lymphoma
145535 terrier X	FN	14.00	Cutaneous lymphoma
143304 Shetland sheepdog	۴N	5.00	Cutaneous vesicular lupus erythematosus
139038 Rough collie	FN	1.08	Demodectic mange
142533 Staffordshire Bull terrier	F	0.50	Demodectic mange
140871 collie X	М	0.33	Demodectic mange adult
200595 German Shepherd dog X	FN	14.00	Demodectic mange adult
200123 Miniature Poodle	MN	9.00	Demodectic mange adult
144550 Newfoundland	М	4.25	Demodectic mange adult
140499 Pekinese	M	7.08	Demodectic mange adult
144153 Akita	Μ	0.67	Erythema multiforme / drug eruption
143866 Golden Retriever	FN	7.08	Exfoliative dermatosis / ichthyosis
145022 Border collie	М	0.83	FAD
139136 Border collie	FN	?	FAD
139340 Border collie	MN	1.25	FAD
142709 Cavalier King Charles spanie	el FN	3.00	FAD

117

selections of the second se

market of the second

and the second sec

and the second sec

Case	Breed	Sex	Age in years	Main diagnosis
144070 0		E	1.00	EAD
144972 0	Joine A Field enaniel		6.58	
1/2/28 (Serman Shanherd dog	Ē	4.00	
1/27/7	Serman Onephero dog Sekinese	- FN	10 92	FAD
138860 9	Semoved	M	7.08	FAD
140200 \$	Shih Tzu	FN	8.50	FAD
1410381	Tibetan spaniel	M	2.00	FAD
143724 \	White Highland White terrier	M	0.50	FAD
141865)	(FN	4.00	FAD
145766 (Serman Shepherd dog X	MN	6.00	FAD / anal furunculosis
144742	Boxer	MN	3.25	Flank alopecia
139616 F	Border collie	MN	7.25	Follicular dysplasia
145427 (German Shepherd dog	M	2.50	Food sensitivity
144901 F	Rough collie	M	1.42	Food sensitivity
141590 (Serman Shepherd dog	FN	1.67	Granulomatous dermatitis
140431 (Collie X	FN	9.92	Hyperadrenocorticism
140500 F	Papillon	M	13.00	Hypothyroidism
145046 L	_akeland terrier	М	7.83	latrogenic cushings
200138 \$	Scottish terrier	М	7.00	Interdigital pyoderma / sterile
				granuloma
144375 6	Beagle	FN	8.00	Interdigital pyoderma / sterile granuloma
143936 [Bulldog	FN	8.00	Interdigital pyoderma / sterile granuloma
143676 (Golden retriever	М	11.50	Keratinsation defect
143938 \	Weimeraner	М	1.00	Microchip related dermatosis
1 4 2184 I	Bearded collie	MN	4.00	Nail dystrophy - symmetrical lupoid onchodystrophy
145126 (Greyhound	М	8.00	Nail dystrophy - symmetrical lupoid onchodystrophy
140457 I	Boxer	FN	5.00	Nail dystrophy - symmetrical lupoid onchodystrophy
142816	_abrador retriever	MN	8.00	Nail dystrophy - symmetrical lupoid onchodystrophy
139525 (German Shepherd dog	FN	4.00	Nodular dermatofibrosis
1440821	Border terrier	MN	5.00	Otitis
144732 (German Shepherd dog	FN	6.92	Otitis
1397401	Lakeland terrier	M	7.75	Otitis
144459	Shar Pei	F	1.50	Otitis

Contract of the second

a survey of the second s

1

1. 18 miles

1.00 C. 1.00 C. 1.00 C.

مىيەت ھۆك ئۇرىيەت بەيدىغ مەلكىمىر مەك مىڭ ئىيە ئىيە تىيە بىيە بى

Appendices

Case Breed	Sex .	Age in years	Main diagnosis
	} <u>P</u> "ki	49.00	Offic (neoudamenes)
143602 Cavaller King Charles spanle		13.00	Ottus (pseudomonas)
142899 Dacrishund (Short Hair)		1.00	Pattern baluness
143703 Akita	FN	2.92	Pempnigus foliaceus
144307 Akita	MIN	2.00	Pempnigus foliaceus
145658 Akita	M	4.92	Pemphigus foliaceus
143066 Dalmatian	MN	2.42	Pemphigus foliaceus
142332 German Shepherd dog	MN	7.08	Pemphigus tollaceus
140456 Collie X	м	1.00	pemphigus vulgaris / bullous
200002 Basset hound	M	1.33	Primary malassezia dermatitis
139283 German Shepherd dog	М	0.67	Primary pyoderma
143587 Border collie	FN	3.00	Primary pyoderma
141885 Boxer	M	2.00	Primary pyoderma
143744 Chow Chow	M	1.00	Primary pyoderma
143290 Deerhound	F	3.08	Primary pyoderma
140764 Dobermann	F	1. 2 5	Primary pyoderma
141464 Dobermann	F	1.25	Primary pyoderma
140281 English Springer spaniel	FN	0.00	Primary pyoderma
142052 German Shepherd dog	F	1.00	Primary pyoderma
142283 German Shepherd dog	M	1.17	Primary pyoderma
145396 Golden retriever	MN	2.00	Primary pyoderma
142196 Irish setter	FN	3.00	Primary pyoderma
125469 Leonberger	M	4.00	Primary pyoderma
138726 Miniature Schnauzer	FN	4.42	Primary pyoderma
139125 Staffordshire Bull terrier	М	1.08	Primary pyoderma
145256 Weimeraner	M	2.00	Primary pyoderma
143372 White Highland White terrier	FN	8.42	Primary pyoderma
145172 X	MN	7.50	Primary pyoderma
145809 X	M	9.00	Primary pyoderma
200017 X	MN	8.00	Primary pyoderma
139371 Cocker spaniel	F	0.92	Primary seborrhoea
140675 Cocker spaniel	FN	11.00	Primary seborrhoea
144018 Labrador retriever	M	3.00	Pyoderma secondary to trauma
143078 English Springer spaniel	M	6.00	Pyotraumatic follicultitis
143536 Golden retriever	F	1.17	Pyotraumatic follicuititis
141765 Rottweiler	M	6.42	Pyotraumatic follicultitis
141065 Rottweiler	M	6.42	Pyotraumatic follicultitis
143590 Collie X	MN	2.50	Sarcoptic mange
143110 English Springer spaniel	М	6.92	Sarcoptic mange
145288 English Springer spaniel	М	13.42	Sarcoptic mange
139800 X	FN	2.00	Sarcoptic mange
139335 Wire haired fox terrier	FN	3.00	Sebaceous hyperplasia

ł

a da serie de series de la companya de la companya

Ì

يتيانك أحكم فاكرته تكرفتنا ويراو فالأفات

and the second second

Appendices

Case	Breed	Sex	Age in years	Main diagnosis
143109 Minia	ture Schnauzer	м	7.00	Schnauzer comedo syndrome
145885 Spine	one	F	7.00	Traumatic wound
145648 Wein	neraner	F	5.00	Urticaria
143560 Cava	lier King Charles spaniel	FN	12.00	Vulval fold dermatitis
140185 Husk	y -	Μ	1.25	Zinc responsive dermatosis

2

2

emotionates activities and the second states the activities and activities activities activities for a task of the states and

1.1.1

11111

Appendix C. Cats referred for investigation of skin disease at the UGVS 2000-2002

Case	Breed	Sex	Age in vears	Diagnosis
145224	Domestic short hair	FN	2.92	Allergic dermatosis
139715	Domestic short hair	FN	2.92	Allergic dermatosis
141263	Domestic short hair	FN	2.92	Allergic dermatosis
142194	Domestic short hair	MN	5.00	Allergic dermatosis
140991	Domestic short hair	MN	7.00	Allergic dermatosis
142894	Domestic short hair	FN	7.00	Allergic dermatosis
139688	Domestic short hair	EN	5.00	Atopic dermatitis
1399 9 2	Domestic short hair	FN	8.00	Atopic dermatitis
145619	Domestic short hair	FN	3.17	Atopic dermatitis
141705	Domestic short hair	FN	5.00	Atopic dermatitis
142380	Domestic short hair	FN	3.00	Atopic dermatitis
140788	Domestic short hair	MN	7.00	Atopic dermatitis
140897	Domestic short hair	NR	3.00	Atopic dermatitis
142430	Domestic short hair	NR	4.00	Atopic dermatitis
141296	Persian (long hair)	F	2.00	Atopic dermatitis
140911	Siamese	MN	7.00	Atopic dermatitis
143545	Domestic long hair	FN	4,92	Atopic dermatitis / FAD
144905	Domestic short hair	FN	4.50	Atopic dermatitis / FAD
140446	Domestic short hair	MN	13.00	Ceruminous gland adenoma
141200	Domestic short hair	MN	6.00	Chin oedema
144022	Domestic short hair	MN	0.75	Cutaneous asthenia
143114	Domestic short hair	FN	4.00	Eosinophilic plaque/eosinophilic ulcers - idiopathic
141558	Ragdoll	MN	2.75	Facial dermatitis - idiopathic
141422	Domestic short hair	۴	10.00	FAD
138942	Domestic short hair	FN	6.42	FAD
139354	Domestic short hair	FN	6.00	FAD
140214	Domestic short hair	FN	11.67	FAD
140907	Domestic short hair	MN	6.00	FAD
144168	Domestic short hair	MN	12.00	FAD
139700	Turkish Van	MN	1.25	FAD
142390	Domestic short hair	FN	1.58	Feline acne / folliculitits
143849	Burmese	MN	0.67	Food sensitivity
143675	Persian (Long Hair)	MN	2.58	Idiopathic facial dermatitis
145816	Persian (Long Hair)	MN	3.00	Idiopathic facial dermatitis
140787	Perslan (Long Hair)	MN	2.00	Idiopathic facial dermatitis
143084	Domestic short hair	MN	9.00	Pemphigus foliaceus
143357	Domestic short hair	MN	10.00	Pemphigus foliaceus
145009	Domestic short hair	MN	2.80	Plasma celì pododermatitis
200759	Siamese	MN	3.42	Psychogenic alopecia
143094	Domestic short hair	FN	0.58	Symmetrical alopecia – idiopathic

and the second se

an Marina da nan sa wasana ana ana ana ana an

100 100 100 100 100 100 100 100 100 100	and marked and the second s	200 2 000000000000000000000000000000000				Appen	dix D. No	rmal dog	is comeo	cyte diam	leter (Jum)	i.		10000		Construction of the	2	10000000000000000000000000000000000000
		2	j r	1		Total		2	3	- - -	5	(otal	i sen Listen	i N	Julio Sector	•	5	total
LBopy	37.94	40.05	42.75	40.00	37.93		38.35	39.63	33,69	46.53	35.60		38.54	35.98	46.49	43.50	38.10	
Labrador	34.37	46.26	188 17 17	38.05	38.90		40.12	39.35	42.01	43.32	45.05		47.75	39,86	41.73	40.50	42.21	
retnever 40 voor	41.08 20.70	51.05 20.05		28.01	14 S		44-US	44.95 20.05	119-115 119-115	40.60	50.02		97.92 1	24.01 26.02	20.62	43.12	90.40 26.00	
EN State	39.84	17 F	45.72	40.33	37.57		37.61	41.80	5 X 7 8	42.77	44 94		4743	20 80 20 80 20 80	40.42	46.46	78°C2	
:	45.04	40.71	46.72	37.52	41.40		41.24	43.38	38.02	36.11	38,66		41.99	33.41	41.37	42.56	35.93	
	40.21	33.11	31.42	39,89	42.58		41.96	37.85	36.51	41.20	38,38		37.79	36.20	45.29	43.61	41,55	
	41.13	40.23	44.74	33.31	40.72		44.61	42.31	41.33	32.64	41.78		40.98	46.05	48.62	39.70	42.47	
	41.18	34.49	38.36	34.78	21.88		42.44	41.84	45.39	38.46	45.01		36.41	40.24	38.64	45.67	42.01	
	39.61	32.05	36.56	36.67	45.37		39.17	40.25	39.47	45.23	41.12		34.66	73.27	40,40	43.20	41.11	
Mean	40.00 5 5 5	35.85	29.55 - 21	37.84	41.09	38,99	40.95	41.24	41.14	40.20	41.48	40.83	4 2 3 3	39.38	42,74	1 3.53	2676E	41,21
50 - 1	21.2	5.34	5.20	23	2.75	4 93 2 4 93	2.34		2.63	3.28	2.89	3.05	4.46	3.85	4,1 1,1	2.67	2.67	0.88 1
mođe	40,00	34.11	38.36	37.84	40.72	39.52	40.95	41.24	41,14	40.65	41.12	41.18	10.51	39.63	41.37	43.20	40.40	41.04
Rddg 2	45.64	40.52	32.10	23.99	40.82		43.67	49.B7	45.61	43.63	46.93		44.52	49.56	49.03	41.31	45.45	
Cocker	39.25	39.45	39.25	34.36	40.58		45.84	46.50	41.5í	44.67	46.67		42,99	46.57	42.33	39.14	46. 2 3	
spaniei	39,44	40.70	33.78	38.29	46.07		46.18	41.73	45,46	45.95	47.74		40.49	43.12	42.81	48.06	45.51	
6 years	38.66	35.83	34.39	38.18	43.34		43,17	4,25	4 .8	42.50	43,75		39.30	4,16	50.28	42.82	39.40	
FN	37.66	41-20	36.03	37.82	32.78		46,65	41.59	41,10	39.53	40.97		13.67	50.92	47.80	43,08	44.27	
	32.19	40,43	36.87	36.12 51 25	43.05		44,86 10	47,80 19	45.25	47.33	37.56		37,78	47.09	55.12	41.57	44.97	
	50.04 20.04 20.04	18.14	4		43.27		53.84 11 20	45.57		40.04	45.03		45.46	44.01	43.75	41.83 44.83	19 19 19 19	
	79.62	43.82			46,73			46.28	41.92	40.86	45.97		46.63	48.19	50.20	43.30	43.87	
	4 1 2 2 2	38.61 1 8.7	41.67	202	42.09		52.61	44 14 14 14	44.03	47.29	46.95			46.16	38.4 5	40.12	39.38	
Hone.		10.25	20.00	\$ 8 8	97.78	07 00	44°14		5. 2	5,5	75.64	20.22	2-14 2-14	48./3 5 4	45.43	61.4	45.F4	07 FT
MBAD.	3.5	07-65	20.01 2 5 0	10.00	42.30	38.76	10.04	43.81 1 8 9 6	43.20	43.78	2.44	22.4	1.14	46.43	46.24	57.73	57.73	04.44
mode	39.44	39.87	37,99	36.95	42.78	39.35	45.84		43.25	43.70 43.70	45.08	44.79	42.98	46.16	45.43	42.73	42.73	44,08
Ndog 3	34.17	39.50	35.99	40.08	40.85								50.04	46.0E	43.52	45.15	49,95	
German	39.61	36.06	42,38	42.15	39.51								47.27	49.87	48.50	48.84	49.47	
shephera	44.30 10.04	37.03	41,42 57,75	29.76	02.75								48.45 4 e c e	48.80	47.61	49.1î E1 12	42.87	
hon /	30.74	20.85	20.05	10.05	20.70								40,04	47.03 E0.11	02.64 C2.02	21.12	19.22	
t years	35.47	40.29	24 EE	5192 26-10	35.47								43.20	45.00	44 61	48.82	10.15 44 75	
	41.58	35.71	31.83	39.56	41.58								48.87	49.07	45.19	52.27	44 85	
	44.00	41.12	38,15	45.43	44.60								51.30	46.57	49.55	47.44	48.12	
	44,86	36.13	40.84	40.16	44.85								49.93	35.39	54.20	44.96	46.72	
	41,38	41.67	N. 85	42.65	41.38								52.50	46,91	48.09	50.35	1 4.85	
Mean	02'6E	38.41	79'6E	39.63	40.31	₩.8							48.41	45.68	48.59	48.63	45.84	47.53
su modo	6/~4	20 00 20 00	200	4.04 10.61	10.4	097.5							90 7 7 90	10. F	587	2,56 10 00	7.38	3.13 40 4 4
ADD	001044	Davac	7.00	20,00	0000								14:01	17:04	40.00	40.02		
, FIBOPN	30.96	40.86	39.64	41,85	42.70								45.23	44.97	41.75	46.31	44.81	
Labrador	37.36	35,59	40.45	36.59	46.25								44.96	40.53	42.75	45.54	45.94	
tehiever C	32.18	41.73	39.55 29.55	45.90	45.24								48.34	30.31	41.08	46.96	41.54	
e years ch	10.00 10.00	13.90 10.00	32.87 20.44	4 1,4 1,4 1,4 1,4 1,4 1,4 1,4 1,4 1,4 1,	36.80 37 EO								50.00 90.00	12.31	91.0E	D6189	21.12	
-	36.00	41.77	38.52		42.36								16.92	39.93	45.78	30.64	43.49	
	39.49	40.24	35.67	37.64	41.46								38.11	46.27	39.51	44.86	46.47	
	37.54	41.30	39.24	39.51	34.15								47,92	40.85	43.46	50.55	45.81	
	20	88	57 15 15 15	42.93	35.33								4,55 1,55	42,93	41.86	40.75	52.49	
	53.73 24 26		30.72	39.9D	32.83	20 47							42,54	39.48 14	43.69	40.35	42.71	64 GV
SD	2.92	3,05	2.64	2.84	447	3.55							4.30	2.19	42.50		08.44	346
mode	35.36	38.84	37.42	39.90	37.59	38.80							44.89	40.85	41,96	44.49	44.98	42.91

and the state of the second

անձատեն ներերը որ ենքեն է։ Դեսելու է երեն նաճենագերեններ առեններու ենիներում։ Արելու հերերությունը հետ նարեներե Առենությունը։ Դենեներում են են երեն եննենագերենները եններու ու հենքներում։ Արելու հետ հետ նարենքեներեն առենում

1

and the second se

4 5 70481 41,84 37.34	42.19 43.39	41.52 35.48	41.62 35.04	43.62 44.93	41,22 35,04	38.90 33.02	47.92 33.89	42.21 39.28	40,35 43,44	42.17 39.37 40.46	2.52 3.74 3.48	41.62 38.89 40.14																										
1 7 38.38	1 38.91	7 43.18	0 43.01	239.94	0 44.51	8 43.29	k0 42.10	8 33.65	37.89	6 40.72	2 3.47	40.72																										
40.6	4	62.2	36.7	37.6	39.0	412	37.8	39.4	41.6	41.1	4.8	39.4																										
37.16	37,32	36.57	38.33	37.86	38.21	39.06	44,0%	42,73	42.79	39.36	2.96	38.33																										
1014										40.82	5.43	41.37											42.78	3.59	42.79											33.57	3.06	33.41
42.64	38.98	38.81	34,55	49.05	44,99	49.42	40.87	44.91	40.68	42,46	4.98	40.87	43.30	43.61	35,83	46.23	42.77	44.91	45,12	41.00	49.29	42.23	43.44	3.76	43.44	48.57	50.41	51.12	47.11	46.25	48.10	42.65	40.22	43.37	43,19	32.89	2.70	32.34
41.52	42.84	56.52	42.65	37.27	38.27	46,12	42.66	37.84	39.75	42,68	5.96	42.66	42.22	43.45	39.72	37.E1	41.25	46.25	45.86	¢1.85	48.97	38.72	42.61	3.81	41,65	43.10	39.10	33.87	46,28	44.06	42.38	40.57	47,59	48.53	46.93	35.66	3,52	33.66
12.9E	44.67	41.26	41.56	35,99	34,11	41.85	39.01	39.60	44.3D	40.28	3.51	40.2B	38.43	48.33	41.13	43.67	45.03	30.65	47.23	42,50	45.37	44.96	44.12	3.00	44.12	42.13	47.24	39.68	46,26	47.19	42.11	44.72	38.72	38.83	43.30	32.64	2.55	32.82
2 41.40	39.75	35.56	46.51	39,26	49,16	34.13	42.14	39.16	44,09	41.08	4.91	39.75	44,43	39.51	42.11	35.52	36.17	67.7C	44.62	43.03	36.74	37.48	39.76	3.11	38.74	45.43	49.50	52,16	51,20	62.20	47.36	46.42	42.00	40.58	43.15	32.41	2,72	31.92
1 27.01	25.33	41.73	41.33	41.62	41,42	37,84	41.61	35.20	41.04	3772	6.31	41.04	48.00	46.31	42.58	43,59	41.28	47.83	42.81	49.13	40.63	39.02	44.12	3.49	42.81	±2.82	38.18	40.74	51.69	40.23	39.65	52.20	51.28	45.19	52.58	34.91	2.73	34.40
Total										33.57	3.05	33.41																										
31.4C	31.90	38.11	31.14	33.44	32.13	35.73	32.34	32,53	28,70	32.89	2.70	32.34																										
34.93	38.39	35.95	31.09	34.05	30,96	39.57	40.61	35.85	33.51	35,66	3.52	35.66																										
27.75	33.38	27.51	35.46	31,54	32.90	30.58	36.10	33.51	32.82	32,64	2.55	32.82																										
33.11	31,55	20°58	33.97	29.47	31.74	35.77	27.64	31.92	35.59	32.41	2.72	31.92																										
1 5.13 35.13	30.87	34,40	38.77	40.76	37.25	34.24	33.10	32.41	33.15	34.91	2.73	34.40																										
Ndog 5	Greyhound	3 years	æ							Mean	SD	mode	Ndog 6	Greyhound	4 years	R							Mean	cs S	араш	Noog 7	Greyhound	5 years	IN							Mean	SD	mode

an a se ban ban a se a ser a se a se la construction e ser a construction de la construction de la construction

;

с 1

14 alter and a stress of a second

											1312.17	1200 61											1673.13	208.67	1690.81											1468,44	1459.29										1433.56	212.74	67.UA46
	1217 1A	1532.01	1188.63	1400.02	1223.10	1208.56	1282.67	1286.66	1108.94	1166.51	1274.71	10.621	1677.56	1704.05	1506.01	1440.86	1453.58	1750.28	1624.36	1800.23	1266.73	1792.11	1602.62	176.60	1613.43		1465.17	1282.07	1531.52	14/67 OC	1783.15	1326.11	1252.99	1370.73	1428.11	1433.21	1426.22	L6 017 F	20.0121	1302.20	1657.28	1286.46	1409.73	1550.89	1818.41	RC 1101	1509.36	171.02	Vernotic
		1500 43	1118.50	1184,85	1306.96	1530.52	1327.90	9:7.87	1425.52	1322.52	1300.40 4 84 94	1314 75	1708.24	1911.46	1841.33	1863.24	1985.37	1844.71	1737.72	1745.49	1580.93	1724.18	1794.27	116.99	1769.88		1390.26	1249.20	1140.22	1480 88	1938.95	1324.31	1359.38	1277,69	1379.90	1430.90	1369.64	G <u>7</u> 797 t	400U C2	1997.92	1975.81	1105.75	1304.79	1460.59	1569.78	1331.00	1608.70	302.87 4 EEE EN	TO DOCE
Ĝnin	100 C10+	1279 68	1310.96	1268.33	1571,96	1207.96	1207,95	114.32	447.24	1400.22	1316.14 122.84	1704 17	1551.09	1538,69	1660.82	1881.17	1571.76	1788,73	1422.14	1875.40	1339.25	1636.91	1630.60	179.65	1010.84		1455.68	1242.63	1206.75	1464 01	1375.32	1623.76	1650,66	1447,84	1464.57	1498.72	1450.93	1070 06	02.0001	1316.74	1243.02	1276.70	1234,46	1231.67	1506.21	1400.00 1504.62	1299.91	136.60	00,6671
	1979 51	1671.98	1003.74	1391.45	1472.14	1034.63	1370.93	12E3.15	1183.25	1420.54	1308.43 202.53	176/202	1576.94	18C6.46	1616.79	2015.26	1613.40	1479.91	1368.14	1174.49	1912.45	1648,30	t641,21	Z59.83	1615.10		1583.52	1509.00	1535.90	1300 10	1650.46	1602.24	1653.05	1479.12	1539.88	1522.01	1528.95	0 - 0 - 7 -	1225.62	1393.65	1253.37	1367.75	1387.07	1035.42	1455.61	1201 78	1344.22	144.26	77/0101
	1100 83	1055 15	1463.38	1473.34	1280.48	1682.34	1/27.71	1523.35	1252.59	1292.24	1364.14	1128.19	1560.62	1727,56	1779.76	1643.88	1203.78	2099.33	2010.48	1909.67	1526,93	1658.43	1721.94	254.80	1691.28		1256.82	1452.02	1340.25 1563 PA	1670.37	1538.50	1520.96	1490.51	1684.33	1088.22	1457.35	1468.94	1240.07	1352.21	1270.32	1376.71	1249.60	1527.13	1325.12	1440.27	10.1021	1405,62	126.13	1000
um²)											1242.46 200.20	1273.11												1515.47	154.30	0																							
ce area (p	075 64	1243.22	1384.88	986.21	1519.96	1292.63	1501.63	1387.07	1317.34	17.096	1251.93	10,022		1403.01	1485.09	1567.78	1519.96	1417.35	1536.11	1563.00	1237.84	1571,96	1655.04	1506.71	125.42	40.73 EI																							
yte surfa	1421 01	1212 74	1419.94	1548.45	1141.61	1190.03	1337.06	1406.20	1283.87	1339.85	1336.07	1336.56		1503.22	1566.50	1618.38	1383,29	1568.37	1120.69	1400.82	1565.98	1277.49	1305.19	1431.00	161.48 ***£ 84	8.0141																							
Thomas	1477 34	1269.52	1302.40	1063.91	1210.95	1234.85	1276.70	1338.65	1357.53	940.79	1246.77	1258.15		1670,73	1577.54	1665.60	1483.98	1390.65	1399.03	1200.59	1444.25	1545,46	1550.04	1490.79	142.41	R ~ 1 1 1 1																							
nal dogs	1374 CO	1093 40	630.81	1116.91	998.17	874.84	913.29	1544.07	1383.48	1130,46	1126.04 ?29.46	1105.16		1737.33	1585.87	1476.13	1314,15	1911.05	1361.37	1453.02	1335.47	1690.51	1444,45	1531.92	196.43 1464.50	00.4044																							
ix E. Non	913 DQ	1322 52	190.63	1245.01	1399.43	1259.56	1323.29	1012.71	1139.02	1703.48	1251.51 217 34	12.11.2		1483.68	1548.65	1518.17	1756,65	1691.10	1583.72	1587.70	1743.30	1649.46	1610.41	1617,90	91,06 1 500 06	01.6661																							
Append											154.16	11.69.45												1158.07	181.91	101.40										1143.33	1129.95										1155.69	179.50	10.01
	1022.07	1135.84	1330.29	1047.58	1024.07	1031.84	1267.13	1286.86	1243.42	1220.91	1161.00	1149.42		1420,74	1285.06	1523.15	1170.50	1091.61	151.97	1333.88	1173.09	1418.55	1039.01	1260.86	160,13 1746 07	16'0121	1145.40	1112.92	805.00	02.020	1032.04	1061.52	1179.47	1022.67	730.79	573.47 466.96	10,798	07 971	1131.05	1071.68	984.02	1253,78	1185,84	1173.08	536.00	1103.42	1110.81	101.54	1 110.00
	1320 13	1135.84	1155.36	:208.15	723.02	1318.53	1131,45	1109.74	1007.33	1170.70	1128,03 170 25	1133.65		1171.50	1241.23	1086.23	1189.83	1364.76	1236.05	1069.89	760,40	885.20	795.54	1082.06	200.16 4084 45	CV-40/5	1136.04	1075.27	77700 77700	12061	1103.76	1089.41	1330.89	1213.94	1123.80	1114.94	1109.35	1570 34	1134.95	1289.21	1471.35	1244.42	1223.90	1181.26	1399.822	1233 26	1287.91	150.24 1770 e4	10.00.01
.	885 21	1052.76	1322.72	1095.99	1161.14	1153.11	1361.77	1306.13	1085.03	1231.07	1156.60 146.12	1163.87		990.59	890.73	1511.13	663.65	1076.86	1059.53	1146.79	1312.56	863.98	1067,30	1018.23	178.31	1020-00	1082.24	1258.37	C7777	1028.61	1115.32	1240.43	1269.72	1109.14	1057.34	1121.83	1112.23	1406.76	110430	1221.51	1124.08	1353.20	1201.78	1396.53	122828	1012 51	1221.67	161.28	1211.00
	17:4 93	1472.74	1074.87	898.55	920.26	1174.29	1042.20	981.63	1389.86	680.98	1085.03 227.09	1058 54		1246,21	1211.15	1078.4C	919.07	1353.40	1340.05	1224.30	1260.56	1432,70	1278.45	1234.44	145.99 1240.22	20.0421	1082.34	1246.81	813.49 1376 -	1275 10	1225.49	1049.37	1199.39	1187.04	800.29 1	1152,23	1169,64	410E 00	205.29	1401.42	1379.30	1276.30	1237,84	1257.97	1126.27	1210.15	1191.92	193.93 1204 E7	0.01
	22 T 76	809.89	1281.88	1053.80	1361.61	1351.21	1529.92	1178.07	1216.33	1175.88	1230.14	00 8241		1201.37	1268.13	1275.70	1248.20	1264.07	857.72	1201.78	1222-10	1322.72	1066.10	1194.79	116.14 1344 ou		1090.01	1284.66	1383.05	12221	1378.11	1483.10	1583.90	1384.48	1308.57	1356.37	1340.84	1000.76	1.000 02 UZU	785.58	1175,88	916.28	963,59	1001.55	839.94 00 00	6072 73	967.14	107.10	17.050
	Ndon 4	l shrador	retriever	10 years	FN					;	Mean	ande Mode		Ndog 2	Cocker	spaniel	6 years	EZ.						Nean	80	Abois	Ndog 3	German	snephero	d veare		ł			;	Mean	mode		Statistics shrathor	retriever	5 years	FN					Mean	5D 2011	anious

and the second second second

anakana kusha sukati se shakarin u du du su du du su su suka kada kada kana kunana u du Marin ana ata ata a se a se

1

يستقصف فسقا فسأشفاث

: الأرض

fotal											1468.44	189.12	1459.29																										
5	1169.51	1618.58	1407.15	937,40	1225.29	1453.02	1019.68	1477.92	1225.29	1483.70	1433.21	150.98	1426.22																										
	1424.33	1347.22	1206.78	1078.06	1282.87	1160.54	12:2.74	1543.27	1061.32	1550.61	1430.90	242.92	1369.64																										
Groh 3	924.45	1506.61	1378.90	1351.61	1316.14	1290.24	1019.28	1190.43	1168.51	1116.91	1498.72	249,31	1450.93																										
.	1123.05	1447.04	1656.83	1234.66	1422.73	985.41	1200.55	1114.12	1514,38	1387.23	1522.01	114.98	1528,96																										
4 4 5	1132.25	1442.06	1328.50	1141.22	1329.89	1269.52	1143.03	1646,87	1194,01	1334,47	1457.36	173.94	1468.94																										
Total											1299.61	222.44	1292.74											1414.55	194.67	1455.21											1524.91	226.44	1504.32
5 2 2 2 2	1397.63	1656.24	1525.14	1416.76	1268,13	1530.52	1285.86	1696.28	1131.66	1304.59	1426.28	172.15	1407.20	1594.67	1505.62	1554.23	1554.83	1017.49	1475.53	1263,15	1544.27	1695.68	1659.82	1486.63	202.44	1524.95	1681.54	1602.64	1469.18	1387.27	1414.96	1507.01	1547.45	1463.78	1436.08	1807.66	1531.76	131.93	1483.09
	1475.33	1658.63	1547.65	1098.58	1298.61	1778,97	1089.41	1163.53	1195.21	1471.35	1377.73	243.49	1338.17	1298.81	1396,63	1403.01	1407.59	1158.15	1366.75	1493.56	971.07	1319.33	1049.97	1286.50	170.90	1309.07	1220.11	1980.65	1439.67	1257.37	1466.76	1995.53	1716.41	1764.42	1619.98	1846.90	1540.98	298.04	1503.87
ltucrae	1210.15	1625.36	1259.56	1456.80	1411.38	1456.76	1395.44	1257.17	1498.04	1327,90	1391.86	128.68	1393.25	1202.50	1584.51	1646.28	1603.84	1283.65	1494.86	1487.49	1282.87	1411.78	1671 38	1467.52	165.32	3477.51	1568.50	1319.73	1552.24	1815.62	1433.27	1339.45	1671.98	1364.96	1564,19	1381.49	1503.64	162.65	1470.96
.	1496.25	1200.59	1330.49	1498.24	1226.69	1236.86	1306.18	1212.94	896.75	1058.53	1251.35	182.28	1239.02	1561.40	1376.11	1076.63	1056.14	1434,89	1571.95	1552.83	1757.45	1547,85	1253 34	1418.87	228.79	1426.83	1508.80	1912.05	1807.66	1674.97	1438.27	1442.66	1335.66	1501.63	932.22	1089.41	1464.43	299.48	1453.55
1	1337.46	998.17	1020.68	981,43	1258.56	1095.39	1040.80	1019.68	742.35	1021.67	1051.62	160.85	1021.18	1488.68	1080.65	1364.96	1235.65	1563.79	1338,46	1351.21	1633.92	1505.02	1569 77	1413.25	171.24	1389.09	(195.81	1836.94	1522.15	1886.95	1624.76	1481.11	1355.39	1770.40	1688.71	14/5.33	1583.76	219.17	1552.95
Total											848.37	115.68	850.04																								•		
C C	838.78	739.36	843.56	668.36	926.44	779.41	780.80	971.67	1028-05	946.56	872.35	93.82	856.21																										
	738.58	867.795	844.56	667,99	809.89	1090.01	990.20	910.10	874,04	855.51	228.87	96.21	880.92																										
. Eu	755.50	701.31	971.27	812.68	654,09	651.70	826.82	698.32	932.02	860.83	786.45	112.97	770.98																										
с. С	708.88	1021.67	778.41	686.16	676.40	740.75	762.47	634,56	670.46	980.23	786.00	130.78	751.61																										
	92B.03	763.70	1023.67	8C6.50	944.97	932.02	8CD.92	916.08	1074.87	900.94	908.17	99.59	912.13																										
	Ndog 5	Greyhound	3 years	Σ							Mean	8	mode	9 BopN	Greyhound	4 years	Σ							Mean	SD	mode	Ndoc7	Greyhound	5 years	Z							Mean	SD	made

ų

والمراجع والمراجع والمراجع

ميكم بنية بولتم تبه يتناعب كريد والكث كثي

. ‡

.

To the state of the

Ĩ

A. Townson

				10 1 10 10	CONTRACT BO	10tal x 10to	にはのアメリンの日本の		New York New York		New Address of the owner own						State Name of State o	
ADdog 1	32.67	38.41	39.20	42.14	44.71		42.42	43.84	38.86	41.55	44.47		36.44	44.40	38.04	40.23	37.23	
Labrador	35,61	36.67	43.16	36.06	41.22		41.69	42.90	46.03	42.61	41.60		28.35	42.90	31.90	43.46	43.74	
retriever	39.40	38.12	43.47	36.24	39.30		47.31	44.15	5.68	44.93	45.02		28.39	35.82	45.19	39,35	38, 19	
1 year	40.39	34.75	46.06	33.05	38.54		41,40	43.73	38,86	28.16	34,10		32.93	35.31	35,50	12.02	42.62	
3 months	36.33	39.42	43,66	42.24	42.11		37.98	40.54	47.49	39.62	18.32		31.97	35.68	37.42	50.95	36.72	
F.	35.79	37.25	33.57	37.03	37.70		38.02	47.53	39.79	37,36	42.59		30.61	41.25	35.60	44.98	41.32	
	34.43	37.78	38.64	37.02	40.77		40.67	39.26	43.71	43.09	40.63		40.57	45.02	38.67	42.57	43.25	
	37.42	44,89	33.71	42.23	4,13		34,33	43.17	40.37	41.17	38.06		39.10	41.31	43,74	43.58	35.21	
	38.17	38.10	37.50	40.15	40.70		45.39	45.37	39.79	35.38	34.51		29.11	39.84	44.64	35.46	43.71	
	37.41	27.95	40.55	33,92	38.45		42.46	47.51	35.22	37,18	43.37		40,47	36.87	39.90	42.12	41.50	
Mean	36.76	37.21	40.03	36,19	40.32	38.68	41.20	43.79	41.29	38.83	37.62	40.69	33.79	39.84	39,06	39,48	40.65	38.56
SD	2.30	4.44	4,49	2.88	2,04	3.54	3.75	2.B0	3.97	5.08	8.18	5.14	4.95	3.71	4.36	10.45	2.79	6.17
mode	36.76	37.25	40.03	37.03	40.32	38.43	41.20	43.73	39.79	38.83	38.06	41.47	32.45	39.84	38,36	41,18	40.98	39.61
ADdeg 2	29,08	38.84	26.80	30.47	37.07		36.43	47.70	46.34	44.29	37.97		41.07	25.66	34.32	29.24	37.43	
Labrador	40.69	33.31	30.63	35.00	34.10		40.00	44.91	44.43	44.50	40.15		32.00	26.96	27.52	35.62	37.47	
Cross	29.25	35.11	32.46	40.40	33.60		38.50	44.39	42.46	41.56	44,72		29.22	29.71	33.95	29.83	35.63	
5 years	32.65	37.17	31.32	30,59	34.78		44.58	45.33	43.94	37.59	41.83		30.38	33.29	30.66	29.00	34,65	
2 months	33 83 28	38.03	37.82	32.43	31.08		40.32	40.60	46.50	45.92	41.45		30.35	35.45	25.48	33.24	32.13	
ž	30.84	37.73	24.70	34,94	8.5		40.80	42.71	42.29	39.19	40.72		38.26	31.36	32.79	30.05	36.40	
	32.46	41,49	30.72	30.84	36.55		42.78	4C.77	46.79	41.49	42.67		31.63	36.45	28.15	34.18	36.97	
	29.48	28.39	26.90	24.06	34.00		43.20	43.21	43.74	40.04	38.89		32.69	24.68	31.23	33.85	37,99	
	30.84	34,59	35.40	36.63	36.18		37.27	41.08	41.53	39.06	41.89		26.37	32.41	34.96	21.13	41.18	
	33.97	33.20	32.78	29.92	40.75		43.23	44,86	45.95	44.64	38.47		34.40	32.74	31.05	28.18	37.32	
Mean	32.31	35.44	31.41	32,75	35.11	33.37	40.71	43.10	44.07	41.65	41.20	42.27	32.64	31.07	31,31	30.44	36.70	32.43
SD	3.46	3.74	3.98	4.58	2.64	4.01	2.74	1.90	1.78	3.10	1.92	2.75	4.33	3.86	2.67	4.17	2.35	4.11
made	32.31	35.11	31.32	32,43	34.78	33.45	40.71	43.10	43.94	41.49	41,20	42.38	31.82	31.21	31.14	29.93	36.84	32.55
ADdog 3	34.61	38,75	44.52	40,05	41.39		39.79	49.98	32.95	38.52	42.73		40.61	46.24	43.75	44.00	40.25	
WHW	33.31	34,39	44.12	39.14	40.41		47.12	44.70	46.17	38.71	41.54		40.84	19.13 1	40.33	48.77	48.53	
1 year	36.30	36,97	44.94	38,39	41.60		42.40	43.82	45.30	47.42	42.17		46.42	49.30	43.40	45.00	46.76	
6 months	40.71	40.45	44.55	34,92	42.17		45,91	30.61	45.86	38.80	38.76		47.06	48,35	42,40	48.91	42.52	
ž	33.40	36.40	40°39	35°28	33.84		41,43	33.57	42.23	36.64	42.72		48.44	36.56	40.36	40.07	47.81	
	33.48	45.98	44.96	32.97	42.61		37.82	44.80	44.70	41,10	43.95		39.03	45.07	42.35	42.31	41.64	
	34.08	43.16	43.76	42.75	41.86		38.26	38.95	46.88	43.82	43.39		47.09	44.94	39.14	41.73	45.43	
	38.08	36.60	39.63	36.01	44.13		42.80	421	43.58	42.39	38.42		47.15	35.06	38.43	-8.35	43.81	
	37.04	39.06	49.78	43.08	44,96		44.61	48.18	32.86	39.78	34.35		40.91	35.69	45.86	41.79	49.92	
	34.76	41,59	39.62	37.38	43.52		42.17	39.15	44.68	40.38	41.24		44,49	37.96	53.10	39.75	42.40	
Mean	35.47	39,40	43.59	37.58	41.68	39.59	42.23	41.57	43.58	41.11	40.72	41.79	45.10	42.19	42.91	44.07	44.91	43.84
SD	2.42	۲ <u>۲</u>	3,19	3.61	3.24	4.17	3.08	4.68	4.26	3.27	3.05	3.96	3.71	5.56	4.24	3.54	3.26	4,13
mode	34.76	39,06	43.76	37.38	41.85	39.84	42.23	41.57	44.68	40.38	41.24	42.31	45.76	42.96	42.37	43.15	4.36	43.78

त्रका करोकारतकी किंदीवर प्रतित के जिल्ही के <mark>किंदीन स्वतंत्रक क्रियंत्र क</mark>ार्यकों की वित्रित के प्रतित के अपने के स

.

4111

1994 - 1911

2

شهدته بالمسيد حامات

- 10-10

otal											45.86	3.84	45.65											40.23	3.45	40.15
- In In	47.95	40,81	42.78	44.28	45.52	44.75	47.74	48.11	45.86	44.93	45.27	2.34	45.10	37.96	44.93	40.41	39.21	42.71	39.68	42,16	40.05	35,92	45.56	40.86	3.01	40.23
	45.79	51.52	47.95	43.87	45.01	44.90	46.60	45.30	41.99	40.80	45.37	3.00	45,15	37.96	44,93	40.41	39.21	42.71	39.68	42.16	40.05	36.92	45.56	40.86	3.01	40.23
Groin 3	46.45	52.06	51.21	41.43	41.36	42.05	41.48	47.36	46.92	52.26	46.26	4.52	46.35	37,96	44,93	40.41	39.21	42.71	39.68	42.16	40.05	35.92	45.56	40.86	3.01	40.23
2	47.70	50.64	45,12	49.83	40.28	47.71	47.51	34.12	45.33	49.75	45.80	5.03	46.65	38.41	36.94	32.75	42,43	31.43	39.88	32.72	41.99	36.87	42.80	37.52	4.28	37.23
	45.27	39.48	47,95	51.27	51.91	50.11	42.67	44.44	43.13	50.07	46.63	4.23	45.95	36.89	∠1,8 3	39.92	36.05	41.66	46.39	43.66	43.42	40.28	40.56	41.07	3.18	40.81
Total											46.11	4.28	45.42											45.33	4.18	45.47
<u>, o</u>	45.22	43.23	35.61	44.72	47.49	42.31	44.85	43.21	42.07	44.44	43.10	3.25	43,25	44,39	39,67	43.08	41.79	50.76	45.17	49.32	47.00	47.43	42.50	45.21	3.67	45.17
4 1 1	53.35	44.39	44,41	54.15	51.18	48.12	51.42	50.72	53,11	50.16	49.74	3.47	50.16	44.38	38.90	46.67	47.03	41.17	40.87	43.03	57.74	48.11	50,64	46.04	5.86	45.04
1003	53.29	44.38	42.46	51.54	43,65	44.70	38.85	44.20	44,69	49.03	44.83	3.64	44.38	45.02	47.90	41.58	45.20	48.36	47,97	50.89	45.93	43.94	47.68	46.60	2.76	46.60
2	47.27	48.32	45.35	47.24	54.39	43.61	48.63	42.84	45.59	47.30	47.04	3.41	47,04	42.74	46.35	43.42	44.73	44.39	49,47	41.11	43.53	47.00	35.80	43.98	3.89	43.98
	38.49	37.17	42.40	47.12	46.24	42.53	46.82	45.49	46.87	50.95	44.42	4.24	45.49	45.46	45.48	45.67	46.97	33.72	46.66	40.69	48.31	46.63	54.06	45.36	5.25	45.67
Total											37.59	5.72	38.21											42.65	5.00	42.02
	51.78	38.68	38.03	40.34	36.86	40.92	43.11	42.18	40.16	44.88	40.57	2.54	40.34	35.33	34.69	40.27	33.46	35.83	33.92	35,11	33.24	41.67	47,55	37.30	4.86	36.11
	38.67	38.40	37.19	33.82	42.06	41.62	35.42	35.77	40.26	35.36	37.76	2.93	37.19	52.61	40.04	44,48	45.72	45.63	39.63	50.19	40.24	51.43	48.63	45.11	4.45	45.11
Ear. 3	45.87	42.12	37.36	32.05	38.40	35.56	39.30	37.61	34.87	48.32	38.42	4.68	37.81	41.65	50.39	40.48	48.66	37.75	39.51	47.48	44.62	43.80	43.24	43.99	4.29	43.80
ંત	41.66	44.97	42.07	37.48	40.45	35.75	39.71	39.89	36.19	39,44	39.55	2.89	39.55	43.55	48.85	47.91	47.94	42.37	40.82	44.27	47.93	39.18	47.16	45.16	3.60	45.16
	29.43	31.86	22.28	35.51	27.06	33.62	22.48	31.07	29.98	31.78	29.50	4.40	29.98	40.99	43.46	40.38	41.47	38.40	39.68	36.07	45.79	40.80	46.58	41.56	2.88	40.80
	ADdog 4	JRT	4 years	FN N							Mean	SD	mode	ADdog 5	WHW	3 years	ž							Mean	SD	mode

and a standard with the second standard of the standard standard standard with the standard standard standard a

and a state of the state of th
3 **		4.0	En .			Append	dix G. Ato	pic dogs	COLINEOC	yte surfa	ce area ()	լ. 		•	Goln :		ente Silve Sala	
ADdog 1	865.40	1138.03	1231.27	1186.24	1503.03		1199.23	1447.51	1272.99	1221.43	1382.97		1239.52	1350.71	1144.15	1182.45	1141.48	
Labrador	963.30	339.76	1144.00	1179.07	1322.72		1254.52	1478.95	1409.90	1417.50	1347.01		612.05	1407.43	769.28	1422.64	1366.53	
retriever	1061.72	1132.25	1510.80	976.85	973.66		1848.69	141.98	953,14	1364.07	1370.03		923.83	773.59	1450.39	1125.86	1204.78	
1 year	953.74	1237,25	1384.68	1069.09	1116.51		1528.69	1195.95	822.51	563.34	826.82		949.32	1204,37	963.71	615.96	1014,47	
3 months	1078.46	927.24	1007.93	1026.66	1043.59		1081,88	1203.04	771.33	1085.58	446.60		726.73	1147.24	1109.22	1198.82	1082.50	
FN	1295.62	1034,03	1184.85	1067.50	1127.27		1089.28	1565.51	1264.80	885.19	1233.75		992.27	1143.63	806.48	1314.12	331.66	
	358.10	1045.18	1350.21	1:34.44	1314.95		1081.06	958,75	1315,15	1520.26	1177.65		1328.10	1415.45	1264.80	1281.86	872.86	
	1050.56	879.42	1415.16	768.05	1338.46		636.92	1715,10	1320.70	1363.86	1129.56		1062.77	1417.71	(277.95	1366.74	1049.61	
	1176.48	962.70	1070.09	1116.31	1063.12		1460.46	1686.33	1115.59	715.43	929.38		613.26	317.67	1361.80	1047.76	1348.24	
	1315.35	1078,86	1277.29	1387.27	1313.35		1292.75	1666.19	943.56	1076.51	1444,32		1204.37	136.78	187,31	1561.16	839.77	
Mean	1063.87	1032.97	1260.56	1030.58	1179.29	1133.56	1247.35	1437.98	1101.52	1110.08	1100.57	1210.99	965.22	1063.21	1138.99	1214.99	1085.60	1099.52
ß	158.63	108.77	169.43	165.67	142.80	169.71	323.74	255.95	237.27	335.17	317.79	303.05	254.25	413.20	232.39	272.78	188.08	273.18
mode	1061.72	1032.97	1260.56	1069.09	1127.27	1116.41	1247.35	1441.96	1101.52	1085.58	1129.56	1244.14	949.32	1147.24	1138.99	1214.99	1049.61	1145.70
ADdog2	786.38	976.85	832.20	341.37	1006.13		1014.09	1533.47	1929.30	1435.67	1309.08		850.53	703.50	953.14	781.60	1083.04	
Labrador	918.47	1380.50	745.74	920.85	913.09		1276.70	1518.11	1629.77	1526.14	1336.24		634.16	783.19	715.65	696.56	1064.31	
Cross	919.47	1004,54	737.96	878.62	803.31		1430.20	1322,24	1327.67	1417.73	1639.17		805.11	694.93	879.02	741.35	905.38	
5 years	1014.30	1352.01	747.93	851,33	729.60		1336.04	1551,64	1292.16	1232.97	1283.17		729.40	899.54	666,44	549.29	1041.00	
2 months	873.05	1009.92	396.75	922.DG	1114.12		1446.95	1177.47	1535.13	1614.11	1427.12		753.11	868.28	977.25	814.47	822.24	
W	744.94	963.62	878.62	1217.92	915.08		1378.64	1351.27	1347.30	1321.63	1236.58		873.25	831.80	707,88	593.32	1010.72	
	783.59	1146.40	681.38	749.12	976.05		1534.28	1155.94	1523.02	1254,55	1222.17		823.63	920.07	807.30	693.93	1444.45	
	798.73	1099.58	639.14	738.35	876.43		1493.35	1637.71	1595.48	1494.81	1345.84		656.28	672.22	875.84	706.09	1017.29	
	892.37	1073.48	597.11	964.89	1031.44		970.01	1099.11	1452.18	1254.55	1364.22		894.76	816.06	830.61	583.96	970,87	
	838.13	1007.33	710.87	914.49	1072.48		1218.20	1710_41	1437.3E	1347.74	1111.85		780.05	824.23	707.68	585.15	1135.04	
Mean	856.95	1117.51	737.28	906,41	936.84	910.17	1307.85	1391.32	1458.90	1390.47	1329.60	1387.97	780.05	812.26	796.41	661.57	£055.70	\$25.10
30	82.07	150.66	99.18	140.43	125.39	164.11	\$91.70	222.68	118.92	129.96	146.58	180.37	87.72	84,51	103.63	88.41	168.28	165.97
mode	856.95	1073.48	737.28	906.41	915.08	904.92	1336.04	1351.27	1452,19	1347.74	1329.60	1357.75	780.05	816.06	796.41	661.57	1017.29	\$15.27
	1038.41	67 2051	1503 83	1243 34	1150 55		1120 GG	1746 B1	80.4.4.4	21 2021	1004 80		424.47	1740.02	1966 70	1505 64	4 E 3 E	
WHW	1179.07	1085.63	1587.70	1145.00	1261.75		1589.69	1373.72	1778.57	1716.41	1366.75		1737.33	1533.71	1097.18	1642-09	1602.84	
1 year	1115,12	1023.07	1452.82	1336.66	1569.37		1521.55	1464.17	1386.67	1957.28	1419.35		1562.40	1659.03	1681.74	1514.58	1576.94	
6 months	1215.33	1050.96	1590.29	919.67	1410.38		1517.17	1245.02	1621.97	1633.91	1393.85		1761.83	1429.11	1585.11	1778.771	1445.65	
Ξ.	943.77	1006.33	1436.48	1171.30	816.06		1342.44	1023.87	1578.73	865.71	1324.31		1587.90	1493.46	1516.18	1346.43	1548.05	
	927.44	1431.10	1517.17	823.83	1156.76		1239.64	891.57	1538.69	1378.11	1169.51		1509.40	1453.22	1168.91	1609.22	(277.89	
	1039.21	1241.43	1331.88	1110.53	1254.98		1511.19	1726.17	1514.38	1741.71	1074.47		1865.63	1102.76	937.80	1372.33	1665.80	
	1470.55	1281.68	1591.88	1228.48	1339.25		1176.88	1338,85	1659,42	1221.51	1412.17		1689.91	1012.11	1633.33	1526.73	1530.12	
	1193.61	1380.50	1680.74	1357.98	1718.20		1710.03	1335.07	1162.93	1441,66	1199.79		1779.16	1172.69	1344.43	1331,68	1555,23	
;	1112.92	1024.28	1497.65	1156.96	1081.05		1250.79	1607.23	1648.87	1250.00	1455.77		1605.63	1142.41	1768.61	1024.07	1067.70	
Mean	1123.54	1152.24	1518.94	1159.37	1276.74	1252.17	1399.20	1333.84	1543.14	1466.59	1313.97	1398.96	1656.34	1333.17	1414.81	1460,86	1474.47	1476.10
SU.	157.11	151.83	56°26	£07.77	254.22	222.61	195,93	261.45	179.08	334.75	134.30	248.13	130.09	227,49	290.97	221.23	168.27	227.48
тоде	21.0111	1085.63	1517.17	1156.96	1261.75	1234.96	1399.ZD	1334.07	1543.14	1441.68	1324.31	1390.15	[656.34	1333.17	14.81	1460.66	1530.12	1524.05

Appendices

Total											1631.91	276.75	1650.46											1130.15	195.66	1133.03
5. S.	1838.54	1353,60	1587.50	1710.83	1690.31	1367,55	1733.14	1680.94	1533.91	1615.00	1585.86	142.48	1587.50	1083.04	1064.31	595.38	1041.00	822.24	1010.72	1444.45	1017.29	970.87	1135.04	1055.70	168.28	1017.29
	1680.94	1183.45	1291.04	1907.87	1529.32	1816.02	1373.72	1162.33	1835.75	1567.38	1518.54	288.16	1518.54	960.31	1226.09	1402,81	1057.74	1319.93	838.58	1144.20	1233.06	1359.78	1274.90	1208.34	173.94	1226.09
Groin G	1921.02	1692.70	1407.79	1262.35	1613.00	1162.93	1770.40	1714.21	1587.50	1913.65	1569.39	245,69	1587.50	1061.52	984.62	1216.13	1365.95	853.92	1070.69	1360.17	1183.45	1199.59	692.74	1103.03	226.19	1103.03
2	2198.95	2415.72	1733.94	1984.97	1337.86	1509.60	948.16	1777.37	1593.28	1706.05	1667.44	408.53	1667.44	1141.02	1172.50	845.95	936.60	996.37	953.74	1378.90	1340.25	1166.32	1373.92	1129.28	204.85	1129.28
t.	2064.67	1476.33	1780.76	1713.82	2000.71	1592.68	1619.98	1467.36	1369.26	1779.56	1688.51	224.54	1619.98	995,58	1116.31	1255.38	922.85	1188.23	1742.90	934.61	1225.89	1215.93	1214.53	1181.22	233.71	1188.23
Total											1636.83	207.55	1648.87											1563.59	203.98	1549.85
С. Э	1729.75	1596.47	1246.01	1766.61	1503.03	1451.03	1340.25	1656.44	1473.54	1485.09	1502.05	157.39	1485.09	1578.14	1574.75	1530.92	1807.26	1699.07	1507.81	1509.00	1690.31	1662.21	1352.80	1592.68	135.84	1574.75
	1691.50	2178.83	1583.32	1463.96	2089.57	1740.51	1755.26	1565.19	1638.71	1882.57	1772.55	239.01	1740.51	1513.37	1450.23	1610.01	1405.20	1841.13	1243.42	1906.08	1526.93	1494.26	1551,64	1558.77	207.07	1526.93
Thoras	1434.49	1321.92	1829.37	1491.67	1641.29	1837.34	1591.68	1307.78	1521.36	1652.25	1577.18	190.20	1577.18	1782.15	1044.79	1853.68	1800.88	1459.39	1461.39	1522.95	1315.55	1406.40	1242.43	1456.38	254.58	1456.38
	1835.75	1921.02	1714.41	1686.72	2041.76	1562.40	1627.33	1516.37	1782.35	1759.24	1723.51	178.12	1714.41	1548.05	1574.35	1532,51	1673.17	1250.03	1717.60	1310.17	1424.53	1337.26	1376.31	1466.21	165.70	1424.53
	1389.26	1430.30	1721.98	1733.34	1843.32	1563.00	1184.05	1732.15	1645.48	1729.36	1597.22	204.30	1645.48	1559.21	1847.30	1775.98	1699.07	2068.65	1602.48	1742.90	1565.19	1699.47	1326.30	1708.66	197.94	1708.66
Total											1112.18	267.84	1143.91											1396.22	259.56	1405.30
-	1584.71	1156.36	1316,54	1414.17	1528.33	1165.52	1301.40	1210.95	1154.56	1219.71	1274.17	129.27	1219.71	1370.93	1075,47	1103.76	845.35	1054.75	1102.17	998,96	1242.83	1073.08	1088.22	1084.95	104.84	1073.05
	1116.51	1326.11	1143.41	1103.96	1343.84	1172.10	995.97	1061.92	1174.09	1068.50	1154.43	\$17,14	143.41	1982.18	1626.55	1452.82	1003.54	1406.00	1816.02	1601.65	1713.02	1143.21	1576.54	1482.13	264.44	1482.13
Ear.	1717.00	1421.14	954.73	1377.11	1000.16	1141.81	1231.67	1366.47	1176.88	1283.27	1219.25	167,20	1219.25	1148.59	1537,49	1316.14	1940.54	1237.84	1183.65	1385.08	1706.64	1309.77	1492.67	1456.65	243.24	1335.08
2.4	1327.30	t426.52	1259.56	1104.95	1111.53	1101.77	1112.92	795.15	1144.40	980.63	1115.27	173.01	1111.53	1427.32	1464.17	2030.40	1587.50	1373,32	1416.16	1439.87	1517.57	1433.89	1286.26	1505.46	214,20	1439.37
	644,13	791.36	593.52	661.86	653.49	710.67	587.34	961.31	755.90	635.56	699.51	112.93	661.86	1262.15	1567.38	1372.13	1440.27	1314.85	1244.02	1404.60	1424.33	1730.95	1538.89	1429,91	149.66	1424.33
hi ki n	A DobdA	JRT	4 years	N							Mean	\$D	mode	ADdog 5	WHW	3 years	R							Mean	SD	mode

Appendices

---- ---- ---- -----

έ÷ ι

å

nan Daar It (Krayk) SH Years N	ខែងរា រៀ រាបថំអ	SH 25 SH 25 Boan Co co co co co	14년 14년 38년 14년 14년 14년 14년 14년 14년 14년 14년 14년 14	Serta Serta ode marka ode marka Neers Nee	SSH SSH SSH Jyears Dean Dean Stars
58.92 58.92 59.37 59.37 55.99 57.91 44.79	44.13 44.13 44.1544.15 44.1544.15 44.15 44.15 44.1544.15 44.15 44.15 44.1544.15 44.15 44.1544.15 44.15 44.1544.15 44.15 44.154	94.20 36.28 41.57 41.57 41.15 41.16 71.33 41.10 77.39 42.56 33.62 33.62 33.62 33.62 33.62	47.18 57.67 26.26 37.67 37.65 37.65 37.73 37.73 27.73 27.73 27.73 27.73 27.73 27.73 27.73 27.74 27.73 27.74 27.74 27.75	2002 2010 20102 2010 2010 2010 2010 2010 2010 2010 2010 2010 2000 2010 2000 2000 2000 2000 2000 200000000	
38.32 38.32 44.83 40.44 39.83	33.7.27 33.7.27 33.7.24 45.3.24 45.62 25.96 25.96 25.96 25.96 25.96 25.96 25.96 25.96 25.96 25.96 25.96 25.777 25.777 25.777 25.777 25.777 25.7777 25.	7.48 2.22 2.22 2.22 2.23 2.23 2.24 2.25 2.25 2.25 2.25 2.25 2.25 2.25	64 64 64 64 64 64 64 64 64 64 64 64 64 6	7285 2017 2017 2017 2017 2017 2017 2017 2017	
4047 35.25 35.25 46.30 46.30 46.30 46.30	42,23 42,29 42,20	867588888888888888888888888888888888888	44444444444444444444444444444444444444	88888888888888888888888888888888888888	
42.65 42.65 42.03 27.77 43.16	41.15 41.15 41.15 41.15 41.15 41.15 41.15 41.15 41.15 41.15	66.49 27.32 27.32 26.47	8289 8488 8488 8588 8588 8588 8588 8588 85	46 46 46 46 47 46 46 46 46 46 46 46 46 46 46 46 46 46	
200 200 200 200 200 200 200 200 200 200	4 4 4 7 3 2 4 4 4 5 5 3 5 4 4 4 5 5 3 5 4 4 4 5 5 5 5	40.58 37,79	003222684444444444444444444444444444444444	4 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	
	5-12 6-13 6-13	38.10 3.23 37.82	43.04 4.70	37.05 5.00 37.64	
16.88 16.88 17.54 17.54 19.01 19.01	24,13 18,41 18,20 19,20 19,20 19,30 19,30 19,30 19,30 19,30 19,30 19,30 19,30 19,30 19,30 19,30 19,30 19,30 19,30 19,30 19,100 19,10 10,1000			22222222222222222222222222222222222222	12.12.00 12.12.00 12.12.00 12.12.00 12.13.00 12.
16.37 16.11 16.11 23.78 16.49	22.46 15.31 17.23 17.48 17.48			R 21 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	12224222222222 12222222222222 1222222222
21.77 21.77 15.77 15.74 15.42 21.54 21.54	2559 2540 25240 25240 2525 2540 2525 2526 2526 2526 2526 2526 2526 252			080,411 080,411 080,412 080,412 080,412 080,441000000000000000000000000000000000	200 200 200 200 200 200 200 200 200 200
21.71 15.74 16.74 18.18 19.42 21.54	16.91 16.15 17.15			7,22,22 7,26,10 7,26,000 7,26,0000 7,26,0000 7,26,00000000000000000000000000000000000	17.37 20.38 15.27 17.52 18.93 18.93 18.93 18.93 18.93 18.93 18.54 18.72 18.54 18.54 18.54 18.54
11.66 11.66 11.66 11.28 17.28	17.11 11.111			12.86 12.86 12.85 12.95 12.20 12.44 12.93	15.80 (15.82 (15.85) (16.57 (16.57 (16.57 (16.54) (16.54) (16.54) (16.54) (16.54) (16.54) (16.54) (16.54) (16.54) (16.54) (16.55) (16.
4 	17.76 3.54 17.33			1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	년 산 25 년 (
			50.13 52.14 52.24 53.24 52.24 52.24 52.24 53.24 52.24 52.24 53.24 52.24 53.24 52.24 53.24 52.24 53.24 52.24	8	
			50.08 51.025	37.20 25.20	
			2012 2012 2013 2013 2013 2013 2013 2013	88848888888888888888888888888888888888	
			ᡩᢄᡷᡷᢧ᠍ᢗᡁᠺᡖᡷ᠀ᢩᡘᡕᢩᠼᡕ ᢊ᠍ᡱᢋᡦᢡᡇᢓᡩᡩ᠖ᢞ ᡭ ᠼᡕ	848 848 85 85 85 85 85 85 85 85 85 85 85 85 85	
			42.45 49.65 49.45 49.45 49.42 49.42 49.45	60.83 60.83	
			N 4 B	ने ⁻ भ	

and the second second

and the second second

المتعمية بمراجع ساليا المرابية ألمان ألمعان مريم لأرامع أنام والمرابع معاريهم منيسيا الارباب والمالية والمناف

1

يترقوها فلاتهم والموسية الألات الأمناء والمراجعة المتطلق للمراجع المراجع والمراجع والمراجع والمراجع والمرابية

	40 10 10	99-62 1	52.83 4.07			53 5 3 57 66 57 68
	8 5 7 7 9 9 7 8 8 8 8 9 9 9 9 9 9 9 9 9 9 9	8, 10, 10, 10, 10, 10, 10, 10, 10, 10, 10	52.76 52.76 52.51 52.51 52.51 52.51 52.51 52.51 52.51 52.51 52.51 52.51 52.51 52.51 55.55 55.555		88888888888888888888888888888888888888	228 228 288 288 288 288 288 288 288 288
	8888 8888 8888 8888 8888 8888 8888 8888 8888	41.03 57.03	59.31 57.66 57.66 52.67		1747 1747 1747 1747 1747 1747 1747 1747	51.75 3.56 3.56
	4844444484 8486888 85888 8888 8888 8888	22.24 22.25 22.25 22.25 22.25 22.25 23.25 23.25 23.25 23.25 23.25 23.25 23.25 24.25 25.25	51.18 51.18 51.56		ស្តេសមេស្ទសទទេស វស្តេសទេសទេសទេស វស្តេសទេសទេសទេស	55.07 3.35 54.57
	44 44 44 44 44 44 44 44 44 44 44 44 44	444 444 444 444 444 444 444 444 444 44	51.15 54.74 5.87 5.87 5.87		2000 2000 2000 2000 2000 2000 2000 200	51.25 2.45 51.76
	2212 2212 2212 2212 2212 2212 2212 221	1977 1978 1978 1978 1978 1978 1978 1978	55,25 56,255		\$ \$ \$ \$ \$ \$ 5 5 6 6 5 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	26.25 26.23 27.23
14.79 3.27 14.90				6.17	24-1 01-4-2	
25.00 25.000 25.000 25.000 25.0000000000				れらたかけれた役 株体務 第44年8月2月2日 第44年8月2日 第44年8月2日 第44年8月2日 第44年8月2日 第44年8月2日 第44年8月2日 第44年8月2日 第44年8月2日 第44年8月21日 第45 8111111111111111111111111111111111111	2.83	
13.01 15.28				11.85 5.52 5.53 5.53 5.53 5.53 5.53 5.53 5.5	3.27	
20.28 20.28 20.29 20.29 20.25				7.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0	13.55 10 10 10 10 10 10 10 10 10 10 10 10 10	
17 17 17 17 17 17 17 17 17 17 17 17 17 1				に 11111 11111 11111 11111 111111	2.80	
25.25.25.25.25.25.25.25.25.25.25.25.25.2				10,82 10,12 10 10,12 10 10,12 10 10,12 10 10,12 10 10,12 10 10,12 10 10,12 10 10,12 10 10,12 10 10,12 10 10 10,12 10 10,12 10 10,12 10 10 10,12 10 10 10 10 10 10 10 10 10 10 10 10 10	3.03 13.10	
42.09 2.43 42.60						
44.13 44.13 42.60 42.60 45.61 45.63 45.64 45.64 45.64 45.64 45.64 45.64 45.64 45.64 45.64 45.64 45.64 45.64 45.64 45.66						
1995年1995年1995年1995年1995年1995年1995年1995						
900 900 900 900 900 900 900 900 900 900						
440 440 440 860 860 860 860 860 860 860 860 860 86						
40.4 40.4 5.5 5.5 5.5 5.5 5.5 5.5 5.5 5.5 5.5 5						
Cartis DSH 7.3 years Mc an Mc	DGH ()) 7.2 years Mr M Ban SD mode	Cart 8 DSH 12.4 ysars MN	Mean SD Riode	Contra 1.9 years Min Maari SD	mode Cat to DSH MN MN	mean SD mode

			1868.75 276.04 1955.87	1464.73 256.85 1361.77	
			1771,98 1546,46 17404,07 1739,52 1739,52 1784,34 177,56 1646,92 177,56 177,56 177,56 177,56	2005.09 1470.81 1780.85 1780.86 1780.86 2035.55 2035.55 1673.57 1566.59 1575.24 2753.24 2753.24 2753.24	
			04.000 16.000 16.000 17.000 17.000 17.000 17.000 17.000 17.000 10.0000 10.00000 10.00000 10.00000 10.00000000	1377.11 1428.11 1428.13 1138.83 1138.83 1138.83 1225.21 1225.10 1237.8.53 1605.43 1705.44 1605.441005.44 1605.44 1605.441000	
			2573.05 2862.40 2962.40 2169.05 2165.73 1655.73 1655.73 1655.73 1655.73 423.24 2005.89 423.24 2005.89	1473.94 1587.89 1587.89 1653.81 1665.81 1665.81 1665.83 1665.83 1665.38 1665.38 1722.38 1655.38 1722.38 1655.38 1665.26 1665.26 1655.2	
			1963.56 2042.15 1964.25 1964.25 2042.15 2042.15 2042.36 2042.36 2079.2 2079.2 2079.2 2079.2 2079.2 2079.2 2079.2 2079.2 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2075.5 2 2 2075.5 2 2075.5 2 2075.5 2 2 2075.5 2 2075.5 2 2075.5 2 2 2075.5 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	1871.5° 1078.28 1279.29 1279.29 1229.24 1472.50 140.55 1172.50 140.55 1222.70 1222.70	
			1943.73 1988.16 22345.19 22345.19 1862.47 1885.36 1885.36 1885.36 1835.36 1835.36 208.86 208.86 208.86 208.86	1125.48 883,43 883,43 10045.73 10045.73 10045.75 1025.38 11225.38 111225.34 111225.55 111225.55	
	227.34 50.13 226.73			177.86 32.78 181.01	192.19 48.26 163.06
ea (µm²)	17.12 245.57 215.57 215.57 245.57 245.57 245.55 245			200.80 126.94 2107.19 2107.19 205.57	178.31 225.33 225.33 225.35 210.79 1582.35 1582.35 234.50 244.50 224.50 224.50 224.50 224.50 224.50 224.50 224.50 22
surface an	207.20 135.68 155.38 150.38 150.38 150.38 187.68 197.68 196.64 196.64 196.64 196.64 186.23 37.09			143.65 168.75 168.75 169.27 155.97 171.94 171.94 185.90 205.51 205.51 205.51 205.51 205.51 205.51 205.51 205.51 205.51 205.51 205.51 205.51 171.94	185.85 205.02 205.02 205.02 205.03 140.86 140.85 245.03 245.05 24
comeacyte	230.22 230.22 210.51 210.51 210.51 225.69 255.69 255.69 255.69 256.69 25			181.30 188.73 188.73 188.73 188.73 188.87 188.87 188.87 188.87 188.87 188.87 188.87 17.88 21.38	153.0° 231.1° 231.1° 238.60 228.66 193.46 193.46 193.46 193.46 193.46 133.43 41.24 133.43 132.46 133.43 132.46 133.43 132.46 133.43 132.46 133.43 132.46 132.46 133.45 145.45 145.45 145.45 145.45 145
ormal cats	180.31 180.31 167.83 167.83 222.53 227.53 254.22 254.22 254.22 255.54 262.59 27.59 2			147.03 207.20 207.20 207.20 207.20 1280.97 1280.97 1280.97 1280.73 120	183.57 195.05 113.57 113.56 151.22 113.56 175.92 251.04 175.92 175.92 175.92 175.92 175.92 175.92 175.92
pendix 1. N	250.04 243.06 243.06 243.06 3325.04 37.0.01 37.0.01 37.0.01 250.24 250.24 250.24 250.24 250.24 250.24 250.24 250.24 250.24			206.01 136.40 152.66 16.215 16.215 16.215 17.63	231.26 97.82 97.82 135.09 135.09 2175.66 202.02 202.02 205.65 175.93 175.93 175.93 155.86 155.86 155.86 155.86
AP S	1200.68 197.07 127.5.04	1121.13 211.12 11136.02	1573,85 257,57 1337,88	1093.85 158.50 1100.47	
	1514.75 1554.90 1354.90 1493.06 1493.06 1443.06 1443.06 154.13 054.13 054.13 054.13 1556.66 1257.66 1257.46 1257.46 1256.10	1020.89 1218.92 1258.37 1258.37 1259.57 1250.61 1250.61 1248.02 1248.02 1248.02 1248.02 1248.02 1248.02 122.63	1261.55 1469.64 1469.64 1515.16 1512.16 1823.55 1823.55 1123.25 1174.09 1153.25 1553.25 1553.25 1553.25	1125.68 930.43 733.58 733.58 733.58 733.58 733.58 965.45 761.08 767.09 767.08 767.08 767.08 767.08 767.09 767.09 767.08 767.08 767.08 767.08 767.09 767.09 767.09 767.09 767.09 767.09 767.09 767.09 767.09 767.09 777.00 7	
	1238.24 1423.53 1681.72 1681.72 1688.14 1146.57 1148.57 1148.79 1231.49 1231.74 1331.74 1331.74 1331.74 1331.74 1331.74	824.23 857.67 11112.55 1165.65 1165.65 755.09 1105.04 1125.55 210.28 210.28 210.28	168.94 1386.53 1386.53 1386.53 1406.74 1571.55 1571.55 1571.54 1520.37 1520.37 1336.44 1336.44 1336.47 1346.57 207.14	1465.18 965.61 972.45 977.45 977.45 977.45 1772.45 1207.33 1207.33 1207.33 1207.33 1207.33 1072.96 1072.96	
	120.05 1140.22 1086.65 1086.65 1088.97 1088.97 1281.94 1135.04 1135.04 1135.04 1135.64 1135.64 1135.64 1135.64 1135.64 1144.65	1269.05 874.05 855.31 855.31 12576.44 12576.62 1353.02 1353.02 1363.25 295.05 295.05 374.05	1153.35 1340,25 1240,25 1242,29 1242,29 1186,59 1186,59 11107,60 11107,60 11107,60 11107,60 11107,60 11107,60 11107,60 11107,60 1122,65 1122,55 1125,55 1125,5	149.38 1407.48 266.72 266.72 266.71 11182.34 11182.34 11182.35 11182.35 11185.34 1065.54 136.10 1065.54 136.10 1065.57 1065.57 1065.57	
	100534 1015.55 970.27 11275.51 11275.51 1044.28 1044.28 1044.28 1055.79 1055.79 1005.55 1205.5	1149.58 974.28 974.28 1058.35 1607.35 11407.00 11409.50 1149.50 24.72 24.72 24.72 24.72 119.50	1479.72 1535.72 1535.72 1535.72 2640.15 1678.55 11168.55 1170.50 1170.50 1170.50 1179.50 1177.	1042.99 1157.80 1110.14 11331.48 1110.14 11139.02 11139.02 11141.22 11288.80 100.79 11141.22 100.74	
	1341.523 1372.53 1372.53 1274.55 860.43 860.43 1665.95 1665.95 1666.93 1666.93 1666.93 1666.93 1666.93 1666.93 1666.93 1675.45 1655.74 1655.74 1655.74 1655.74	834,79 991,39 994,58 984,57 1024,58 1024,58 945,77 945,77 1060,02 1054,60 1054,60 1054,60 1054,60 1054,60	1208.37 1105.55 131.57 131.57 133.109 145.55 1445.25 985.98 1445.25 1445.25 1445.25 1445.25 1100.97 1100.97 1100.97 1120.81 1100.97 1120.83 1120.83	1305.19 1267.13 855.73 855.75 11218.12 1145.26 1175.26 1175.25 1175.25 1175.53 1175.53 1175.54 1171.50	
	Meen SH 11 years Min Ma SD SD SD SD SD SD SD SD SD SD SD SD SD	Catility Cat	Caracter D1.H 5 Years Mn Mean 3D 3D mode	C.## D.5H B. 9 years F.N Maan Maan Maan Maan Maan Maan Maan	CALSS CISH DISH Bury years FN Maan Maan Maan Maan Maan

S.

and the second of the second second

The survey of the strength

		1227.36 195.21	40,4121	2077.76 268.70 2132.41					2161.95 257.55	2160.60
	1137.03 1001.92 1102.96 1148.09 1204.97 1311.96 1232.47 1415.55 1146.00	1694.69 7257.95 196.67	194,97 1948,7: 1972,222 2990,78 1932,58 1932,58 2118,46	2463,13 2436,19 1627,78 1743,50 2063,71 172,52 2053,71			3364.27 2506.37	2126.65 2498.40 1825.00 2408.35 2013.05 2013.05 2013.45 2177.45 2177.45	2177.B1 236.62	2126.63
	1319.73 1512.89 1565.15 1085.38 1085.38 1085.38 1387.41 1587.41 1587.41 1587.41	1570.17 1327.88 176.57	722025.03 2025.03 2306.54 2007.18 25644.42 25644.42	1684.14 2021.63 1978.18 1872.18 286.27 286.27 2087.18			1713.22	2004.50 2005.89 2016.89 2018.05 2130.26 2380.26 2380.26 2380.26	2139.09	2110.89
	1150.38 922.26 11116.90 1158.65 1413.17 1536.30 1455.61 1255.06 1255.06	928.43 1 199.26 234.60	1158.55 1818.47 1850.97 2026.81 2026.81 1902.69 1902.69	2183.41 2174.45 2147.35 2147.35 2021.59 181.27 2021.59			2077.22 2136.38	2368.50 2152.52 2152.52 2266.25 2193.57 2018.02 2218.02 1546.51	2175.93 153.73	2175.93
	256.22 1197,02 256,22 256,22 256,22 1086,20 100,20 100	1127.47 1170.70 153.42	1730.35 2524.00 2524.00 2346.56 1984.54 2346.56 2346.56	1486.20 2261.11 2204.73 1566.95 2003.97 387.64 2003.97 2003.97			1691.50 1878.33	2278.04 2355.15 2004.51 2004.51 2004.51 1955.29 2286.21 1757.65 2786.65	2695.67	2093.67
	1529:52 924:25 1285:66 1(90.83 1383.16 1447,84 815.84 815.84 115:417 703:42	1223.10	2327.06 2425.48 245.39 2141.57 2141.57 2141.57	2456.56 2475.09 2229.43 1584.71 2238.97 273.52 273.52 273.52			2176.43 2380.45	2104.51 2265.88 2316.30 2317.2.85 2049.05 2049.05 22127.43 2127.43 2127.43	2209.50	2172.85
28281 28281 2011						135.46 42.33 134.68				
135.28 125.12 1125.12 1125.12 1125.12 1125.12 1125.05 1177.96 177.96 177.96					139.46 157.59 157.59 141.26 141.26 144.44 172.52 164.65 164.63 164.63	98,441 77,34 70,961				
112.77 185.29 209.40 201.78 201.78 201.78 201.78 153.24 153.24 153.24 152.48					74.55 165.65 208.66 208.66 208.66 201.65 152.23 152.23 152.23 152.23 152.23 152.23 153.23 153.23 153.23 153.23 153.53 155	138.51 46.30 132.29				
208.80 143.25 143.25 143.25 143.25 143.25 147.17 147.19 147.17 143.25 143.78 143.78					8972 8972 1998 1997 1998 1998 1998 1997 1998 1997 1997	67.921 52.06 11.721				
127.77 157.40 246.45 106.19 1135.48 1135.48 1135.48 1135.48 1135.48 1135.48 1136.08 1136.08 1136.08 1136.08 1136.08 1136.08 1136.08 1136.08 1136.08 1136.08 1136.08 1137.00 11000000000000000000000000000000000					22122 2212 2212 2212 2	127.85 42.82 10.121				
130.50 151.22 151.22 151.22 148.15 148.15 148.15 148.15 161.75 161.75 161.75 161.75 161.75 161.75					167,78 157,78 157,78 130,07 73,92 175,72 100,21 100,21 100,58 160,58	102.41 38.45				
1315.3Z 221.25 221.25										
1068./3 1547.85 1255.45 11151.53 11547.85 1341.45 1174.49 986.7.50 1174.49 986.7.50 1455.27 1455.27 1455.27 1238.57 180.27 1204.97										
1235.25 1273.57 1573.57 1533.52 1539.25 1650.50 1540.55 1400.50 1360.50 1400.50 1400.50 1400.50 1400.57 147.75										
1367.25 1354.00 1355.84 1728.84 1728.85 2365.84 2355.21 2357.45 024.45 024.45 024.45 2381.41 2										
11736.17 1726.17 1422.14 1551.44 1551.44 1513.52 1334.20 1271.32 1335.27 1335.27 1533.25 1335.25 1335.25 1337.74										
1026.66 1224.66 392.50 1071.58 1334.55 1334.55 1337.55 888.50 1481.11 1401.25 1401.55										
CERTS	Carly DGIT 7.2 years MAN	Mean SD mode	Cat 3 DSH 13.4 Years f.in	Mean SD mode	Carlowson DSH 	Mean SD mode	Cartan () DSH	16 years MN	Maan SD	abom

Appendices

Appendices

Appendix J. Scripts used in image analysis studies.

Script 1, 400 x 400 pixel box. macro 'box 400 x 400'; {Produces a centrally located box 400 x 400 pixels. Copies selection to new window.}

```
var
w,h:integer;
begin
GetPicSize(w,h);
MakeRoi((w-400)/2,(h-400)/2, 400, 400);
copy;
paste;
end;
```

```
Script 2. 200 x 200 pixel box.
```

macro 'box 200 x 200';

{Creates a 200x200 selection centred on the image. Copies selection to new window.}

var

```
w,h:integer;
```

begin

```
GetPicSize(w,h);
```

```
MakeRoi((w-200)/2,(h-200)/2, 200, 200);
```

copy;

paste;

end;

Script 3. Count bacteria

.macro 'count bacteria';

{Opens user define ROI. Copies selection to new window.

Opens threshold plugin. Find objects gives count. Opens object classification.

Open dialog box with result. Closes all windows}

begin

GetUserRoi('Outline the area to add to the object set', true);

copy;

close;

paste;

PlugIn('C:\\Program Files\\UTHSCSA\\ImageTool\\Plug-Ins\\bilevel');

FindObjects;

NextWindow;

```
PlugIn('C:\\Program Files\\UTHSCSA\\ImageTool\\Plug-Ins\\objclass.dll');
```

ShowMessage('count is: ', GetResults('count', 0));

CloseAll;

end;

REFERENCES

AARESTRUP FM, Jacobeson MJ (1993). Bacterial paronychia caused by *Staphylococcus felis* in cats, *Dansk Veterinaertidsskrift* 76: 1066-1067

ALLAKER RP, Lloyd DH, Bailey RM (1992). Population sizes and frequency of staphylococci at mucocutaneous sites on healthy dogs, *Veterinary Record* 130: 303-304

ALLAKER RP, Lloyd DH, Simpson AI (1992). Occurrence of *Staphylococcus intermedius* on the hair and skin of normal dogs, *Research in Veterinary Science* 52: 174-176

ALLAKER RP, Noble WC (1993). Microbial interactions on skin, in *The Skin Microflora and Microbial Skin Disease*, ed. Noble WC, Cambridge University Press, Cambridge p 331-354

ALLEN TD, Potten CS (1974). Fine-structural identification and organization of the epidermal proliferative unit, *Journal of Cell Science* 15: 291-319

ALY R, Bibel DJ (1993). Adherence of skin micro-organisms and the development of skin flora from birth, in *The Skin Microflora and Microbial Skin Disease*, ed. Noble WC, Cambridge University Press, Cambridge p 355-372

ALY R, Shinefield HI, Strauss WG, Maibach HI (1977). Bacterial adherence to nasal mucosal cells, *Infection and Immunity* 17: 546-549

AMER M, Mostafa FF, Tosson Z, Nasr AN (1996). Corneocytes in scaly parakeratotic diseases, *International Journal of Dermatology* 35: 417-421

BAKER BB, Maibach HI, Park RD, McFarland LZ, O'Brien TR (1973). Epidermal cell renewal in the dog, *American Journal of Veterinary Research* 34: 93-94

BARTELT MA, Duncan JL (1978). Adherence of group A streptococci to human epithelial cells, *Infection and Immunity* 20: 200-208

BARTHELSON R, Hopkins C, Mobasseri A (1999). Quantitation of bacterial adherence by image analysis, *Journal of Microbiological Methods* 38: 17-23

BEACHEY EH (1981). Bacterial adherence: adhesin-receptor interactions mediating the attachment of bacteria to mucosal surface, *Journal of Infectious Diseases* 143: 325-345

and the second second

in survey

Ş

Contract of the Contract of th

BERG JN, Wendell DE, Vogelweid C, Fales WH (1984). Identification of the major coagulase-positive *Staphylococcus* spp. of dogs as *Staphylococcus intermedius*, *American Journal of Veterinary Research* 45: 1307

BIBEL DJ (1982). Bacterial interference, bacteriotherapy and bacterioprophylaxis, in *Bacterial Interference*, ed. Aly R and Shinefield HR, CRC Press, Boca Raton, Florida p 1-12

BIBEL DJ, Aly R, Lahti L, Shinefield HR, Maibach HI (1987). Microbial adherence to vulvar epithelial cells, *Journal of Medical Microbiology* 23: 75-82

BIBEL DJ, Aly R, Shinefield HR, Maibach HI, Strauss WG (1982). Importance of the keratinized epithelial cell in bacterial adherence, *Journal of Investigative Dermatology* 79: 250-253

BIBEL DJ, Lebrun JR (1975). Changes in cutaneous flora after wet occlusion, *Canandian Journal of Microbiology* 21: 496-500

BIBERSTEIN EL, Jang SS, Hirsh DC (1984). Species distribution of coagulase-positive staphylococci in animals, *Journal of Clinical Microbiology* 19: 610-615

BLAIR C (1968). Morphology and thickness of the human stratum corneum, *British Journal of Dermatology* 80: 430-436

BLAND JM, Altman DG (1986). Statistical methods for assessing agreement between two methods of clinical measurement, *Lancet* 1: 307-310

BLAND JM, Altman DG (1999). Measuring agreement in method comparison studies, *Statistical Methods in Medical Research* 8: 135-160

BURKETT G, Frank LA (1998). Comparison of production of Staphylococcus intermedius exotoxin among clinically normal dogs, atopic dogs with recurrent pyoderma, and dogs with a single episode of pyoderma, *Journal of the American Veterinary Medical Association* 213: 232-234

CHARBONEAU D, Rubins JB (1998). Interaction of Bacteria and Their Products with Tissue Culture Cells, in *Methods in Microbiology. Volume 27. Bacterial Pathogenesis*, ed. Williams P, Ketley J, and Salmond G, Academic Press, San Diago p 113-122

CHESNEY CJ (1997). The microclimate of the canine coat: The effect of heating on coat and skin temperature and relative humidity, *Veterinary Dermatology* 8: 183-190

a a a comparte de sector a comparte de secondo de secondo de secondo de secondo de secondo de secondo de second

CHO SH, Strickland I, Boguniewicz M, Leung DY (2001). Fibronectin and fibrinogen contribute to the enhanced binding of *Staphylococcus aureus* to atopic skin, *Journal of Allergy and Clinical Immunology* 108: 269-274

CHRISTOPHERS E (1971). Cellular architecture of the stratum corneum, *Journal of Investigative Dermatology* 56: 165-169

CHRISTOPHERS E, Kligman AM (1964). Visualization of the cell layers of the stratum corneum, *Journal of Investigative Dermatology* 42: 407-409

COLE GW, Silverberg NL (1986). The adherence of *Staphylococcus aureus* to human corneocytes, *Archives of Dermatology* 122: 166-169

COOPER MD, Jeffery-Wiseman C, Bortner CA, Wheat LJ, Wadstrom T (1990). *Pathogenesis of wound and biomaterial-associated infections*, Springer-Verlag, London p 397-407

CORCUFF P, Lotte C, Rougier A, Maibach HI (1991). Racial differences in corneocytes. A comparison between black, white and oriental skin, *Acta Dermato-venereologica.* 71: 146-148

COX HU, Hoskins JD, Newman SS, Foil CS, Turnwald GH, Roy AF (1988). Temporal study of staphylococcal species on healthy dogs, *American Journal of Veterinary Research* 49: 747-751

COX HU, Hoskins JD, Newman SS, Turnwald GH, Foil CS, Roy AF, Kearney MT (1985). Distribution of staphylococcal species on clinically healthy cats, *American Journal of Veterinary Research* 46: 1824-1828

COX HU, Hoskins JD, Roy AF, Newman SS, Luther DG (1984). Antimicrobial susceptibility of coagulase-positive staphylococci isolated from Louisiana dogs, *American Journal of Veterinary Research* 45: 2039-2042

COX HU, Newman SS, Roy AF, Hoskins JD (1984). Species of Staphylococcus isolated from animal infections, *The Cornell Veterinarian* 74: 124-135

CREE RG, Noble WC (1995). In vitro indices of tissue adherence in *Staphylococcus intermedius*, *Letters in Applied Microbiology* 20: 168-170

DAHL MV (1983). *Staphylococcus aureus* and atopic dermatitis, *Archives of Dermatology* 119: 840-846

DE PAEPE K, Janssens K, Hachem JP, Roseeuw D, Rogiers V (2001). Squamometry as a screening method for the evaluation of hydrating products, *Skin Research and Technology* 7: 184-192 $\sim 1.5 \, {\rm eV}_{\rm e}^{\rm A}$ to be the results of setting the state $\sim 1.5 \, {\rm eV}_{\rm e}^{\rm A}$ to be the set of the se

DEVRIESE LA, Nzuambe D, Godard C (1984). Identification and characterization of staphylococci isolated from cats, *Veterinary Microbiology* 9: 279-285

DUNCAN-HEWITT W (1990). Nature of the hydrophobic effect, in *Microbial cell surface hydrophobicity*, ed. Doyle RJ and M.Rosenberg, American society of microbiology, Washington p 39-73

DYCE KM (1987). Textbook of Veterinary Anatomy, W.B. Saunders Co., Philadelphia

EL GAMMAL C, Kligman AM, el Gammal S (1997). Anatomy of the skin surface, in *Bioengneering of the skin. Skin surface imaging and analysis*, ed. Wilhelm K-P, Elsner P, Berardesca E, and Maibach HI, CRC Press, Washington, D.C. p 3-37

EL GAMMAL C, Pagnoni A, Kligman AM, el Gammal S (1996). A model to assess the efficacy of moisturizers--the quantification of soap-induced xerosis by image analysis of adhesive-coated discs (D-Squames), *Clinical and Experimental Dermatology* 21: 338-343

ESPINOLA MB, Lilenbaum W (1996). Prevalence of bacteria in the conjunctival sac and on the eyelid margin of clinically normal cats, *Journal of Small Animal Practice* 37: 364-366

EVANS HE, Christensen GC (1993). *Miller's Anatomy of the Dog*, W. B. Saunders Co., Philadelphia p 98-121

EVANS-HURRELL JA, Adler J, Denyer S, Rogers TG, Williams P (1993). A method for the enumeration of bacterial adhesion to epithelial cells using image analysis, *FEMS Microbiology Letters* 107: 77-82

FEINGOLD DS (1986). Bacterial adherence, colonization, and pathogenicity, *Archives of Dermatology* **122**: 161-163

FLUHR JW, Kao J, Jain M, Ahn SK, Feingold KR, Elias PM (2001). Generation of free fatty acids from phospholipids regulates stratum corneum acidification and integrity, *Journal of Investigative Dermatology* 117: 44-51

FORSYTHE PJ, Hill PB, Thoday KL, Brown J (2002). Use of computerized image analysis to quantify staphylococcal adhesion to canine corneocytes: does breed and body site have any relevance to the pathogenesis of pyoderma?, *Veterinary Dermatology* 13: 29-36

FOSTER TJ, Höök M (1998). Surface protein adhesins of *Staphylococcus* aureus, *Trends in Microbiology* 6: 484-488

i,

1.00

140

FRETER R (1978). Association of enterotoxigenic bacteria with the mucosa of the small intestine-mechanisms and pathogenic implications, in *Cholera and Related Diarrhea: Molecular Aspects of a Global Health Problem.*, ed. Ouchterlony O. and Holmgren J., S. Karger, Basel p 155-170

FRETER R (1980). Prospects for preventing the association of harmful bacteria with host mucosal surfaces, in *Bacterial Adherence (Receptors and Recognition, Vol. 6)*, ed. Beachey EH, Chapman and Hall, London p 439-458

FRETER R, O'Brien PCM, Halstead SA (1978). Adhesion and chemotaxis as determinants of bacterial association with mucosal surfaces, in *Secretory Immunity and Infection*, ed. McGhee JR, Mestecky J, and Babb JL, Plenum Press, New York p 429-438

GARTHWAITE G, Lloyd DH, Thomsett LR (1983). Location of immunoglobulins and complement (C3) at the surface and within the skin of dogs, *Journal of Comparative Pathology* 93: 185-193

GIBBONS RJ, Houte JV (1971). Selective bacterial adherence to oral epithelial surfaces and its role as an ecological determinant, *Infection and Immunity* 3: 567-573

GIBBONS RJ, Houte JV (1975). Bacterial adherence in oral microbial ecology, *Annual Review of Microbiology* 29: 19-44

GIBBONS RJ, Spinell DM, Skobe Z (1976). Selective adherence as a determinant of the host tropisms of certain indigenous and pathogenic bacteria, *Infection and Immunity* 13: 238-246

GOERTTLER K, Reuter M, Stahmer HE (1973). [Morphologic investigations of the proliferation kinetics of the mouse epidermis. Standardized methods and results (author's transl)], *Zeitschrift fur Zellforschung und mikroskopische Anatomie* 142: 131-146

GOLDSCHMIDT H, Kligman AM (1967a). Desquamation of the human horny layer, *Archives of Dermatology* 95: 583-586

GOLDSCHMIDT H, Kligman AM (1967b). Exfoliative cytology of human horny layer. Methods of cell removal and microscopic techniques, *Archives of Dermatology* 96: 572-576

GRANT DI (1991). Skin diseases in the dog and cat, Blackwell Scientific Publications,

GREENE C, McDevitt D, Francois P, Vaudaux PE, Lew DP, Foster TJ (1995). Adhesion properties of mutants of *Staphylococcus aureus* defective in fibronectin-binding proteins and studies on the expression of fnb genes, *Molecular Microbiology* 17: 1143-1152

GREENE RT, Lammler C (1993). *Staphylococcus intermedius*: current knowledge on a pathogen of veterinary importance, *Zentralbl Veterinarmed B* 40: 206-214

GRONE A (2002). Keratinocytes and cytokines, Veterinary Immunology and Immunopathology 88: 1-12

HAIDL G, Plewig G (1988). Exfoliative cytology of stratum corneum and the effects of topical retinoids on the physical properties of corneocytes, *Journal of the Society of Cosmetic Chemists* 39: 53

HALLIWELL REW, Schwartzman RM (1971). Atopic disease in the dog, Veterinary Record 89: 209-213

HANIFIN JM, Homburger HA (1986). Staphylococcal colonization, infection, and atopic dermatitis-- association not etiology, *Journal of Allergy and Clinical Immunology* 78: 563-566

HARVEY RG (1993). Distribution of propionibacteria on dogs: A preliminary report of the findings on 11 dogs, *Journal of Small Animal Practice* 34: 80-84

HARVEY RG (1994). Nasal carriage of *Staphylococcus intermedius* in humans in contact with dogs, *Microbial Ecology in Health and Disease* 7: 225

HARVEY RG, Lloyd DH (1994). The distribution of *Staphylococcus intermedius* and coagulase-negative staphylococci on the hair, skin surface, within the hair follicles and on the mucous membranes of dogs, *Veterinary Dermatology* 5: 75-81

HARVEY RG, Lloyd DH (1995). The distribution of bacteria (other than staphylococci and *Propionibacterium acnes*) on the hair. at the skin surface and within the hair follicles of dogs, *Veterinary Dermatology* 6: 79

HARVEY RG, Noble WC (1994). A temporal study comparing the carriage of *Staphylococcus intermedius* on normal dogs with atopic dogs in clinical remission, *Veterinary Dermatology* 5: 21-25

HASHIMOTO K, Kanzaki T (1975). Surface ultrastructure of human skin, *Acta Dermato-venereologica*. 55: 413-430

HASTY DL, Ofek I, Courtney HS, Doyle RJ (1992). Multiple adhesins of streptococci, *Infection and Immunity* 60: 2147-2152

HERRMANN S, Scheuber E, Plewig G (1983). *Exfoliative cytology: effects of the seasons, in Stratum Corneum.*, Springer, Berlin p 181

HIGGINS R, Gottchalk M (1991). Isolation of *Staphylococcus felis* from cases of external otitis in cats, *Canadian Veterinary Journal* 32: 312-313

HILL PB, Moriello KA (1994). Canine pyoderma, *Journal of the American Veterinary Medical Association* 204: 334-340

HILL S, Edwards C (2002). A comparison of the effects of bath additives on the barrier function of skin in normal volunteer subjects, *The Journal of Dermatological Treatment* 13: 15-18

HOLLAND KT, Roberts CD (1974). A technique for sampling microorganisms from the pilo-sebaceous ducts, *Journal of Applied Bacteriology* 37: 289-296

HOLZLE E, Plewig G (1977). Effects of dermatitis, stripping, and steroids on the morphology of corneocytes. A new bioassay., *Journal of Investigative Dermatology* 68: 350

IGIMI S, Kawamura S, Takahashi E, Mmitsuoka AJ (1989). Staphylococcus felis, a new species from clinical specimens from cats, International Journal of Systematic Bacteriology 39: 373-377

IHRKE PJ (1987). An overview of bacterial skin disease in the dog, *The British Veterinary Journal* 143: 112-118

IHRKE PJ, Franti CE (1985). Breed as a risk factor Associated with skin diseases in dogs seen in Northern California, *California Veterinarian* 39: 13-28

IHRKE PJ, Halliwell REW, Deubler MJ (1977). Canine pyoderma, in *Current veterinary therapy VI. Small animal practice*, ed. Kirk RW, WB Sauders Co, Philadelphia p 513-519

IHRKE PJ, Schwartzman RM, McGinley K, Horwitz LN, Marples RR (1978). Microbiology of normal and seborrheic canine skin, *American Journal of Veterinary Research* 39: 1487-1489

IHRKE PJ, Stannard AA, Ardans AA, Griffin CE (1985). Pemphigus foliaceus in dogs: a review of 37 cases, *Journal of the American Veterinary Medical Association* 186: 59-66

- a No o M Montol Market

JEMEC GB, Serup J (1992). Scaling, dry skin and gender. A bioengineering study of dry skin, *Acta Derm Venereol.Suppl (Stockh)* 177: 26-28

JOHANSSON A, Flock JI, Svensson O (2001). Collagen and fibronectin binding in experimental staphylococcal osteomyelitis, *Clinical Orthopaedics* 1: 241-246

JONES ME, Karlowsky JA, Draghi DC, Thornsberry C, Sahm DF, Nathwani D (2003). Epidemiology and antibiotic susceptibility of bacteria causing skin and soft tissue infections in the USA and Europe: a guide to appropriate antimicrobial therapy, *International Journal of Antimicrobial Agents* 22: 406-419

JONES ME, Schmitz FJ, Fluit AC, Acar J, Gupta R, Verhoef J (1999). Frequency of occurrence and antimicrobial susceptibility of bacterial pathogens associated with skin and soft tissue infections during 1997 from an International Surveillance Programme. SENTRY Participants Group, *European Journal of Clinical Microbiology & Infectious Diseases* 18: 403-408

JORDENS JZ (1990). *Pathogenesis of wound and biomaterial-associated infections*, Springer-Verlag, London p 325-327

KANZAKI H, Morishita Y, Akiyama H, Arata J (1996). Adhesion of *Staphylococcus aureus* to horny layer: role of fibrinogen, *Journal of Dermatological Science* 12: 132-139

KASHIBUCHI N, Hirai Y, O'Goshi K, Tagami H (2002). Three-dimensional analyses of individual corneocytes with atomic force microscope: morphological changes related to age, location and to the pathologic skin conditions, *Skin Research and Technology* 8: 203-211

KATZ HI, Prawer SE, Hien NT, Mooney JJ (1985). Skin-surface touch print for diagnosing fungal infections, *American Family Physician* 31: 189-194

KAWAKAMI S, Callicott RH, Zhang N (1998). Trace analysis of benzalkonium chloride on skin by flow injection ionspray mass spectrometry-mass spectrometry, *Analyst* 123: 489-491

KEDDIE F, Sakai D (1965). Morphology of the horny cells of the superficial stratum corneum: Cell membrances and melanin granules, *Journal of Investigative Dermatology* 44: 135-138

KEDDIE F, Shadomy J, Shadomy S (1963). Intrafollicular tinea versicolor demonstrated on monomer plastic strips, *Journal of Investigative Dermatology* 41: 103-106

KRAL F, Schwartzman RM (1964). *Veterinary and Comparative Dermatology*, J. B. Lippincott Co., Philadelphia

KRISTENSEN S, Krogh HV (1978). A study of skin diseases in dogs and cats. III. III. Microflora of the skin of dogs with chronic eczema, *Nordisk Veterinaermedicin* 30: 223-230

KROGH HV, Kristensen S (1976). A study of skin diseases in dogs and cats. II. Microflora of the normal skin of dogs and cats, *Nordisk Veterinaermedicin* 28: 459-463

KROGH HV, Kristensen S (1981). A study of skin diseases in dogs and cats. VI. Microflora of the major canine pyodermas, *Nordisk Veterinaermedicin* 33: 17-22

KWOCHKA KW, Rademakers AM (1989). Cell proliferation of epidermis, hair follicles, and sebaceous glands of beagles and cocker spaniels with healthy skin, *American Journal of Veterinary Research* 50: 587-591

LAI A FAT RF, Van Furth R (1974). The role of the skin & adjacent mucous membranes in host defense, *Journal of Cutaneous Pathology* 1: 61-70

LEE SH, Lee S (1986). Regional and sexual differences in corneocytes among young Korean adults, *Yonsei Medical Journal* 27: 213-218

LEIGH IM (1994). *The keratinocyte Handbook*, Cambridge University Press, New York

LEVER WF, Schaumburg-Lever G (1990). *Histopathology of the Skin*, Lippincott Co., Philadelphia

LEYDEN JJ, Marples RR, Kligman AM (1974). *Staphylococcus aureus* in the lesions of atopic dermatitis, *British Journal of Dermatology* 90: 525-530

LLOYD DH (1976). The cutaneous defence mechanisms, *Veterinary Dermatology Newsletter* 1: 9

LLOYD DH (1980a). Skin surface immunity, Veterinary Dermatology Newsletter 5: 10

LLOYD DH (1980b)The inhabitants of the mammalian skin surface. Proceedings of the Royal Society of Edinburgh. P25-42 المحاطية والمراجع المحسر مسر

والمراجع والمعادية مستنا والقوافين والمرا

LLOYD DH (1995) The skin as a microbial habitat. Proceedings of the 12th annual congress of the European Society of Veterinary Dermatology. Barcelona. p60-63

LLOYD DH, Allaker RP (1991). Carriage of *Staphylococcus intermedius* on the ventral abdomen of clinical normal dogs and those with pyoderma, *Veterinary Dermatology* 2: 161-164

LLOYD DH, Dick WD, Jenkinson DM (1979). Structure of the epidermis in Ayrshire bullocks, *Research in Veterinary Science* 26: 172-179

LLOYD DH, Garthwaite G (1982). Epidermal structure and surface topography of canine skin, *Research in Veterinary Science* 33: 99-104

LLOYD DH, Jenkinson DM (1980). The effect of climate on experimental infection of bovine skin with *Dermatophilus congolensis*, *The British Veterinary Journal* 136: 122-134

LLOYD DH, Sasiak AB, Kitson S, McEwan JD, Elder HY (1993). Temporal changes in the granulocytic responses to experimental infection of the skin of mice and sheep with *Dermatophilus congolensis*, *Revue d'elevage et de medecine veterinaire des pays tropicaux* 46: 271-276

LOOKINGBILL DP (1985). Principles of clinical diagnosis, in *Dermatology*, ed. Moschella S.L. and Hurley H.J., WB Saunders Co, Philadelphia p 126-200

LOVELL JE, Getty R (1957). The hair follicle, epidermis, dermis, and skin glands of the dog, *American Journal of Veterinary Research* 18: 873

LOWY FD (1998). Staphylococcus aureus infections, New England Journal of Medicine 339: 520-532

LUND EM, Armstrong PJ, Kirk CA, Kolar LM, Klausner JS (1999). Health status and population characteristics of dogs and cats examined at private veterinary practices in the United States, *Journal of the American Veterinary Medical Association* 214: 1336-1341

LYNE AG, Short BF (1965). *Biology of the Skin and Hair Growth*, American Elsevier Publishing Co., New York

MACKENZIE IC (1969). Ordered structure of the stratum corneum of the mammalian skin, *Nature* 222: 881-883

;

a Warner and a sur-

2

÷.,

a the second strained that we are a second

ł

MACKENZIE IC (1972). The ordered structure of mammalian skin, Epidermal Wound Healing, ed. H maibach and D Rovee, Yearbook Medical Publishers, Chicago p 5-25

MACKENZIE IC, Linder JE (1973). An examination of cellular organization within the stratum corneum by a silver staining method, *Journal of Investigative Dermatology* 61: 245-250

MAKKI S, Barbenel JC, Agache P (1979). A quantitative method for the assessment of the microtopography of human skin, *Acta dermato-venere* 59: 285-291

MARKS R (1972). Histochemical applications of skin surface biopsy, *British Journal of Dermatology* 20-26

MARKS R (1980). Epidermal and corneocyte size changes with age, *British Journal of Dermatology* 102: 738

MARKS R, Barton S (1983). The significance of the size and shape of corneocytes, Springer-Verlag, Berlin p 161

MARKS R, Dawber RPR (1971). Skin surface biopsy: A improved technique for the examination of the horny layer, *British Journal of Dermatology* 117-123

MARKS R, Dawber RPR (1972). In situ microbiology of the stratum corneum: An application of skin surface biopsy, *Archives of Dermatology* 105: 216-221

MARPLES RR (1965). The effect of hydration on the bacterial flora of the skin, in *Skin Bacteria and their Role in Infection*, ed. Maibach HI and Kildick-Smith G, McGraw-Hill, New York

MASENGA J, Garbe C, Wagner J, Orfanos CE (1990). *Staphylococcus aureus* in atopic dermatitis and in nonatopic dermatitis, *International Journal of Dermatology* 29: 579-582

MASON IS, Mason KV, Lloyd DH (1996). A review of the biology of canine skin with respect to the commensals *Staphylococcus intermedius*, *Demodex canis*, and *Malassezia pachydermatitis*, *Veterinary Dermatology* 7: 119-132

MASON IS (1999) Pathogenesis of canine pyoderma. Proceeding of the British Veterinary Dermatology Study Group. P37

MASON IS, Lloyd DH (1989). The role of allergy in the development of canine pyoderma, *Journal of Small Animal Practice* 30: 216-218

MASON IS, Lloyd DH (1990). Factors influencing the penetration of bacterial antigens through canine skin, in *Advances in Veterinary Dermatology*, ed. von Tscharner C and Halliwell REW, Bailliere Tindall, London p 370-374

MASON IS, Lloyd DH (1993). Scanning electron microscopic studies of the living epidermis and stratum corneum in dogs, in *Advances in Veterinary Dermatology*, ed. Ihrke PJ, Mason IS, and White SD, Oxford p 131

MASON KV, Stewart LJ (1993). *Malassezia* and canine dermatitis, in *Advances in Veterinary Dermatology*, ed. Ihrke PJ, Mason IS, and White SD, p 399-402

MCBRIDE ME (1993). Physical factors affecting the skin flora and skin desease, in *The Skin Microflora and Microbial Skin Disease*, ed. Noble WC, Cambridge University Press, Cambridge p 73-101

MCBRIDE ME, Duncan WC, Knox JM (1977). The environment and the microbial ecology of human skin, *Applied and Environmental Microbiology* 33: 603-608

MCDONALD J, Kunkle G (1983). Skin disease of selected dog breeds, *Proceedings of American Animal Hospital Association* 50: 133-135

MCEWAN JENKINSON D. (1993). The basis of the skin surface ecosystem, in *The Skin Microflora and Microbial Skin Disease*, ed. Noble WC, Cambridge University Press, Cambridge p 1-32

MCEWAN NA (1990). Bacterial adherence to canine corneocytes, in *Advances in Veterinary Dermatology*, ed. von Tscharner C and Halliwell REW, Bailliaere-Tindall, London p 454

MCEWAN NA (2000). Adherence by *Staphylococcus intermedius* to canine keratinocytes in atopic dermatitis, *Research in Veterinary Science* 68: 279-283

MCEWAN, NA (2002a) An Epidemiological study of canine skin disease with special reference to atopic dermatitis and lethal acrodermatitis. Glasgow, DVM 38-80.

MCEWAN, NA (2002b) An Epidemiological study of canine skin disease with special reference to atopic dermatitis and lethal acrodermatitis. Glasgow, DVM 45.

MCEWAN, NA (2002c) An Epidemiological study of canine skin disease with special reference to atopic dermatitis and lethal acrodermatitis. Glasgow, DVM .

÷

2

.

MCGINLEY KJ, Leyden JJ, Marples RR, Kligman AM (1975). Quantitative microbiology of the scalp in non-dandruff, dandruff, and seborrheic dermatitis, *Journal of Investigative Dermatology* 64: 401-405

MEDLEAU L, Blue JL (1988). Frequency and antimicrobial susceptibility of Staphylococcus spp isolated from feline skin lesions, *Journal of American Veterinary Medical Association* 193: 1080-1081

MEDLEAU L, Long RE, Brown J, Miller WH (1986). Frequency and antimicrobial susceptibility of Staphylococcus species isolated from canine pyodermas, *American Journal of Veterinary Research* 47: 229-231

MENTON DN, Eisen AZ (1971a). Structural organization of the stratum corneum in certain scaling disorders of the skin, *Journal of Investigative Dermatology* 57: 295-307

MENTON DN, Eisen AZ (1971b). Structure and organization of mammalian stratum corneum, *Journal of Ultrastructure Research* 35: 247-264

MIHARA M (1997). Scanning electron microscopy, in *Bioengineering of the skin-Skin surface imaging and analysis*, ed. Wilhelm K-P, Elsner P, Berardsca E, and Maibach HI, CRC Press, London p 73-83

MILLS OH, Jr., Kligman AM (1983). The follicular biopsy, *Dermatologica* 167: 57-63

MIYAKE Y, Kohada A, Fujii I, Sugai M, Suginaka H (1989). Aminoglycosides enhance the adherence of Staphylococcus aureus to HeLa cells, *The Journal of Antimicrobial Chemotherapy* 23: 79-86

MIYAKE Y, Sugai M, Kohada A, Minagi S, Suginaka H (1990). Adherence of Staphylococcus aureus to cultured epidermal cells during differentiation, *Journal of Medical Microbiology* 32: 9-14

MONTAGNA W (1967). Comparative anatomy and physiology of the skin, Archives of Dermatology 96: 357

MORIELLO KA, Mason IS (1995). *Handbook of Small Animal Dermatology*, Pergamon Veterinary,

MOURICOUT M, Petit JM, Carias JR, Julien R (1990). Glycoprotein glycans that inhibit adhesion of Escherichia coli mediated by K99 fimbriae: treatment of experimental colibacillosis, *Infection and Immunity* 58: 98-106

MURPHY CT (1975). Nutrient materials and the growth of bacteria on human skin, *Transactions of the St.John's Hospital Dermatological Society* 61: 51-57

NAGATA M, Sakai T (1999). Clinical survey of canine dermatosis in Japan, *Journal of the Japanese Veterinary Medical Association* 52: 775-779

NESBITT GH (1983). Canine and feline dermatology: a systematic approach, Lea & Febiger, Philadelphia

OFEK I, Beachey EH (1980). Bacterial adherence, *Advances in Internal Medicine* 25: 503-532

OFEK I, Doyle RJ (1994). *Bacterial adhesion to cells and tissues*, Chapman & Hall, New York p ix, 578

PATEL, A (2002) A study on feline staphylococci in England. Dissertation for the RCVS Diploma in Veterinary Dermatology.

PATEL A (2003) Feline Bacterial Skin Disease. Proceedings. Spring. British Veterinary Dermatology Study Group. p5-10

PATEL A, Lloyd DH, Howell SA, Noble WC (2002). Investigation into the potential pathogenicity of Staphylococcus fells in a cat, *Veterinary Record* 150: 668-669

PEACOCK SJ, Day NP, Thomas MG, Berendt AR, Foster TJ (2000). Clinical isolates of *Staphylococcus aureus* exhibit diversity in fnb genes and adhesion to human fibronectin, *Journal of Infectious Diseases* 41: 23-31

PHILLIPS WE, Jr., Kloos WE (1981). Identification of coagulase-positive *Staphylococcus intermedius* and *Staphylococcus hylcus* subsp. *hylcus* isolates from veterinary clinical specimens, *Journal of Clinical Microbiology* 14: 671-673

PIERARD GE, Pierard-Franchimont C (1993). Dihydroxyacetone test as a substitute for the dansyl chloride test, *Dermatology* 186: 133-137

PIMENTEL GC, McClellan AL (1960). *The Hydrogen Bond*, W.H. Freeman, San Francisco

PINCUS H (1951). Examination of the epidermis by the strip method of removing the horny layer, *Journal of Investigative Dermatology* 16: 383-386

PLEWIG G (1970). Regional differences of cell sizes in the human stratum corneum. II. Effects of sex and age, *Journal of Investigative Dermatology* 54: 19-23

PLEWIG G, Marples RR (1970). Regional differences of cell sizes in the human stratum corneum. I, *Journal of Investigative Dermatology* 54: 13-18

PRICE PB (1938). The bacteriology of normal skin: A new quantitative test applied to a study of the bacterial flora and the disinfectant action of mechanical cleansing, *Journal of Infectious Diseases* 63: 301-318

PRIESTLEY GC (1993). *Molecular Aspects of Dermatology*, John Wiley and Sons, New York

RAYCHAUDHURI SP, Raychaudhuri SK (1993). Relationship between kinetics of lesional cytokines and secondary infection in inflammatory skin disorders: a hypothesis, *International Journal of Dermatology* 32: 409-412

ROMERO-STEINER S, Witek T, Balish E (1990). Adherence of skin bacteria to human epithelial cells, *Journal of Clinical Microbiology* 28: 27-31

ROTH RR, James WD (1989). Microbiology of the skin: resident flora, ecology, infection, *Journal of the American Academy of Dermatology* 20: 367-390

SADER HS, Pfaller MA, Jones RN, Doern GV, Gales AC, Winokur PL, Kugler KC (1999). Bacterial Pathogens Isolated from Patients with Bloodstream Infections in Latin America, 1997: Frequency of Occurrence and Antimicrobial Susceptibility Patterns from the SENTRY Antimicrobial Surveillance Program, *The Brazilian Journal of Infectious Diseases* 3: 97-110

SAIJONMAA-KOULUMIES LE, Lloyd DH (1995). Carriage of bacterial antagonists towards *Staphylococcus intermedius* on canine skin and mucosal surfaces, *Veterinary Dermatology* 6: 187

SAIJONMAA-KOULUMIES LE, Lloyd DH (1996). Colonisation of canine skin with bacteria, *Veterinary Dermatology* 7: 153-162

SAIJONMAA-KOULUMIES LE, Lloyd DH (2002). Adherence of *Staphylococcus intermedius* to canine corneocytes *in vitro*, Veterinary Dermatology 13: 169-176

SARKANY I (1962). A method for studying the microtopography of the skin, *British Journal of Dermatology* 254-259

SCHÄTZLEIN A, Cevc G (1998). Non-uniform cellular packing of the stratum corneum and permeability barrier function of intact skin: a high-resolution confocal laser scanning microscopy study using highly deformable vesicles (Transfersomes), *British Journal of Dermatology* 138: 583-592

SCHUMMER A (1981). The Circulatory System, the Skin, and the Cutaneous Organs of the Domestic Mammals, Verlag Paul Parey, Berlin

SCHWARTZMAN RM, Orkin M (1962). A comparative study of skin diseases of dog and Man, Charles C Thomas, Springfield, CT

SCHWARZ R (1979). Micromorphology of the skin (epidermis, dermis, subcutis) of the dog, *The Onderstepoort Journal of Veterinary Research* 46: 105

SCOTT DW (1980). Feline dermatology 1900-1978, *Journal of American* Animal Hospital Association 16: 331-332

SCOTT DW (1981). Observations on canine atopy, *Journal of American* Animal Hospital Association 17: 91-100

SCOTT DW (1988). *Large animal dermatology*, W. B. Saunders, Philadelphia p 7-8

SCOTT DW (1992). Bacteria and yeast on the surface and within noninflamed hair follicles of skin biopsies from dogs with non-neoplastic dermatoses, *The Cornell Veterinarian* 82: 379-386

SCOTT DW, Miller WH, Griffin CE (2001a). Structure and function of the skin, in *Muller & Kirk's Small Animal Dermatology*, 6th Edition, ed. Scott DW, Miller WH, and Griffin CE, W.B. Saunders, Philadelphia p 1-70

SCOTT DW, Miller WHJr, Griffin CE (2001b). *Muller and Kirk's Small* Animal Dermatology, 6th Edition, W. B. Saunders Co., Philadelphia p 275-277

SCOTT DW, Paradis M (1990). A survey of canine and feline skin disorders seen in a university practice: Small animal clinic, University of Montreal, Saint-Hyacinthe, Qubec, *Canadian Veterinary Journal* 31: 830-835

SCOTT DW, Walton DK, Slater MR, Smith CA, Lewis RM (1987a). Immune-mediated dermatoses in domestic animals: Ten years after- part I, *Compendium on Continuing Education for the Practising Veterinarian* 9: 423-437 SCOTT DW, Walton DK, Slater MR, Smith CA, Lewis RM (1987b). Immune-mediated dermatoses in domestic animals: Ten years after- part II, *Compendium on Continuing Education for the Practising Veterinarian* 9: 539-554

SCOTT SW (1984). Feline dermatology 1979-1982: Introspective retrospections, *Journal of American Animal Hospital Association* 20: 537

SELLWOOD R, Gibbons RA, Jones GW, Rutter JM (1975). Adhesion of enteropathogenic Escherichia coli to pig intestinal brush borders: the existence of two pig phenotypes, *Journal of Medical Microbiology* 8: 405-411

SERUP J (1992a). A double-blind comparison of two creams containing urea as the active ingredient. Assessment of efficacy and side-effects by non-invasive techniques and a clinical scoring scheme, *Acta Derm Venereol.Suppl (Stockh)* 177: 34-43

SERUP J (1992b). A three-hour test for rapid comparison of effects of moisturizers and active constituents (urea). Measurement of hydration, scaling and skin surface lipidization by noninvasive techniques, *Acta Derm Venereol.Suppl (Stockh)* 177: 29-33

SERUP J, Winther A, Blichmann C (1989). A simple method for the study of scale pattern and effects of a moisturizer--qualitative and quantitative evaluation by D-Squame tape compared with parameters of epidermal hydration, *Clinical and Experimental Dermatology* 14: 277-282

SISCHO WM, Ihrke PJ, Franti CE (1989). Regional distribution of ten common skin diseases in dogs, *Journal of the American Veterinary Medical Association* 195: 752-756

SOMERVILLE-MILLAR DA, Noble WC (1974). Resident and transient bacteria of the skin, *Journal of Cutaneous Pathology* 1: 260-264

STRICKLAND JH, Calhoun ML (1963). The integumentary system of the cat, *American Journal of Veterinary Research* 24: 1018-1029

SUEKI H, Gammal C, Kudoh K, Kligman AM (2000). Hairless guinea pig skin: anatomical basis for studies of cutaneous biology, *European Journal* of *Dermatology* 10: 357-364

SULZBURGER MB (1960). On the biology and function of the skin surface, *Japanese Journal of Dermatology* 70: 105-122

SUTER MM (1987). Monoclonal antibodies: Cell surface markers for canine keratinocytes, *American Journal of Veterinary Research* 40: 1367

SUTER MM, Crameri FM, Olivry T, Mueller E, von Tscharner C, Jensen PJ (1997). Keratinocyte biology and pathology, *Veterinary Dermatology* 8: 67-100

TAVENDALE A, Jardine CK, Old DC, Duguid JP (1983). Haemagglutinins and adhesion of Salmonella typhimurium to HEp2 and HeLa cells, *Journal of Medical Microbiology* 16: 371-380

THODAY AL, Friedmann PS (1986). Scientific Basis of Dermatology. A Physiological Approach., Churchill Livingstone, New York

THOMSON ML (1953). The identification and enumeration of active sweat glands in man from plastic impressions of the skin, *Transactions of the Royal Society of Tropical Medicine and Hygiene* 47: 412-417

TOYOSHIMA H (1989). Aging of epidermis. I. Morphological changes and their quantitative analysis, *Nippon Hifuka Gakkai zasshi.The Japanese Journal of Dermatology*. 99: 1-7

TRING FC, Murgatroyd LB (1974). Surface microtopography of normal human skin, *Archives of Dermatology* 109: 223-226

TRING FC, Murgatroyd LB (1975). Psoriasis: Changes in surface microtopography., *Archives of Dermatology* 476-480

TZANCK A (1947). Le cytodiagnostic immediat en dermatologie, Bulletin de la Societe francaise de dermatologie et de syphiligraphie 7: 68

VALENTIN-WEIGAND P, Chhatwal GS, Blobel H (1987). A simple method for quantitative determination of bacterial adherence to human and animal epithelial cells, *Microbiology and Immunology* 31: 1017-1023

VALENTIN-WEIGAND P, Chhatwal GS, Blobel H (1988). Adherence of streptococcal isolates from cattle and horses to their respective host epithelial cells, *American Journal of Veterinary Research* 49: 1485-1488

VAN BELKUM A, Kools-Sijmons M, Verbrugh H (2002). Attachment of *Staphylococcus aureus* to eukaryotic cells and experimental pitfalls in staphylococcal adherence assays: a critical appraisal, *Journal of Microbiological Methods* 48: 19-42

VAN WAMEL WJ, Vandenbroucke-Grauls CM, Verhoef J, Fluit AC (1998). The effect of culture conditions on the in-vitro adherence of methicillin-resistant *Staphylococcus aureus*, *Journal of Medical Microbiology* 47: 705-709

VERBRUGH HA (1990). Pathogenesis of wound and biomaterialassociated infections, Springer-Verlag, London p 361-365 Number of Advenue

WEBB AJ, Calhoun ML (1954). The microscopic anatomy of the skin of mongrel dogs, *American Journal of Veterinary Research* 15: 274-280

WHITE MI, Noble WC (1980) Skin response to protein A. 1980. Proceedings of the Royal Society of Edinburgh. P43-46

WHITE SD (1991). Pyoderma in five cats, *Journal of the American Animal Hospital Association* 27: 141-146

WHITE SD, Ihrke PJ (1987). Pyoderma, in *Dermatology*, ed. Nesbitt G.H., John Wiley & Sons, New York p 376-381

WHITE SD, Ihrke PJ, Stannard AA, Sousa C, Reinke S, Rosser EJ, Jr., Jang S (1983). Occurrence of Staphylococcus aureus on the clinically normal canine hair coat, *American Journal of Veterinary Research* 44: 332-334

WILKINSON GT (1985). A color altas of small animal dermatology, Williams & Wilkins, Baltimore

WILLEMSE T (1986). BSAVA Education Committee commissioned article. Atopic skin disease: a review and a reconsideration of diagnostic criteria, *Journal of Small Animal Practice* 27: 771-778

WOLDEHIWET Z, Jones JJ (1990). Species distribution of coagulasepositive staphylococci isolated from dogs, *Veterinary Record* 126: 485

WYATT JE, Poston SM, Noble WC (1990). Adherence of *Staphylococcus aureus* to cell monolayers, *Journal of Applied Bacteriology* 69: 834-844

YAGER JA, Scott DW (1993). *The skin and appendages*, Academic Press, New York p 531

YOON HS, Baik SH, Oh CH (2002). Quantitative measurement of desquamation and skin elasticity in diabetic patients, *Skin Research and Technology* 8: 250-254

ľ	GLASCO	
I	UNIVERSITY	:
ļ	UBRARY	

ų,