

https://theses.gla.ac.uk/

Theses Digitisation:

https://www.gla.ac.uk/myglasgow/research/enlighten/theses/digitisation/

This is a digitised version of the original print thesis.

Copyright and moral rights for this work are retained by the author

A copy can be downloaded for personal non-commercial research or study, without prior permission or charge

This work cannot be reproduced or quoted extensively from without first obtaining permission in writing from the author

The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the author

When referring to this work, full bibliographic details including the author, title, awarding institution and date of the thesis must be given

Enlighten: Theses <u>https://theses.gla.ac.uk/</u> research-enlighten@glasgow.ac.uk



FRONTISPIECE A 3-day-old Zea mays seedling

ProQuest Number: 10647023

All rights reserved

INFORMATION TO ALL USERS The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



ProQuest 10647023

Published by ProQuest LLC (2017). Copyright of the Dissertation is held by the Author.

All rights reserved. This work is protected against unauthorized copying under Title 17, United States Code Microform Edition © ProQuest LLC.

> ProQuest LLC. 789 East Eisenhower Parkway P.O. Box 1346 Ann Arbor, MI 48106 – 1346

. ٢

• •

Thesis 6556 Copy 2 ABGOW ONIVERSITY LIBRARY

ABSTRACT

The metabolism of exogenous indole-3-acetic acid (IAA) was examined in root and coleoptile segments taken from dark-grown seedlings of Zea The rate of disappearance of IAA, and the chemical mays L. cv. Fronica, nature of the total spectrum of products in methanolic extracts of plant material which had been incubated with IAA for 2h were investigated. The difference in IAA metabolism in separated cortex plus epidermis, and stelar tissues of the root was monitored. Two methods of supplying IAA to the segments were employed, and the effect of the mode of application on the metabolism pattern noted. The metabolism of IAA, transported through coleoptile and root segments was also studied. In each experiment, the uptake of exogenous IAA was noted and compared with published measurements of endogenous IAA levels. Finally, experiments were carried out to investigate the effect of IAA on coleoptile and root elongation.

Segments were usually supplied with IAA by floating them in aqueous solutions of the radioactively-labelled compound. Alternatively, agar blocks containing ¹⁴C-IAA were placed at the apical ends of coleoptile segments, or on protruding portions of stele at the basal ends of root segments. All incubations were carried out in darkness. Crude methanolic extracts of plant material were analysed using a gradient-elution, reversephase high-performance liquid chromatograph with an on-stream homogeneous radioactivity monitor. Sample purification was kept to a minimum to avoid selective loss of metabolites. Control experiments showed that degradation of IAA did not occur during sample preparation or analysis. Information on the chemical nature of metabolites was obtained from high-performance liquid chromatograph retention times, use of IAA labelled with ¹⁴C at different positions on the molecule, examination of the products of base-catalysed hydrolysis and methylation with diazomethane, UV spectrophotometry, and

i

co-chromatography with standards of IAA derivatives. The identity of the IAA remaining at the end of experiments was confirmed by combined gas chromatography-mass spectrometry.

IAA was metabolised rapidly by both roots and coleoptiles to a large number of products; at least 11 in roots. All metabolites, in methanolic extracts of tissue incubated for 2h, were more polar than IAA and did not involve decarboxylation. The main product in both roots and coleoptiles appeared to be oxindole-3-acetic acid. Other tentative identifications included 5-hydroxyindole-3-acetic acid, indole-3-acetyl glycine and an ester of IAA possibly with a sugar or <u>myo</u>-inositol. The appearance of several products, in similar quantities, after very short incubation times (10 min), indicated that metabolism probably did not take a single, linear pathway.

Experiments using sterile tissues confirmed that metabolism was taking place within the plant cells. At the lowest external concentrations used, the amounts of ¹⁴C-IAA taken into the plant represented only a small fraction of endogenous levels (measurements obtained from published data). Increasing the concentration of IAA over two orders of magnitude did not alter the pattern of metabolism. The products observed were thus unlikely to represent results of a detoxification process.

Comparison of IAA metabolism by cortical and stelar tissues of the root indicated that the majority of metabolism took place in the cortex. IAA present in the stele, the site of the majority of endogenous IAA in the root, appeared to be protected from metabolism.

The mode in which IAA was supplied to roots had a substantial effect on the metabolism pattern. IAA applied in agar blocks to the stele at the basal end of segments was metabolised more slowly than that taken up from solution. The relative proportions of products also differed.

Although substantial metabolism took place in all experiments, radioactivity transported through both root and coleoptile segments, and collected

ii

in agar blocks, remained exclusively associated with the IAA molecule. This provided further evidence for a transport system specific for IAA.

Preliminary experiments were also carried out to study the effects Coleoptile segments incubated in 10^{-2} mol m⁻³ of IAA on segment elongation. On the other hand, IAA $(10^{-3} \text{ mol m}^{-3})$ IAA showed maximal growth stimulation. was shown to inhibit root elongation. Uptake at this concentration was of the same order of magnitude as endogenous levels. IAA at lower concentrations could not be shown to promote growth. The effect of IAA on roots was dependent on the presence of oxygen. Growth experiments were always carried out using lh incubation times, as growth of segments decreased rapidly after this time. The proportion of exogenous IAA which was metabolised also increased with time.

These results have several implications for research on IAA physiology. The complexity of the metabolism pattern observed highlights the need for definitive identification of the total spectrum of metabolites. Cochromatography with standards in thin-layer or paper systems was clearly shown to be inadequate for this purpose. The lack of decarboxylation products calls into question the role of IAA-oxidases in the <u>in vivo</u> metabolism of IAA. The fast rate of metabolism, the substantial effect of mode of IAA application on its metabolism, and the rapid decrease in growth rate of segments, all confound research on the effects, transport and binding of exogenous IAA.

iii

ACKNOWLEDGEMENTS

Thanks are due to the following persons for their help and advice during the compilation of this thesis.

Dr. J. R. Hillman for supervision of research and criticism of the manuscript.

Professor M. B. Wilkins for use of the research facilities in the Botany Department of Glasgow University, and also for helpful advice concerning the project.

Dr. V. B. Math and Mr. R. McGowan carried out GC-MS analyses and synthesised the derivatives of indole-3-acetic acid.

Dr. A. Crozier and Mrs. A. Sutcliffe for assistance with the HPLC analyses.

Mr. J. McMurray for technical assistance.

My father Dr. D. C. Nonhebel and several members of Strathclyde University Chemistry Department for advice and criticism of the chemistry involved in the project.

Miss M. G. Cuthill for typing the thesis.

Mr. T. N. Tait for photography.

The work reported in this thesis was carried out with the support of a scholarship from the Carnegie Trust for the Universities of Scotland.

Finally, my parents are thanked for their moral and financial support during my university career.

ABBREVIATIONS

A.E.S.	Automatic External Standard
BSTFA	N,O-bis-(trimethy1si1y1)trifluoroacetamide
cpm	Counts per minute
<u>et al</u> .	<u>et alia</u>
f.s.d.	Full scale deflection
f.wt.	Fresh weight
GC -MS	Gas chromatography-mass spectrometry
HPLC	High-performance liquid chromatography
IAA	Indole-3-acetic acid
I.D.	Internal diameter
<u>m/e</u>	Mass-to-charge ratio
Me-IAA	Methyl indole-3-acetate
NMR	Nuclear magnetic resonance
ODS	Octadecylsilane
PPO	2,5-diphenyloxazole
R _f	Retention factor
TLC	Thin-layer chromatography
T _R	Retention time
UV	Ultra-violet
v/v	Volume for volume
TMSi	Trimethylsilyl derivative

Where practicable S1 units have been employed, however, for measurement of time, minutes, hours and days were used.

,

ŧ

÷

CONTENTS

	Page No.
Abstract	i
Acknowledgements	iv
Abbreviations	v
Contents	vi
INTRODUCTION	1
MATERIALS AND METHODS	40
RESULTS	57
SECTION 1 - METABOLISM OF IAA	58
A. Comparison of Separatory Techniques Used in the analysis	
of IAA Metabolites in Plant Extracts	
A.1 TLC analysis of IAA metabolites	58
A.2 HPLC analysis of IAA metabolites	61
A.3 Use of HPLC and TLC in a preparative mode	61
B. Quantitative Fate of IAA-2- ¹⁴ C Supplied to Zea mays	
Root Segments	
B.1 Metabolism of IAA-2- ¹⁴ C by root segments	66
during a 2h incubation period	
B.2 Metabolism of IAA by sterile roots	77
B.3 Effect of exposure to light on the metabolism	79
of IAA by Zea mays root segments	
B.4 Effect of varying the IAA concentration in the	79
incubating solution on its metabolism by root segments	
B.5 Use of Sep-pak C ₁₈ cartridges in preparation of	80
samples for HPLC	
C. Time-Course of IAA Metabolism in Root Segments	87
D. A Comparison of the Metabolism of IAA by Cortical and	94
Stelar Tissues of the Root	

.....

METABOLISM OF INDOLE-3-ACETIC

.

ACID IN SEEDLINGS OF ZEA MAYS L.

By

Heather Margaret Nonhebel

Thesis submitted to the University of Glasgow

for the degree of Doctor of Philosophy

Department of Botany University of Glasgow

September 1982

the St	ele at the Basal End of Root Segments	
F Metabo	lism of ¹⁴ C-IAA by Coleontile Segments	
F.1	HPLC analysis of IAA-2- ¹⁴ C metabolites from	
	coleoptile segments after 2h incubation	
F.2	Metabolism of IAA by coleoptile segments exposed	
	to light during sectioning	
F.3	Effect of varying the external concentration of IAA	
	on its metabolism by coleoptile segments	
F.4	Time-course of IAA metabolism by Zea mays	
	coleoptile segments	
G. Metabo	lism of IAA-2- ¹⁴ C Transported through Zea mays	
Coleop	tile Segments	
H. The Ch	emical Nature of IAA Metabolites	
<u>H. The Ch</u> H.1	emical Nature of IAA Metabolites Experiment to investigate the number of radioactive	
H. The Ch H.1	emical Nature of IAA Metabolites Experiment to investigate the number of radioactive compounds represented by each peak	
<u>H. The Ch</u> H.1 H.2	emical Nature of IAA Metabolites Experiment to investigate the number of radioactive compounds represented by each peak Polarity of IAA metabolites	
<u>H. The Ch</u> H.1 H.2 H.3	emical Nature of IAA Metabolites Experiment to investigate the number of radioactive compounds represented by each peak Polarity of IAA metabolites Experiment to investigate whether carbon-1 of the	
<u>H. The Ch</u> H.1 H.2 H.3	emical Nature of IAA Metabolites Experiment to investigate the number of radioactive compounds represented by each peak Polarity of IAA metabolites Experiment to investigate whether carbon-1 of the side chain has been lost	
<u>H. The Ch</u> H.1 H.2 H.3 H.4	emical Nature of IAA Metabolites Experiment to investigate the number of radioactive compounds represented by each peak Polarity of IAA metabolites Experiment to investigate whether carbon-1 of the side chain has been lost UV Spectrometry	
H. The Ch H.1 H.2 H.3 H.4 H.5	emical Nature of IAA Metabolites Experiment to investigate the number of radioactive compounds represented by each peak Polarity of IAA metabolites Experiment to investigate whether carbon-1 of the side chain has been lost UV Spectrometry Methylation of IAA metabolites	
H. The Ch H.1 H.2 H.3 H.4 H.5 H.6	emical Nature of IAA Metabolites Experiment to investigate the number of radioactive compounds represented by each peak Polarity of IAA metabolites Experiment to investigate whether carbon-1 of the side chain has been lost UV Spectrometry Methylation of IAA metabolites Hydrolysis of IAA metabolites	
H. The Ch H.1 H.2 H.3 H.4 H.5 H.6 H.7	emical Nature of IAA Metabolites Experiment to investigate the number of radioactive compounds represented by each peak Polarity of IAA metabolites Experiment to investigate whether carbon-1 of the side chain has been lost UV Spectrometry Methylation of IAA metabolites Hydrolysis of IAA metabolites Co-chromatography of metabolites with standards	

154
154
156
159
159
162
165
167

REFERENCES

Ø......

INTRODUCTION

Intercellular communication is a property of fundamental importance to multicellular organisms. As the physiological and morphological complexity of the organism increases, there is a corresponding requirement for efficient and adaptive coordination mechanisms. The plant or animal must be able to integrate directly the activities of cells which may be considerable distances apart.

Specific Requirements for Intercellular Communication in Higher Plants

The sedentary habit of vascular plants requires them to be adaptable to changing environmental conditions. As they are nearly all autotrophic, plants have relatively simple nutrient requirements. They are, however, dependent on having an adequate supply of water, light and mineral nutrients and must respond to a changing external situation in a way which will ensure that this supply continues. Plants must also be responsive to seasonal changes in temperature and daylength, and should be able to react to adverse circumstances such as allelochemicals and predators in a way which will ensure their survival. For any such reaction to environmental conditions the sensing mechanisms of the plant must be able to communicate with the other organs and tissues.

Intercellular communication is also required to integrate the celldivision activity of separate meristems and to control subsequent cell growth and differentiation. The localisation of cell-division activity in these embryonic regions, which are present throughout the life of the organism, is a unique feature of plants. At any time the majority of these areas are dormant, and their activity governs the morphological organisation of the plant. In addition certain plant cells have the ability to dedifferentiate and may regain the capacity for cell division.

The existence of mechanisms which control cell division, growth, and differentiation may be demonstrated using undifferentiated callus tissue, which can be produced from a variety of plant species. This tissue can be induced to form meristems from which whole plants may be regenerated.

Once an organism has reached a certain size, diffusion is no longer an adequate means of distributing nutrients and metabolites. Thus higher plants, in common with other complex multicellular organisms, must possess one or more efficient transportation systems. The movement of water and nutrients within the plant, while a facet of communication in its own right, may also carry information concerning the activity of one part of the plant to another. For example, if the rate of photosynthesis in the leaves is reduced, the level of sugars in the phloem sap supplying the root system is also likely to be depressed. In this thesis the transport of nutrients will only be discussed with respect to this latter rôle.

Possible Modes of Intercellular Communication

Research into methods of communication in higher animals has revealed two main systems (e.g. Metzler, 1977). The message may take the form of electrical impulses as in the nervous system, or it may be chemical in nature. Hormones are examples of chemical messengers.

Evidence for these two modes of communication has also been investigated by plant physiologists. Electrical impulses or action potentials have been observed in certain plants which exhibit very rapid responses to environmental stimuli (e.g. <u>Dionaea</u> and <u>Mimosa pudica</u>; see Bentrup, 1979). The generality of such nerve-like electrical signals is uncertain, however, and there is considerable scope for more research.

A wide variety of "chemical messengers" are theoretically possible. The active molecules might vary in size from simple compounds such as ethylene to complex proteins which could carry a large amount of information.

As in animals there is the possibility that two or more substances with opposite effects might interact. The active compounds could be present in minute quantities or may be relatively abundant and ubiquitous nutrient molecules or ions. It is also feasible that the gradient of a substance across a cell or tissue may be more important than its absolute amount.

There are several possible routes for the transmission of a chemical message. For rapid communication across considerable distances the mass flow transportation systems of the xylem and phloem might be used. For shorter distances, the substance could move by diffusion in the symplast or apoplast or even through air spaces. There is also evidence that specific active-transport mechanisms for certain compounds (e.g. indole-3-acetic acid; see Goldsmith, 1977) may exist at cell interfaces.

The Discovery of Phytohormones

Despite the large variety of possible forms of chemical message, the majority of research on the means of intercellular communication in plants has concentrated on a few small organic molecules which show The earliest physiological activity when present in minute quantities. evidence for the movement of an "influence" from one part of the plant to another, came from the work of the Darwins (1880). They discovered that a coleoptile could be prevented from responding phototropically by covering the tip with an opaque cap, although the growing cells producing the curvature were located further down the organ. Indications that the "influence" was chemical in nature came from the work of Boysen Jensen (1910), who showed that the stimulus could travel through a block of gelatin, and from Páal (1919). When Páal excised the tips of Avena coleoptiles and replaced them asymmetrically, he observed a greater rate of growth in cells directly below the apex, producing a curvature

resembling that induced by a unilateral light stimulus. He concluded that the coleoptile tip secreted a growth-promoting substance, the distribution of which could be controlled in response to light. This work was further confirmed by Went (1928) who obtained growth-active diffusate from coleoptile tips. Curvature could be induced by placing a block of agar, containing diffusate, asymmetrically on a decapitated The angle of curvature thus obtained was proportional to coleoptile. the number of apices which had been placed on the agar block. Went also showed that when an excised coleoptile tip was given a directional light stimulus, more growth-promoting activity diffused out of the side away from the light source. While studying geotropism, Dolk (1930) demonstrated a redistribution of growth-promoting substance in a horizontal coleoptile tip. He further showed that the total growth and total amount of growth-promoting substance did not change in response to the gravity stimulus.

The first studies of the chemical identity of the growth-promoting substance, which became known as auxin, were carried out on extracts of human urine which had been shown to possess considerable auxin activity in bioassay. In 1934, Kögl <u>et al</u>. isolated crystalline indole-3-acetic acid from this source and demonstrated that this compound had considerable auxin activity. Indole-3-acetic acid (IAA) was subsequently extracted from <u>Rhizopus suinus</u> (Thimann, 1935) and corn seeds (Haagen-Smit <u>et al</u>., 1946). It was not until 1972, however, that IAA was conclusively identified from Zea mays coleoptile tissue (Greenwood et al.).

IAA has now been identified from a large number of plant species using a range of analytical techniques, the accuracy of which is discussed extensively by Reeve and Crozier (1980). Table 1 gives a selection of the plant tissues from which IAA has been identified conclusively by obtaining a full-scan mass spectrum.

Since the discovery of IAA, a number of other groups of compounds, capable of affecting growth and development when present in small

Table 1:A Selection of Plants from which IAA has been Extractedand Identified by Full-Scan Mass Spectra

Species	Tissue	Selected Reference
<u>Zea mays</u>	coleoptile tips roots : stele & cortex roots (sterile)	Greenwood <u>et al</u> . (1972) Bridges <u>et al</u> . (1973) Elliott and Greenwood (1974)
Ricinus communis	xylem & phloem sap	Hall and Medlow (1974)
<u>Avena sativa</u>	etiolated seedlings	Bandurski and Schulze (1974)
Prunus cerasus	seeds	Hopping and Bukovac (1975)
<u>Citrus</u> unshiu	fruit	Igoshi <u>et al</u> . (1971)
Gossypium	ovules	Shindy & Smith (1975)
Phaseolus vulgaris	shoots	McDougall & Hillman (1978)
<u>Picea</u> <u>sitchensis</u>	cambium	Little <u>et al</u> . (1978)
<u>Pinus</u> sylvestris	seedlings	Sandberg et al. (1981)
<u>Fragaria</u> x <u>ananassa</u>	immature fruits	Swartz and Powell (1979)
<u>Malus pumila</u>	shoot tips	Swartz and Powell (1979)
<u>Undaria pinnatifida</u>	young thalli	Abe <u>et al</u> . (1972)

.

•

quantities, have also been extracted from vascular plants. There are five generally accepted groups of "phytohormones" or "plant growth substances": the auxins, the gibberellins, the cytokinins, abscisic acid and ethylene (for a discussion of the terminology for such compounds see Weyers, 1978; and Hewitt, 1980). Several other substances whose physiological activity is less well established, have also been studied (e.g. various phenolics, fatty acids and terpenoids). A selection of the better documented responses to phytohormone application is shown in Table 2.

The Rôle of Phytohormones in Higher Plant Physiology

The term hormone was first used by Bayliss and Starling (1902 and 1904) to describe the action of secretin, a molecule secreted by the duodenum that stimulates the activity of the pancreas. They defined hormones as "chemical messengers produced in one organ and transported in the bloodstream to a site of action in another". Plant growth substances have traditionally been assumed to operate in a similar manner. Thimann (1948) defined a phytohormone as "an organic substance produced naturally in higher plants, controlling growth or other physiological functions at a site remote from its site of production, and active in minute amounts". A physiological response was believed to be induced by a change in the amount of a substance binding to receptor sites in the cells of a target tissue.

Early experiments on the rôle of auxin in the geotropism and phototropism of <u>Avena</u> coleoptiles supported this theory. Auxin appeared to be synthesised in one place, i.e. the coleoptile tip, and transported to a site of action in another, i.e. the zone of curvature. There is, however, an increasing body of evidence that the function of phytohormones may be more complex than the traditional theories suggest, and several alternative hypotheses have been proposed.

Table 2:Physiological Responses to Plant Growth Substances(Generally Species-Specific)

a. Effects on the Fundamental Processes of Cell Division, Growth and Differentiation

Phenomenon	Growth Substance	Effect	Example Reference
Cell Division	Cytokinins	Stimulation	Letham (1963)
	IAA	Stimulation of cambial cell division	Snow (1935)
Cell Growth	IAA	Stimulation of shoot growth	Went & Thimann (1937)
		Inhibition of root growth	Went & Thimann (1937)
	Gibberellins	Promotion of growth of green shoots	Brian and Hemming (1955)
	Cytokinins	Promotion of cotyledon enlargement	Esashi and Leopold (1969)
	Abscisic acid	Growth inhibition	Addicott <u>et al</u> . (1964)
	Ethylene	Growth inhibition	Pratt and Goeschl (1969)
Cell	IAA	Stimulation of xylem	Jacobs (1952)
Differentiation		and phloem differentiation	LaMotte and Jacobs (1963)
Organ Morpho- genesis	IAA	Control of callus tissue morphogenesis	Miller and Skoog (1953)
		Promotion of root initiation	Avery and Johnson (1947)
	Cytokinins	Control of callus tissue morphogenesis	Skoog and Miller (1957)

b. Some Examples of Responses to Plant Growth Substances concerning Aspects of Correlative Control of Plant Development and Response to the Environment

Phenomenon	Growth Substance	Response	Example Reference
Seed Dormancy	Gibberellins	Break dormancy	Suzuki and Takahashi (1968)
	Cytokinins	Break dormancy	Miller (1956)
	Abscisic acid	Inhibits germination	Aspinall <u>et al</u> . (1967)
	Ethylene	Breaks dormancy	Vacha and Harvey (1927)
Seedling Morphology	Ethy lene	Maintains plumular hook	Kang <u>et al</u> . (1967)
Geotropism	IAA	IAA transported to lower side of shoot	Shaw <u>et al</u> . (1973)
Phototropism	IAA	Transported to darkened side of shoot	Gard.ner. <u>et al</u> . (1974)
Bud Dormancy	Gibberellins	Break dormancy	Brian et al. (1955)
	Cytokinins	Break dormancy	Benes et al. (1965)
	Ethy1ene	Breaks dormancy	Pratt and Goeschl (1969)
Apical Dominance	IAA	Replaces apical bud	Thimann and Skoog (1933)
	Cytokinins	Release lateral buds from apical dominance	Sachs and Thimann (1964)
Juvenility	Gibberellins	Reduce period of juvenility	Pharis and Morf (1968)
Stomatal Movement	Abscisic acid	Causes stomatal closure	Mittelheuser and Van Steveninck (1969)
	Cytokinins	Promote stomatal opening	Kuiper (1972)
Leaf	IAA	Delays senescence	Sacher (1959)
Senescence	Gibberellins	Delay senescence	Fletcher and Osborne (1965)
	Cytokinins	Delay senescence	Richmond and Lang (1957)
	Ethylene	Accelerates senescence	Burg (1968)
Leaf Abscission	IAA	Inhibits abscission	La Rue (1935)
	Abscisic acid	Promotes abscission in explants	Bornman <u>et al</u> . (1967)
	Ethylene	Promotes abscission	Zimmerman et al. (1931)

Phenomenon	Growth Substance	Response	Example Reference
Flowering	IAA	Promotes flowering in Bromeliads	Clark and Kerns (1942)
	Gibberellins	Induces flowering in long day and cold requiring plants	Lang (1956)
	Cytokinins	Promotes flowering, some short-day plants	Ogawa (1961)
	Abscisic acid	Promotes flowering, some short-day plants	E1-Antably and Wareing (1966)
	Ethylene	Promotes flowering in Bromeliads	Rodriguez (1932)
Fruit Setting	Gibberellins IAA Cytokinins)) ^{Allows} development of) parthenocarpic fruit	Crane (1964) Crane (1965) Crane (1965)
Fruit Ripening	Ethylene	Stimulates fruit ripening	Lyons and Pratt (1964)
	IAA	Defers ripening	Kendrell (1969)
	Gibberellins	Retards ripening	Coggins and Lewis (1962)
Fruit Abscission	Abscisic acid	Stimulates abscission	Addicott <u>et al</u> . (1964)
	Ethylene	Stimulates abscission	Pratt and Goeschall (1969)

r

-

In the specific example of coleoptile phototropism and geotropism, Firn and Digby (1980) and Trewavas (1981) have pointed out that amounts of IAA moving laterally across the organ may not be sufficient to produce the observed change in growth rate. The effect of IAA on coleoptile elongation is proportional to the log of its concentration (Bentley and Bickles, 1952) with relatively large changes in hormone levels being required to invoke a noticeable response. However, the amount of exogenous IAA transported laterally in response to a gravity or light stimulus (Shaw et al., 1973 and Gardner et al., 1974) was no more than Curry (1969) also showed that the tip was 56% and 15.3% respectively. not the only region of the coleoptile capable of detecting a directional light stimulus, although it was somewhat more sensitive than regions lower down. Similarly, geosensitivity is found along the length of the organ (Firn and Digby, 1980). These results have led the above authors to question whether IAA has any rôle in shoot phototropism or geotropism. In the case of geotropism they suggested that the epidermal cells of the elongation zone may both detect the gravity stimulus and respond directly by changing their rate of growth.

The logarithmic relationship between exogenous concentration and effect is common to the majority of bioassays. On the other hand, the largest observed endogenous changes in amounts of phytohormones are the 40-50 fold increases in the abscisic acid content of some leaves subjected to water stress; a two- to five-fold increase is more commonly recorded (see Trewavas, 1981). On this basis Trewavas proposed that changes in levels of plant growth substances are not normally responsible for physiological changes, but that they might serve to integrate physiological processes, ensuring that growth and differentiation occur in an evenly distributed fashion throughout a group of cells. It must be recognised, however, that measurements of plant hormone levels are frequently made on whole organs or large areas of tissue. Different conclusions might be

reached if the movement of substances between individual cells or even between different compartments within cells was considered. It has been shown (Loveys, 1977; and Weyers, 1978) that movement of abscisic acid from the leaf mesophyll to the guard cells of the epidermis occurs in response to water stress and precedes the increase in total abscisic acid levels (Walton <u>et al.</u>, 1977). Furthermore, the relationship between exogenous hormone concentration and response may not reflect accurately the effect of the endogenous substance. The influence of rates of uptake and metabolism on the activity of the applied compound must be considered.

The effective levels of a plant growth substance might also be altered by regulating the activity of putative receptor sites. Trewavas (1981) replotted the results of Wright (1961 and 1966) on the change in sensitivity to auxin. and the growth rate of wheat coleoptiles, as a function of the developmental age. The graph showed a substantial increase in sensitivity which preceded the rise in growth rate. Osborne (1978) discussed a similar development in the flower buds of Ecballium Only buds which had passed a certain stage in development elaterium. could be induced by ethylene to separate from the pedicel. However, as the ovary is not usually shed until after the ovule has been pollinated and the fruit ripened, development of ethylene sensitivity cannot be the normal trigger for abscission.

Another alternative theory suggests that gradients of plant growth substances may in some cases be responsible for inducing cell differentiation. Wetmore and Rier (1963) presented evidence that gradients of IAA and sucrose within a cylinder of callus induced the formation of nodules of vascular tissue. The relative proportions of IAA and sucrose also affected the ratio of xylem and phloem.

While phytohormones have many dramatic effects on excised tissues and organs they are frequently ineffective when applied to intact plants. For example, the application of IAA to intact coleoptiles led to growth increases of only some 25% (Went and Thimann, 1937). This could be due

to problems of penetration and/or metabolism before the compound reaches its site of action but might also suggest that phytohormones, while necessary for plant growth and development, are not usually present in limiting amounts.

Table 2 shows that a large variety of physiological events appear to be controlled by the five known groups of phytohormones. Individua1 compounds all influence at least one of the fundamental processes of cell division, growth and differentiation; IAA appears to effect all three. Phytohormones also have a wide variety of specific effects in different tissues at various stages of the plants' development, with each process usually being affected by at least two substances. It is difficult to envisage how simple changes in the levels of these five groups of compounds alone can supply enough information to regulate such a wide variety of growth and developmental phenomena. The problem might be partly solved if it could be shown that a unique combination of substances was required for a specific process. Osborne (1978) suggested that the variety of responses might be attributable to there being more than one class of target cell on which a plant growth substance might act, with different results.

In addition, there are several situations in which the existence of a transported message is implied which cannot be explained in terms of the known phytohormones. For example, in many plants where flowering is controlled by photoperiod, it can be shown (see Zeevaart, 1976) that daylength is detected by the mature leaves although the responding tissue is the apex. Grafting experiments implicate the existence of a translocated signal, however all attempts to isolate "florigen" have failed. The nature of the inhibitor responsible for preventing the outgrowth of lateral buds, in the presence of an intact apex, remains unsolved. Similarly, the identities of "senescence factor" and the dormancy-inducing stimulus are unknown.

In the case of the flowering stimulus there is a need for genes to be

activated and deactivated, in order to effect the fundamental changes taking place when a plant is converted from the vegetative to the reproductive phase. There is evidence that steroid hormones in animals have this property (e.g. O'Malley <u>et al.</u>, 1977) but the alteration of gene activity is not thought to be the initial response to plant growth substances. At least some effects of phytohormones are much too rapid to involve the mediation of gene-activated protein synthesis. For example, IAA can alter the rate of coleoptile elongation within 15 min (Ray and Ruesink, 1962). It is therefore possible that other substances have yet to be discovered.

In conclusion, the exact rôle of phytohormones remains unclear. Ιt has been shown that these compounds have many effects when applied to plant tissues. The levels at which growth substances are present in intact plants are also important in certain situations. There are several examples of disease symptoms being caused by excessive hormone production by, or in response to, a pathogenic organism. The "foolish seedling" disease of rice is caused by the synthesis of large quantities of the gibberellin GA_z by the fungus <u>Fusarium moniliforme</u> (Cross <u>et al.</u>, 1961). Dwarfism in certain varieties of Zea is believed to be a result of inadequate gibberellin production (Phinney, 1960). Nevertheless, it is by no means clear whether changes in the levels of plant growth substances are normally responsible for physiological changes, or whether the translocation of a phytohormone constitutes a chemical message. Although they are necessary, phytohormones may not normally be limiting factors controlling plant growth and development.

Rationale Behind the Present Study on the Metabolism of Indole-3-

acetic Acid

If an understanding of the rôle of plant growth substances is to be achieved, certain fundamental questions concerning the biochemistry of these compounds must be answered, viz.:

1. Where is each substance synthesised?

2. How are they distributed within the plant at a cellular and intracellular level, and how does this distribution change at various stages of the plants' development?

3. How are the levels of substances controlled? Several methods are possible:

- (a) Control of synthesis and catabolism.
- (b) Reversible activation and inactivation.
- (c) Intra- and intercellular movement.
- 4. What is the nature of the "receptor sites"?
- 5. What are the primary effects of each substance?

For the purposes of this present investigation, one of these aspects was chosen. The metabolism of indole-3-acetic acid in the root and coleoptile tissues of dark-grown Zea mays seedlings was studied. The rates of catabolism and conjugation of IAA may influence the levels present in the tissue. It is also possible that IAA metabolism might be directly linked in some way to its physiological activity. Research on the effects, transport and binding of IAA has often assumed that metabolism of the applied IAA during the course of the experiment is negligible (e.g. Batra et al., 1975; Moloney and Pilet, 1981) although adequate evidence to support this contention is rarely provided. A detailed investigation of the extent to which exogenous IAA is metabolised is therefore essential.

The Plant System

Zea mays seedlings have been used extensively for research on the metabolism, transport, receptor sites and effects of IAA (e.g. Reinecke and Bandurski, 1981; Wilkins and Scott, 1968; Moloney and Pilet, 1981; Shaw et al., 1973). Zea is the third most important cereal crop in the world, with an annual production of 362,971,000 tonnes in 1978 (Encyclopedia Britannica 1980 Year Book). Commercially available varieties have the advantage of rapid and even germination, reaching a size suitable for experimentation within three to five days. Both the coleoptile and the primary root are unusually large in diameter when compared with other cereal species. When grown in the dark, the lack of chloroplastic components, and products of photosynthesis, reduces the complexity of extracts. While IAA causes substantial stimulation of coleoptile growth, similar amounts inhibit root growth (e.g. Pilet, 1976). Thus the project allows a comparison of IAA metabolism in organs with different responses to the phytohormone. As Zea mays roots can easily be divided into cortex and stele (Greenwood et al., 1973) the metabolism of IAA by these tissues could also be compared.

IAA in Zea mays Seedlings

The main source of IAA in young Zea mays seedlings appears to be the endosperm. Bandurski and co-workers have shown that this tissue contains large quantities of IAA, most of which is esterified to either <u>myo</u>-inositol or high molecular weight glucans (Ueda <u>et al.</u>, 1970; Piskornik and Bandurski, 1972; Bandurski and Schulze, 1977). They have further demonstrated that a sufficient quantity of IAA-<u>myo</u>-inositol is exported to the shoot where it can be hydrolysed to supply the coleoptile with its full requirement for IAA (Nowacki and Bandurski, 1980). IAA-<u>myo</u>-inositol is also transported to the root (Epstein <u>et al.</u>, 1980). On the other hand, there is some evidence that the root may be able to synthesise IAA (Elliott,

1977; Feldman, 1980).

The movement of IAA in Zea mays roots, as in many other plants, is predominantly acropetal (e.g. Scott and Wilkins, 1968; Shaw and Wilkins, 1974). The majority of IAA in the root is found in the stele, which also shows more efficient transport of IAA (Bowen <u>et al.</u>, 1972). Quantities of IAA measured using mass spectrometry in the variety Giant White Horsetooth were as follows (Bridges et al., 1973):

stele	53.3	μg	IAA/ Kg	fresh	weight
cortex	4.8	μg	IAA/ Kg	fresh	weight
root tip	29.0	μg	IAA/Kg	fresh	weight

Pilet and Elliott (1981) have, however, warned that quantities of IAA might vary substantially between varieties.

In <u>Zea mays</u> shoots, the transport of IAA is polarised in a basipetal direction (Hertel and Leopold, 1963). The quantity of IAA present in coleoptiles from the variety Anjou 21 was measured as 10 pmol /2.5 mm tip by Weiler et al. (1981) using a radioimmunoassay.

IAA has been shown on many occasions to cause substantial stimulation of coleoptile elongation (e.g. Thimann, 1969). Its effect on root growth is, however, less clear. Exogenous IAA concentrations of around 10^{-2} mol m⁻³ which produce maximum growth rates in mesocotyl and coleoptile segments inhibit the growth of root sections (e.g. Batra et al., 1975; Greenwood and Yčas, 1975). While some workers have found no evidence of IAA-induced promotion of root growth (e.g. Greenwood and Yčas, 1975) others claim to have observed a small and often transient stimulation at very low concentrations of between 10^{-4} mol m⁻³ and 10^{-7} mol m⁻³ (e.g. Batra et al., 1975; Edwards and Scott, 1977; Elliott, 1977; Evans et al., 1980). Pilet and co-workers (Pilet et al., 1979; Pilet and Elliott, 1981) presented evidence that root segments, depleted of IAA by diffusing onto agar for periods of up to 8h, will show significant IAA-induced growth promotion. They suggested that in the intact plant the roots have a saturating level of IAA.

It has been shown (Rayle & Gleland, 1970) that low pH can stimulate shoot growth in a similar manner to IAA. This led Marré and co-workers (1974) to put forward the "acid-growth" theory, in which they suggested that IAA stimulates growth by causing protons to be pumped out of the cell into the wall, where the lowered pH causes wall-loosening. In support of this hypothesis was the observation that fusicoccin, a fungal toxin, could stimulate shoot growth by such a proton-pump mechanism (e.g. Lado <u>et al.</u>, 1973). Edwards and Scott (1974) and more recently Moloney <u>et al.</u> (1981) showed that the growth of roots could also be stimulated by low pH. Furthermore, fusicoccin can cause proton extrusion in roots and thereby promote elongation (Lado <u>et al.</u>, 1976). Pilet (1976), however, has pointed out that fusicoccin and IAA cannot be acting by a common mechanism in the root, as the optimum concentration of fusicoccin for growth promotion was 10^{-2} mol m⁻³, with stimulation also found at 10^{-3} , 10^{-4} and 10^{-1} mol m⁻³. IAA at any of these concentrations produced growth inhibition.

Batra, Edwards and Scott (1975) used a different approach to investigate the relationship between acid growth and the effect of IAA on root elongation. They studied the effect on growth, of IAA in buffered agar of varying pH. An IAA concentration of 10^{-4} mol m⁻³ caused growth inhibition at pH 4 while promoting extension at pH 7. In a later paper, Edwards and Scott (1977) found that 10^{-4} mol m⁻³ IAA would promote growth only in a buffered solution at pH 7 and not in an unbuffered solution at the same pH. The authors suggested that their observations might be explained in terms of pH effects on IAA transport as they are not easily compatible with the acid-growth theory. The results are also complicated by the influence of the buffering substance itself on growth (Edwards and Scott, 1976; Gabella and Pilet, 1978). Thus under most experimental conditions, IAA appears to inhibit root growth. Evidence for growth promotion is inconclusive, but the acid-growth theory is not thought to Evans and co-workers (Evans and Mulkey, 1980; Evans et al., 1980) apply. have recently presented evidence that IAA may inhibit root growth by

causing protons to be pumped into the cells, in the reverse direction to shoots, thereby raising the wall pH. Such a change in pH was measured in the bathing solution surrounding maize root segments incubated in 2×10^{-3} mol m⁻³ IAA. The start of the pH change almost coincided with the time at which the growth rate began to fall rapidly.

On the other hand, the effect of exogenous IAA may not be a reliable indication of the rôle of the endogenous compound. The use of segments necessarily involves the presence of cut surfaces. Both wounding, and the application of auxin can induce ethylene production (e.g. Burg, 1962; Abeles, 1966). Chadwick and Burg (1967) and more recently Mulkey <u>et al</u>. (1981) have suggested that the inhibition caused by exogenous IAA is due to this stimulation of ethylene biosynthesis.

In conclusion, the existence, origin and transport of IAA in Zea mays seedlings are well established, as is the effect of the exogenous compound on the growth of the coleoptile and mesocotyl. However, much work is needed before its rôle in the control of root elongation is understood.

Metabolism of IAA

The importance of IAA metabolism was realised as early as the 1950s when Kamerbeek (1956) and Galston (1957) hypothesised that there might be a causal relationship between dwarfism, endogenous levels of IAA, and the activity of IAA-oxidising enzymes. Since then dwarfism has been found to be more readily explained by other means, such as gibberellin content or sensitivity to plant growth substances (see Jones, 1973). Interest in IAA metabolism as a possible means of regulating the levels of this phytohormone has remained, however, and much research has been carried out on this topic.

Three main approaches have been used in the study of IAA metabolism: 1) The partial characterisation of enzyme preparations capable of oxidising IAA, and identification of the reaction products.

2) The application of IAA, usually labelled with 14 C or 3 H to plant tissues, followed by the extraction and analysis of its metabolites.

3) The extraction, and identification with varying degrees of certainty, of endogenous indolic compounds.

The substances found as a result of these lines of investigation are listed in Table 3 and their chemical structures shown in Fig. 1. Table 4 compares the detection limits, specificity and information content of the various techniques used to identify IAA and its metabolites. According to Reeve and Crozier (1980) the conclusive identification of an unknown component of a complex plant extract requires either a minimum of 140 bits of information, or a demonstration that the specificity of the detector is adequate, using the successive approximation method. A comparison of Tables 3 and 4 indicates that in many cases identification of compounds has not been sufficiently rigorous.

Products formed by IAA-oxidase preparations fall into two groups: 3-met hylene oxindole and related compounds, and indole-3-aldehyde. In both cases the reaction involves decarboxylation with loss of carbon-1 of the side chain as CO_2 . Various pat hways leading to these compounds have been proposed (e.g. Hinman and Lang, 1965; BeMiller and Colilla, 1972; Ricard and Job, 1974; Suzuki and Kawarada, 1978). Fig. 2 shows a simplified scheme from Sembdner <u>et al.</u> (1980). The same enzyme can produce either 3-met hylene oxindole or indole-3-aldehyde; Ricard and Job (1974) demonstrated that the proportions of these two substances varied with the relative amounts of enzyme and substrate, and with the pH of the medium.

In vivo studies of IAA metabolism give a completely different picture, however. The most common and abundant metabolites appear to be IAA conjugates with amino acids, sugars or <u>myo</u>-inositol (e.g. Feing <u>et al.</u>, 1976; Kopcewicz <u>et al.</u>, 1974). Reports of decarboxylation products such as 3-methyleneoxindole or indol e-3-aldehyde have usually been based on somewhat inadequate chromatographic and colorimetric identifications.

Table 3: IAA Metabolites

a. Products of In Vitro IAA Metabolism

Compound	Enzyme	Selected Reference	Method of Identification			
3-methylene- oxindole	Horseradish peroxidase	Hinman and Lang (1965)	1 b, 6			
	Zea mays IAA- oxidase	BeMiller and Colilla (1972)	la, 6			
	Wheat peroxidase B _l	Zmrhal and Mackáčková (1978)	1a, 6			
	Turnip peroxidases P ₁ and P ₇	Ricard and Job (1974)	6			
3-hydroxymethy1- oxindole	Horseradish peroxidase	Suzuki and Kawarada (1978)	6, 7, 8			
	Zea mays IAA- oxidase	BeMiller and Colilla (1972)	la, 6			
indole-3-aldehyde	Horseradish peroxidase	Ricard and Job (1974)	6			
	Turnip peroxidases P ₁ and P ₇	Ricard and Job (1974)	6			
	Zea mays IAA- oxidase	BeMiller and Colilla (1972)	1a			
indole-3-methanol	Wheat peroxidase ^B 1	Zmrhal and Mackáčková (1978)	1a, 6			
3,3' diindoly1- methane	Zea mays IAA- oxidase	BeMiller and Colilla (1972)	1a, 5, 7			
	Horseradish peroxidase	Suzuki and Kawarada (1978)	6,7,8			
3-acetoxyindole	Horseradish peroxidase	Suzuki and Kawarada (1978)	6, 7, 8			
3-(indol-3-yl- methyl)oxindole						
3 (2"-indol-3"-y1met indol-3-y1methy1	hyl) oxindole					
3-hydroxymethy1-3(3-y1methy1)oxindd	indo1- le					
3-hydroxymethy1-3- 3"-y1methy1)indo1-	3-hydroxymethy1-3-[(2 ["] -indo1- 3"ylmethy1)indo1-3'ylmethy1]oxindole					
2-(indol-3-ylmethy	v1)indoly1-3-acetic a	acid				

Compound	Tissue	Species	Selected Reference	Method of Identification
Indol-3-acety1 aspartate	crown gall tissue	Parthenocissus tricuspidata	Feung <u>et</u> <u>a1</u> . (1976)	1c, 3, 7
	shoot bark	<u>Pinus pinea</u>	Riov and Gottlieb (1980)	8,9
	stem sections	<u>Pisum sativum</u>	Andreae and Good (1955)	2a, 3, 6
	callus	<u>Citrus</u> <u>sinensis</u>	Epstein <u>et</u> <u>al</u> . (1977)	1c
	stem segments	Phaseolus vulgaris	Davies (1972)	1c, 4
	leaves	<u>Scrophularia</u> arguta	Miginiac <u>et al.</u> (1978)	1c
	leaves	<u>01ea</u> europa	Epstein and Lavee (1977b)	2c
Indole-3-acety1 glycine	crown gall callus tissue	Parthenocissus tricuspidata	Feung <u>et al.</u> (1976)	1c, 3, 7
Indole-3-acetyl alanine				
Indole-3-acety1 valine				
Indole-3-acetyl glutamate				1c, 3
Indole-3-acetyl glucose	needles	<u>Pinus pinea</u>	Riov and Gottlieb (1980)	1c, 8
	epicoty1s	Pisum sativum	Zenk (1961)	2a,2c,3,5,6
	caryopses	Zea mays	Kopcewicz <u>et al</u> . (1974)	1a,1c,3,5
	whole plants	Orobanche ramosa	Magnus <u>et al</u> . (1982)	la, 3
Indole-3-acetyl myo-inositol	caryopses	Zea mays	Kopcewicz <u>et al</u> . (1974)	la,1c,3,5
Indole-3-aldehyde	stem sections	<u>Pisum</u> <u>sativum</u>	Magnus <u>et</u> <u>al</u> . (1971)	1a
	leaves	<u>Olea</u> europa	Epstein and Lavee (1977b)	2c, 4
	leaves	<u>Scrophularia</u> arguta	Miginiac et <u>al</u> . (1978)	1c
	inflorescence	Orobanche <u>lutea</u>	Magnus <u>et</u> <u>al</u> . (1982)	la, 6, 8
	whole plants	Orobanch ^e ramosa	Magnus <u>et</u> <u>al</u> . (1982)	1a, 6, 8

b. Products of the In Vivo Metabolism of Exogenous IAA

		_		
Compound	Tissue	Species	Selected Reference	Method of Identification
Indole-3-carbo- xylic acid	stem sections	<u>Pisum sativum</u>	Magnus <u>et</u> <u>al</u> . (1971)	1a, 5
	leaves	<u>01ea europa</u>	Epstein and Lavee (1977b)	2c, 4
	whole plants	Orobanche ramosa	Magnus <u>et</u> <u>a1</u> . (1982)	1a, 6, 8
Indole-3- methanol	stem sections	<u>Pisum</u> <u>sativum</u>	Magnus <u>et al</u> . (1971)	1a, 5
	inflorescence	Orobanche lutea	Magnus <u>et</u> <u>al</u> . (1982)	1a, 5
3-Methylene " oxindole	callus	app1e	Epstein <u>et</u> <u>al</u> . (1975)	2a,2b,2c,5
	coleoptiles	<u>Avena sativa</u>	Menschick and Hild (1976)	1c
	callus	<u>01ea</u> europa	Lavee and Epstein (1976)	2c, 4
3-Hydroxymethy1- oxindo1e	etiolated seedlings	<u>Pisum</u> sativum	Tuli and Moyed (1967)	1b, 1d, 6, 5
	stem segments	<u>Pisum sativum</u>	Davies (1972)	1c, 4
	stem segments	Phaseolus vulgaris	Davies (1972)	1c, 4
	callus	apple	Epstein <u>et al.</u> (1975)	2a,2b,2c,4
	callus	<u>Olea</u> europa	Lavee and Epstein (1976)	2c, 4
3-Methy1- oxindole	in c ubating solution surrounding callus	Parthenocissus tricuspidata	Hamilton <u>et al</u> . (1976)	1c
Oxindole-3- acetic acid	endosperm	Zea <u>mays</u>	Reinecke and Bandurski (1981)	1c,7,9,10

c.	Endogenous	Compounds	which	may	be	IAA	Metabolites

Compound	Tissue	Species	Reference	Method of Identification			
Indole-3-acetyl aspartate	seedlings	Phaseolus vulgaris	Tillberg (1974)	la, 3			
Indole-3-acetyl aspartate	seed	Glycine max	Cohen (1981)	7			
2- 0 -(indole-3- acetyl)-D-) glucose							
4-0-(indole-3- acety1)-D-) glucose	caryopses	Zea mays	Ehmann (1974)	7			
6-0-(indole-3- acety1)-D-) glucose							
Indole-3-acetyl <u>myo</u> -inositol	caryopses	<u>Zea mays</u>	Ehmann and Bandurski (1974)	7,1a,5,9			
	shoots	Zea <u>mays</u>					
	seeds	<u>Oryza</u> sativa	Hall (1980)	la,5,7,9			
IAA- <u>myo</u> -inositol- arabinosides	caryopses	Zea mays	Ueda <u>et al</u> . (1970)	la, 7, 9			
IAA-myo-inositol- galactosides	caryopses	Zea mays	Ueda <u>et al</u> . (1970)	la, 7, 9			
Methyl(oxindole- 3-acetate)	rice bran	<u>Oryza</u> <u>sativa</u>	Kinashi <u>et</u> <u>al</u> . (1976)	6, 7, 8			
Methyl(3-Hydroxyoxindole- 3-acetate)							
Methy1(5-hydroxy- oxindo1e-3-acetat	e)						
3,5-Dihydroxyoxind 3-acetic acid	ole-						
Methyl(3,5-Dihydro 3-acetate	xyoxindole-						
Oxindole-3- acetate	endosperm	<u>Zea</u> <u>mays</u>	Reinecke and Bandurski (1981)	7, 9, 10			
- d. Key to Techniques Used in the Identification of Metabolites
- Co-chromatography with standards in thin-layer system(s); detection of spots by (a) colour reactions
 - (b) UV absorbance
 - (c) radioactivity
 - (d) bioassay

2. Co-chromatography with standards in paper system(s); detection of

- spots by (a) colour reactions
 - (b) UV absorbance
 - (c) radioactivity
 - (d) bioassay

3. Susceptibility to, and products of hydrolysis.

- 4. Comparison of metabolism of IAA labelled in different positions.
- 5. Other chemical tests.
- 6. UV Absorption spectroscopy
- 7. Mass spectrometry (full scan).
- 8. NMR spectroscopy.
- 9. Co-chromatography with standards on gas chromatograph.
- 10. Co-chromatography with standards on high-performance liquid chromatograph.

(a) Oxidation Products



3-Methyleneoxindole



3-Methyloxindole



3-Hydroxymethyloxindole



Indole-3-aldehyde



Indole-3-methanol



Indole-3-carboxylic acid



3,3'-di-indolylmethane





3-(Indol-3'-ylmethyl)oxindole



3-Hydroxymethyl-3-(indol-3-ylmethyl)oxindole

(a) Oxidation Products







2-(Indol-3-ylmethyl)indole-3-acetic acid



Methyl 3,5-Dihydroxyoxindole-3-acetate

Methyl Oxindole-3-acetate



НО ОН СООН

Methyl 3-Hydroxyoxindole-3-acetate

3,5-Dihydroxyoxindole-3-acetate



Methyl 5-Hydroxyoxindole-3-acetate



3-Hydroxymethyl-3-[(2"-indol-3"-ylmethyl)indol-3-ylmethyl]oxindole

(b) IAA Conjugates

.



Indole-3-acetyl Aspartate



1-0-(Indole-3-acetyl)-\$-D-Glucose



2-0-(Indole-3-acetyl)-<u>myo</u>-inositol

Table 4:Assessment of Methods Used to Identify IAA and its Metabolitesin Complex Plant Extracts(Modified from Reeve & Crozier, 1980)

Technique	Detection Limit	Reference	Selectivity	Information Content (bits)
Bioassay e.g. <u>Avena</u> straight growth assay	20ng	Eliasson (1969)	medium	4
Colour Reactions				
Ehrlich's	50ng	Stahl (1965)	medium	4
Salkowski's	2 ,4 g	Tang and Bonner (1947)	medium	4
Spectrofluorimetry				
fixed	10pg	Reeve and Crozier	high	4
scan	lng	(19 80)	medium	32
Electrochemical	50pg	Sweetzer and Swartfager (1978)	medium	4
GLC Detectors				
Flame ionisation detection	10ng '	Reeve and Crozier (1980)	low	4 .
Alkali flame ionisation detection	5ng	Swartz and Powell (1979)	medium	4
Electron capture detection	10pg	Seeley and Powell (1974)	high	4
Mass Spectrometry				
scan	100ng	Reeve and Crozier (1980)	low	2,000
single ion monitoring	100pg	н н	high	4
Infra-Red Spectroscopy	10 , µg	Powell (1967)	1ow	320
Spectrofluorometry of indole apyrone derivative	of lng	Stoess1 and Venis (1970)	high	4
Radioimmunoassay	100pg	Weiler <u>et al</u> . (1981)	high	4
UV Spectrometry (scan)	100ng	Reeve and Crozier (1980)	medium	24
Liquid Scintillatior Spectrometry	n depends on specific activity		high	4
$1_{\rm H}~{\rm NMR}$ spectroscopy	10 µg	Reeve and Crozier	low	124

,

Note: Where these techniques are used to detect compounds following chromatographic analysis, the information content quoted does not include that provided by the chromatogram. Although not easily quantified, this will be high in the cases of HPLC and capillary column GC.

Fig.2. Proposed Pathway of IAA Oxidation Catalysed by IAA Oxidases (Sembdner <u>et al.</u>, 1980)



3-Methyloxindole

Moreover, Epstein <u>et al</u>. (1980) have failed to find any evidence for the evolution of ${}^{14}\text{CO}_2$ from carboxy1-labelled IAA. Instead Reinecke and Bandurski (1981) have identified oxindole-3-acetic acid as the major product of IAA catabolism in <u>Zea mays</u> caryopses. This compound must be formed by a quite different oxidation reaction and is unlikely to be catalysed by any of the IAA-oxidases studied. An alternative sequence of reactions to the decarboxylation pathway (see Fig. 3) was proposed by Kinashi <u>et al</u>. (1976) who extracted the methyl esters of oxindole-3-acetic acid and several related compounds from rice bran.

Analyses of endogenous indolic compounds have indicated that IAA conjugates may be important metabolites, on a quantitative basis. A large proportion of the IAA present, particularly in tissues such as seeds, with a storage function, is conjugated (e.g. Bandurski and Schulze, 1977). Dry Zea caryopses contained 70-90 mg IAA/g tissue as "alkali-labile" esters (Ueda and Bandurski, 1969) while the amounts of "free" IAA varied between 1 and 10% of this figure. The seedlings also contained the majority of their IAA in the form of esters (Bandurski and Schulze, 1974). In the caryopses about half of the esterified IAA was conjugated to high molecular weight glucans, most of the other half being esters of IAA with myo-inositol. The structures of the IAA-myo-inositol isomers have been verified using mass spectrometry (Ueda et al., 1970). Several esters of IAA with glucose were also identified from the same tissue (Eh mann 1974). Kopcewicz et al. (1974) fed Zea mays caryopses with labelled IAA and showed that it became esterified to the same compounds found in the tissue. An enzyme preparation, which was later purified (Michalczuk and Bandurski, 1980), was found to catalyse the esterification of IAA by the following route.

IAA + UDP-Glucose IAA-Glucose + UDP IAA-Glucose + myo-Inositol - IAA-myo-Inositol + Glucose IAA-myo-Inositol + UDP-Galactose - IAA-myo-Inositol-Galactoside + UDP Several functions for the esters of IAA have been proposed. The large



соон

decrease in high molecular weight esters of IAA during germination, accompanied by a small increase in free IAA, observed by Ueda and Bandurski (1969), suggested that these compounds have a storage function. Quantitative studies of the turnover of indolic compounds in <u>Zea</u> caryopses (Epstein <u>et al.</u>, 1980) strongly indicated that high molecular weight esters are the principal source of IAA in the young seedling. Nowacki and Bandurski (1980) further demonstrated that IAA-<u>myo</u>-inositol was transported into the shoot, where it could be hydrolysed at a rate sufficient to fulfil the requirements for IAA. An enzyme preparation capable of catalysing this reaction has recently been extracted from <u>Zea mays</u> shoots (Hall and Bandurski, 1981).

In other species, labelled IAA has been shown to form conjugates with, the amino acids glycine, valine, alanine, aspartic acid and glutamic acid (Feung <u>et al.</u>, 1976). Endogenous IAA-aspartate has also been extracted recently from seeds of <u>Glycine max</u> (Cohen, 1981). The function of these conjugates is thought to be similar to that of esters of IAA, i.e. storage. Feung <u>et al</u>. (1977) and Hangarter and co-workers (Hangarter <u>et al</u>., 1980; Hangarter and Good, 1981) studied the physiological activity of various amino acid conjugates. The majority of compounds were apparently growth active although slightly less so than IAA. Activity of the conjugates outlasted that of IAA and was also related to the rate of decarboxylation, suggesting that these compounds were acting as slow release sources of IAA. Conjugates with amino acids or sugars have also been identified as products of the metabolism of the synthetic auxin 2,4-dichlorophenoxyacetic acid (Scheel and Sanderman, 1981).

The lack of evidence for decarboxylation of IAA <u>in vivo</u> casts some doubt on whether traditional IAA-oxidases are involved in the catabolism of endogenous IAA. It is now known that plants often contain several peroxidase isoenzymes which frequently possess varying degrees of IAAoxidase activity and have different cofactor requirements (e.g. McCune,

1961; Sequeira and Mineo, 1966; Ockerse <u>et al.</u>, 1975). Schneider and Wightman (1976) have described four classes of IAA-oxidase/peroxidase enzymes:

- 1. Typical peroxidases that catalyse peroxidation reactions at a very high rate while also oxidising IAA more slowly (e.g. Horseradish peroxidase).
- Enzymes that have both high peroxidase and high IAA-oxidase activity (e.g. Turnip peroxidase P₇, Ricard <u>et al.</u>, 1972).
- 3. Peroxidases that show no IAA-oxidase activity (e.g. from pea roots, Collet, 1968).
- 4. IAA-oxidases that display no peroxidase activity (e.g. Sequeira and Mineo, 1966; Bryant and Lane, 1975). The non-peroxidative enzyme isolated by Bryant and Lane did not possess a heme group.

Conflicting opinions exist as to whether IAA-oxidase activity and peroxidase activity can be separated (c.f. Lee, 1976; Laurema , 1974). Stonier <u>et al.</u> (1979) proposed that high molecular weight IAA-oxidase inhibitors, known as auxin protectors, which do not inhibit peroxidase activity, may contaminate enzyme preparations to varying degrees giving the appearance that IAA-oxidase and peroxidase activity is due to different enzymes. Siegel and Galston (1967) suggested that there might be different active sites for IAA-oxidase and peroxidase activity on the same enzyme, a view also shared by £utz (1957). Removal of the heme prosthetic group with a mixture of cold acetone and HC1 destroyed the peroxidase activity, although the apoenzyme retained its IAA-oxidase activity. This result was not, however, obtained by Ku <u>et al</u>. (1970) who found that both IAA-oxidase and peroxidase activity stopped when the heme group was removed by a different method.

Auxin protectors are reported (Stonier, 1976) to be derivatives, complexes or polymers of <u>o</u>-dihydroxyphenyl propanoids. <u>O</u>- and <u>p</u>-dihydric phenols and polyphenols are also generally found to be inhibitory to <u>in vitro</u> IAA oxidation (Ray, 1958; Sirois and Miller, 1972). The latter compounds

affect the oxidation rate, rather than induce a lag period (Stonier, In contrast, certain monophenols and m-dihydric phenols e.g. 2,4-1970). dichlorophenol, are often required as cofactors (Ray, 1958). Mn²⁺ ions are also frequently needed. Maclachlan and Waygood (1956 a,b) have suggested that IAA may not be directly oxidised by the enzyme, but indirectly by Mn³⁺ formed in the mixture. Earlier, Kenten (1955) had demonstrated that peroxidase could oxidise Mn^{2+} to Mn^{3+} in the presence of certain Maclachlan and Waygood were able to decarboxylate IAA using a phenols. Mn^{3+} - EDTA solution. This was subsequently confirmed by BeMiller and Colilla (1972) who claimed that the products of this reaction were identical to those of the enzymic process. Some highly purified enzymes may, however, oxidise IAA in the absence of cofactors and several publications now favour a more intimate association of the enzyme with IAA, at least in the case of horseradish peroxidase (e.g. Gelinas, 1973; Yamazaki and Yamazaki, 1973; Nakajima and Yamazaki, 1979). The exact mechanism of the reaction is not fully understood, with several conflicting hypotheses being put forward on the basis of kinetic studies. However, it seems clear that the reaction involves the formation of free radicals (e.g. Parups, 1969; Meudt, 1971; Horng and Yang, 1975). Hydrogen peroxide is usually a requirement for peroxidase-catalysed IAA oxidation (e.g. Kokkinakis and Brooks, 1979) and, where H_2O_2 is not needed, it is thought that a bound form of peroxide is produced and used during the reaction (Mackáčková et al., 1975).

To summarize, much work has been confined to the details of the mechanism of IAA oxidation by a few enzymes, such as horseradish peroxidase, although the <u>in vivo</u> significance of this reaction is questionable. In addition to the lack of evidence for IAA decarboxylation products <u>in vivo</u>, the results of these studies show several points which bring further doubt on the rôle of IAA-oxidases:

1. There is a general lack of specificity shown by IAA-oxidising peroxidase enzymes (Butt, 1980).

- 2. The intimacy of the relation between substrate and enzyme has been questioned.
- 3. Certain inorganic species and UV radiation will catalyse IAA oxidation by the same pathway as IAA-oxidases.

(see Sembdner et al. 1980)

- 4. Peroxidase enzymes are known to be increased by injury and application of abnormal levels of growth substances (e.g. Birecka and Miller, 1974). Briggs <u>et al</u>. (1955) found that application of a drop of 0.005 M KCN to the cut ends of <u>Osmunda cinnamomea</u> fronds considerably increased the yield of diffusable auxin. It is possible that the IAA-oxidase/ peroxidase system is produced in response to injury and its activity is predominantly artifactual.
- 5. Finally, the activity of tissue homogenates might sometimes result from the meaningless mixing of enzyme, phenolic compounds and inactive protein from different cells or organelles, and which are not normally in contact with one another.

On the other hand, there is some evidence to indicate that IAA-oxidases may have a rôle in vivo. A few reports have been published which suggest that IAA levels may be regulated by IAA-oxidase activity. An inverse relationship between enzyme activity and endogenous IAA content has been reported during abscission in bean seedlings (Jain et al., 1969) and lightinduced hook opening of pea seedlings, by precise kinetic studies (De Greef et al., 1977). More recently, Saleh (1981) found that kinetin treatment increased the levels of extractable IAA in the roots of young Phaseolus vulgaris, Zea mays and Avena sativa plants. This treatment also decreased the amount of IAA-oxidase activity detectable. Nevertheless, the quantitative correlation was not good. The fall in IAA-oxidase activity was greatest in oat roots, where kinetin had very little effect on IAA levels, but was rather small in bean roots, where kinetin treatment

significantly raised the amount of IAA. Lee (1980) has published evidence that certain phenolic compounds which act as cofactors or inhibitors of IAA-oxidase preparations may have similar effects on the <u>in vivo</u> catabolism of IAA. Although Epstein <u>et al</u>. (1980) were not able to detect any $^{14}CO_2$, several other workers, using a variety of plant material, have detected substantial quantities of labelled CO_2 following application of IAA-1- ^{14}C (eg.Wilkins <u>et al</u>.,1972b,Zea mays roots; Epstein <u>et al</u>. 1977, orange callus; Epstein and Lavee, 1975, apple callus; Epstein and Lavee 1977a, olive leaves; Hamilton <u>et al</u>., 1976, <u>Parthenocissus tricuspidata</u> callus tissue). It must be borne in mind that where experiments were performed in the light, or involved contact of IAA with cut surfaces, these results could be artifactual.

In conclusion, in order to understand whether IAA-oxidase has a rôle in the catabolism of endogenous IAA, there must be further definitive studies of IAA metabolism <u>in vivo</u>. In the past such work has been seriously hampered by the use of inadequate identification procedures and lack of controls, together with a disregard for the quantitative importance of the products studied. It is therefore proposed that ideally in future, the following precautions should be considered.

1. Where exogenous compounds are used, their initial purity must be proved.

- 2. Care should be taken to include controls with dead tissue and to check for breakdown of compounds during purification and analysis of extracts.
- 3. Identification of products must be rigorous and as far as possible quantitative (see Reeve and Crozier, 1980).
- 4. Exogenous IAA should not be added to <u>in vivo</u> systems in amounts which far exceed natural levels.
- 5. The site of application must be such that entry to the tissue mimics as closely as possible the endogenous system.
- 6. Contact of the IAA with wounded tissue is to be avoided as far as possible and must be considered when interpreting results.

- It must be determined whether or not isotopic discrimination occurs in labelling experiments.
- The possibility that surface microorganisms may be metabolising significant quantities of the applied substance should be borne in mind (see Libbert and Rische, 1969).
- 9. Experiments are based on the assumption that IAA itself is a biologically important compound. The growth activity and general physiology of metabolites needs to be considered.

With these guidelines in mind, this thesis presents a study of the metabolism of radioactively-labelled IAA in the root and coleoptile tissues of dark-grown Zea mays seedlings. Initial experiments were performed using segments floated in solutions of 14 C-IAA. Methanolic extracts of plant material were analysed by the traditional method of thin-layer chromatography in three solvent systems and by the more efficient separatory technique of gradient-elution reverse-phase high-performance liquid chromatography. Virtually no sample purification was carried out, to allow analysis of the whole spectrum of IAA metabolites. Controls using boiled tissue were included, and small-scale experiments were carried out using sterile roots to ensure that the metabolism observed was not due to contamination. The effect of external IAA concentration on the metabolism pattern was monitored. uptake of exogenous IAA was compared with endogenous levels. The changes in the pattern of metabolites were followed during time-course experiments and quantities of label remaining associated with the IAA at each time were estimated. Root segments could be divided into epidermis plus cortex and stele, and the metabolism of the two different tissues compared. In later experiments IAA was supplied to root segments via the stele from agar blocks. This method was also used to supply IAA to the apical ends of coleoptile segments. In this way, metabolism of IAA from a source more closely resembling the endogenous supply could be monitored. The metabolism of the IAA transported through the segments was also studied. Finally, some information

on the chemical structures of metabolites was gained from comparison of 14 C-IAA labelled in different positions, tests for methylation and hydrolyis, co-chromatography with standards of putative IAA metabolites, and UV spectrometry.

A few preliminary experiments were carried out to study the effect of IAA on root and coleoptile elongation.

MATERIALS AND METHODS

1. PLANT MATERIAL

Dark-grown seedlings of <u>Zea mays</u> L. cv. Fronica were used for the majority of experiments although preliminary studies were carried out using the variety Kelvedon 33. Caryopses were obtained from Sinclair McGill (Scotland) Ltd. and stored at 4°C prior to sowing. Plants were grown on damp tissue paper in plastic seed trays covered with a glass sheet, in darkness at 22°C. After 3-4 days, seedlings with roots between 35 and 55mm in length were used as a source of root tissue (Frontispiece), whereas coleoptiles between 15 and 25mm in length were selected from 5-day-old plants.

For metabolism experiments root segments, 20mm in length, were taken from 1mm behind the tip unless stated otherwise. Sections 5mm long were used for experiments on the effect of IAA on growth. This region contains the elongation zone, but excludes the root cap and the majority of the meristem (Erickson and Sax, 1956; and Pilet and Senn, 1980). Coleoptile segments, 10mm in length, were taken from 2mm behind the apex and the leaf tissue was removed from the centre. Sections were cut either in dim green light or in the laboratory, as stated in each experiment.

The physiological safe light was obtained by covering a 40W Tungsten bulb with red and green Cinnamoid filter. This allowed light of between 520 and 590 nm to be transmitted. Tests for physiological activity of this light, using Avena sativa and Zea mays coleoptiles proved negative.

Sterile Plants

Seeds were surface-sterilised by soaking in a 10% Chloros solution (Industrial Grade; ICI, Durham Chemicals Distributors Ltd.) for 30min. After washing thoroughly with autoclaved water, they were transferred to sterile glass petridishes lined with moist sterile filter paper and allowed to grow for 4 days. Manipulation of seeds and cutting of root segments were carried out in a sterile air-flow cabinet. To test for fungal and bacterial contamination, portions of the roots were incubated at 20°C for 2 weeks on either nutrient agar (pH 7.4) or potato dextrose agar (pH 6).

41

Nutrient agar was prepared by dissolving 'Oxoid' nutrient agar (14g) in distilled water (500 cm³). The recipe for potato dextrose agar was as follows: potato (100g) was boiled in water (300 cm³) and strained through muslin. Dextrose (10g) was added, and the volume brought up to 500 cm³ before dissolving agar (10g). Both media were autoclaved for 15min at 120°C and 103 KPa. After cooling, 50mg of Streptomycin was added to the molten potato dextrose agar.

•

«a

2. <u>CHEMICALS</u>

Chemicals were obtained from the following sources:

a. Solvents

methanol* Lothian Chemicals, Edinburgh diethyl ether*, chloroform May and Baker Ltd. propan-2-ol*, toluene A & J Beveridge Ltd. methyl acetate, methoxyethanol, ethyl acetate* British Drug Houses Ltd. acetic acid, formic acid, xylene Hopkin and Williams Ltd. Solvents denoted * were routinely glass distilled before use.

b. Gases

Oxygen free	nitrogen,				
2% propan	e/98% Argon	British	Oxygen	Company	Ltd.

c. Bulk Chemicals

2,5-diphenyloxazole,
2-phenylethylamine, ammonia
Koch Light Laboratories Ltd.
Triton-X 100
A & J Beveridge Ltd.
agar (technical), nutrient agar
NaOH, KOH, HCl
May and Baker Ltd.
Hopkin and Williams Ltd.

d. Fine Chemicals

indole-3-acetic acid, indole-3-aldehyde,				
indole-3-carboxylic acid	Sigma	Chemical	Company	Ltd.
<u>N-methyl-N-nitro-p-toluene</u> sulphonamide				
<u>N,O</u> -bis-(trimethylsilyl)				
trifluoroacetamide	Pierc	e & Warri	ner Ltd.	

e. Radiochemicals

All radioactively labelled chemicals were obtained from Amersham International Ltd.

compound	specific activity
$IAA-1-^{14}C$	2.18 GBq mmo1 ⁻¹
IAA-2- ¹⁴ C	2.11 GBq $mmol^{-1}$
Na ¹⁴ CO ₃	2.21 GBq mmol^{-1}
¹⁴ C-hexadecane	18.6 KBq g ⁻¹

Solutions of IAA

Unlabelled IAA was dissolved in a small volume of methanol and diluted to the required concentration with distilled water. For growth experiments the same amount of methanol (i.e. 2 mm^3 per 100 cm³ of solution) was added to each IAA concentration and to the control. Aqueous solutions of IAA were stored in darkness at 4°C for no more than 48h.

 14 C-IAA was stored in methanolic solution at -15°C. Aqueous solutions were made up when required.

3. EXPERIMENTAL PROCEDURES

(a) Metabolism experiments

Incubation

50mm diameter glass petridishes containing 2 cm³ aqueous IAA solution. Coleoptile segments were incubated in groups of 20 or 40. All experiments were carried out in darkness at 22°C. After incubation the tissue from each dish was washed 5 or 6 times with 25 cm³ distilled water for 1min each, by which time the efflux of radioactivity had reached a steady value (see Fig. 5).

Blocks of agar containing IAA for experiments E and G were prepared as follows. Approximately 2 cm³ of a hot 1.5% agar solution with 100 μ l of methanol containing IAA-2-¹⁴C was poured into moulds (25 x 21 x 3 mm) and allowed to set. The agar was divided into 8 blocks of approximate dimensions 12 x 5 x 3 mm, each of which was used to supply IAA to 5 root or coleoptile segments.

Preparation of Extracts

Segments were extracted overnight with redistilled methanol at 4°C (25 cm³ methanol per 50 segments). Extracts were then filtered. Two alternative methods of sample preparation were used:

1. Small root extracts were reduced to dryness using a Buchii rotary evaporator. After being redissolved in 1 cm³ of methanol and centrifuged at 10,000 x g, samples were stored at -15° C in darkness until required for analysis.

2. The majority of coleoptile extracts, and root extracts from more than 100 segments were reduced <u>in vacuo</u> to the aqueous phase. They were then diluted with ammonium acetate buffer (pH 3.5, 20 mol m⁻³) and loaded onto C_{18} Sep-pak cartridges (Waters Associates Inc.) using a 5 cm³ glass syringe. Cartridges were wetted prior to use by pumping through 5 cm³ methanol followed by 2 cm³ buffer. Radioactive compounds in the extracts were eluted from the cartridges with 60% methanol in ammonium acetate buffer (2 cm³). The methanol was blown off under a stream of N₂ and the samples freeze dried. Extracts were redissolved in 1 cm³ methanol for storage until required for analysis.

Immediately prior to HPLC analysis all samples were blown dry under N2



Fig. 5. Radioactivity present in successive washings of $\frac{\text{root segments incubated for 2h in IAA-2-}^{14}C}{(10^{-3} \text{ mol m}^{-3})}$

and dissolved in 100 mm^3 10% methanol in ammonium acetate buffer (pH 3.5, 20 mol m⁻³).

(b) Growth Experiments

Incubation

Root segments,5mm in length,were incubated in groups of 10,in 50mm diameter glass petridishes containing 2 or 5 cm³ aqueous IAA or control solution. Incubation usually took place in darkness at 22°C. Roots in some experiments were incubated in the laboratory at 18-20°C.

Measurement of Growth

Tissue segments were measured using the shadowgraph method. Sections were placed on a glass plate and a photographic enlarger used to produce images of x5 magnification which were then measured using a ruler with 0.5mm divisions. Growth of intact plants was determined by placing spots of indian ink lmm and 6mm from the root tip. The distance between the spots after incubation was measured using a ruler.

4. LIQUID SCINTILLATION SPECTROMETRY

The amount of radioactivity present in samples was assayed using Packard model 3380 liquid scintillation spectrometers. Samples were placed in 20 cm³ plastic vials with 10 cm³ scintillation cocktail. They were routinely counted for 5min.

Scintillation cocktails

A. For non-aqueous samples:

4 g cm⁻³ PPO (2,5-diphenyloxazole) in toluene

B. For aqueous samples:

2 parts A plus 1 part Triton-X 100

C. For both aqueous and non-aqueous samples in later experiments:

Packard Emulsifier-Scintillator 299

- D. For trapping 14 CO₂ in sample oxidiser
 - 7g PPO 50 cm³ H_2O 220 cm³ methanol 330 cm³ 2-phenylethylamine 400 cm³ toluene

Counting Efficiency

The instruments used comprised an external standard radiation source, the counting efficiency for which was automatically recorded for each sample. Actual counting efficiency for 14 C was obtained from a calibration graph in which the counting efficiency for a known quantity of 14 C hexadecane was plotted against the A.E.S. (Automatic External Standard) ratio. A quantity of around 20 mm³ of labelled hexadecane was weighed accurately in a scintillation vial and 10 cm³ of scintillation cocktail added. The number of counts and the A.E.S. ratio were recorded as an appropriate "quenching agent" was added dropwise to the vial. Separate calibration curves were prepared for different scintillants. Three quenching agents were used: water, a methanolic root extract and a methanolic coleoptile extract.

Oxidation of Tissue for Radioassay

After methanolic extraction, the quantity of radioactivity remaining in the tissue was measured. The plant material was combusted using an Intertechnique IN 4101 Sample Oxidiser. The ${}^{14}\text{CO}_2$ produced was trapped by phenylethylamine in the scintillation cocktail and the radioactivity assayed by liquid scintillation spectrometry.

5. THIN-LAYER CHROMATOGRAPHY

Glass TLC plates, precoated by Camlab (Macharey-Nagel and Co.) with a 0.25mm layer of silica gel were used. Chromatograms were developed for a distance of 15cm in one of three solvent systems:

1. Methyl acetate : propan-2-o1 : 25% ammonia (45 : 35 : 20 v/v)

2. Chloroform : methanol : glacial acetic acid (75 : 20 : 5 v/v)

3. Chloroform : ethyl acetate : 90% formic acid (35 : 55 : 10 v/v)

Solvents used for TLC were either redistilled in glass, or Analar grade. The position of radioactive compounds was found by scanning the plates using a Panax thin-layer radiochromatogram scanner. Alternatively the silica gel from each of ten zones was scraped off, and the quantity of radioactivity determined by liquid scintillation spectrometry.

6. HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY

The HPLC system used is shown in diagrammatic form in Fig. ^{5a.} Solvents were delivered at flow rates of either 1 or 0.75 cm³ min⁻¹ by an Altex Model 332 gradient liquid chromatograph (Altex Scientific Inc.). A Shandon reversephase column (I.D. 250 x 5mm) packed with 5 μ m ODS Hypersil (Shandon Southern Products Ltd.) was used. Samples were introduced off column <u>via</u> a Perkin-Elmer Model 7105 sample valve. The mobile phase consisted of a mixture of HPLC grade methanol (Rathburn Chemicals Ltd.) and ammonium acetate buffer (pH 3.5, 20 mol m⁻³) The buffer was cleaned prior to use by passing through a 250 x 10mm I.D. column packed with 10 μ m ODS silica gel. Three solvent gradients were used:

- A. 10 to 60% methanol over 20 min
- B. 10 to 60% methanol over 30 min
- C. 50 to 100% methanol over 20 min.

The percentage of methanol was then increased to 100% over 5min and the column cleaned for a further 10min. The solvent composition was returned to the initial value and the column equilibrated for 10min prior to injecting the next sample.

Two detectors were used in series. The column was connected to a Perkin-Elmer 650-10S Fluorescence Spectrophotometer fitted with a 16 nm^3 flow cell. The excitation and emission wavelengths were set at 280 and 350 nm respectively (the fluorescence λ_{max} for IAA; Crozier <u>et al.</u>, 1980) Eluant leaving the fluorimeter was mixed with scintillant and passed through a 400 mm.³ spiral glass flow cell in an adapted ICN Tracerlab manual scintillation spectrometer (see Reeve and Crozier, 1977). The scintillant comprised 10g 2,5-diphenyloxazole, 330 cm³ Triton X-100, 670 cm³ distilled xylene and 150 cm³ methanol. A 3:1 scintillant/eluant ratio was compatible with all HPLC mobile phase conditions and gave a <u>circa</u> 65% counting efficiency for ¹⁴C solutes.

During analysis of unlabelled indole standards the fluorescence detector

Fig, 5a. HPLC Apparatus



was substituted with a Pye Unicam LC 871 UV absorbance detector.

The HPLC column gave 15,000 theoretical plates with an ODS test mixture comprising benzamide, acetophenone, benzophenone and biphenyl. With IAA this value was considerably lower (<u>circa</u> 4,000 plates). This is the result of ionisation of a proportion of the molecules at the surface of the stationary phase.

The resolution of radioactive peaks was reduced due to mixing in the connecting tubing (which was kept to a minimum) and the fluorimeter flow cell, and also due to the size of the radioactivity monitor flow cell. The width of the fluorescence peak attributable to IAA, eluting isocratically in 50% methanol at a flow rate of 1 cm³ min⁻¹ was approximately 28s compared with 57s for the radioactive peak.

7. PREPARATION OF DERIVATIVES

(a) Trimethylsilylation

Samples were dried thoroughly, dissolved in 50 or 100 μ l <u>N,O</u>-bis-(trimethylsilyl) trifluoroacetamide and heated at 60°C for 2h. This procedure was shown by McDougall (1978) to give 99% yield of the bistrimethylsilyl derivative of IAA.

(b) Methylation

Compounds were methylated using an etherial solution of diazomethane prepared using a modification of the method of Schlenk and Gellerman (1960). Approximately 0.5g <u>N-methyl-N-nitro-p</u> toluene sulphonamide was placed in a 100 cm³ conical flask with side arm attachment. To this was added 10 cm³ diethyl ether, 1 cm³ methoxyethanol, and 3 cm³ concentrated potassium hydroxide. The flask was stoppered and heated in a hot water bath. The open end of the side arm was placed under <u>ca</u>. 10 cm³ diethyl ether in a flask cooled in ice. On heating the reaction mixture, diazomethane gas and

diethyl ether distilled over and dissolved in the cold ether. When an intense yellow colour was observed, the solution was ready for use.

Samples were dissolved in $100 \,\mathrm{mm}^3$ methanol. After addition of 0.5 cm³ of diazomethane solution they were stored at 4°C for at least 2h before blowing dry under a stream.of nitrogen. This method converted ¹⁴C-IAA to a single peak which co-chromatographed with IAA-methyl ester on the HPLC column.

8. COMBINED GAS CHROMATOGRAPHY-MASS SPECTROMETRY

The GC-MS instrumentation comprised a single beam, double focusing AEI-MS 30 mass spectrometer coupled <u>via</u> a single stage glass jet separator to a Pye 104 gas chromatograph. For selected ion monitoring an AEI Multipeak Selectro WF-055 multipeak monitor acted as an accelerating voltage alternator. Up to 6 ions could be monitored at one time. The tuning of the instrument was checked using a perfluorokerosene standard which gave a reference peak at $\underline{m/e}$ 181. Gas chromatograph and mass spectrometer operating conditions are stated separately for each experiment.

For selected ion-monitoring detection of IAA the instrument was tuned to detect ions of $\underline{m/e}$ 202, the most prominent peak in the mass spectrum of bis-trimethylsilyl IAA (see Fig. 6).

9. UV SPECTROPHOTOMETRY

UV absorbance spectra were obtained using a Unicam SP 8000 UV recording spectrophotometer. Samples were dissolved in methanol, placed in lcm quartz cuvettes and scanned from 200-450 nm.

.

.

$$\begin{bmatrix} CH_{2}CO_{2}Si(CH_{3})_{3} \\ Si \\ (CH_{3})_{3} \end{bmatrix}^{\frac{m}{e} 319} (M^{+})$$

,

$$\underline{m}/\underline{e}$$
 304 = $(\mathbf{M} - \mathbf{CH}_3)^*$

$$\underline{m/e} \quad 202 =$$

$$\underline{m}/\underline{e} \quad 73 = \overset{\dagger}{\text{Si}} (CH_3)_3$$

$$\underline{m}/\underline{e} \quad 75 = (CH_3)_2 \quad Si = OH$$



10. SYNTHESIS OF IAA DERIVATIVES

Conjugates of IAA with glycine, valine and alanine

Conjugates of IAA with the amino acids glycine, valine and alanine were prepared as follows using a modification of the methods of Mollan <u>et al</u>. (1972) and Hangarter <u>et al</u>. (1980). The tetramethylammonium salts were prepared by adding 50 or 100 mmol of each amino acid to 25% aqueous solutions of tetramethylammonium hydroxide. After dissolving, using additional water when necessary, the salts were freeze dried. Each compound was then suspended in dimethyl formamide (5 cm³) and the mixed anhydrides prepared. IAA (0.88g) was dissolved in a mixture of tetrahydrofuran (20 cm³) and triethylamine (0.7 cm³) and stirred in a flask cooled in an ice bath. Over a period of 30s, ethyl chloroformate (0.48 cm³) was added dropwise with rapid stirring. Stirring was continued for 10 min. The white precipitate was then filtered off and the filtrate added dropwise to the suspension of the amino acid salt. Stirring was continued for 2h at room temperature.

Water (30 cm^3) and diethyl ether (30 cm^3) were added to the reaction mixture and the whole was shaken. The ether layer was separated off and the aqueous phase acidified to pH 5 with HCl. This was then extracted twice with ether before further acidification to pH 2.

The glycine and L-valine conjugates were precipitated on acidification to pH 2. The L-alanine conjugate was obtained by freeze drying the acidified (pH 2) aqueous phase. The residue was dissolved in methanol or ethyl acetate, filtered and evaporated to dryness. All products were analysed by GC-MS and found to contain the amino acid conjugate plus some unreacted IAA (see Experiment H.8).

Oxindole-3-acetic Acid

This was prepared according to the methods of Hinman and Bauman (1964) and Reinecke and Bandurski (1981). N-Bromosuccinimide (0.712 g) was added to a solution of IAA (0.7 g) in tertiary butyl alcohol (25 cm³) at room temperature and stirred. After 2h the solution was concentrated under reduced pressure at room temperature, to give a thick syrup. Water (30 cm³) was then added and the mixture extracted three times with ethyl acetate (8 cm³). The combined extracts were washed with saturated salt solution, dried over Na_2SO_4 and evaporated at room temperature. Residual ethyl acetate was entrained by repeated evaporation with acetone <u>in vacuo</u>. The residue was taken up in a mixture of acetone (0.5 cm³) and benzene (8 cm³). This was left for 2 days at -20°C. The resulting yellow solid was filtered and recrystallised twice from acetone/benzene before drying in a vacuum desiccator. The identity of the product was confirmed by mass spectrometry (see Experiment H.8).

RESULTS

This part of the thesis is divided into two sections:

- 1. An investigation of the metabolism of IAA in coleoptile and root tissues of dark-grown Zea mays seedlings.
- A study of the effect of exogenous IAA on the growth of these tissues.

The aim of the first section was to assess quantitatively the metabolic fate of exogenous IAA, supplied to root and coleoptile segments. Reversephase high-performance liquid chromatography combined with a continuous radioactivity monitor was used for most analyses, and was compared with the older technique of thin-layer chromatography. A number of methods, including combined capillary-column gas chromatography-mass spectrometry were used to obtain information on the chemical structures of the products. Measurements of the uptake of IAA and its effect on growth confirmed that the amount of phytohormone entering the tissue was comparable to the endogenous content and was physiologically active.

SECTION 1 - METABOLISM OF IAA

A. Comparison of Separatory Techniques Used in the Analysis of IAA Metabolites in Plant Extracts

Traditionally, the techniques of thin-layer and paper chromatography have been used extensively in the separation and characterisation of IAA metabolites (e.g. Tuli and Moyed, 1967; Davies, 1972; Epstein and Lavee, 1977b; Magnus <u>et al.</u>, 1982). Recently various forms of high-performance liquid chromatography (HPLC) have been used to analyse phytohormones in plant extracts (e.g. Durley and Kannangara, 1976; Crozier and Reeve, 1977; Sweetser and Swartzfager, 1978) and also to purify their metabolites (e.g. Summons <u>et al.</u>, 1980; and Reinecke and Bandurski, 1981). HPLC has the advantage of potentially much greater resolution than thin-layer or paper chromatography. The following three experiments illustrate the usefulness of a reverse-phase HPLC system in the separation of IAA metabolites, when compared with thin-layer chromatography (TLC).

A.1 TLC Analysis of IAA Metabolites

Methanolic extracts containing the products of ${}^{14}C$ -IAA metabolism were prepared from groups of 50 root segments which had been incubated in aqueous solutions of IAA-2- ${}^{14}C$ ($10^{-2}mol m^{-3}$) for 2h in darkness. After drying <u>in</u> <u>vacuo</u> and redissolving in a small volume of methanol, extracts were spotted onto silica gel TLC plates and developed in three different solvent systems: 1. Methyl acetate : propan-2-ol : 25% ammonia (45 : 35 : 20 v/v) 2. Chloroform : methanol : acetic acid (75 : 20 : 5 v/v)

3. Chloroform : ethyl acetate : 90% formic acid (35 : 55 : 10 v/v)

The positions of radioactive compounds were detected using a thinlayer radiochromatogram scanner (Fig. 7). The number of distinct peaks varied from two to five according to the solvent system. The most



Fig. 7. TLC analysis of IAA-2-¹⁴C metabolites from extracts of Zea mays roots. Root segments were incubated in IAA-2-¹⁴C (10⁻² mol m⁻³) for 2h. Traces represent typical results from 3 replicate experiments.


Fig. 7. Cont'd

.

prominent peak always had the same Rf value as IAA.

A.2 HPLC Analysis of IAA Metabolites

Extracts were prepared in the same manner, from root segments which had been incubated with IAA-2- 14 C (10⁻³mol m⁻³) for 2h. Samples, dissolved in 100 mm³ of 10% methanol in ammonium acetate buffer (pH 3.5, 20 mol m^{-3}), were injected onto the reverse-phase HPLC column. A solvent gradient of 10 to 60% methanol was run over 30 min. Fig. 8 shows the profile of radioactive peaks eluting from the column (typical result from 6 replicate experiments). A large peak with a retention time identical to that of IAA was apparent, together with at least 11 metabolite peaks all eluting before IAA. When six identical extracts were analysed during a single day these 12 peaks were consistently present and the variation in retention time of each peak was small (Table 5) giving a strong indication that the same compounds were consistently present. The metabolite peaks shown in Fig. 8 were thus labelled 1 to 11 for identification. In order to achieve an objective comparison of extracts from different experiments, in the absence of definitive identification of compounds, samples from roots incubated in 10^{-3} mol m⁻³ IAA for 2h were used as standards for numbering of peaks throughout the thesis. Peaks 7 and 8, and 1 and 2 eluted close together and were not always resolved. When this was the case the peaks were labelled 7/8 and 1/2.

A.3 Use of HPLC and TLC in a Preparative Mode

Although the column used for HPLC was of a narrow bore it could be used in a preparative mode to purify metabolite peaks. Samples of peaks 4-11 and IAA were collected after separation on the column, dried and rechromatographed. Peaks 7, 8, 9, 10 and IAA remained unchanged (Table 6). Peaks 4 and 5 appeared to be interconverted, while a proportion of radioactivity from peak 6 appeared with the retention time of peak 7. Some of peak 11 was converted to a new product.



Fig. 8. HPLC Analysis of a Zea mays root extract containing metabolites of IAA-2-¹⁴C. Root segments were incubated in 10⁻³ mol m⁻³ ¹⁴C-IAA for 2h and extracted in methanol. The trace represents a typical result from 6 replicate experiments. Solvent gradient : 10-60% methanol over 30 min. Flow rate : 0.75 cm⁻¹. Detector : homogeneous radioactivity monitor; 30 cps full scale deflection; 10s time constant.

l4 _{C-IAA metabolites in}
ks representing ¹
f radioactive pea
mes (min) o
m in retention tin
5 : Variatio
Tab le

extracts of Zea mays roots, chromatographed on the same day

;			Replic	a t e			Mean ± standard
Peak No	1	2	34	4	5	6	error
П	5.7	6.0	6.2	5.8	6.2	5.9	6.0 ± 0.08
7	7.4	7.2	7.0	6.8	7.2	6.8	7.1 ± 0.10
ю.	11.2	11.4	11.0	10.8	11.0	10.6	11.0 ± 0.12
4	13.4	13.2	12.4	12.7	12.6	12.3	12.8 ± 0.18
Ω	14.7	14.8	14.0	14.1	14.0	13.8	14.2 ± 0.17
6	15.9	15.8	15.3	15,3	15.3	I5.2	15.5 ± 0.12
7	17.6	17.4	16.8	17.0	17.0	16.6	17.1 ± 0.15
8	18.2	18.2	17.6	17.6	17.7	17.6	17.8 ± 0.12
ი	21.0	21.0	20.1	20.4	20.4	20.2	20,5 ± 0,16
10	24.3	24.3	23.4	23.8	23.8	23.7	23.9 ± 0.14
11	25.8	25.8	25.0	25.2	25.2	25.1	25.4 ± 0.15
IAA	32.1	32.1	31.2	31.5	31.4	31.5	31.6 ± 0.15

Table 6 : Degradation of IAA and its metabolites during preparativepurification by HPLC.Peaks 7 and 8 were not completelyresolved and had to be collected together.

Peak No	Initial T _R on HPLC (min)	T _R during rechromatography after purification by HPLC (min)
4	9,9	9.4 (44%) + 11.1 (56%)
5	11.2	9.8 (28%) + 11.2 (72%)
6	12.6	12.4 (56%) + 14.0 (44%)
7/8	14.2 + 15	14.4 + 15.1
9	17.8	18.1
10	18,5*	18.7*
11	24	24.6 (69%) + 25.6 (31%)
IAA	30	29.6

* Peak 10 was analysed on a separate day

On the other hand, TLC was found to cause substantial sample degradation. Individual peaks, collected from the liquid chromatograph were loaded onto silica gel TLC plates, which were then developed in chloroform : ethyl acetate : 90% formic acid (35 : 55 : 10 v/v). Zones of radioactivity were collected and eluted, and the compounds rechromatographed on the HPLC. In each case at least some degradation had occurred. Peaks 4, 10 and IAA had undergone a small amount of breakdown, while peaks 5, 6, 7, 8 and 9 appeared to have changed completely. Peak 11 was not analysed.

A. Conclusions

The resolving power of HPLC is of considerable value in separating the large number of IAA metabolites present in methanol extracts of <u>Zea mays</u> root segments which had been incubated in IAA-2-¹⁴C. Although it is possible that a single peak on the HPLC trace might represent more than one compound, TLC was clearly unable to separate individual metabolites. Furthermore, while degradation of compounds occurred during TLC separation, most metabolites could be recovered unchanged from the HPLC column (degradation of metabolites during TLC analysis might be reduced in other solvent systems). The reverse-phase HPLC column was ideal as all the metabolites appeared to be more polar than IAA.

)

B. Quantitative Fate of IAA-2-¹⁴C Supplied to Zea mays Root Segments

In the past, metabolites of IAA which were analysed sometimes represented only a small fraction of the IAA supplied to the tissue (e.g. Tuli and Moyed, 1967; Magnus et al., 1971). The aim of this study was to quantify the amount of IAA metabolised and to record the relative production of the whole spectrum of IAA-2-¹⁴C metabolites produced by Zea mays root segments. Control experiments were designed to confirm that the reactions were carried out by the living plant cells and to assess the effect of experimental conditions on the metabolism pattern. To answer the possibility that the compounds might represent products of a detoxification process unrelated to the normal pathways of endogenous IAA metabolism, the amount of IAA taken up by the roots was compared with published measurements of endogenous quantities (Bridges et al., 1973). The effect of varying the amount of IAA supplied, on the pattern of metabolism was also studied.

B.1 Metabolism of IAA-2-¹⁴C by Root Segments during a 2h Incubation Period

Groups of 50 root segments from 3 to 4-day-old, dark-grown Zea mays seedlings, were incubated in aqueous solutions of IAA-2-¹⁴C (10^{-3} mol m⁻³; 2h; darkness). All manipulations of plant material were carried out under dim green light. After following the washing procedure, the tissue was extracted overnight with methanol and samples prepared for HPLC analysis. When the column was routinely cleaned with tetrahydrofuran, it was possible to analyse relatively crude extracts, thereby eliminating the need for lengthy purification procedures, during which products might be lost. Root extracts were simply filtered, dried <u>in vacuo</u>, redissolved in 1 cm³ methanol and centrifuged at 10,000 x g (method 1). Samples were stored at-15°C in this form until required. Prior to analysis, each sample was dried under a stream of N₂ and redissolved in 100 mm³ of 10% methanol in

ammonium acetate buffer (pH 3.5, 20 mol m^{-3}). The average dry weight of an extract prepared in this manner was 25.4 mg. Aliquots for measuring radioactivity were taken from extracts at two stages: after filtration and after drying in vacuo. Substantial and variable losses (up to 36% of the sample; Table 7) occurred during drying in vacuo. Large losses at this stage were also noted by Mann and Jaworski (1970). The amounts of radioactivity lost during drying under N_2 and redissolving in 100 mm $^3\,\mathrm{10\%}$ methanol in buffer, and also due to centrifugation, were small. Radioactivity present in the incubating solutions at the end of the experiment, and in the washings was also measured by counting aliquots. The amount of label remaining in the roots after methanol extraction was assessed by combusting the tissue in a sample oxidiser and counting the $^{14}CO_2$ trapped by phenylethylamine.

The fate of IAA-2-¹⁴C supplied to the root segments is presented in Table 7 (column 1). An average of 31% of the radioactivity supplied (44-45 Bq segment⁻¹) was taken into the tissue. Methanol extraction removed 93% of the ¹⁴C from the roots (similar extraction efficiencies were found in all subsequent experiments). Losses of radioactivity during incubation were small (4.2% on average).

Fig. 9 shows the profile of radioactive compounds eluting from the HPLC column during the analysis of a typical methanolic root extract. Twelve peaks were consistently present (Table 8). The largest in each case co-chromatographed with IAA and also produced a fluorescence peak when the excitation and emission wavelengths of the fluorescence spectrophotometer were set at 280 and 350 nm respectively. Further evidence for the identity of this peak is presented in experiment H. After 2h incubation, an average of 46% of the radioactivity present in the methanol extracts remained associated with the IAA peak (calculated from the peak integrator on the chart recorder). Eleven metabolite peaks were present in each replicate and were numbered for identification as described in experiment A.2. The largest was always peak 10, which accounted for a mean of 25% of the radio-

	(50 segments) were incub	ated for $2h$ in 10^{-3} mol	m ⁻³ IAA-2- ¹⁴ C.	Extracts designated (S)	were purified using
	Sep-pak C ₁₈ cartridges.	Radioactivity is vy	ressed in Bq. Th	e experiment was carri	ed out in triplicate.
TRE	ATMENT	Root segments cut in dim green light	Root segments c in the laborato	rt Sterile root ry segments	Dead root segments
• ~	Initial Radioactivity	7130 7190 7530	7210 7840 7900	32,000 21,400 23,500	6070 6350
ν,	Radioactivity remaining in incubating solution	3430 3620 3660	3500 3430 3490	17,000 13,100	3410 3300
ч.	Radioactivity in methanol extract	2110 2060 2130	2230 2250 2210	5,340 2,190 2,260	241 156
4.	Radioactivity remaining in tissue after extraction	154 161 132	171 177 153	161 55 221	14 12
л.	Radioactivity in washings	1180 1110 1160	1130 1600 1370	1	2520 2360
6.	Final total of radioactivity	6880 6950 7080	7030 7450 7220	3	6190 5840
7.	Loss of radioactivity during incubation	244 240 447	175 387 677	1	518

Root tissues

Analysis of the Fate of IAA-2-¹⁴C supplied in Aqueous Solution to Zea mays Root Segments.

Table 7 :

as a if 1	3.4% 3.3% 5.9%	2.4% 4.9% 8.6%	1	- 8.2%
(mg)	40.1 38.4 39.7	33.6 36.5 37.5	62.6 30.7 50.1	32.3 32.5
ity	45 44 45	48 47	ł	5.1 3.4
×	93% 93% 94%	93% 93% 94%	97% 98% 91%	95% 93%
hanol ation	1440 1310 1370	1900 1680 1830	- 1640(S) 2020	229 149
	68% 64% 64%	85% 75% 83%	- 75% 89%	95% 96%
le of ifter	2760 2940 3030	2820 2800 3020	I	2780 2920
	80% 81% 83%	81% 82% 87%	3	82% 82%



Table 8 : IAA Metabolites pro	esent in methanol extracts	s of root segments incub	ated in IAA-2- ¹⁴ C for 2	<u>i</u> . Metabolite peaks in
extracts of non-st	erile roots were numbered	I-11 (see Fig. 8) and u	sed as standards for th	e other extracts (1).
The experiment was	carried out in triplicate	ċ		
TREATMENT	Root segments cut in dim green light	Root segments cut in the laboratory	Sterile root segments	Dead root segments
Radioactive peaks in methanol extracts in order of decreasing heigit (1)	IAA,10,1,7,9,6,4,5,8, 11,2,3 IAA,10,5,6,1,9,7,3,4, 8,2,11	IAA,10,9,7,6,5,4,3, 1,11,8,2 IAA,10,9,6,1,5,7,8, 4,3,2,11	10,8,IAA,2,4,5,7,1, 11,9,3,6 7,8,5,4,10,IAA,6, 9,2,1,3	IAA only IAA only
	IAA,10,1,4,7,3,8,11,5, 9,6,2	IAA,10,9,6,7,5,1,4, 3,8,2,11	10,IAA,8,7,5,4,9,1, 11,3,6,2	
% radioactivity in methanol extract, associated with IA4 peak	49 40 48	34 29 30	11 11 14	100 100
% radioactivity in methanol extract, associated with peak 10	21 31 23	21 19 18	16 12 23	0
% radioactivity in incubating solution, associated with IAA peak	98 93 97	9 9 5 5	- 6	98 95

activity in each sample. The relative prominence of the other peaks varied.

Samples of the incubating solution were also analysed to determine the amount of IAA breakdown occurring during the experiment. Solutions were dried <u>in vacuo</u> and taken up in 1 cm³ of methanol. Before analysis they were dried again under a stream of N₂ and redissolved in 100 mm³ of 10% methanol in ammonium acetate buffer (pH 3.5, 20 mol m⁻³). An average of 95% of radioactivity remained associated with the IAA peak (Table 8, Fig. 10) indicating that very little degradation had taken place.

IAA Breakdown during Sample Preparation

Three control experiments were carried out to confirm that the compounds obtained were in fact products of IAA metabolism by the plant tissue. To test whether any breakdown of IAA was occurring during the preparation of extracts, six groups of 50 root segments were extracted in methanol, aliquots of IAA-2-¹⁴C were added and samples processed for HPLC analysis. Samples 1-3 were prepared according to method 1,while Sep-pak C₁₈ cartridges (method 2) were used to purify samples 4-6. Typical HPLC traces from each group are presented in Fig. 11. The amount of breakdown in samples 1-3 was negligible although a small amount of peak 10 was observed in samples 4-6. Losses of radioactivity during sample preparation varied between 17 and 33% (Table 9) and were similar to those for extracts containing metabolites of IAA-2-¹⁴C.

Metabolism of IAA by Killed Root Segments

Groups of 50 root segments which had been boiled for 2min were incubated for 2h in aqueous solutions of IAA-2-¹⁴C ($10^{-3} \text{ mol m}^{-3}$). After following the washing procedure, the tissue was extracted with methanol and samples prepared for HPLC analysis using method 1. No metabolism of IAA was apparent (Fig. 12; Table 8). A single radioactive peak with the same retention time as IAA was present and coincided with a peak for fluorescence.



Fig. 10. HPLC analysis of radioactive compounds remaining in the bathing solution after a 2h incubation of root segments in an aqueous solution of IAA-2-14C (10⁻³mol m⁻³). The trace represents a typical result from 3 replicate experiments. Solvent gradient : 10-60% methanol over 30 min. Flow rate : 0.75 cm min⁻¹. Detector : homogeneous radioactivity monitor; 30 cps full scale deflection; 10s time constant.



Fig. 11. Control experiment to test for IAA degradation during sample preparation. IAA-2-14C was added to methanolic root extracts and samples prepared for HPLC analysis. (a) Extracts prepared by filtration and drying in vacuo (method 1). (b) Samples purified using Sep-pak C₁₈ cartridges (method 2). Each trace represents a typical result from 3 extracts. Solvent gradient : 10-60% methanol over 20 min. Flow rate : 1 cm min⁻¹. Detector : homogeneous radioactivity monitor; 30 cps full scale deflection; 10s time constant. Table 9 :Loss of IAA-2-14C during preparation of control root extractsfor HPLC analysis.Samples 1-3 were filtered, dried in vacuo,redissolved in 1 cm3 methanol and centrifuged (method 1) whilesamples 4-6 were purified using Sep-pak C18 cartridges (method 2).

REPLICATE	1	2	3	4	5	6
1. Initial radioactivity in extract (Bq)	1180	1150	1100	759	859	821
2. Radioactivity in extrac after preparation for HPLC analysis (Bq)	ct 784	885	849	500	614	679
3. 2 expressed as a percentage of 1	67%	77%	76%	66%	71%	83%



Fig. 12. Metabolism of IAA-2-¹⁴C by boiled root segments (2h incubation). The trace represents a typical result from 4 replicate experiments. Solvent₃gradient : 10-60% methanol over 30 min. Flow rate : 0.75 cm min⁻¹. Detector : homogeneous radioactivity monitor; 30 cps full scale deflection; 10s time constant.

Efficiencies of extraction and preparation procedures were comparable to those for living tissue although uptake of radioactivity was reduced to <u>circa</u> 10% (Table 7).

B.2 Metabolism of IAA by Sterile Roots

Libbert al (1966) demonstrated that surface bacteria are often capable of large-scale production of IAA. Much of the earlier research on the biosynthesis of IAA in higher plants is of questionable significance because account was not taken of the contribution of epiphytic microorganisms. On the other hand, Libbert and Rische (1969) have published results indicating that degradation of IAA by bacteria on pea plants is negligible. To confirm that the metabolism of IAA by Zea mays seedlings observed in previous experiments was in fact due to plant cells, the fate of IAA-2-¹⁴C supplied to sterile roots was studied.

Sterile plants were grown in groups of 10 in glass petridishes. Roots from each dish were incubated in IAA-2- 14 C (2cm³ of 10⁻³ mol m⁻³) for 2h, washed and extracted separately. The sterility of plants in each dish was tested by incubating portions of root for 2 weeks on potato dextrose agar and nutrient agar. Extracts from plants for which tests for fungal and bacterial contamination proved negative were then combined, and samples prepared for HPLC analysis according to methods 1 or 2.

The metabolism pattern observed in three replicate experiments (Fig. 13; Table 8) was closely similar to that for non-sterile roots. The majority of peaks 1 to 11 were present in each case, strongly indicating that metabolism of IAA was taking place in the plant tissue. The amount of IAA remaining in the extracts was reduced (an average of 12% compared with 31% for non-sterile root segments cut in the light). Uptake, extraction efficiences and losses of radioactivity were all similar to those in experiments with non-sterile tissue (Table 7).



Fig. 13. Metabolism of IAA-2-¹⁴C by sterile root segments (2h incubation). The trace represents a typical result from 3 replicate experiments. Solvent gradient :₃ 10-60% methanol over 30 min. Flow rate : 0.75 cm min⁻¹. Detector : homogeneous radioactivity monitor; 30 cps full scale deflection; 10s time constant.

B.3 Effect of Exposure to Light on the Metabolism of IAA by Zea mays Root Segments

Root segments for sterile experiments, and for preparation of largescale extracts, were cut in the laboratory. The effect of exposure to light on the metabolism of IAA was therefore studied.

Groups of 50 root segments were incubated in aqueous IAA $(10^{-3} \text{ mol m}^{-3})$ for 2h. Plant material was both grown and incubated in darkness, but section cutting was performed in the laboratory.

Analysis of methanolic extracts, prepared using method 1, revealed a pattern of metabolism closely similar to that obtained in experiment B.1 with root segments cut under dim green light (Table 8; compare Figs. 8 and 9). The same peaks were present in each of three replicates although the proportion of radioactivity associated with the IAA peak was somewhat reduced (31% compared with 46%). The exposure to light had no significant effect on uptake, extraction efficiency or the amount of radioactivity lost during incubation (Table 7).

B.4 Effect of Varying the IAA Concentration in the Incubating Solution on its Metabolism by Root Segments

The rationale for this experiment was two-fold. Firstly, when applying plant growth substances to plant tissue the question always arises as to how much apparent metabolism actually represents detoxification of abnormally large amounts of the exogenous compound. A partial answer to this question might be gained by examining the metabolism of different external concentrations of IAA. Secondly, the metabolism of higher concentrations of IAA than those previously used was investigated with a view to preparing large extracts for studies on the chemical structures of metabolites.

Root segments (cut in the laboratory) were incubated in groups of 50 in four different concentrations of IAA-2- 14 C (10⁻⁴, 10⁻³, 10⁻², 10⁻¹ mol m⁻³) for 2h. After following the washing procedure, tissue was extracted overnight

Samples were prepared according to method 1 and analysed by in methanol. HPLC. The pattern of metabolism in segments incubated at the three lowest concentrations was similar, with the majority of peaks 1-11 present in each case (Fig. 14; Table 10). The proportion of radioactivity in extracts, remaining associated with the IAA peak was also similar (25%, 25% and 29% for incubating solutions of 10^{-4} , 10^{-3} and 10^{-2} mol m⁻³ IAA respectively). The percentage of IAA metabolised in tissue incubated in 10^{-1} mol m⁻³ IAA was reduced and only metabolite peaks 4, 5 and 10 were consistently present. The average uptake of radioactivity from a 14 C-IAA concentration of 10^{-4} mol m⁻³ was equivalent to 2.3 pmol segment⁻¹. Given that the mean fresh weight of a root segment was 17.6mg and that an average of 25% of the radioactivity in the tissue remained associated with the IAA peak, the approximate amount of exogenous IAA present in the roots after 2h incubation could be calculated as $5 \cdot i \mu g kg^{-1}$. This is substantially less than the endogenous IAA content of Zea mays (cv. Giant White Horsetooth) root tips measured as 29 kg^{-1} by Bridges et al. (1973).

B.5 Use of Sep-pak C₁₈ Cartridges in Preparation of Samples for HPLC

When large extracts were prepared for HPLC analysis, it was necessary to employ a purification step which would reduce the dry weight of samples, with minimal loss of IAA metabolites. Sep-pak C_{18} cartridges were found to be suitable for this purpose. An experiment was carried out to record the effect of this procedure on the profile of radioactive peaks observed during HPLC analysis of root extracts containing ¹⁴C-IAA metabolites.

Root tissue (2 groups of 50 segments) which had been incubated in IAA-2- 14 C (10⁻² mol m⁻³) for 2h was washed and extracted in methanol. Extracts were filtered and reduced to the aqueous phase <u>in vacuo</u>. After diluting with 10 cm³ ammonium acetate buffer (pH 3.5, 20 mol m⁻³) samples were loaded onto wetted Sep-pak cartridges. Radioactivity was eluted with 2 cm³ 60% methanol in ammonium acetate buffer. Both the buffer passing

Fig. 14. Metabolism of TAA-2-¹⁴C supplied at 4 different concentrations

to Zea mays root segments (2h incubation). (a) 10^{-1} mol m⁻³. (b) 10^{-2} mol m⁻³. (c) 10^{-3} mol m⁻³. (d) 10^{-4} mol m⁻³.

Each trace represents a typical result from 3 replicate experiments HPLC Conditions: Solvent gradient; 10-60% methanol over 30 min. Flow rate; 0.75 cm³min⁻¹. Detector: homogeneous radioactivity monitor; 30cps full scale deflection; 10s time constant.







Table 10:Uptake and MetabcMetabolite peaksFig. 8) and used	olism of IAA-2- ¹⁴ C supplied in extracts of roots suppl as standards for other ext	1 at four concentrations t lied with IAA at 10 ⁻³ mol m tracts (1). The experime	to <u>Zea mays</u> root segments - ³ were numbered 1-11 (se ent was carried out in tri	(2h incubation). se Experiment A.2, iplicate.	
Concentration of IAA in incubating solution	10 ⁻⁴ mol m ⁻³	10 ⁻³ mol m ⁻³	10 ⁻² mol m ⁻³	10 ⁻¹ mol m ⁻³	
Uptake of IAA_1 (pmol segment ⁻¹)	2.7 1.9	17 23 22	129 90 103	1000 740 765	
Radioactive peaks in methanol extract in order of decreasing height (1)	IAA,10,7/8,9,1,6,4,3, 11,5,2 IAA,10,1,9,2,5,6,7/8, 4,3,11	IAA,10,7/8,9,1/2,4,5, 6,11,3 IAA,10,4,5,6,7,9,8, 1,3,2 IAA,10,9,6,7,4,5,8	IAA,10,4,5,7/8,6,9 IAA,10,4,5,8,9,7,6,11 1,3 IAA,10,9,4,7/8,5, 6,1,11	IAA,11,10,4,5 IAA,10,4,5,9,8,7,6 IAA,10,11,9,7/8,5,4	
Percentage radioactivity associated with IAA peak	24 26	23 23 29	33 26 27	62 57 66	
Percentage radioactivity associated with peak 10	20 12	23 20 22	23 19 24	10 11 11 11	
					i .

through the cartridge and the 2 cm ³ of 60% methanol were dried and analysed by HPLC.

A substantial proportion of peak 1 had been lost from the purified extract (Fig. 15a). Most of this was found in the buffer passing through the cartridge (Fig. 15b).

Conclusions

Metabolism of IAA-2-¹⁴C by root segments taken from dark-grown Zea mays seedlings is rapid and extensive. Non-sterile roots, cut under dim green light and incubated for 2h in darkness in 10^{-3} mol m⁻³ ¹⁴C-IAA, had metabolised an average of 54% of the radioactivity extracted with methanol. At least 11 products were formed, one of which, peak 10, was always more prominent than The efficiency of the extraction procedure was high, at least the others. 93%, thus only a small fraction of the radioactivity escaped analysis in However, substantial and variable amounts of label, up to 36% this way. of samples, were lost during drying of extracts in vacuo. Control experiments confirmed that no significant breakdown of IAA occurred during sample Sterile roots yielded an apparently similar range of products preparation. to non-sterile indicating that metabolism was taking place in the plant tissue.

Evidence that the compounds observed probably did not represent products of a detoxification mechanism to eliminate abnormally high hormone levels was obtained from experiments in which the external IAA concentration was varied. The pattern and rate of IAA metabolism did not change significantly when uptake of IAA increased from 2.3 to 107 pmol segment⁻¹. Furthermore, the amount of exogenous IAA present in the tissue after a 2h incubation in 10^{-4} mol m⁻³ IAA-2-¹⁴C was substantially less than published measurements of the endogenous IAA content.

Exposure of root segments to light for a short period resulted in a decrease in the amount of radioactivity associated with the IAA peak, but



Fig. 15. HPLC of root extract purified using Sep-pak C₁₈ cartridges.
(a) Analysis of radioactive compounds eluted from the cartridge in 2 cm³ 60% methanol in ammonium acetate buffer (pH 3.5, 20mol m⁻³). (b) Radioactive compounds passing through cartridge during loading of extract in 10 cm³ ammonium acetate buffer (pH 3.5, 20mol m⁻³). Solvent gradient : 10-60% methanol over 30 min. Flow rate : 0.75 cm³min⁻¹. Detector : homogeneous radioactivity monitor; 30 cps full scale deflection; 10s time constant.

apparent stimulation of IAA metabolism. The increased proportion of IAA metabolised by sterile roots could be due to the increased exposure to light during section cutting in the sterile air-flow cabinet.

C. Time-Course of IAA Metabolism in Root Segments

Research, using root tissues, on the effects, transport and binding of IAA to putative receptor sites has often been performed with little consideration of the metabolism of exogenous IAA during the course of the experiment (e.g. Batra <u>et al.</u>, 1975; Moloney and Pilet, 1981). The results in section B of this thesis suggest that <u>Zea mays</u> root segments are capable of extensive metabolism of IAA. A knowledge of the amount of exogenous IAA remaining in the tissue after varying incubation times is therefore essential. It was also hoped that this study would yield information on the sequence in which metabolite peaks were produced.

Seven groups of 50 root segments (cut in the laboratory) were placed in aqueous solutions of IAA-2- 14 C (10⁻³ mol m⁻³) and incubated for various times : Table 11.

Dish	Time incubated in IAA-2- ¹⁴ C	Time incubated in distilled water	Total incubation time
1 and 2	10 min	-	10 min
3	1 h	-	1 h
4	1 h	1 h	2 h
5	1 h	3 h	4 h
6	1 h	7 h	8 h
7	1 h	23 h	24 h

Table 11 : Incubation protocol for time-course experiment of IAA metabolism

in root segments

Root segments incubated for periods longer than lh were washed thoroughly (Materials and Methods) before transferring to dishes of distilled water. Methanol extracts were prepared for HPLC analysis according to method 1.

Metabolism of IAA was rapid; after incubating segments for only 10min,

an average of 43% of the radioactivity was no longer associated with the IAA peak (Table 12; Fig. 16). The amount of IAA in extracts continued to decline until only around 1% of the radioactivity co-chromatographed with IAA after 24h. The extraction efficiency declined only slightly during the course of the experiment (from 93% for a 10min incubation to 88% after 24^h)

At least 3 and sometimes up to 8 metabolite peaks were apparent after 10 min (the variation appeared to result from difficulty in distinguishing small metabolite peaks from fluctuations in background radiation). The appearance of several products after such a short time indicates that there is probably not a single linear pathway of IAA metabolism. The apparently complex nature of the reaction sequence presented difficulties in interpreting relationships between different peaks. Some consistent trends were visible, The most prominent metabolite at short incubation times was peak 9. however. This declined as there was an increase in the height of peak 10 which became the largest metabolite after 2h and 4h before decreasing in size. After 24h the most prominent compounds were the components of peaks 7 and 8, and peaks 1 and Precise information on the interconversions of individual metabolites, 2. however, could not be deduced from these observations. This might be achieved by supplying labelled IAA metabolites to root tissue and extracting and analysing the radioactive products.

In a single experiment, root tissue remained in IAA-2-¹⁴C for the whole incubation time, up to 24h. Although radioactivity continued to be taken up during this period and there was no significant breakdown of IAA in the incubating solution, the proportion of IAA in the tissue decreased at the same rate as in root segments transferred to water after 1h (Table 13). This would indicate that radioactivity taken up after longer periods was more rapidly metabolised.

Tissue was incubated in	
Uptake and metabolism of IAA-2- ¹⁴ C by Zea mays root segments over a time-course.	
Table 12 :	

÷
products
44
comparison o
for
standards
ЗS
used
and
.
Fig.
1.2,
experiment .
(see
1-11

INCUBATION TIME	10 min	1 h	2 h	4 h	Ч 8	24 h
Uptake of radioactivi' (Bq segment ¹)	ty 8.6 9.6 6.3	19 23 18	19 22 15	18 27 16	19 25 20	21 31 16
Radioactive peaks in methanol extract in order of decreasing height (1)	IAA,9,7/8,4,5, IAA,9,7/8,4,5, I,10,11,8 IAA,9,7/8,10,	IAA,9,7/8,1,10,5, 6,4,11,3 IAA,9,7/8,10,1,5, 4,11,6,3 IAA,9,7/8,10,4,1, 11,5,6	IAA,9,10,7/8,6,1, 11,3,5,4,2 IAA,10,9,7/8,1,6, 4,3,5,2 IAA,10,7/8,6,9	10,2,1AA,7/8,4,1 , 9,6,4,5,11 10,8,2,1AA,3,7,11, 6,9,1,4,5 10,2,1AA,3,11,7/8 9,6,1,4,5	8,2,7,10,3,11,9, 6,1AA,4,5 2,8,10,3,7,11,9,6, 1AA,4,9,5 2,10,7/8,11,3,9,4, 6,1AA,5	2,7/8,3,9,10,11,IAA, 4,5,1b,6 1,7/8,10,1b,3,9,4, 6,11,IAA,5 8,2,7,3,10,11,9, 8,2,7,3,10,11,9,
<pre>% radioactivity in methanol extract associated with IAA p</pre>	64 56 sak 50	28 35 30	31 25 17	14 9 11	2.8 3.5 3.0	110
<pre>% radioactivity in methanol extract associated with peak 1</pre>	10 2.9 10 -	88 87 7	9 11 6	16 20 14	10 14 11	0.04



Fig. 16. Time-course of IAA-2-¹⁴C metabolism by Zea mays root segments. Tissue was incubated in ¹⁴C-IAA (10⁻³ mol m⁻³) for up to 1h, then washed and transferred to distilled water. (a) 10 min incubation, (b) 1h, (c) 2h, (d) 4h, (e) 8h, (f) 24h. Each trace represents a typical result from 3 replicate experiments. ³Solvent gradient: 10-60% methanol over 30 min. Flow rate: 1 cm³ min⁻¹. Detector: homogeneous radioactivity monitor; 30 cps full scale deflection; 10s time constant.



.

Fig. 16 cont'd



Fig. 16 cont'd

ī

92

۰.

Incubation time	Radioactivity taken up by tissue (Bq segment ⁻¹)	% of radioactivity in methanol extract associated with IAA peak
10 min	7.5	58
30 min	20	54
1 h	29	40
2 h	36	29
4 h	76	4
8 h	91	3
24 h	103	No detectable IAA

Table 13 : Metabolism of IAA-2- 14 C supplied to root segments for up to 24h

Conclusions

IAA is metabolised rapidly by <u>Zea mays</u> root tissue, along an apparently non-linear and complex reaction pathway. After a 24h incubation, very little IAA-2-¹⁴C remained in the tissue. Similar results were obtained from experiments in which roots were removed from IAA-2-¹⁴C after 1h and transferred to distilled water, as for tissue incubated in ¹⁴C-IAA for the full incubation time. It appears that IAA-2-¹⁴C taken up after longer periods is more rapidly metabolised.

D. A Comparison of the Metabolism of IAA by Cortical and Stelar Tissues of the Root

Measurements of the endogenous content of IAA in Zea mays roots have indicated that the amount present in the stele, per gram fresh weight of tissue, is approximately 10 times that in the cortex (Bridges <u>et al.</u>, 1973). Wilkins and co-workers (Bowen <u>et al.</u>, 1972; Cane and Wilkins, 1970; Shaw and Wilkins, 1974) have also demonstrated that the polar acropetal transport of IAA in Zea mays roots takes place predominantly in the stele. The products of IAA-1-¹⁴C metabolism in separated cortical and stelar tissues have been examined using TLC (Greenwood <u>et al.</u>, 1973). These studies revealed a slower rate of IAA metabolism in the stele. The major products formed by the two tissues also appeared to differ. It was decided to follow up this work by investigating the metabolism of IAA applied to separated cortical and stelar tissues, using HPLC analysis.

As the apical 5mm of the root cannot be divided into cortex and stele, segments 20mm in length were taken from 5mm behind the tip. Separated tissues were incubated in groups of 50 in aqueous solutions of IAA-2- 14 C (10⁻³ mo1 m⁻³) for 2h, 4h, 8h or 24h after which they were washed (Materials and Methods) Samples were prepared for HPLC analysis using and extracted in methanol. The results (Fig. 17 : Table 14) confirmed the observations of method 1. Greenwood et al. (1973) that metabolism took place more slowly in the stele. After 2h incubation an average of 96% of the radioactivity present in stelar extracts was associated with the IAA peak. Extracts of cortical tissue contained an apparently similar range of metabolites to those from complete root segments; only 8% of the ¹⁴C appeared to represent IAA. Stele segments incubated in IAA for longer times metabolised the majority of IAA entering the tissue. As IAA was taken up from solution for up to 8h, and there was no significant breakdown of IAA in the incubating solutions, this represents an increase in the rate of IAA metabolism. This may be a result Without conclusive identification of the of senescence of the tissue.


Fig. 17. Comparison of ¹⁴C-IAA metabolism in separated cortical and stelar tissues of Zea mays roots - <u>HPLC Analysis</u>. Tissues were incubated in 14C-IAA (10⁻³mol m⁻³) for 2h. The traces represent typical results from 3 replicate experiments. Solvent gradient : 10-60% methanol over 20 min. Flow rate : 1 cm min⁻¹. Detector : homogeneous radioactivity monitor; 30 cps full scale deflection; 10s time constant.

Tissues (50 segments p 24h. Cuantitative re:	er dish) were incub: sults were obtained	ated in aqueous so in 2 replicate ext	lutions of ¹⁴ C-IAA beriments.	. (10 ⁻³ mol m ⁻³) foo	: 2h, 4h, 8h or
		• • •			
TREATMENT	cortical segments 2h incubation	stelar segments 2h incubation	stelar segments 4h incubation	stelar segments 8h incubation	stelar segments 24h incubation
Uptake of radioactivity (Bq segment-1)	39 41	18 18	21 44	54 77	25 70
Dry weight of tissue (mg)	28.0 25.2	7.5 8.0	7.2 8.8	8.4	5.5 8.9
Percentage radioactivity in extract, associated with IAA peak	∞ Ο	99 94	61 55	14 1.8	10 . 1
Percentage radioactivity in extract, associated with peak 10	16 13	6 1	15 15 15	30 31	30 35

Uptake and metabolism of ¹⁴C-IAA by separated cortical and stelar tissues of Zea mays roots over a time-course.

Table 14:

products, it was not possible to know whether the same compounds were produced in both tissues. However, a substance with the same retention time as peak 10 was the most prominent metabolite in both cortical and stelar extracts.

Conclusion

Although results in sections B and C indicate that exogenous IAA taken up from solution. was rapidly metabolised, during the first 2h at least most of this appears to take place in the cortical tissues, where only a small proportion of the endogenous IAA is located.

E. Metabolism of IAA-2-¹⁴C Supplied from Agar Blocks to the Stele at the Basal End of Root Segments

In order to obtain information concerning the metabolism of endogenous IAA, it is necessary to supply labelled IAA in a manner which reproduces as closely as possible its natural source in the root. Bridges <u>et al</u>. (1973) showed that the majority of endogenous IAA is located in the stele. Wilkins and co-workers (e.g. Bowen <u>et al</u>., 1972) also demonstrated that IAA is transported acropetally predominantly in the stele. In this experiment the metabolism of IAA-2-¹⁴C supplied from agar blocks to the stelar tissue at the basal end of root segments was studied. As the results in experiment D showed that metabolism of IAA by the stele during a 2h incubation was slow, this means of donating the IAA would eliminate the possibility that a substantial proportion of metabolism was taking place during entry at the cut surfaces.

Sections of tissue, 13mm in length, were taken from 10mm behind the root tips of 3-day-old, dark-grown Zea mays seedlings. The cortex was removed from the basal 3mm of each segment, and the protruding stele placed in contact with a block of 1.5% agar containing IAA-2- 14 C (see Fig. 18). Blocks of plain agar at the opposite end of each root were present to protect the tissue from drying, and to collect radioactive compounds diffusing from the segment. Eight groups of 20 segments were enclosed in glass petridishes The protruding section of stele plus 1mm of tissue and incubated for 2h. from each end of the segments was then removed and the roots extracted with methano1. Extracts were prepared for HPLC analysis according to method 1. Radioactivity present in both the donor and receiver blocks of agar was also extracted into methanol. The distribution of radioactivity at the end of the experiment, which was only performed once, was as follows:





donor block	338,000	Bq
root tissue	3,482	вq
receiver block	1,480	Bq

The approximate amount of radioactivity present in each segment after a 2h incubation was therefore equivalent to 19.8 pmol IAA.

The proportion of IAA which had been metabolised by the root tissue was significantly less than in experiments in which segments were floated in aqueous solutions of IAA; approximately 50% of the radioactivity remained associated with the IAA peak (Fig. 19a) compared with 31% in experiment B.5. Six of the same metabolite peaks were apparently present (peaks 5, 6, 8, 9, 10 and 11). However, there was a noticeable decrease in the proportion of peak 10 which only accounted for approximately 9% of the radioactivity and was no longer the most prominent metabolite. The percentage of IAA metabolised was considerably greater than that in stelar tissues alone (experiment D), indicating that some IAA had probably moved into the cortex.

Although approximately 50% of the radioactivity in the root tissue extracts represented metabolites of IAA, extracts of the receiver block yielded only a single radioactive peak with the same retention time as IAA (Fig. 19b). This was accompanied by a fluorescence peak. When methylated using diazomethane, and analysed again by HPLC, a single peak was obtained with the same retention time as IAA - methyl ester (Fig. 20). Thus there is strong evidence that radioactivity in the receiver block was exclusively associated with the IAA molecule. HPLC analysis of donor block extracts indicated that breakdown of IAA during the experiment was negligible.

A similar experiment to that described above was carried out in which root segments were incubated for 24h. HPLC analyses of tissue extracts showed that very little IAA remained in the tissue after this time; in two out of three experiments no 14 C-IAA was detectable (Fig. 21). Four metabolite peaks were consistently present : peaks 3 and 10 and unresolved compositpeaks 1/2 and 7/8. However, stelar tissue alone also metabolised virtually all of the IAA taken up after 24h. These results may reflect the senescence



Fig. 19. HPLC analyses of extracts from root tissue and "receiver" blocks of agar. IAA-2-"C was supplied to the stele at the basal end of root segments, from agar blocks (2h incubation). Traces represent results from single extracts analysed in duplicate. Solvent_gradient : 10-60% methanol over 30 min. Flow rate : 0.75 cm min⁻¹. Detector: a. (solid line) homogeneous radioactivity monitor; 30 cps full scale deflection; 10s time scale. b. (dashed line) fluorescence spectrophotometer.



Fig. 20. HPLC analysis of radioactivity extracted from "receiver" block, after methylation using diazomethane. Trace represents results from a single extract, analysed in duplicate. Solvent gradient : 50-100% methanol over 20 min. Flow rate : 0.75 cm min⁻¹. Detector : homogeneous radioactivity monitor; 20 cps full scale deflection; 10s time constant.





of the tissues and their significance in relation to elucidating the normal pathways of endogenous IAA metabolism is questionable.

Conclusions

IAA-2-¹⁴C supplied to the stele at the basal end of root segments for 2h is metabolised to at least 6 products which appear to be similar to those in extracts of tissue floated in aqueous solutions of IAA. The rate of metabolism was reduced, however, and the relative proportions of peaks differed. Radioactivity diffusing from the apical end of the segments appeared to be exclusively associated with the IAA molecule, thus providing evidence for a transport mechanism specific for IAA.

F. Metabolism of ¹⁴C-IAA by Coleoptile Segments

Coleoptiles of dark-grown Zea mays seedlings have been used extensively for research on the rôle of IAA in geotropism and phototropism (e.g. Goldsmith and Wilkins, 1964; de la Fuente and Leopold, 1968; Wilkins and Whyte, 1968; Cane and Wilkins, 1969; Hertel et al., 1969; Shaw et al., 1973; Gardner et al., 1974), on the polar transport of IAA (e.g. Edwards and Goldsmith, 1980), and on binding of IAA to putative receptor sites (e.g. Hertel et al., 1972; Batt et al., 1976; Ray et al., 1977; Cross and Briggs, 1979; Moloney and Pilet, 1981). As with similar experiments on roots, the results of these studies have often been interpreted on the basis that metabolism of applied ¹⁴C-IAA during the course of the experiment is negligible. In some cases this was checked using TLC (e.g. Cane and Wilkins, 1969; Shaw et al., 1973; Gardner et al., 1974; Edwards and Goldsmith, 1980). Where significant metabolism had occurred this was taken into account, although in each instance the majority of radioactivity cochromatographed with the IAA standard. The results in section A of this thesis, however, demonstrated that TLC was not able to resolve the large number of metabolites present in methanolic extracts of Zea mays root tissue which had been incubated in IAA-2- 14 C. The metabolism of IAA-2- 14 C in coleoptile tissues was therefore investigated using HPLC analysis.

F.1. HPLC Analysis of IAA-2-¹⁴C Metabolites from Coleoptile Segments after 2h Incubation

Coleoptile sections from 5-day-old, dark-grown Zea mays seedlings were cut under dim green light and incubated in groups of 10 or 20 in aqueous solutions of IAA-2- 14 C (2 cm³; 10⁻² mol m⁻³) for 2h in darkness. After following the standard washing procedure, tissue was extracted overnight with methanol and samples prepared for HPLC analysis. In one replicate the extract was prepared according to method 1, while two samples were

purified using Sep-pak C₁₈ cartridges (method 2).

In each experiment a substantial proportion of IAA was metabolised (Fig. 22; Table 15, column 1); an average of 47% of the radioactivity remained associated with a peak which shared the same retention time as IAA. A corresponding fluorescence peak was also observed when the excitation and emission wavelengths were set at 280 and 350 nm respectively. All the major metabolites observed corresponded closely to peaks present in root extracts. Peak 10 was the most prominent metabolite and comprised a mean of 35% of the label. Other compounds were proportionately reduced and could not always be distinguished from background radiation. Although only metabolites 3,4,5,10 and 11 appeared consistently, peak 2 alone was absent from all coleoptile extracts.

The average amount of radioactivity taken up by the tissue was 121 Bq segment⁻¹ (equivalent to approximately 57 pmol IAA). The extraction efficiencies for all coleoptile experiments were high (at least 90%; Tables 16, 18 and 20). A small and variable amount of radioactivity remained unaccounted for at the end of each experiment. Label was also lost during preparation of samples for HPLC (up to 22%). Very little breakdown of IAA took place in the incubating solution, the amount of radioactivity associated with the IAA peak never being less than 94% (Table 15).

IAA Breakdown during Sample Preparation

As with experiments on the metabolism of IAA in root tissue, controls were included to test whether breakdown occurred during the preparation of extracts for analysis. Aliquots of IAA-2-¹⁴C were added to 3 separate methanolic extracts of 20 coleoptile segments each. Samples were prepared for HPLC analysis using Sep-pak C₁₈ cartridges (method 2). A small amount of peak 10 was present in each trace (Fig. 23). The average proportion of radioactivity lost during sample preparation was 26% (Table 17).



Fig. 22. Metabolism of IAA-2-¹⁴C by coleoptile tissues of Zea mays <u>seedlings : HPLC analysis.</u> 20 segments (cut in dim green light) were incubated in 10⁻² mol m⁻³ IAA-2-¹⁴C for 2h. Trace represents a typical result from 3 replicate experiments. Root extracts containing IAA metabolites were used as standards for numbering of peaks. Solvent gradient : 10-60% methanol over 30 min. Flow rate : 0.75 cm min⁻¹. Detector : homogeneous radioactivity monitor; 30 cps full scale deflection; 10s time constant.

Table 15 :	IAA Metabolites present in methanol extracts of coleoptile segments incubated in IAA-2- ¹⁴ C for 2h.
	Metabolite peaks were numbered, using root extracts as standards; see Expt A.2, Fig. 8 (1).
	The exneriment was nerformed in trinlicate.

FREATMENT	Coleoptile segments cut in dim green light	Coleoptile segments cut in the laboratory	Boiled coleoptile segments	
Radioactive peaks present in	IAA,10,4,7/8,11,5,6,9,1,3	IAA,10,11,4,5,3,1,7/8	IAA,10	
methanol extracts in order of	IAA,10,11,4,5,3	IAA,10,3,4,5,1	IAA,10	
decreasing height (1)	IAA,10,11,5,4,3		IAA, IO	
<pre>% radioactivity in methanol</pre>	63	52	96	
extract remaining associated	39	44	95	
vith IAA peak	38		66	
¢ radioactivity in methanol	32	36	4	
sxtract associated with peak 10	36	44	ъ	
	38		0.5	
¢ radioactivity in incubating	1	66	1	
solution associated with IAA peak	98	67	95	
	67		96	

nalysis of the fate of IAA-2- ¹⁴ C supplied in aqueous solution to Zea mays coleoptile segments.	oleoptiles (10, 20 or 40 segments from which leaves had been removed) were incubated for 2h in	0^{-2} mol m $^{-3}$ IAA-2- 14 C. Extracts designated (S) were purified using Sep-pak C $_{18}$ cartridges.
Table 16 : Ana	Col	10

The experiment was carried out in triplicate. Radioactivity is expressed in Bq.

TRI	ATAENT	Coleoptile segments cut in dim green light	Coleoptile segments cut in the laboratory	Boiled coleoptile segments
1.	Initial Radioactivity	50300 49000 47800	50300 49300	45400 43300 42700
2.	Radioactivity remaining in incubating solution	45500 40700 32500	42200 42600	25800 38000 35500
м.	Radioactivity in methanol extract	949 2620 2500	1800 2060	. 6980 3280 3610
4.	Radioactivity remaining in tissue after extraction	43 55 72 .	46 52	103 35 63
<u>ب</u> ا	Radioactivity in washings	- 5170 6810	8160 436 0	11800 4750 7160
6.	Final total radioactivity	- 48500 41900	52200 49000	44700 46100 · 46400
7.	Loss of radioactivity during incubation	- 451 5900	225	671 - -

TOA

.

	7 expressed as a percentage of 1	- 0.9% 12%	- 0.5%	1.5%
14	umber of segments	10 20 20	20 20	40 20 20
)ry weight of tissue (mg)	- 16.1 16.9	- 15.1	64.2 12.4 13.0
	Jptake (Bq segment ⁻¹)	99 134 129	92 105	199 164 184
	Extraction efficiency	96% 98% 97%	9 8% 9 8% 8	909 809 88
	Radioactivity in methanol extract after preparation for HPLC	- 2230(S) 2180(S)	1500 - (S)	58.30 (S) 2580(S) 2830 (S)
	13 expressed as a percentage of 3	- 85% 87%	83% -	84% 79% 78%
	kadioactivity in incubating solution after preparation for HPLC	30400 33300 28300	39300 35600	20600 32400 29200
-	5 expressed as a percentage of 2	67% 828 87%	93% 84%	80% 85% 82%

.

110

.

.



Fig. 23. IAA degradation during preparation of coleoptile extracts for <u>HPLC analysis</u>. IAA-2-14C was added to methanolic coleoptile extracts and samples prepared for HPLC analysis using Sep-pak C₁₈ cartridges (method 2). The trace represents typical results from 3 extracts. Solvent gradient : 10-60% methanol over 20 min. Flow rate : 1 cm min⁻¹. Detector : homogeneous radioactivity monitor; 30 cps full scale, 10s time constant.

Table 17 :	Loss o	of IAA-2- ^{14}C	during	preparation	\mathbf{of}	contro1	coleor	otile
	Contraction of the local division of the loc				the second s	the second s	the second s	The second s

PLICATE	1	2	3
Initial radioactivity in extract (Bq)	887	931	993
Radioactivity in extract after preparation for HPLC analysis (Bq)	626	686	761
2 expressed as a percentage of 1	71%	74%	77%
	PLICATE Initial radioactivity in extract (Bq) Radioactivity in extract after preparation for HPLC analysis (Bq) 2 expressed as a percentage of 1	PLICATE 1 Initial radioactivity in extract (Bq) 887 Radioactivity in extract after preparation for HPLC analysis (Bq) 626 2 expressed as a percentage of 1 71%	PLICATE12Initial radioactivity in extract (Bq)887931Radioactivity in extract after preparation for HPLC analysis (Bq)6266862 expressed as a percentage of 171%74%

extracts for HPLC analysis

Metabolism of IAA-2-¹⁴C by Boiled Coleoptile Segments

Coleoptile tissue (groups of 20 or 40 segments) which had been boiled for 2 min was incubated in aqueous solutions of IAA-2- 14 C (10⁻² mol m⁻³) for 2h.

After following the standard washing procedure, plant material was extracted overnight with methanol, and samples prepared for HPLC analysis using Sep-pak C_{18} cartridges (method 2).

As with the previous control, the majority of the radioactivity remained associated with the IAA peak (Fig. 24; Table 15) although a small amount of peak 10 was present, representing a mean of 3% of the radioactivity in the extract. Losses of radioactivity and extraction efficiencies were similar to those for living tissue (Table 16). In contrast to the root experiments the uptake of radioactivity had increased by a mean of 51%.



Fig. 24. Metabolism of IAA-2-¹⁴C by boiled coleoptile segments (2h incubation). Trace represents a typical result from 3 replicate experiments. Solvent gradient : 10-60% methanol over 30 min. Flow rate : 0.75 cm min⁻¹. Detector : homogeneous radioactivity monitor; 300 cps full scale deflection; 10s time constant.

113

• •

F.2. Metabolism of IAA by Coleoptile Segments Exposed to Light during Sectioning

Coleoptiles for large-scale experiments were sectioned in the laboratory. An experiment was therefore carried out to determine the effect of this short exposure to light on the metabolism of IAA. Groups of 20 coleoptile segments were incubated in aqueous $IAA-2-{}^{14}C$ (10^{-2} mol m⁻³) for 2h. Plant material was both grown and incubated in darkness, but sections were cut in the laboratory. This took approximately 15 min.

Analysis of methanolic extracts, prepared according to either method 1 or 2 revealed a pattern of metabolism closely similar to that obtained in experiment F.1 with coleoptile segments cut in dim green light (Table 15). No significant differences could be found in either the amount of IAA metabolised (a mean of 52% compared with 53% for experiment F.1) or the products obtained. Uptake and extraction efficiencies were also not significantly affected (Table 16).

F.3. Effect of Varying the External Concentration of IAA on its Metabolism by Coleoptile Segments

Coleoptile segments (cut in the laboratory) were incubated in groups of 20 in three different concentrations of IAA-2-¹⁴C (10^{-3} , 10^{-2} and 10^{-1} mol m⁻³) for 2h. After washing (Materials and Methods), tissue was extracted with methanol and samples prepared for HPLC analysis using Sep-pak C₁₈ cartridges (method 2).

No consistent and significant differences could be found in the metabolism patterns obtained at different concentrations (Fig. 25, Table 18). The average proportions of radioactivity remaining associated with the IAA peak were 57%, 45% and 58% for external IAA concentrations of 10^{-3} , 10^{-2} and 10^{-1} mol m⁻³ respectively. Peak 10 was always the most prominent metabolite with peak 4 also consistently present. Variation in the minor components observed, appeared to be due to the limitations of the detector sensitivity. The



Fig. 25. Metabolism of IAA-2-¹⁴C supplied at 3 different concentrations to Zea mays coleoptile segments (2h incubation). a. 10⁻³ mol m⁻³, b. 10⁻² mol m⁻³, c. 10⁻¹ mol m⁻³. Each trace represents typical results from 3 replicate experiments. Solvent gradient: 10-60% methanol over 20 min. Flow rate: 1 cm⁻³ min⁻¹. Detector: homogeneous radioactivity monitor; 30 cps full scale deflection (a. and c.), 100 cps (b); 10s time constant.



Fig. 25 cont'd

.

Table 18:	Uptake and metabolisn	m of IAA-2- ¹⁴ C supplied a	t 3 different concentrations	to Zea mays coleoptile
	segments (2h incubati incubated for 2h in 1	<u>ion)</u> . Metabolite peaks 10 ⁻³ mol m ⁻³ IAA-2- ¹⁴ C as	were numbered using extracts standards: see Expt. A.2, 1	from root segments ig. 8 (1).
	The experiment was ca	arried out in triplicate.		
Concentrat: in incub:	ion of IAA ating solution	10 ⁻³ mol m ⁻³	10 ⁻² mol m ⁻³	10 ⁻¹ mol m ⁻³
Untake of	IAA	7.5	38	320
(pmol set	gment_1)	6.8	53	654
		6.8	65	551
Radioactive	e peaks in methanol	IAA, 10, 4, 5, 3	IAA, 10, 4, 5, 3	IAA, 10, 4, 5, 9
extract : decreasir	ın order of ng height (1)	IAA, 10, 11, 4, 5	IAA, 10, 11, 4, 5, 3	IAA, 10, 4, 5
		IAA, 10, 11, 4, 3	IAA, 10, 11, 4, 3, 5	IAA, 10, 11, 4, 5
% radioacti	ivity in methanol	57	39	64
extract ; TAA neak	associated with	59	50	63
		56	47	48
% radioacti	ivity in methanol	31	38	13
with peak	k 10	24	25	28
		25	29	27

lack of peak 1, which was present in some extracts in Experiments F.1 and F.2, was probably a result of its loss during purification of samples using Sep-pak C_{18} cartridges (see Experiment B.5). The different external IAA concentrations also had no effect on the extraction efficiency which was always at least 95%.

At the lowest concentration, i.e. $10^{-3} \text{ mol m}^{-3}$ IAA, the average amount of radioactivity taken up over 2h was equivalent to 7.0 pmol segment⁻¹. Given that approximately 57% of this remained associated with the IAA peak. the amount of exogenous IAA in the tissue after this time appeared to be <u>circa</u> 4.0 pmol segment⁻¹.

F.4. Time-Course of 4AA Metabolism by Zea mays Coleoptile Segments

Four groups of 20 coleoptile segments (cut in the laboratory) were placed in aqueous solutions of IAA-2- 14 C (10⁻² mol m⁻³) and incubated for various times: Table 19.

Dish	Time incubated in ¹⁴ C-IAA	Time incubated in distilled water	Total incubation time
1	10 min	_	10 min
2	1 h	-	1 h
3	1 h	3 h	4 h
4	1 h	23 h	24 h

Table 19 : Incubation protocol for time-course experiments on IAA

metabolism in coleoptile segments

Sections incubated for longer than 1h were washed thoroughly (Materials and Methods) before transferring to dishes of distilled water for the remainder of the incubation time. Methanol extracts were prepared for HPLC analysis using Sep-pak C_{18} cartridges (method 2).

Metabolism of IAA was slower than in root tissues, with no products apparent after a 10 min incubation (Fig. 26; Table 20). However, in common with root experiments very little IAA was detectable after 24h (IAA was only observed in one replicate). After 4h the IAA peak was approximately 15% of the size of peak 10. Peak 10 was the most prominent metabolite at each time. Other peaks were small and variable, although peaks 7 and 8 had consistently increased in size after a 24h incubation. As with root experiments, there were no clear indications of conversions of one metabolite into another.

Conclusions

Metabolism of IAA-2-¹⁴C by Zea mays coleoptile segments was rapid with approximately 47% of radioactivity remaining associated with the IAA peak after 2h incubation. During short incubation times the amount of IAA metabolised by coleoptiles was significantly less than in roots. The products of IAA metabolism in roots and coleoptiles appeared to be similar although peak 10 had greater prominence in coleoptile extracts. Contro1s for metabolism by dead tissue, and for breakdown of IAA during sample preparation indicated that metabolism was genuinely taking place in the living tissue. Metabolism of IAA in sterile coleoptiles was not investigated. The exposure of coleoptiles to light for a short period during section cutting did not affect the rate or pattern of IAA metabolism. Changes in the exogenous IAA concentration also did not alter its metabolism.



Fig. 26. Time-course of IAA-2-¹⁴C metabolism by Zea mays coleoptile segments. Tissue was incubated in IAA-2-¹⁴C (10⁻² mol m⁻³) for up to 1h, then washed and transferred to distilled water. (a) 10min incubation, (b) 1h, (c) 4h, (d) 24h. Each trace represents typical results from 3 replicate experiments. Solvent gradient: 10-60% methanol over 30 min. Flow rate: 1 cm min⁻¹. Detector: homogeneous radioactivity monitor; 30 cps full scale deflection; 10s time course.



Table 20 :Metabolism of IAA-2-14C by Zea mays coleoptile segments over
a 24h time course. Tissue was incubated in IAA-2-14C (10⁻²
mol m⁻³) for up to 1h, then washed and transferred to distilled
water. Extracts designated (S) were purified using Sep-pak C18
cartridges. The experiment was carried out in triplicate.
Metabolite peaks were numbered using a root extracts as standards;
see Expt. A.2 Fig. 8 (1).

Incubation Time	Metabolite peaks present in methanol extract in order of decreasing height (1)
	IAA only
10 min	IAA only (S)
	IAA only (S)
	IAA, 10, 11
1 h	IAA, 10, 11 (S)
	IAA, 10, 11 (S)
	10, 3, IAA, 11, 7/8
4 h	10, IAA, 11, 7/8, 3 (S)
	10, 11, IAA, 7/8, 3 (S)
	10 8 7 3 1 9 /
24 h	10, 8, 7, 3, 1, 5, 4
N T 11	10, 11, 8, 7, 9, TAA (S)
	· · · · · · · · · · · · · · · · · · ·

In the previous experiment it was shown that <u>Zea mays</u> coleoptile tissue is capable of rapid metabolism of IAA. This has potentially important consequences for the interpretation of published results on the transport of IAA. An experiment was therefore carried out to investigate the metabolism of IAA-2-¹⁴C moving basipetally along coleoptile segments.

Labelled IAA was supplied from agar blocks (donor blocks) to the apical ends of sections. Radioactivity diffusing from the basal ends was collected in agar (receiver blocks) which also served to prevent drying of the tissue. Coleoptile segments were thus incubated in petridishes, in four groups of 20, for 4h in darkness. After removing 1mm from each end of the sections, both tissue and agar blocks were extracted with methanol. Coleoptile extracts were purified using Sep-pak C_{18} cartridges (method 2) prior to HPLC analysis. Samples from agar blocks were prepared according to method 1.

Substantial metabolism of IAA took place within the plant tissue (Fig. Approximately 25% of the radioactivity remained associated 27: Table 21). with the IAA peak after 4h. Metabolites 3 and 10 are seen in Fig. 26. In a repeat experiment small amounts of peaks 6, 7 and 8 were also visible. In contrast, only a single peak, which co-chromatographed with IAA, was observed in extracts from receiver blocks. When methylated using diazomethane a compound was formed which co-chromatographed with IAA-methyl Further proof that only IAA had reached the ester on the HPLC (Fig. 28). receiver block of agar came from GC-MS analysis. When used in the selected ion-monitoring mode and tuned for ions with m/e 202 a peak was observed with the same retention time as IAA (Fig. 29).

Only a very small proportion of the IAA-2-¹⁴C remaining in the donor blocks had undergone modification.



Fig. 27. HPLC analysis of methanol extract from coleoptile segments supplied with IAA from agar blocks, and of extracts from agar receiver blocks. 4h incubation. HPLC conditions: 10-60% methanol over 30 min. Flow rate: 0.75 cm min⁻¹. Homogeneous radioactivity monitor; 30cps full scale deflection; 10s time constant

<u>Table 21</u>: <u>Metabolism of IAA-2-¹⁴C transported down Zea mays coleoptile</u> <u>segments</u>. Segments were supplied with IAA-2-¹⁴C from agar blocks placed at the apical ends and incubated for 4h. The experiment was repeated in duplicate.

	Amount of Radioactivity (Bq)	% of radioactivity remaining associated with IAA peak
Donor block	12,600	92
	19,400	98
Coleoptile extract	1,014	26
	2,560	25
Receiver block	917	97
	1,880	100
·		



Fig. 28.	HPLC Analysis of radioactivity collected in agar "receiver
	block", after methylation using diazomethane. Solvent
	gradient: 50-100% methanol over 20 min. Flow rate:
	0.75 cm ³ min ⁻¹ . Detector: homogeneous radioactivity
	monitor; 10 cps full scale deflection; 10s time constant.



Fig. 29.GC-MS analysis of the IAA peak from methanolic extracts of
"receiver blocks", after purification by HPLC, and derivatisation
using BSTFA.
GC column: 5ft x ¼" (1.52m x 6.35mm) 3% Dexil-300 on Supelcoport
100-120. Temperature: 190°C isothermal. Flow rate: 20 cm ³ min⁻¹.

100-120. Temperature: 190°C isothermal. Flow rate: 20 cm⁻ min⁻. MS Conditions: 24 eV; Source temperature 280°C; Separator temperature 250°C. (a) extract only. (b) extract plus coinjection of IAA-TMSi standard. Conclusions

IAA-2-¹⁴C supplied to coleoptile segments from apically placed agar blocks was substantially metabolised by the tissue. A comparison with Fig. 26c indicated that metabolism was somewhat slower than that taking place in coleoptiles supplied with IAA from the incubating solution.

Analyses of radioactivity in receiver blocks indicated that the IAA metabolites formed were not mobile. IAA alone was transported.

H. The Chemical Nature of IAA Metabolites

A series of experiments were carried out to investigate the chemical structures of the products of IAA metabolism. For this purpose large extracts were prepared from 10 or 15 groups of 50 root segments incubated for 2h in 10^{-2} mol m⁻³ IAA-2-¹⁴C. After washing (Materials and Methods), the tissue was extracted with methanol. For the majority of studies the extracts were purified using Sep-pak C₁₈ cartridges (Method 2). An aliquot of each sample was analysed using HPLC and the retention times of metabolite peaks 4-11 and the IAA peak were measured. The remainder of the extract was then loaded onto the column and metabolites collected. Peaks 1, 2 and 3 could only be collected at a very low yield and were not used. The samples were dried by first blowing off the methanol under a stream of nitrogen, then freeze-drying.

H.1. Experiment to Investigate the Number of Radioactive Compounds Represented by each Peak

Samples of individual peaks were prepared as described above. Each was then reanalysed by HPLC, firstly on a solvent gradient of 10-60% methanol over 30min to check for sample degradation, and secondly by isocratic elution. The experiment was carried out in triplicate. The results of the gradient analysis have been presented in Experiment A.3 (Table 6). The compounds giving rise to peaks 4 and 5 appeared to be interconverted. A proportion of peaks 6 and 11 had been modified while the other compounds remained unchanged.

The results of the isocratic analyses are given in Table 22. In each case no further resolution of products was observed, indicating that the radioactivity in each peak is probably associated with a single compound.

Table 22 : Results of isocratic HPLC analysis of IAA-2-14 C metabolites fromZea mays root tissue. The experiment was carried out in
duplicate.

Metabolite peak	% Methanol in buffer used for HPLC analysis	No. of Peaks observed
4	12%	2 Peaks (peaks 4 and 5)
5	12%	2 Peaks (peaks 4 and 5)
6	12%	2 Peaks (peak 6 and break- down product)
7/8	18%	2 Peaks (peaks 7 and 8)
9	23%	Single peak
10	23%	Single peak
11		Not analysed
IAA	50%	Single peak
H.2. Polarity of IAA Metabolites

In a reverse-phase HPLC system compounds of greatest polarity are normally eluted first. It can be seen from the traces obtained (e.g. Figs. 8 and 22) that all metabolites from both root and coleoptile segments were eluted before IAA. This is thus a good indication that the products are all more polar than IAA itself. It must be remembered, however, that the relative polarities of compounds may be altered by the solvent system. Furthermore, in addition to the main process of solvent partitioning, other effects such as adsorption may play a part in the separation.

H.3. Experiment to Investigate whether carbon-1 of the Side Chain has been lost

An experiment was carried out to compare the metabolism of IAA-1- 14 C with that of IAA-2- 14 C. Groups of 50 root or 20 coleoptile segments (cut in the laboratory) were incubated for 2h in 2 cm³ volumes of 10⁻³ and 10⁻² mol m⁻³ IAA respectively. After following the standard washing procedure, tissues were extracted with methanol. Samples were prepared for HPLC analysis using either method 1 or method 2 as indicated in Table 23.

HPLC analyses of both root and coleoptile extracts showed metabolism patterns closely resembling those obtained in experiments using IAA-2-¹⁴C (Figs. 30 and 31, Table 23). The range of metabolite peaks present appeared to be the same, as did the proportion of radioactivity remaining associated with the IAA molecule; means of 38% compared with 31% for roots and 51% compared with 48% for coleoptiles. Similar amounts of radioactivity were also associated with peak 10. No significant differences between experiments utilising IAA-1-¹⁴C and IAA-2-¹⁴C were found in the quantities of radioactivity lost during incubation (Table 24). It would thus appear that all the major metabolites of IAA, observed in the present experiment₅ on <u>Zea mays</u> seedlings, do not represent products of a decarboxylation reaction.

leoptile segments from Zea mays seedlings.
o p
tan
r00
C by
[AA-2- ¹⁴ (
n pu
4 C :
IAA
of
metabolism
the
Ъ
Comparison
 10
Table 2

٠

Segments (50 root or 20 coleoptile segments) were jncubated for $2h_{14}^{j}n_{14}C_{-1AA}$ (10^{-3} or 10^{-2} mol m⁻³). Metabolite peaks in extracts of roots incubated for 2h in 10^{-3} mol m⁻¹ IAA-2- $1^{4}C$ were numbered 1-11 (see Experiment A.3) and used as standards for the other extracts. Samples designated (S) were purified using C_{18} Sep-pak cartridges. The experiment was carried out in triplicate.

•

•	•			
	IAA-2- ¹⁴ C supplied to roots	IAA-1- ¹⁴ C supplied to roots	IAA-2- ¹⁴ C supplied to coleoptiles	IAA-1- ¹⁴ C supplied to coleoptiles
Radioactive peaks present in methanol extracts in	IAA,10,9,7,6,5,4,3, 1,11,8,2	IAA,10,7,9,8,6,1,11, 5,4,2,3	IAA,10,11,4,5,3,1, 9,7/8	IAA,10,4,5,1,3,11, 9, 7/8
order of decreasing height	IAA,10,9,6,1,5,7,8, 4,3,2,11	IAA,10,7/8,9,4,6,1, 5,3,11,2	IAA,10,3,4,5,1(S)	IAA,10,11,4,5,1,3(S)
	IAA,10,9,6,7,5,1,4, 3,8,2,11	IAA,10,1,3,9,5,4,6, 7,11,8,2		IAA,10,11,4,5,3,1(S)
% Radioactivity in methanol	34	42	52	48
extract associated with	29	39	44	57
IAA peak	30	32		49
<pre>% Radioactivity in methanol</pre>	21	21	17	36
extract associated with	19	16	44	31
peak 10	18	21		32
% Radioactivity in incubating	93	98	66	97
solution associated with IAA peak	se 98	06	67	100
	95	100		. 94
			•	

.

.





Fig. 31. Metabolism of IAA-1-¹⁴C by coleoptile segments (2h incubation)

A typical result from 3 replicate experiments. HPLC Conditions: Solvent gradient; 10 - 60 % methanol over 30 min. Flow rate; 0.75 cm³min⁻¹. Detector: homogeneous radioactivity monitor; 30cps full scale deflection; 10s time constant.



ings.
seed1
ays s
of 26
nts (
segme
ile
€opt
col
and
root
by
4
- ,
9
-FA-I
and
14 C
IA
n of
lis
tabo
e me
f th
o uo
aris
Comp
24
able
-

Segments (50 root or 20 coleoptile segments) were incubated for 2h in 14 C-IAA (10^{-3} or 10^{-2} mol m⁻³). The results represent the means of 3 replicate experiments \pm standard error (2 experiments in the case of coleoptile segments incubated in IAA-2- 14 C). Radioactivity is expressed in Bq.

-

Ì	TREATMENT	IAA-2- ¹⁴ C supplied to roots	IAA-1- ¹⁴ C supplied to roots	IAA-2- ¹⁴ C supplied to coleoptiles	IAA-l-14C supplied to coleoptiles
	Initial radioactivity	7650 ± 221	5760 ± 21	49800 ± 513	44200 ± 3660
2.	Radioactivity remaining in incubating solution	3470 ± 22	2310 ± 43	42400 ± 174	37700 ± 3150
3.	Radioactivity in methanol extract	2230 ± 11	1890 ± 12	1930 ± 130	1730 ± 30
4.	Radioactivity remaining in tissue after extraction	167 ± 7.2	114 ± 5.2	49 ± 3	69 ± 31
5.	Radioactivity in washings	1360 ± 136	900 ± 34	6257 ± 1900	4900 ± 1340
6.	Final total radioactivity	7230 ± 122	5210 ± 44	50600 ± 1600	44400 ± 1810
7.	Loss of radioactivity during incubation	413 ± 145	545 ± 43	-858 ± 1080	-268 ± 1850
8.	7 expressed as a percentage of 1	5.3%	9.5%	-1.7%	•0.6%
<u>б</u> .	Uptake o£ radigactivity (Bq se;ment ⁻¹)	48 ± 0.4	40 ± 0.3	98 ± 6.5	104 ± 14
10.	. Extraction efficiency	$93\% \pm 0.3\%$	94% ± 0.2%	98% ± 0%	97% ± 1 %
11.	. Radicactivity in methanol extract after preparation for HPLC	1800 ± 63	1590 ± 170	I	1397 ± 99
12.	. Il expressed as a percentage of 3	81%	84%	ı	81%

.

H.4. UV Spectrometry

The chromophore responsible for the UV absorbance of IAA is the indole nucleus. As this is not altered directly by the conjugation of the molecule with amino acids, sugars or <u>myo</u>-inositol these compounds should have UV absorption spectra very similar to that of IAA (see Fig. 32). On the other hand, if the molecule is oxidised, this will alter the chromophore and thus change the spectrum very significantly. This can be seen by a comparison of the spectra of indole-3-aldehyde, oxindole-3-acetic acid and 5-hydroxyindole-3-acetic acid with that of IAA (Fig. 33).

Thus if a metabolite is found to have an absorption spectrum identical to that of IAA this would be a clear indication that it represents a conjugate of IAA. Unfortunately most of the metabolites were not sufficiently purified to give a definitive UV spectrum. However, the UV spectrum of the combined peaks 7 and 8 was very similar to that of IAA, with a broad peak at 272-282 nm and shoulders at 268 and 289 nm (Fig. 34). The sample also contained at least one other UV absorbing substance. This result was obtained in three separate experiments. Thus, at least one component of this peak appears to be a conjugate of IAA.

H.5. Methylation of IAA Metabolites

The IAA metabolites were derivatised using diazomethane which reacts rapidly with organic acids to form the corresponding methyl esters. Diazomethane also reacts with phenols to give the corresponding methyl ethers. Both types of product will be less polar than the starting material.

After methylation each IAA metabolite was analysed by HPLC. The results are summarised in Table 25. The experiment was performed once.





Fig. 33. UV absorbance spectra of IAA and its oxidation products.

IAA ----- 5-Hydroxyindole-3-acetic acid 0xindole-3-acetic acid ----- Indole-3-aldehyde



Fig. 34. UV absorbance spectrum of metabolite peak 7/8 after purification by HPLC, and compared with IAA standard

Peak No	Product of methylation					
4	inconclusive					
5	inconclusive					
6	inconclusive					
7/8	unreacted starting material plus one peak with a T _R between that of IAA and methyl-IAA.					
9	single peak with T _R between that of IAA and methyl-IAA.					
10	not analysed					
11	single peak with same T _R as methyl-IAA					
IAA	single peak with same T _R as standard methyl-IAA					

The radioactive compound represented by peak 9 and a component of peak 7/8 appeared to methylate, forming products considerably less polar than the starting material. In view of the apparently high polarity of these metabolites prior to methylation, they probably possess a carboxyl group. Treatment of the peak containing putative ¹⁴C-IAA with diazomethane resulted in a product which co-chromatographed with authentic IAA-methyl ester. This provided further evidence that the only radioactive component of this peak Methylation of peak 11 also resulted in a radioactive product which was IAA. had the same retention time as methyl IAA. However, the compound represented by peak 11 was somewhat labile and appeared to revert to IAA under certain conditions (see Expt H.6). It is likely that this metabolite had reverted to IAA prior to methylation.

H.6. Hydrolysis of IAA Metabolites

This work is of a preliminary nature. Two different hydrolysis conditions were used, as described by Bandurski and Schulze (1977). They reported that esters of IAA with glucose, <u>myo</u>-inositol and oligosaccharides could be hydrolysed using 10^3 mol m^{-3} NaOH for 1h at 22 to 25°C but that hydrolysis of peptidyl IAA required more rigorous conditions of 7 x 10^3 mol m^{-3} NaOH at 100°C for 3h. Hydrolysis of the compounds represented by peaks 4-11 was attempted. The effect of these conditions on IAA was also monitored. After each treatment, the reaction mixture was neutralised, and the radioactive products analysed by HPLC. The results are presented in Table 26. The experiment was performed once.

Peak No	Products of 10 ³ mol m ⁻³ NaOH, 22°C	Hydrolysis 7 x 10 ³ mol m ⁻³ NaOH, 100°C
4	Peak with same T _R as peak 1	-
5	Peak with same T _R as peak 1/2	-
6	Peak with same T _R as peak 1/	2 -
7/8	No effect	Peak with same T as peak 1/2
9	No effect	Peak with same T _R as peak 1/2
10	Several peaks	-
11	Peak with same T _R as IAA	-
IAA	No effect	Peak with same T _R as peak 1/2

Table 26 : Results of Hydrolysis of Metabolites

Only one metabolite (represented by peak 11) appeared to be hydrolysed to IAA. After treatment with 10^3 mol m⁻³ NaOH at room temperature this metabolite was converted to a product with the same retention time as IAA. Peak 11 thus probably represents an ester of IAA with a sugar or myo-inositol. The radioactive components of peaks 4, 5 and 6 were also altered by mild hydrolysis However, the products were highly polar compounds which were conditions. eluted close to the void volume. There is no need to assume that the products of each compound were necessarily identical as the separation of compounds eluting at the start of a gradient is poor. It is possible that the compounds represented by peaks 4, 5 and 6 are esters of IAA oxidation products. This interpretation is also supported by the lack of clear cut products of methylation in experiment H.5.

Unfortunately treatment with 7 x 10^3 mol m⁻³ NaOH at 100°C caused breakdown of IAA itself. The effect of these conditions on IAA metabolites was therefore conjectural.

H.7. Co-chromatography of Metabolites with Standards

The following putative IAA metabolites were synthesised for use as standards: IAA-glycine, IAA-alanine, IAA-valine and oxindole-3-acetic acid. The identities of these compounds were confirmed by GC-MS. The mass spectra obtained are presented in Fig. 35-38. 5-Hydroxyindole-3-acetic acid was also obtained from Aldrich Chemical Company. The HPLC retention times of the standards were compared with those of radioactive IAA metabolites. Standards were monitored using a UV absorbance detector, the wavelength of which was set at the absorbance maximum for each compound, measured from spectra shown in Figs. 32 and 33. The UV absorbance detector replaced the fluorimeter in Fig. 5a. The results are presented in Table 27.









Table 27:Comparison of the retention times of IAA metabolites with those of
standards. HPLC Conditions: Flow rate 1 cm³ min⁻¹. Solvent
gradient 10-60% methanol over 20 min. T_R values for standards are
corrected for the time difference between detectors. Average peak
width 1min. The experiment was carried out once.

Peak No	T _R	Standard .	Τ _R
7	13.2 min	5-Hydroxyindole-3-acetic acid	12.9 min
10	17.8 min	Oxindole-3-acetic acid	17.8 min
8*	13.8 min	IAA-glycine	14.2 min

* Peak 8 was analysed on a separate occasion

Peak 10, representing the most prominent metabolite, co-chromatographed with oxindole-3-acetic acid. Peaks 7 and 8 had the same retention times as 5-hydroxyindole-3-acetic acid and IAA-glycine respectively, within experimental error. IAA-valine and IAA-alanine did not co-chromatograph with any of the metabolites.

H.8. GC-MS Analysis

Only the sample of the IAA peak was sufficiently pure to permit identification using GC-MS. The instrument was used in the selected ionmonitoring mode and was tuned to detect ions with $\underline{m/e}$ 202, the most prominent fragment in the mass spectrum of the trimethylsilyl derivative of IAA (IAA-TMSi) (see Fig. 6). The IAA peak was derivatised using BSTFA. A peak was present at the same retention time as trimethylsilyl IAA. This was confirmed by coinjection (Fig. 39).

Conclusions

The product represented by peak 10, i.e. the most prominent metabolite



Fig. 39.GC-MS analysis of the IAA peak from methanolic extracts of root
segments incubated in solutions of 10⁻⁵ mol m⁻³ IAA for 2h.
The extract was purified by HPLC, and derivatised using BSTFA.
(a) extract only (b) extract plus coinjection of IAA-TMSi standard.
GC Column: 5ft x ¼" (1.52m x 6.35mm) 3% Dexil-300 on Supelcoport
100-200. Temperature: 190°C isothermal. Flow rate: 20 cm³ min⁻¹.
MS conditions: 24eV; Source temperature 280°C; Separator
temperature 250°C.

in both root and coleoptile extracts, appeared to be oxindole-3-acetic acid. Evidence was provided by co-chromatography with authentic oxindole-3-acetic Further indication that this metabolite was an oxidation acid on the HPLC. product came from control analyses in Experiments B.1 and F.1. Contro1 extracts usually contained a small amount of peak 10 (see also Experiment I). Peak 7 co-chromatographed on HPLC with 5-hydroxyindole-3-acetic acid, a compound not previously identified in plant tissue. Peak 8 had a retention time the same as that of IAA-glycine, within experimental error. Further evidence that peak 7/8 contained an IAA conjugate was provided by the UV absorption The metabolite represented by peak 11 was converted to IAA under spectrum. mild hydrolysis conditions. It is likely that this compound is an ester of IAA with a sugar and/or myo-inositol. The radioactive components of peaks 4, 5 and 6 could not be hydrolysed to IAA and are unlikely to represent sugar, myo-inositol or related esters of IAA. The compounds represented by peaks 4 and 5 were interconvertible and are probably isomeric. The metabolite represented by peak 9 could be methylated to form a substantially less polar This compound probably possessed a carboxyl group. product.

The identity of the IAA peak was confirmed by methylation and cochromatography with IAA-methyl ester and by GC-MS (selected ion-monitoring) of the trimethylsilyl derivative.

I. HPLC Analysis of IAA Metabolites Present in the Incubating Solutions

The metabolites of IAA, present in bathing solutions after 2h incubation with Zea mays root and coleoptile segments were investigated. Samples of incubating solutions containing IAA-1- 14 C and IAA-2- 14 C (10⁻² mol m⁻³ for coleoptiles: 10^{-3} mol m⁻³ for roots) were prepared for HPLC analysis. Aqueous solutions were usually dried in vacuo at 45°C and redissolved in 1 cm³ methanol. Immediately prior to analysis, samples were dried under a stream of nitrogen and taken up in 100 mm³, 10% methanol in ammonium acetate buffer (pH 3.5, 20 mol m^{-3}). Losses of radioactivity during this procedure are given in Tables 7 and 16. Two samples (root incubating solutions) were prepared by first acidifying the incubating solution (diluted to 10 cm³ with water) to pH3 with HC1 and partitioning three times with equal volumes of diethyl ether. The ether was then dried in vacuo and the sample dissolved in 1 cm³ of methanol. When freshly distilled diethyl ether was used, this procedure had no significant effect on the spectrum of compounds observed. However, when diethyl ether which had not been recently distilled was used, this resulted in the oxidation of the majority of the IAA to a compound which co-chromatographed with peak 10. This was presumably due to the presence of peroxides.

The results of the analysis, which was carried out in triplicate for each treatment, are presented in Figs. 40 and 41. As stated previously (see Tables 8 and 15), in each case the majority of the radioactivity co-chromatographed with IAA. Small amounts of some of the metabolites found in methanol extracts of roots and coleoptiles were also present; peaks 3 and 10 were consistently observed. The presence of one or more compounds which were less polar than IAA, and only appeared when methylenelabelled IAA was supplied, was of particular interest. These compounds





monitor; 30cps full scale deflection; 10s time constant.

152

.

are probably decarboxylation products of IAA. They did not co-chromatograph with indole-3-aldehyde or indole-3-carboxylic acid, but might represent other products such as 3-methyleneoxindole.

Conclusions

Solutions of IAA, in which <u>Zea mays</u> root and coleoptile segments had been incubated for 2h, contained small amounts of metabolites which appeared to be decarboxylation products. As these compounds were not present in methanol extracts of plant material they may represent the products of IAA oxidation at cut surfaces. The formation of large amounts of peak 10, when aqueous solutions of IAA were partitioned against diethyl ether which had not been redistilled, provided further evidence that this compound was an oxidation product.

SECTION 2 - THE EFFECT OF EXOGENOUS IAA ON THE ELONGATION OF ZEA MAYS ROOT AND COLEOPTILE SEGMENTS

The purpose of this Section of the thesis was two fold. Firstly, it was important to confirm whether the amounts of IAA supplied to root and coleoptile segments during metabolism experiments were physiologically active with regard to cell elongation. Secondly, published data on the effect of IAA on root growth are conflicting (see Introduction); it was hoped to clarify the effect of applied IAA on the elongation of Zea mays root segments.

A. Effect of IAA on Coleoptile Elongation

The stimulation of coleoptile elongation by IAA has been well established since 1934 (Kögl <u>et al.</u>). A single experiment was carried out to investigate the external concentration required to give maximum growth stimulation under the experimental conditions used for metabolism studies. Coleoptile segments (5mm in length, taken from 2mm behind the apex) were incubated for 1h (darkness; 22°C) in 5 cm³ volumes of aqueous IAA solution (0, 10^{-5} , 10^{-4} , 10^{-3} , 10^{-2} and 10^{-1} mol m⁻³). Growth was measured using the shadowgraph method.

IAA at the lowest three concentrations had no significant effect on growth when compared with the control (Fig. 42). Maximum stimulation occurred in 10^{-2} mol m⁻³ IAA.

Conclusions

Under the experimental conditions used in metabolism experiments, maximum growth was obtained in 10^{-2} mol m⁻³ IAA. This concentration was used in the majority of experiments on IAA metabolism in coleoptile tissue.



Fig. 42. Effect of IAA supplied at 5 different concentrations on the growth of Zea mays coleoptile segments during 1h after excision. Each point represents the mean of 10 segments ± standard error. The experiment was performed once.

B. Effect of IAA on Root Elongation

B.1. Comparison of the Growth Rate of Intact Roots with that of Segments

As other scientists (e.g. Edwards and Scott, 1977; Pilet <u>et al.</u>, 1979) appeared to have difficulty in obtaining significant effects of IAA on root growth, a preparatory experiment was carried out to investigate the optimum incubation time for such an experiment. The growth rate of intact roots was compared with that of segments at various times after excision.

Root segments 5mm in length were taken from 1mm behind the tip. Erickson and Sax (1956) and Pilet and Senn (1980) reported that this region contains the elongation zone. This was confirmed for the variety Fronica. Groups of 10 sections were incubated in water (2 cm³) for $\frac{1}{2}h$, 1h, 2h, 4h, 8h and 24h, and growth measured using the shadowgraph method. This was compared with growth of roots on intact plants, measured by recording the distance between spots of indian ink initially spaced 1mm and 6mm from the root tip.

Growth of the segments decreased rapidly with very little elongation taking place after 8h (Fig. 43). The rate of growth remained similar to that of intact plants for only circa lh.

In a similar experiment addition of IAA at concentrations of 10^{-7} , 10^{-6} and 10^{-5} mol m⁻³ (amounts which have been reported to stimulate root elongation; e.g. Batra <u>et al.</u> 1975; Pilet <u>et al.</u> 1979) had very little effect on the growth of segments at any time (Fig. 44). Addition of IAA certainly did not allow them to maintain the rate of elongation found in intact plants.

It could thus be concluded that the decrease in growth rate of segments was due to factors other than lack of IAA, such as shortage of nutrients and other growth substances or the presence of inhibitory compounds such as ethylene. For the purposes of investigating the rôle of IAA in root growth, the tissue was therefore incubated for no longer than lh, during which the growth rate was similar to that of intact roots.

1,56



Fig. 43. A comparison of the growth of 5mm root segments (taken from 1mm behind the root tip) during 24h after excision (•) with that of the same portion of root on the intact plant (o). Each point represents the mean of 30 plants ± standard error. The same result was obtained in 3 replicate experiments.



Fig. 44. Effect of IAA at three different concentrations on the growth of root segments during a 24h time course, and compared with that of intact roots. Control (•), 10^{-7} mol m⁻³ IAA (o), 10^{-6} mol m⁻³ IAA (**A**), 10^{-5} mol m⁻³ IAA (**A**), intact plants (•).

B.2. The Effect of IAA and Oxygen on Growth of Root Segments during a lh Incubation

An experiment was carried out to study in greater detail the effect of different concentrations of IAA on the growth of root segments during the first 1h after excision. Root segments were incubated in groups of 10 in the 1aboratory in solutions of 0, 10^{-9} , 10^{-7} , 10^{-5} and 10^{-3} mol m⁻³ IAA (5 cm³ per dish). Growth was measured using the shadowgraph method. Evans et al. (1980) have suggested that a supplementary source of oxygen is required for the optimum response to IAA. For this reason solutions were supplied with oxygen, nitrogen (20 cm³ min⁻¹) or air (120 cm³ min⁻¹).

The results (Fig. 45) showed no stimulation of growth by IAA at any concentration. Incubation in $10^{-3} \text{ mol m}^{-3}$ IAA inhibited growth and this occurred only when solutions were supplied with O_2 or air. Roots incubated under N_2 did not respond significantly to IAA supplied at any concentration. The bubbling of solutions with O_2 and air caused significant increases in growth of control roots and those incubated in low concentrations of IAA. An analysis of variance was carried out on measurements of roots supplied with various concentrations of IAA under O_2 or N_2 (Table 28). This revealed a highly significant interaction between the O_2 availability and the effect of IAA on growth.

Although the experiments presented were carried out in the laboratory similar experiments in which roots were incubated in darkness showed this to have no significant effect on the growth rate.

B.3. The Effect on IAA-2-¹⁴C Metabolism of Bubbling the Incubating Solution with O₂

A single experiment was carried out to discover whether the effect of addition of 0_2 on the growth and response to IAA of Zea mays root segments could be attributable to an alteration in the pattern of IAA metabolism.

Root segments (50, 20mm in length) were incubated in the laboratory in



Fig. 45. Growth of root segments incubated in the laboratory for <u>1h in 4 different concentrations of IAA and compared with</u> <u>controls</u> Roots were incubated in the absence - 0 - (4 <u>dishes per concentration</u>) or presence of supplementary oxygen - ● - (3 dishes), nitrogen - △ - (2 dishes) or air - ▲ - (2 dishes). Each dish contained 10 segments. Results represent the means ± standard error.

Table 28 : Analysis of variance on results obtained when root segments were incubated at 5 different concentrations

of IAA in the presence of O_2 or N_2

	S.Sq	D.f	Mean Sq.	F
Between concentrations of IAA	112	4	28	
Between 0 ₂ availability	176	1	176	
Interaction	103	4	25.75	32.8
Residual	186	237	0.785	

.

.

an aqueous solution of IAA-2-¹⁴C (5 cm³ of $10^{-3} \text{ mol m}^{-3}$) which was bubbled with 0_2 at a rate of 20 cm³ min⁻¹. After 2h the segments were washed (Materials and Methods), extracted overnight in methanol, and a sample prepared for HPLC analysis. The results (Fig. 46) revealed a metabolism pattern similar to that for roots incubated in aqueous IAA with no supplemantary source of O_{λ} (see Fig. 8). Peaks 1,2,3,4,5,6,7,8,9,10,11 and a peak corresponding to IAA were all present with peak 10 representing the most prominent metabolite. A detailed quantitative comparison of results is not meaningful without further repetition of the experiment and conditions were not identical to those in Section 1. The proportion of radioactivity remaining in the IAA peak was 13% compared with a value of 31% obtained in Section 1, Experiment B.3.

B.4. Changes in pH of IAA Solutions during Incubation with Root Tissue

Evans <u>et al</u>. (1980) have reported that the inhibitory effect of 2 x 10^{-3} mol m⁻³ IAA on root growth was associated with an increase in the pH of the incubating medium. They suggested that IAA causes H⁺ influx into the cells; i.e. a proton pump operating in the reverse direction to that in shoots might be activated. An experiment was carried out to try to repeat these results.

Groups of 10 root segments were incubated in darkness in 5 cm³ volumes of solution (0, 10^{-9} , 10^{-7} , 10^{-5} or 10^{-3} mol m⁻³ IAA) for 1h. The pH of the medium was measured before and after incubation. The experiment was carried out in triplicate.

An increase of approximately 0.5 pH units was observed after 1h incubation (Table 29); a value similar to that obtained by Evans <u>et al.</u> (1980). However, this increase occurred irrespective of the IAA concentration and O_2 availability, and was even observed in control dishes lacking IAA. Bubbling the solution with O_2 in the absence of root tissue produced no observable change in pH.





Table 29:Change in pH of bathing solution during incubation of root segmentswith IAA.Results represent the means of 3 dishes ± standard error.

		IAA conce	entration	$(mo1 m^{-3})$	
	0	10 ⁻⁹	10 ⁻⁷	10 ⁻⁵	10 ⁻³
Mean pH before incubation	6.1	6.1	6.1	6.1	6.1
Mean pH after incubation	6.6	6.5	6.6	6.6	6.6
pH change	+0.5	+0.4	+0.5	+0.5	+0.5
Standard error	0.07	0	0,07	0.07	0

(a) Solution bubbled with air (120 $\text{cm}^3 \text{min}^{-1}$)

(b) No supplementary gas

α το προστρατικό το προστρατικό το προστρατικό το προστρατικό το προστρατικό το προστρατικό το ποιο το ποιο το Το προστρατικό το ποιο		IAA concentration (mol m^{-3}			
	0	10 ⁻⁹	10 ⁻⁷	10 ⁻⁵	10^{-3}
Mean pH before incubation	5.8	5.8	5.8	5.8	5.8
Mean pH after incubation	6.3	6.3	6.3	6.3	6.3
pH change	+0.5	+0.5	+0.5	+0.5	+0.5
Standard error	0.09	0.08	0.09	0.09	0.06

(c) Solution bubbled with oxygen (20 $\text{cm}^3 \text{min}^{-1}$)

		IAA conce			
	0	10 ⁻⁹	10 ⁻⁷	10 ⁻⁵	10 ⁻³
Mean pH before incubation	5.7	5.8	5.8	5.8	5.8
Mean pH after incubation	6.3	6.3	6.3	6.4	6.4
pH change	+0.6	+0.5	+0.5	+0.6	+0.6
Standard error	0.17	0.15	0.09	0.07	0.12

(d) Control : Solution bubbled with oxygen (20 $\text{cm}^3 \text{min}^{-1}$), no plant tissue

	IAA concentration (mol m^{-3})				
	0	10 ⁻⁹	10 ⁻⁷	10 ⁻⁵	10 ⁻³
Mean pH before incubation	6.0	5.9	5.9	5.9	5.9
Mean pH after incubation	6.0	5.9	5.9	5.9	5.9
pH change	0	0	0	0	0
Standard error	0	0.1	0.15	0.1	0

B.5. Uptake of IAA by Root Tissue

. .

In order to assess the significance of experimental observations on the effects of exogenous IAA, the amounts of IAA entering the tissue must be compared with the endogenous levels. An experiment was carried out to measure the uptake of IAA from five different external concentrations. Groups of 10 root segments were incubated in 14 C-IAA solutions for 1h. They were then washed (Materials and Methods) and extracted in 1 cm³ methanol. The amount of radioactivity taken up by each group of segments was assayed by liquid scintillation spectrometry and the approximate uptake of IAA calculated (Table 30).

Table 30.	Uptake of	¹⁴ C-IAA	into	5mm	root	segments	incubated	in	five
	different	concent	ratior	ns fo	or 1h				

concentration of IAA (mol m ⁻³)	Bq segment ⁻¹	Uptake of µg kg ⁻¹ fresh weight of tissue	IAA µg kg ⁻¹ fresh weight taking into account IAA metabolism
5.9×10^{-6}	0.088	2.1	0.64
1.4×10^{-5}	0.21	48	0,15
1.0×10^{-4}	0.82	19	5.9
9.4 x 10^{-4}	5.3	124	38
1.0×10^{-5}	2.7	628	195

These values can be compared with measurements of the endogenous content of IAA in Zea mays roots: 30 μ g/kg f.wt. (Bridges <u>et al.</u>, 1973) and 76 μ g/kg f.wt. (Rivier and Pilet, 1974).

Conclusions

The growth rate of Zea mays root segments decreased rapidly after excision
due to factors other than lack of IAA. IAA supplied at concentrations of 10^{-9} , 10^{-7} and 10^{-5} mol m⁻³ had no effect on elongation of root segments during 1h incubation. In the presence of 0_2 , IAA at 10^{-3} mol m⁻³ had an . inhibitory effect on growth. This was not apparent under N_2 . Supplementary 0_2 increased the growth rate of control segments and those in low concentrations of IAA. This addition of oxygen did not have a significant effect on IAA metabolism or on the pH of the incubating solution.

A decrease of approximately 0.5 units in the pH of the bathing solution surrounding root segments was observed during 1h incubation. This change took place both in the presence and absence of various concentrations of IAA. Measurements of IAA uptake from solutions containing different concentrations of ¹⁴C-IAA were compared with published values for the endogenous IAA content of <u>Zea mays</u> roots. The amount of exogenous IAA present in the tissue after 1h incubation was of the same order of magnitude as the endogenous content, where the external concentration was 10^{-3} mol m⁻³. Where the IAA concentration was $5*9\times 10^{-6}$ mol m⁻³ the quantity of IAA taken up was <u>circa</u> 1% of published measurements of the endogenous levels.

DISCUSSION

A. The Pathway of IAA Metabolism in Zea mays Seedlings and its Rôle in the Regulation of IAA Levels within the Plant

If a chemical substance is to be used to regulate physiological processes, it is imperative that the organism can control accurately the amounts of that substance which are present and active at the responding tissue. There are several possible means by which it might do this. The amount of the compound itself could be regulated by one or more of 4 general ways: by controlling the rate of biosynthesis, by regulating its catabolism, by reversible inactivation of the molecule or by transport both between cells and tissues, and between different compartments within the cell. Alternatively, the effectiveness of a physiologically active compound might be controlled by changes in the levels or activity of hypothetical receptor sites. Thus to understand fully how a plant growth substance operates it is necessary to investigate the details of its metabolism and transport, as well as possible binding sites.

In this thesis the metabolism of exogenous radiolabelled IAA in the roots and coleoptiles of dark-grown <u>Zea mays</u> seedlings has been studied. Products were analysed using the high resolution technique of reverse-phase HPLC in combination with an on-stream radioactivity monitor.

In initial experiments IAA was supplied by floating root and coleoptile segments in aqueous solutions of the plant growth substance. Metabolism of IAA was rapid and extensive with a large number of products being formed. Non-sterile root segments, incubated for 2h in darkness in 10^{-3} mol m⁻³ IAA-2-¹⁴C metabolised on average approximately 59% of the radioactive IAA taken up by the tissue. At least 11 products were formed and some information about their chemical nature has been obtained.

IAA metabolites described in the literature can be divided into three groups (see Introduction):

- IAA conjugates with amino acids, sugars, <u>myo</u>-inositol or high molecular weight compounds.
- Oxidation products formed by decarboxylation, with loss of carbon-1 of the side chain (e.g. 3-methyleneoxindole, indole-3-aldehyde and indole-3-carboxylic acid).
- 3. Products formed by oxidation of the indole nucleus (e.g. oxindole-3-acetic acid, 3-hydroxyoxindole-3-acetic acid, 5-hydroxyoxindole-3-acetic acid, and 3,5-dihydroxyoxindole-3-acetic acid).

Conditions reported by Bandurski and Schulze (1977) to hydrolyse esters of IAA with sugars and <u>myo</u>-inositol were used to test IAA metabolites. Only a single metabolite, represented by peak 11 was hydrolysed to IAA under these conditions. This compound may be an ester of IAA with glucose or <u>myo</u>inositol. It is unlikely that the other metabolites are esters of IAA. UV absorbance data provided some evidence for the presence of a second IAA conjugate, as a component of peak 7/8. Peak 8 co-chromatographed with IAAglycine. A comparison of HPLC retention times of IAA metabolites with those of standards indicated that IAA-valine and IAA-alanine were not present in the plant extracts.

All the major metabolites appeared to be more polar than IAA. A comparison of the metabolism of IAA-1- 14 C and IAA-2- 14 C also showed that the products had not been formed by decarboxylation. They therefore did not include 3-methyleneoxindole, indole-3-aldehyde, indole-3-carboxylic acid or related compounds.

It was thought probable that several of the metabolites might represent compounds from group 3, formed by oxidation of the indole nucleus. These compounds are likely to be more polar than IAA. A standard of oxindole-3acetic acid was synthesised as this compound had been found in <u>Zea mays</u> endosperm by Reinecke and Bandurski (1981). When analysed by HPLC peak 10, the most prominent metabolite, co-chromatographed with oxindole-3-acetic acid. This identification was supported by evidence from IAA breakdown controls that

peak 10 represented an oxidation product of IAA. Peak 7 also co-chromatographed with 5-hydroxyindole-3-acetic acid. This compound has not been previously identified in plant extracts.

Coleoptile segments, incubated under the same conditions, had metabolised a somewhat smaller proportion of the IAA-2-¹⁴C taken up by the tissue than root segments. An average of 57% of the radioactivity remained associated with the IAA peak after 2h incubation. The products obtained were apparently similar to those formed in root tissue. However, there was a much greater predominance of peak 10 (presumptive oxindole-3-acetic acid).

The products observed in either tissue did not appear to be artifacts of the sample preparation procedure. Only a very small amount of breakdown was found in control experiments to test for IAA degradation during sample preparation. Similarly boiled segments did not metabolise IAA significantly. Further tests confirmed that modification of IAA and its metabolites during HPLC analysis was small. Only two metabolites (peaks 6 and 11) showed any sign of breakdown on rechromatography after collection from the column and drying. Experiments employing sterile roots indicated that metabolism was genuinely taking place within the plant tissue and was not due to epiphytic microorganisms.

When IAA metabolism has been studied in the past, work has usually concentrated on the identification of one or two products which often represented only a small fraction of the total amount of IAA metabolised (e.g. Tuli and Moyed, 1967; Magnus <u>et al.</u>, 1971). In the present study relatively crude extracts were analysed, eliminating purification steps during which components might be selectively lost. This was made possible by the relatively low dry weight of extracts from dark-grown tissue and by the high selectivity of a radioactivity detector. Methanolic extracts of root segments were simply filtered, dried <u>in vacuo</u> and redissolved in a small volume of methanol. In most cases they were then centrifuged.

The efficiency of the methanol extraction procedure was always high: usually at least 90%. Thus a relatively small proportion of IAA metabolites

escaped analysis in this way. Similarly, recovery of radioactivity after HPLC analysis was good (ca.95%). In spite of the minimum amount of sample preparation involved, however, a significant and variable amount of radioactivity (up to 36% of the sample) was lost during preparation. Most of this occurred during drying of the sample under reduced pressure (see Mann and Jaworski, 1970). It was not possible to be sure whether some metabolites were selectively lost. This was not thought to be probable, however, for three reasons. Firstly, as the samples were dissolved in the same solvent (i.e. methanol) after drying, differential solubility was unlikely to cause Secondly, although losses were variable, the same products were losses. Thirdly, similar losses occurred during present in each replicate extract. the preparation of control extracts which contained IAA alone. It was thus tentatively concluded that HPLC traces gave a reliable representation of the total spectrum of ¹⁴C-IAA metabolites extracted into methanol from Zea mays root segments which had been incubated in 10^{-3} mol m⁻³ IAA-2-¹⁴C for 2h.

During preparation of the majority of coleoptile samples, reduction of the extract to dryness by rotary evaporation was avoided. Samples were reduced to the aqueous phase then diluted with buffer and purified using C₁₈ Sep-pak cartridges. However, as shown in experiment B.5, peaks 1 and 2 were selectively lost during this procedure.

In addition to eliminating selective purification procedures it was imperative to use an analytical technique which had high resolution. TLC was found to be inadequate for the separation of the large number of metabolites. Reverse-phase HPLC was, however, able to resolve the extract into 12 major components. Some evidence that these peaks might represent individual metabolites was obtained by isocratic chromatography. This could only be confirmed by the use of another analytical technique dependent on different properties of the molecules, e.g. ion-exchange chromatography. These results highlight the inconclusive nature of "identifications" of IAA metabolites based on co-chromatography with standards on paper or TLC systems (see Table 3).

Time-course experiments were carried out to study the interconversions of

products. However, after only 10min incubation at least 3 and sometimes up to 8 radioactive compounds, in roughly similar quantities, were observed in methanolic root extracts. These comprised a mean total of 43% of the radioactivity present in the extracts. The appearance of several products apparently simultaneously indicated that the path of metabolism in this tissue was probably complex and non-linear. Metabolism took place more slowly in coleoptile segments and also appeared less complex. After 1h only two products, represented by peaks 10 and 11 (presumptive oxindole-3-acetic acid and probably an ester of IAA with sugar or <u>myo</u>-inositol) were observed.

Without conclusive identification of metabolites and demonstrations of their interconversions, deductions concerning the pathway of IAA metabolism in <u>Zea mays</u> seedlings remain tentative. However, it would appear that root tissues are capable of very rapid metabolism of IAA by a non-linear and predominantly oxidative reaction pathway. Oxidation did not involve loss of the carboxyl group. Instead, the major product appeared to be oxindole-3-acetic acid. 5-Hydroxyindole-3-acetic acid may also have been formed. There was some evidence for the formation of two conjugates with IAA. One may have been an ester of IAA with a sugar or <u>myo</u>-inositol while the other co-chromatographed with IAA-glycine.

Coleoptiles appeared to metabolise IAA more slowly, at least at short incubation times. Although most of the products found in root extracts were apparently present in coleoptile samples, a single compound which cochromatographed with oxindole-3-acetic acid was predominant. Again none of the major products involved decarboxylation of the IAA.

Certain questions must be answered before these results can be related to the metabolism of endogenous IAA. Firstly, how does the manner in which IAA is supplied to the plant affect its metabolism? When segments are floated on aqueous solutions of IAA, much of the plant growth substance is taken up at cut surfaces. Degradation might occur at this site and have very little relation to endogenous IAA metabolism. The remainder of the IAA will enter

across the epidermis and move towards the centre of the organ. In the case of roots this is likely to give rise to a distribution rather different from that in the intact plant, where the majority of IAA enters the root from, and is located in, the central stele (see Bridges <u>et al.</u>, 1973; Bowen <u>et al.</u>, 1972).

Experiments were carried out to investigate the metabolism of separated cortical and stelar segments. The results agree with those of Greenwood <u>et al</u>. (1973) in that very little IAA metabolism was found to take place in the stele during a 2h incubation. Products from cortical extracts were similar to those from whole segments and metabolism was rapid. Thus the IAA present in the stele, i.e. the bulk of the endogenous compound, would appear to be protected from metabolism. The rate at which IAA, taken up from solution, is metabolised is thus likely to be considerably greater than the overall rate of endogenous IAA metabolism.

An experiment was therefore carried out in which IAA was supplied to protruding portions of stele at the basal ends of root segments. In this way a closer approximation to the natural site of entry of IAA was obtained. As very little IAA was metabolised by stelar segments at short incubation times, this method of donation also avoided the possibility that much of the metabolism was taking place during entry at cut surfaces. IAA supplied in the above manner was metabolised more slowly than IAA taken up from solution; the proportions of radioactivity, in methanol extracts, remaining associated with the IAA peak were 50% and 31% respectively. The distribution of metabolites also differed. Peaks 5,6,8,9,10 and 11 were present as well as a small amount of a new product which was less polar than peak 11. In addition, peak 10 (oxindole-3-acetic acid) was no longer the most prominent metabolite.

Experiments were also carried out in which IAA was supplied to coleoptile segments from agar blocks placed at their apical ends. The apex is believed to be the major source of free IAA to the intact coleoptile (see Cohen and Bandurski, 1982) and transport is polarised in a basipetal direction (e.g.

Hertel and Leopold, 1963). This mode of donation is therefore a closer approximation to the natural source of IAA than that in experiments in which IAA was taken up from a bathing solution. The quantity of IAA metabolised by the coleoptiles supplied by the two different methods could not be directly compared as the experimental procedure was not identical. In both cases the most prominent metabolite was peak 10 (presumptive oxindole-3-acetic acid).

From these observations it can be deduced that the site at which exogenous IAA is supplied to the tissue may have a marked effect on its metabolism, although this was much more pronounced in the root than in the coleoptile. The results of experiments in which IAA was supplied from agar blocks were believed to give a more reliable approximation of the metabolism of endogenous IAA.

ſ

A second question which must be answered when relating the metabolism of exogenous IAA to that of the endogenous compound is, to what extent does the observed metabolism represent the detoxification of abnormally large amounts plant growth substance? To answer this question in part, experiments \mathbf{of} were carried out to look at metabolism in the presence of different external Coleoptiles incubated in solutions of 10^{-3} , 10^{-2} and IAA concentrations. 10^{-1} mol m⁻³ IAA all metabolised similar proportions of the IAA taken up by the tissue. The nature and distribution of metabolites was also not Furthermore, when the effect of different IAA significantly altered. concentrations on segment elongation was studied, maximum growth took place in 10^{-2} mol m⁻³ IAA. The amount of IAA taken up from 10^{-3} mol m⁻³ IAA solution and present in the tissue after 2h incubation was approximately 4 pmol segment⁻¹ (0.20 pmol mg⁻¹ fresh weight). Coleoptile segments supplied with IAA-2-¹⁴C from agar blocks over 4h contained approximately 2.2 pmol segment⁻¹ (0.11 pmol mg⁻¹ fresh weight) of exogenous IAA. Published measurements of the endogenous IAA content of Zea mays coleoptiles (Weiler et al., 1981; cv Anjou 21) using a radioimmunoassay are as follows: 10 pmol extractable "free" IAA per 2.5mm tip; ca. 3 pmol "diffusible" IAA

per mg fresh weight of coleoptile taken from between 2.5 and 10mm from the tip.

The amounts of IAA supplied were therefore of the same order of magnitude or lower, than the endogenous levels. Thus three lines of evidence indicated that the metabolism of IAA observed in the present experiments was unlikely to be a detoxification reaction to reduce abnormally high levels of the compound.

The same results were obtained in experiments with root tissues. Segments incubated in external concentrations of 10^{-4} , 10^{-3} and 10^{-2} mol m⁻³ IAA metabolised comparable proportions of the IAA-2-¹⁴C taken up and contained a similar pattern of products, despite a large increase in uptake of IAA at higher concentrations. After a 2h incubation in 10^{-4} mol m⁻³ IAA solution, the amount of exogenous IAA present in the tissue was calculated to be approximately 5.7 µg kg⁻¹ fresh weight. Measurements of endogenous quantities of IAA in similar portions of root by Bridges <u>et al</u>., (1973) for the variety "Giant White Horsetooth" and by Rivier and Pilet (1974) for the variety Kelvedon 33, were 29 and 76 µg kg⁻¹ fresh weight respectively. Thus the amounts of IAA taken up were an order of magnitude less than endogenous quantities.

A third problem concerns whether metabolism of IAA changes due to the ageing of excised segments. Results of time-course experiments on IAA metabolism in roots indicated that IAA taken up after longer incubation times was more rapidly metabolised. Growth experiments indicated that after 1h there was a progressive decrease in the rate of elongation. It would therefore appear that short-term experiments are more likely to give an accurate representation of IAA metabolism in the intact plant. More work is required to discover whether more immediate wound responses might also have a significant effect on IAA metabolism.

Fourthly, it should be considered whether the 14 C label will have a significant effect on the distribution of products. However, with 14 C at

room temperature the maximum isotope effect is ca. 10% (e.g. Alder et al., 1971).

In summary, it is difficult to relate the results of studies on exogenous IAA metabolism in excised plant tissues to the endogenous metabolism. However, the results obtained in the present study did not appear to be artifacts produced at cut surfaces or consequences of detoxification of abnormally large amounts of hormone. Experiments in which IAA was supplied from agar blocks placed in a situation to mimic the endogenous source of IAA were most likely to give reliable information on endogenous metabolism. In future experiments it would be possible to supply labelled IAA to the endosperm of intact seedlings. As IAA is transported acropetally along the root this is likely to provide an adequate means of donation to this organ. Hall and Bandurski (1978) have shown, however, that when labelled IAA was supplied in this way, 98% of the radioactivity moving into the shoot had been metabolised. This method would therefore not be a feasible means of supplying IAA to the coleoptile. Instead, IAA could be applied to the coleoptile tip of intact seedlings using a micropipette.

Without the results of such studies, considerations of the potential of pathways of IAA metabolism to regulate its endogenous levels are speculative. Nevertheless, it would appear that both root and coleoptile tissues have the capacity for rapid metabolism of IAA. It is likely that the free IAA in the plant has a relatively rapid turnover. Whether or not the metabolism of IAA is a limiting step in regulating its levels, this property would allow the plant to change the amount of IAA present within a short time.

If the level of IAA is controlled by the rate of its metabolism, the latter might be expected to rise as the amount of exogenous IAA taken up is increased. In fact the proportion of IAA metabolised by both root and coleoptile segments was independent of the amount of exogenous IAA taken up over a range of two orders of magnitude. Thus there is some evidence that solely IAA levels might not be controlled by the rate of metabolism. The results of experiments D and E indicated that IAA present in the stele of the root is

largely protected from metabolism, while IAA in the cortex disappears rapidly. The levels of IAA in the root might therefore be regulated by the amount of lateral movement from the stele into the cortex.

Bandurski and coworkers (e.g. Cohen and Bandurski, 1978) have proposed that the reversible conjugation of IAA with sugars or <u>myo</u>-inositol may be important in regulating IAA levels in <u>Zea mays</u> seedlings. The results of this thesis indicated that IAA conjugates comprised only a small fraction of the total spectrum of metabolites of exogenous IAA. It is possible that conjugation may be more important in storage tissues such as the endosperm.

Experiments in which the metabolism of IAA-1- 14 C and IAA-2- 14 C were compared indicated that none of the major products had been formed by decarboxylation. Reinecke and Bandurski (1981) also presented evidence that the initial oxidation of IAA in <u>Zea mays</u> endosperm does not involve the loss of carbon-1 of the side chain. These results call into question the rôle of traditional IAA-oxidases in the <u>in vivo</u> oxidation of IAA in <u>Zea mays</u> seedlings, as these enzyme preparations have been shown to catalyse the decarboxylation of IAA to form either 3-methylen eoxindole or indole-3aldehyde (e.g. Hinman and Lang, 1965; Ricard and Job, 1974).

IAA-oxidase preparations, usually peroxidases, (see Introduction), frequently show low substrate specificity (Butt, 1980). They have been extracted from a wide range of plant species (Schneider and Wightman, 1974). The possibility that these enzymes might be important in regulating IAA levels in higher plants has been studied extensively. Various types of correlations between IAA-oxidase activity and endogenous IAA levels have been reported (e.g. Frenkel, 1972; Atsumi and Hayashi, 1978; Saleh, 1981; Rao <u>et al.</u>, 1982). None of these investigations has shown conclusively that IAA-oxidase can regulate endogenous IAA levels. Briggs <u>et al</u>. (1955) reported no relation between extractable IAA-oxidase activity and the disappearance of IAA <u>in vivo</u> in tissues of the fern <u>Osmunda cinnamomea</u> L.

Despite the lack of evidence for decarboxylation in the present study,

other researchers have found evidence for the decarboxylation of IAA. Substantial quantities of ${}^{14}\text{CO}_2$ may be produced when IAA-1- ${}^{14}\text{C}$ is supplied to a variety of plant species (e.g. Wilkins <u>et al.</u>, 1972b; Grochowska, 1974; Ryugo and Breen, 1974; Epstein and Lavee, 1975; Hamilton <u>et al.</u>, 1976; Menschick and Hild, 1976; Epstein and Lavee, 1977a).

There are at least three possible explanations for this discrepancy. Firstly, decarboxylation might be occurring at cut surfaces catalysed by wound-produced peroxidase enzymes. Evidence for products of decarboxylation in the incubating medium rather than the plant tissue have been found by Andreae and Collet (1968) and Hamilton <u>et al.</u> (1976). In the present study there was some evidence for the presence of decarboxylation products in the incubating solution. Une or sometimes two peaks, which were less polar than IAA, and only appeared when methylene-labelled IAA was supplied, were observed on HPLC traces.

Secondly, it is probable that there may be differences in IAA metabolism between species. Evidence for such differences in the nature of the IAA conjugates formed is substantial. Bandurski and Schulze (1977) studied the endogenous IAA conjugates present in the seeds from several plant species. They discovered that in cereals much of the IAA was esterified, while in legumes peptidyl IAA was more abundant. More conclusive identifications of endogenous IAA conjugates indicate the same trend. Esters of IAA with sugars and <u>myo-inositol have been extracted from Zea mays</u> (Ehmann, 1974; Ehmann and Bandurski, 1974) and <u>Oryza sativa</u> (Hall, 1980) while IAA-glycine was found in soybean (Cohen, 1981). Similar variation may be found in oxidation pathways.

Thirdly, while the initial oxidation of IAA may not involve decarboxylation, this must occur at some stage in its catabolism. In most of the publications quoted, the majority of ${}^{14}\text{CO}_2$ evolution occurred after incubating the tissue for longer than 2h. Indeed preliminary experiments in the present study (results not given) indicated that ${}^{14}\text{CO}_2$ was evolved by Zea mays root segments and intact seedlings incubated in aqueous solutions of IAA-). ${}^{-14}\text{C}$ for longer periods.

In conclusion, until it is shown conclusively that products such as indole-3-aldehyde and 3-methyleneoxindole are actually formed <u>in vivo</u> the rôle of IAA-oxidases remains conjectural.

B. Implications of Results on IAA Metabolism for Research into other Aspects of IAA Physiology

In this thesis it has been shown that exogenous IAA is rapidly and extensively metabolised by the roots and coleoptiles of <u>Zea mays</u> seedlings. As a large proportion of research into the rôle of IAA in higher plants is carried out using exogenous IAA, it is imperative that metabolism during the course of such experiments is taken into consideration.

A large variety of responses may be elicited when IAA is applied to various plant tissues (Table 1). It is possible, however, that metabolites of IAA may also have physiological activity. Indeed, IAA itself might not be the active molecule. A series of papers by Tuli and coworkers (e.g. Tuli and Moyed, 1967 and 1969) presented evidence that 3-methyleneoxindole was responsible for at least some effects of IAA. However, a number of other scientists (e.g. Skytt Andersen <u>et al.</u>, 1972; Evans and Ray, 1973; Roberts, 1974; Evans, 1976 a and b; Hamilton <u>et al.</u>, 1976; Sabater <u>et al.</u>, 1976; Lau <u>et al.</u>, 1978) have since found 3-methyleneoxindole to be devoid of auxin activity and the theory has now been discounted. Nevertheless, other IAA metabolites may well be physiologically active. Kinashi <u>et al</u>. (1976) suggested that the growth-inhibitory substance " β -acid" found in rice bran is



 β -acid

an oxidation product of IAA. Hangarter and coworkers (e.g. Hangarter <u>et</u> <u>al</u>, 1980, and Hangarter and Good, 1981) and Feung <u>et al</u>. (1977) have examined the physiological activity of various amino acid conjugates of IAA: however, results indicated that activity was dependent on their hydrolysis.

The metabolism of IAA must also be considered when planning the duration of experiments on the effects of IAA. After 24h, Zea mays root and coleoptile segments incubated in IAA-2-¹⁴C had metabolised at least 99% of the radioactive IAA taken up by the tissue. The metabolism of IAA in roots was markedly affected by the site of its donation. This may also have a bearing on experiments on the effects of exogenous IAA. When root segments are supplied with IAA from solution, most will enter the tissue through the cortex where it is rapidly metabolised. This must be borne in mind when interpreting the results of experiments such as those of Steen and Hild (1980) and Schurzman and Hild (1980) in which IAA was supplied from agar blocks placed on the side of maize roots for a 4h or 6h period. Firn (1982) noted that IAA, supplied to sunflower hypocotyl sections via the cut surfaces, produced a very weak response. On the other hand, IAA entering through the epidermis induced characteristic rapid growth promotion. The difference in effectiveness of IAA supplied at two different sites could reflect alterations in the rate or pattern of metabolism.

Zea mays roots and coleoptiles have been used extensively in experiments on various aspects of the physiology of IAA. These studies are frequently based on the premise that applied IAA is not metabolised significantly during the experiment. In particular, Zea mays coleoptiles and sometimes roots have been used to investigate IAA binding sites (e.g. Hertel <u>et al.</u>, 1972; Batt <u>et al.</u>, 1976; Ray <u>et al.</u>, 1977; Cross and Briggs, 1979; Moloney and Pilet, 1981). None of these studies mention any form of test to ensure that radioactivity from labelled IAA supplied to cell preparations remains associated with the IAA molecule at the end of the experiment.

It was proposed by Went (1928) that geotropic and phototropic curvatures

in plants were mediated by the lateral transport of growth promoting substance from one side of the organ to the other. After the proposal that the chemical identity of auxin was IAA, several studies were carried out using radiolabelled IAA to investigate its lateral movement in Zea coleoptiles (e.g. Goldsmith and Wilkins, 1964; de la Fuente and Leopold, 1968; Wilkins and Whyte, 1968; Cane and Wilkins, 1969; Hertel et al., 1969; Shaw et al., 1973; Gardner et al., 1974). In some cases, tests for metabolism were carried out using TLC analysis of radioactive compounds (e.g. Cane and Wilkins, 1969; Shaw et al., 1973; Gardner et al., 1974). In the present study TLC was found not to have sufficient resolution to separate the large number of It was therefore thought possible that the amount of IAA IAA metabolites. metabolised may have been underestimated. However, Shaw et al. (1973) found that about 40% of the radioactivity extracted from coleoptile tissue after lh, had been metabolised; i.e. a little more than recorded in this thesis. Given the differences in experimental conditions the results are in reasonable agreement. As in the present experiments Shaw et al. (1973) found only a single peak of radioactivity, corresponding to IAA, on analysis of extracts from receiver blocks of agar. This indicated that IAA metabolites were not mobile and gave further evidence for the specific transport of IAA.

Zea coleoptiles have also been used in the study of polarised basipetal transport of IAA (e.g. Hertel and Leopold, 1963; Edwards and Goldsmith, 1980; Sussman and Goldsmith, 1981). In later papers TLC analysis of extracts indicated a slow rate of metabolism (circa 5-10% of IAA supplied, per hour); in comparison with present results this would appear to be an underestimate.

In <u>Zea mays</u> roots, transport of IAA appears to be polarised in an acropetal direction (e.g. Wilkins and Scott, 1968; Wilkins <u>et al.</u>, 1972a; Martin <u>et al.</u>, 1978) and takes place predominantly in the stele (e.g. Cane and Wilkins, 1970; Bowen <u>et al.</u>, 1972; Shaw and Wilkins, 1974). Some papers (e.g. Wilkins <u>et al.</u>, 1972a and Shaw and Wilkins, 1974) have tested for metabolism of IAA using TLC analysis. The results indicated that although

substantial metabolism took place in the plant tissue, only IAA itself was collected in receiver blocks of agar. This result has been confirmed in this thesis using the higher resolution technique of HPLC.

The results in this thesis thus confirm the validity of the published work on IAA transport. It is suggested, however, that in future more rigorous, quantitative identification of the radioactivity at the end of experiments should be carried out.

C. The Effect of IAA on Root Elongation

As detailed in the Introduction, data on the effect of IAA on root elongation are contradictory. Some investigators have only found evidence for an inhibition of growth by IAA (e.g. Greenwood and Yčas, 1975). Others have observed a small growth promotion at very low concentrations (e.g. Batra <u>et al.</u>, 1975; Edwards and Scott, 1977; Elliott, 1977; Evans <u>et al.</u>, 1980; Mulkey et al., 1981; Pilet and Elliott, 1981). Experiments on the effects of IAA on root growth, described in this thesis, are of a preliminary nature. Nevertheless, some interesting observations have been made.

The growth rate of root segments decreased rapidly after excision. Experiments in this thesis were carried out using short incubation periods (1h). During this time the growth rate of segments was comparable to that of intact roots. Metabolism experiments which showed exogenous IAA to be rapidly metabolised provided a further reason for using short incubation periods.

Other scientists have frequently used longer incubation times; e.g. Batra <u>et al.</u>, 1975 (6 and 2h); Pilet <u>et al.</u>, 1979 (4, 8 and 12h); Pilet and Elliott, 1981 (8h). The deceleration of growth should be considered when interpreting these results. The data of Pilet <u>et al.</u> (1979), may be treated as an example. In this paper the effect of IAA on the growth of root segments which had been stood on agar for 8h was studied. The results were compared with those for freshly cut segments supplied immediately with IAA. Roots left on agar for 8h showed a substantial decrease in endogenous IAA. Furthermore,

low concentrations of IAA, inhibitory to the growth of freshly cut segments, were reported to stimulate growth of roots which had been left on agar for 8h. The authors hypothesised that intact Zea maysroots contained a saturating or supra-optimal level of IAA. A growth-promotory effect was therefore only observed when the endogenous IAA content was depleted. However, root segments left for 8h would show very little growth. It is noteworthy that Baehler and Pilet (1981) also noticed a decrease in the growth rate of segments after Although this appeared to be somewhat slower than reported in excision. this thesis, growth had decreased substantially 8h after excision. It is thus difficult to see how a valid comparison between growth of segments left for 8h and that of freshly cut segments, could be carried out. Furthermore, substantial changes other than a decrease in the IAA levels, will have taken place during 8h after excision. Results presented in this thesis indicated that the decrease in growth was not principally due to lack of IAA. Incubation with various concentrations of exogenous IAA over a 24h period did not maintain elongation at the same rate as in intact roots. It was therefore concluded that the decrease in growth was due to lack of other substances (e.g. nutrients, other plant growth substances) or to the presence of inhibitory compounds such as ethylene. The same conclusion was reached by Greenwood and Yčas (1975). They monitored the growth of root segments after excision. Whole root segments were compared with those from which the stele (the main site of IAA in the root) had been removed. The growth of both groups decreased at the same rate. It is also worth pointing out that if the IAA levels in the root are limiting growth then IAA should not be present in saturating amounts.

More detailed studies of the effect of various concentrations of IAA on growth of root segments were carried out with greater replication. The availability of oxygen was also varied. A promotory effect of IAA on growth was not observed under any circumstances. On the other hand, IAA at a concentration of 10^{-3} mol m⁻³ and in the presence of _{oxygen} inhibited root elongation. To assess the relevance of this work with respect

to the rôle of endogenous IAA, it was important to compare the amounts of IAA taken up from different external concentrations with endogenous quantities. Data on uptake of IAA-2-¹⁴C indicated that with an external concentration of 10^{-3} mol m⁻³ IAA (i.e. inhibitory to growth), the amount of exogenous IAA present in the tissue after 1h incubation was approximately 38 µg per kg fresh weight of tissue. This value was of the same order of magnitude as published measurements of endogenous levels in similar portions of Zea mays root: 29 µg kg⁻¹ for the variety Giant White Horsetooth (Bridges <u>et al.</u>, 1973); 76 µg kg⁻¹ for the variety Kelvedon 33 (Rivier and Pilet, 1974). Thus it is possible that endogenous IAA may have an inhibitory effect on growth.

A comparison of the IAA concentrations reported to promote root growth (e.g. $10^{-4} \text{ mol m}^{-3}$, Edwards and Scott, 1977; 10^{-5} and $10^{-7} \text{ mol m}^{-3}$, Mulkey <u>et al.</u>, 1981; $10^{-5} \text{ mol m}^{-3}$ Pilet and Elliott, 1981) with the results presented in this thesis indicated that the amount of IAA taken up from growth stimulating concentrations was equivalent to only a small fraction of the endogenous content. The amount of exogenous IAA present in segments incubated in 5.9 x $10^{-6} \text{ mol m}^{-3}$ IAA for 1h was approximately 0.64 µg per kg fresh weight of tissue; i.e. <u>circa</u> 1% of endogenous levels.

Accordingly, IAA uptake should be viewed as a critical factor in assessing the significance of experiments on the effects of low concentrations of IAA on root growth. Nevertheless, it is possible that IAA may be located in more than one discrete compartment and that measurements of the total IAA content of the tissue may mask the levels of "effective" IAA. It must also be noted that published measurements of endogenous IAA were carried out using different varieties of <u>Zea mays</u> from that employed in the present study. Pilet and Elliott (1981) suggested that there may be substantial differences in the endogenous IAA levels of different cultivars.

Significant interaction was found between oxygen availability and the effect of IAA. IAA did not inhibit the growth of segments incubated under nitrogen. The effect of oxygen was to increase the growth of roots incubated in control solution and in low concentrations of IAA. IAA at a concentration

of 10^{-3} mol m⁻³ then reduced the growth rate to that observed for segments incubated under nitrogen at any concentration of IAA.

This dependence of the inhibitory effect of exogenous IAA on the presence of supplementary oxygen may indicate that root segments incubated in aqueous solutions rapidly become depleted of oxygen, and that the intact roots possess an oxygen-dependent component of growth which may be inhibited by IAA. Segments incubated in solutions not bubbled with gas showed a small growth inhibition in the presence of 10^{-3} mol m⁻³ IAA. On the other hand, the levels of oxygen supplied to roots bubbled with pure oxygen will be substantially higher than those encountered <u>in vivo</u> and the resulting high growth rate may be artifactual.

To summarise, the results presented in this thesis indicated that exogenous IAA is only able to inhibit <u>Zea</u> root growth. Several publications reporting a growth promotory effect used long incubation times (2h or more) during which ageing of the tissue, shortage of essential substances and possibly a build-up of wound-produced inhibitory compounds may have affected the response to IAA. In addition, calculations of uptake of ¹⁴C-IAA from various concentrations indicated that the amount of exogenous IAA taken up from the low concentrations reported to promote growth, may be only a small fraction of the endogenous content.

On the other hand, it was suggested by Chadwick and Burg (1967), that the inhibition of root growth, induced by exogenous IAA may be a result of IAA-stimulated ethylene biosynthesis. This was investigated by Mulkey <u>et al</u>. (1981). Roots were pretreated with the ethylene biosynthesis inhibitor aminoethoxyvinylglycine before being placed in solutions of IAA. This treatment increased growth of controls and permitted growth enhancement by 10^{-3} and 10^{-4} mol m⁻³ IAA; higher concentrations remained inhibitory.

The site at which IAA is supplied to root segments might also affect results. Changes in the method of applying IAA-2- 14 C resulted in variation in the rate and pattern of its metabolism. Firn (1982) also noticed that the site of IAA uptake into sunflower hypocotyls substantially affected the

growth response.

Evans <u>et al.</u> (1980) reported that the inhibitory effect of IAA on root growth was accompanied by an increase in the pH of the medium. This was not found in the present study. A pH increase occurred in all solutions containing root segments, irrespective of the presence or concentration of IAA. It is possible that the discrepancy is a result of small differences in experimental procedure; the above authors used buffered incubating solution, and a larger number of sections for the volume of medium.

D. Future Experiments

The present study has emphasised the need to investigate the whole spectrum of IAA metabolites produced within a single species. Conclusive identification of these products is now required. A comparison of mass spectra of metabolites with those of pure compounds is likely to prove the most suitable method of definitive characterisation. The following compounds may prove to be useful standards, in addition to those already synthesised: 3-hydroxyoxindole-3-acetic acid, 5-hydroxyoxindole-3-acetic acid, 3,5dihydroxyoxindole-3-acetic acid, and conjugates of IAA with <u>myo</u>-inositol and glucose.

The sequence in which metabolites are produced also requires investigation. This is likely to prove easier in the coleoptile where the metabolic pathway appears to be less complex than in the root. Radioactively-labelled IAA metabolites could be supplied to the plant tissue and the products isolated and identified.

The overall aim of a study on IAA metabolism should be to obtain information leading to an understanding of the role of IAA in the intact plant. The site at which IAA was supplied to plant tissue was found to have a substantial effect on the metabolism pattern in roots. It is therefore suggested that in future experiments, the metabolism of IAA supplied to the intact plant by means, and in amounts, which mimic the endogenous source should be investigated. Application to the endosperm

might serve as a suitable site for root studies, while IAA could be supplied to the coleoptile tips from micropipettes. Extractions should also be carried out to determine whether the exogenous IAA metabolites are naturally present within the plant. Possible diurnal variations in the rate and/or path of IAA metabolism should be considered (see Sandberg et al., 1982).

Once the pathway of IAA metabolism has been established it should be possible to identify the enzymes catalysing these reactions. Proteins catalysing the esterification of IAA with glucose and myo-inositol have already been isolated (Michalczuk and Bandurski, 1980). Enzymes involved in the oxidation of IAA to compounds such as oxindole-3-acetic acid have not The rôle of "IAA-oxidases" requires further investigation, vet been studied. It would appear unlikely that this type of enzyme is normally responsible for the in vivo oxidation of IAA in Zea mays seedlings. However, further study of IAA metabolism in other species might provide evidence for an IAAoxidase-catalysed decarboxylation reaction. An investigation of the enzymes involved in IAA metabolism might reveal correlations between their activity and endogenous levels of IAA and thus provide information as to how endogenous quantities of IAA are controlled. The subcellular location of IAA metabolism and the products should also be examined. This might be achieved by the use of autoradiography and cellular fractionation techniques. In order to understand how the activity of IAA is controlled, the physiological activities of IAA metabolites should be studied. Hangarter et al. (1980) and Feung et al. (1977) have investigated the physiological activity of several amino acid conjugates. 3-Methyleneoxindole has also been studied extensively in this context (e.g. Tuli and Moyed, 1969). The activity of compounds such as oxindole-3-acetic, however, has not been tested. Experiments using esters of IAA with myo-inositol and glucose would be complicated by the ease with which these compounds are hydrolysed.

Bandurski and co-workers (e.g. Cohen and Bandurski, 1978) have suggested that IAA may be oxidised as a result of its producing a physiological response. This hypothesis might be tested by examining possible correlations

between the rate of IAA oxidation and tissue sensitivity.

Experiments on the effect of IAA on root elongation have highlighted several points which should be borne in mind in future experiments: It is essential that the quantity of exogenous IAA entering the root is of the same order of magnitude as the endogenous content. The level of IAA in the tissue and <u>Zea mays</u> variety used should therefore be measured and compared with the uptake of IAA during growth experiments. The importance of subcellular compartmentation should also be investigated.

IAA should be applied to the root in a manner which most closely mimics the natural source of the compound. Donation to the endosperm or to a portion of stele protruding from the basal end of segments might be suitable.

The metabolic fate of IAA should be correlated with growth effects and the bioactivity of metabolites considered. The rate of IAA metabolism would be expected to influence choice of the duration of experiments.

The use of buffers is to be discouraged, as these may affect the growth in their own right (e.g. Edwards and Scott, 1976). Nevertheless the pH of solutions may be important in determining uptake of IAA (see Edwards and Goldsmith, 1980). The levels of oxygen available to the tissue presents another variable to be investigated. The effect of oxygen on IAA-controlled growth needs further study.

Finally the role of ethylene in the response to IAA requires clarification. This would involve the use of inhibitors of its biosynthesis and mode of action, or removal of ethylene (e.g. by mercuric perchlorate). The production of ethylene in the presence and absence of IAA should be studied, as should the response to exogenous ethylene.

The results presented in this thesis must be viewed in the context of the role of endogenous IAA in the correlative control of growth and differentiation in Zea mays seedlings. Early work (e.g. Went, 1928) provided evidence that auxin acted as a chemical messenger within the plant. Confirmation of this theory requires unequivocal proof that IAA is synthesised at a discrete site, from where it is transported to specific site(s) of An understanding of the role of plant growth substances is most action. likely to be achieved by the detailed study of a relatively simple plant system. Young Zea mays seedlings represent a compact experimental system, exhibiting most of the correlative features of vegetative plant growth and have been used extensively for research on the rôle of IAA. It is now beyond dispute that endogenous IAA is present in extracts of roots, endosperm, mesocotyls and coleoptiles of dark-grown Zea mays seedlings. Furthermore, the zone of cell expansion in the coleoptile is at least one site of action. Levels of exogenous IAA comparable with the endogenous pool size inhibit Exogenous IAA is metabolised rapidly by coleoptiles and root elongation. roots. Moreover, cortical and stelar tissues of the root metabolise IAA The exogenous compound is specifically at substantially different rates. transported in a polarised fashion in both roots and coleoptiles. Recent advances in analytical technology have made possible the conclusive identification of minute quantities of plant growth substances in complex plant extracts. Methods also exist for determining the intracellular These technological developments, coupled with an location of compounds. increasing understanding of the effects, transport and metabolism of exogenous IAA provide encouraging signs for clarifying the role of endogenous IAA in the life of the intact plant.

REFERENCES

- Abe, H., Uchiyama, M. and Sato, R. (1972). Isolation and identification of native auxin in marine algae. Agr. Biol. Chem., <u>36</u>, 2259-2260.
- Abeles, F.B. (1966). Auxin stimulation of ethylene evolution. Plant Physiol., <u>41</u>, 585-588.
- Addicott, F.T., Carns, H.R., Lyon, J.L., Smith, O.E. and McMeans, J.L. (1964).
 On the physiology of Abscisins. In: Colloques Internationaux 123.
 Regulateurs naturels de la croissance végétale. pp 687-703. C.N.R.S. Paris.
- Alder, R.W., Baker, R. and Brown, J.M. (1971)*. Mechanism in Organic Chemistry. p 17. Wiley, Chichester.
- Andreae, W.A. and Collet, G. (1968). The effect of phenolic substances on the growth activity of indoleacetic acid applied to pea root or stem sections. In: Biochemistry and Physiology of Plant Growth Substances. Proceedings 6th International Conference on Plant Growth Substances: Eds. F. Wightman and G. Setterfield. pp 553-561. Runge Press, Ottawa.
- Andreae, W.A. and Good, N.E. (1955). The formation of indoleacetylaspartic acid in pea seedlings. Plant Physiol., <u>30</u>, 380-382.
- Aspinall, P., Paleg, L.G. and Addicott, F.T. (1967)*. Abscisin II and some hormone-regulated plant responses. Aust. J. Biol. Sci., <u>20</u>, 869-882.
- Atsumi, S. and Hayashi, T. (1978). The relationship between auxin concentration, auxin protection and auxin destruction in crown gall cells cultured <u>in vitro</u>. Plant Cell Physiol., <u>19</u>, 1391-1397.
- Avery, G.S. and Johnson, E.B. (1947)*. Hormones and Horticulture. McGraw-Hill, New York.
- Baehler, W. and Pilet, P.E. (1981). Kinetic analysis of root growth and georeaction. Physiol. Plant., 51, 27-32.
- Bandurski, R.S. and Schulze, A. (1974). Concentrations of indole-3-acetic acid and its esters in <u>Avena and Zea</u>. Plant Physiol., <u>54</u>, 257-262.
 Bandurski, R.S. and Schulze, A. (1977). Concentrations of indole-3-acetic acid and its derivatives in plants. Plant Physiol., 60, 211-213.

- Batra, M.W., Edwards, K.L. and Scott, T.K. (1975). Auxin transport in roots: Its characteristics and relationship to growth. In: The Development and Function of Roots. Eds. J.G. Torrey and D.T. Clarkson. pp 299-325. Academic Press.
- Batt, S., Wilkins, M.B. and Venis, M.A. (1976). Auxin binding to corn coleoptile membranes, kinetics and specificity. Planta, <u>130</u>, 7-13.
- Bayliss, W.M. and Starling, E.H. (1902). On the causation of the so-called peripheral reflex secretion of the pancreas (preliminary communication). Proc. Roy. Soc., 69, 352-353.
- Bayliss, W.M. and Starling, E.H. (1904). The chemical regulation of the secretory process. Proc. Roy. Soc., 73, 310-322.
- BeMiller, J.N. and Colilla, W. (1972). Mechanism of corn indole-3-acid oxidase in vitro. Phytochem., 11, 3393-3402.
- Benes, J., Veres, K., Chvojka, L. and Friedrich, A. (1965). New types of kinins and their action on fruit tree species. Nature, 206, 830-831.
- Bentley, J.A. and Bickle, A.S. (1952). Studies on plant growth hormones. II. Further biological properties of 3-indolyl-acetonitrile. J. Exp. Bot., 3, 406-423.
- Bentrup, F.W. (1979). Reception and transduction of electrical and mechanical stimuli. In: Encyclopedia of plant physiology, Vol. 7. Physiology of Movements. Eds. W. Haupt and M.E. Feinleib. pp 42-70. Springer.
- Birecka, H. and Miller, A. (1974). Cell wall and protoplast isoperoxidases in relation to injury, indoleacetic acid and ethylene effects. Plant Physiol., 53, 569-574.
- Bornman, C.H., Spurr, A.R. and Addicott, F.T. (1967)*. Abscisin, auxin and gibberellin effects on the developmental aspects of abscission in cotton (Gossypium hirsutum). Am. J. Bot., 54, 125-135.
- Bowen, M.R., Wilkins, M.B., Cane, A.R. and McCorquodale, I. (1972). Auxin transport in roots. VIII. The distribution of radioactivity in the tissues of Zea root segments. Planta, 105, 273-292.

- Boysen-Jensen, P. (1910)*. Über die leitung des phototropischen reizes in avenakeimpflanzen. Berichte der Deutschen botanischen Gesellschaft, <u>28</u>, 118-120.
- Brian, P.W. and Hemming, H.G. (1955)*. The effect of gibberellic acid on shoot growth of pea seedlings. Physiol. Plant., 8, 669-681.
- Brian, P.W., Hemming, H.G. and Radley, M. (1955). A physiological comparison of gibberellic acid with some auxins. Physiol. Plant., 8, 899-912.
- Bridges, I.G., Hillman, J.R. and Wilkins, M.B. (1973). Identification and localisation of auxin in primary roots of <u>Zea mays</u> by mass spectrometry. Planta, 115, 189-192.
- Briggs, W.R., Steeves, T.A., Sussex, I.M. and Wetmore, R.H. (1955). A comparison of auxin destruction by tissue extracts and intact tissues of the fern Osmunda cinnamomea L. Plant Physiol., 30, 148-155.
- Bryant, S.D. and Lane, F.E. (1975). A study of pea indole-3-acetic acid oxidases and horseradish peroxidase. Plant Physiol., 56, Suppl. 17.
- Burg, S.P. (1962). The physiology of ethylene formation. Ann. Rev. Plant Physiol., 13, 265-302.
- Burg, S.P. (1968). Ethylene, plant senescence and abscission. Plant Physiol., 43, 1503-1511.
- Butt, V.S. (1980). Direct oxidases and related enzymes. In: The Biochemistry of Plants, A Comprehensive Treatise. Vol. 2. Metabolism and Respiration. Ed. D.D. Davies. pp 81-123. Academic Press.
- Cane, A.R. and Wilkins, M.B. (1969). Independence of lateral and differential longitudinal movement of indole-3-acetic acid in geotropically stimulated coleoptiles of Zea mays. Plant Physiol., <u>44</u>, 1481-1487.
- Cane, A.R. and Wilkins, M.B. (1970). Auxin transport in roots. VI. Movement of IAA through different zones of <u>Zea</u> roots. J. Exp. Bot., <u>21</u>, 212-218.

Chadwick, A.V. and Burg, S.P. (1967). An explanation of the inhibition of root growth caused by indole-3-acetic acid. Plant Physiol., <u>42</u>, 415-420.

Clark, H.E. and Kerns, K.R. (1942). Control of flowering with phytohormones. Science, 95, 536-537.

- Coggins, C.W. and Lewis, L.N. (1962). Regreening of Valencia orange as influenced by potassium gibberellate. Plant Physiol., <u>37</u>, 625-627.
- Cohen, J.D. (1981). Indolylic auxins of soybean seed. Plant Physiol., <u>67</u>, Suppl. 2.
- Cohen, J.D. and Bandurski, R.S. (1978). The bound auxins : protection of indole-3-acetic acid from peroxidase-catalysed oxidation. Planta, 139, 203-208.
- Cohen, J.D. and Bandurski, R.S. (1982). Chemistry and physiology of bound auxins. Ann. Rev. Plant Physiol., 33, 403-454.
- Collet, G.F. (1968)*. Metabolisme intracellulaire et extracellulaire de l'acide 3-indolylacétique dans les culture aseptiques ou non aseptiques de points de racine du <u>Pisum sativum</u>. Can. J. Bot., <u>46</u>, 969-978.
- Crane, J.C. (1964)*. Growth substances in fruit settling and development. Ann. Rev. Plant Physiol., 15, 303-326.
- Crane, J.C. (1965). The chemical inductions of parthenocarpy in the Calimyrna fig and its physiological significance. Plant Physiol., <u>40</u>, 606-610.
- Cross, J.W. and Briggs, W.R. (1979). Solubilised auxin-binding protein. Subcellular localisation and regulation by a soluble factor from homogenates of corn shoots. Planta, <u>146</u>, 263-270.
- Cross, B.E., Grove, J.F., McClosky, P., MacMillan, J., Moffatt, J.S. and Mulholland, T.P.C. (1961). The structures of fungal gibberellins. Adv. Chem. Ser., <u>28</u>, 3-17.
- Crozier, A., Loferski, K., Zaerr, J.B. and Morris, R.O. (1980). Analysis of picogram quantities of indole-3-acetic acid by high performance liquid chromatography-fluorescence procedures. Planta, 150, 366-376.
- Crozier, A. and Reeve, D.R. (1977). The application of high performance liquid chromatography to the analysis of plant hormones. In: Plant Growth Regulation. Ed. P.E. Pilet. pp 67-76. Springer, Berlin.

- Curry, G.M. (1969). Phototropism. In: The Physiology of Plant Growth and Development. Ed. M.B. Wilkins. pp 245-277. McGraw-Hill, London.
- Darwin, C. and Darwin, F. (1880)*. The Power of Movement in Plants. Murray, London.
- Davies, P.J. (1972). The fate of exogenously applied indoleacetic acid in light grown stems. Physiol. Plant., 27, 262-270.
- De Greef, J.A., Van Hoof, R. and Caubergs, R. (1977). Light induced changes in auxin metabolism during hook opening of etiolated bean seedlings. Biochem. Soc. Transact., <u>5</u>, 1049-1051.
- De La Fuente, R.K. and Leopold, A.C. (1968). Geotropism and lateral movement of auxin. In: Biochemistry and Physiology of Plant Growth Substances. Proceedings 6th International Conference on Plant Growth Substances. Eds. F. Wightman and G. Setterfield. pp 1129-1144. Runge Press, Ottawa.
- Dolk, H.E. (1930). Geotropie en groeistoff. Doctoral dissertation, University of Utrecht. pp 129.
- Durley, R.C. and Kannangara, J. (1976). Use of high pressure liquid chromatography for the analysis of plant growth regulators in <u>Sorghum</u> <u>bicolor</u> L. Moench. In: Abstracts 9th International Conference on Plant Growth Substances. pp 81-83.
- Edwards, K.L. and Goldsmith, M.H.M. (1980). pH dependent accumulation of indoleacetic acid by corn coleoptile sections. Planta, 147, 457-466.
- Edwards, K.L. and Scott, T.K. (1974). Rapid growth responses of corn root segments : Effect of pH on elongation. Planta, 119, 27-37.
- Edwards, K.L. and Scott, T.K. (1976). Rapid growth responses of corn root segments : Effect of citrate-phosphate buffer on elongation. Planta, <u>129</u>, 229-233.
- Edwards, K.L. and Scott, T.K. (1977). Rapid growth responses of corn root segments : Effect of auxin on elongation. Planta, <u>135</u>, 1-5.
- Ehmann, A. (1974). Isolation of 2-<u>O</u>-(indole-3-acetyl)-D-glucopyranose, 4-<u>O</u>-(indole-3-acetyl)-D-glucopyranose and 6-<u>O</u>-(indole-3-acetyl)-Dglucopyranose from kernels of <u>Zea mays</u> by gas liquid chromatography-mass spectrometry. Carbohydr. Res., <u>34</u>, 99-114.

- Ehmann, A. and Bandurski, R.S. (1974). The isolation of di-<u>O</u>-(indole-3acetyl)-<u>myo</u>-inositol and tri-<u>O</u>-(indole-3-acetyl)-<u>myo</u>-inositol from mature kernels of <u>Zea mays</u>. Carbohydr. Res., <u>36</u>, 1-12.
- El-Antably, H.M.M. and Wareing, P.F. (1966). Stimulation of flowering in certain short-day plants by abscisin. Nature, 210, 328-329.
- Eliasson, L. (1969)*. Growth regulators in <u>Populus tremula</u>. I. Distribution of auxin and growth inhibitors. Physiol. Plant., 22, 1288-1301.

Elliott, M.C. (1977). Auxins and the regulation of root growth. In:

Plant Growth Regulation. Ed. P.E. Pilet. pp 100-108. Springer, Berlin.

Elliott, M.C. and Greenwood, M.S. (1974). Indol-3yl-acetic acid in roots of Zea mays. Phytochem., 13, 239-241.

Encyclopedia Britannica. 1980 Year Book. p 164.

- Epstein, E., Cohen, J.D. and Bandurski, R.S. (1980). Concentration and metabolic turnover of indoles in germinating kernels of <u>Zea mays</u> L. Plant Physiol., 65, 415-421.
- Epstein, E., Klien, I. and Lavee, S. (1975). Uptake and fate of IAA in apple callus tissues : metabolism of IAA-2-¹⁴C. Pl. Cell Physiol., 16, 305-311.
- Epstein, E., Kochba, J. and Neumann, H. (1977). Metabolism of indoleacetic acid by embryog^enic and non-embryogenic callus lines of 'Shamouti' orange (<u>Citrus sinensis</u> Osb.). Z. Pfl. physiol., <u>85</u>, 263-268.
- Epstein, E. and Lavee, S. (1975). Uptake and fate of IAA in apple callus tissue using IAA-1-¹⁴C. P1. Cell Physiol., <u>16</u>, 553-561.
- Epstein, E. and Lavee, S. (1977a). Uptake, translocation and metabolism of IAA in the olive (<u>Olea europa</u>). I. Uptake and translocation of $[1-^{14}C]$ IAA in detached 'Manzanilla' olive leaves. J. Exp. Bot., <u>28</u>, 619-628.
- Epstein, E. and Lavee, S. (1977b). Uptake, translocation and metabolism of IAA in the olive (<u>Olea europa</u>). II. The fate of exogenously applied [2-¹⁴C]IAA in detached 'Manzanilla' olive leaves. J. Exp. Bot., 28, 629-635.

- Erickson, R.O. and Sax, K.B. (1956). Rates of cell division and cell elongation in the growth of the primary root of <u>Zea mays</u>. Proc. Am. Phil. Soc., <u>100</u>, 499-514.
- Esashi, Y. and Leopold, A.C. (1969). Cotyledon expansion as a bioassay for cytokinins. Plant Physiol., 44, 618-620.
- Evans, M.L. (1976a). Effects of auxin, acidic pH and 3-methyleneoxindole on intact root elongation. In: Abstracts 9th International Conference on Plant Growth Substances. pp 91-92.
- Evans, M.L. (1976b). Comparative effects of indole-3-acetic acid and 3-methyleneoxindole on intact lentil root elongation. Pl. Sci. Lett., <u>7</u>, 447-453.

Evans, M.L. and Mulkey, T. (1980). Correlation between auxin effects on H^+ movement and growth in roots. Plant Physiol., <u>65</u>, Suppl. 130.

Evans, M.L., Mulkey, T.J. and Vesper, M.J. (1980). Auxin action on proton influx in corn roots and its correlation with growth. Planta, <u>148</u>, 510-512.
Evans, M.L. and Ray, P.M. (1973). Inactivity of 3-methyleneoxindole as a

mediator of auxin action on cell elongation. Plant Physiol., <u>52</u>, 186-189. Feldman, L.J. (1980). Auxin biosynthesis and metabolism in isolated roots of

Zea mays. Physiol. Plant., 49, 145-150.

- Feung, C.S., Hamilton, R.H. and Mumma, R.O. (1976). Metabolism of indole-3acetic acid. III. Identification of metabolites isolated from crown gall callus tissue. Plant Physiol., <u>58</u>, 666-669.
- Feung, C.S., Hamilton, R.H. and Mumma, R.O. (1977). Metabolism of indole-3acetic acid. IV. Biological properties of amino acid conjugates. Plant Physiol., 59, 91-93.
- Firn, R.D. (1982). Auxin uptake, proton uptake and cell elongation in sunflower hypocotyl sections. In: Abstracts 11th International Conference on Plant Growth Substances. p 10.
- Firn, R.D. and Digby, J. (1980). The establishment of tropic curvatures in plants. Ann. Rev. Plant Physiol., <u>31</u>, 131-148.

- Fletcher, R.A. and Osborne, D.J. (1965)*. Regulation of protein and nucleic acid synthesis by gibberellin during leaf senescence. Nature, <u>207</u>, 1176-1177.
- Frenkel, C. (1972). Involvement of peroxidase and indole-3-acetic acid oxidase isozymes from pear, tomato and blueberry fruit in ripening. Plant Physiol., 49, 759-763.
- Gabella, M. and Pilet, P.E. (1978). Effects of pH on georeaction and elongation of maize root segments. Physiol. Plant., <u>44</u>, 157-160.
- Galston, A.W. (1957). Studies on indoleacetic acid oxidase and its inhibitor in light grown peas. Plant Physiol., 32, Suppl. XXI.
- Gardner, G.K., Shaw, S. and Wilkins, M.B. (1974). IAA transport during the phototropic responses of intact Zea and <u>Avena</u> coleoptiles. Planta, <u>121</u>, 237-251.
- Gelinas, D.A. (1973). Proposed model for the peroxidase-catalysed oxidation of indole-3-acetic acid in the presence of the inhibitor ferulic acid. Plant Physiol., <u>51</u>, 967-972.
- Goldsmith, M.H.M. (1977). The polar transport of auxins. Ann. Rev. Plant Physiol., <u>28</u>, 439-478.
- Goldsmith, M.H.M. and Wilkins, M.B. (1964). Movement of auxin in coleoptiles of Zea mays L. during geotropic stimulation. Plant Physiol., <u>39</u>, 151-162.
- Greenwood, M.S., Hillman, J.R., Shaw, S. and Wilkins, M.B. (1973). Localisation and identification of auxin in roots of <u>Zea mays</u>. Planta, <u>109</u>, 369-374.
- Greenwood, M.S., Shaw, S., Hillman, J.R., Ritchie, A. and Wilkins, M.B. (1972). Identification of auxin from <u>Zea</u> coleoptile tips by mass spectrometry. Planta, <u>108</u>, 179-183.
- Greenwood, M.S. and Yčas, J. (1975). The role of stelar auxin in the development of the primary root of Zea mays. Planta, 122, 311-314.
- Grochowska, M.J. (1974). Photolytic decarboxylation of carboxyl ¹⁴C-labelled indol-3yl-acetic acid in leaves of the apple tree. J. Exp. Bot., <u>25</u>, 638-645.

- Haagen Smit, A.J., Dandliker, W.B., Wittner, S.H. and Murneek, A.E. (1946). Isolation of 3-indoleacetic acid from immature corn kernels. Am. J. Bot., <u>33</u>, 118-120.
- Hall, P.J. (1980). Indole-3-acetyl-<u>myo</u>-inositol in kernels of <u>Oryza sativa</u>. Phytochem., <u>19</u>, 2121-2123.
- Hall, P.J. and Bandurski, R.S. (1978). Movement of indole-3-acetic acid and tryptophan-derived indole-3-acetic acid from the endosperm to the shoot of Zea mays L. Plant Physiol., 61, 425-429.
- Hall, P.J. and Bandurski, R.S. (1981). Hydrolysis of ³H-IAA-<u>myo</u>-inositol by extracts of Zea mays. Plant Physiol., <u>67</u>, Suppl. 2.
- Hall, S.M. and Medlow, G.C. (1974). Identification of IAA in phloem and root pressure saps of <u>Ricinus communis</u> L. by mass spectrometry. Planta, <u>119</u>, 257-261.
- Hamilton, R.H., Meyer, H.E., Burke, R.E., Feung, C.S. and Mumma, R.O. (1976). Metabolism of indole-3-acetic acid. II. Oxindole pathway in Parthenocissus tricuspidata crown-gall tissue cultures. Plant Physiol., <u>58</u>, 77-81
- Hangarter, R.P. and Good, N.E. (1981). Evidence that IAA conjugates are slow release sources of free IAA in plant tissues. Plant Physiol., <u>68</u>, 1424-1427.
- Hangarter, R.P., Peterson, M.D. and Good, N.E. (1980). Biological activities of indoleacetylamino acids and their use as auxins in tissue culture. Plant Physiol., <u>65</u>, 761-767.
- Hertel, R., De La Fuente, R.K. and Leopold, A.C. (1969). Geotropism and the lateral transport of auxin in the corn mutant amylomaize. Planta, <u>88</u>, 204-214.
- Hertel, R. and Leopold, A.C. (1963).* Versuche zur analyse des auxintransports in der ^koleoptile von <u>Zea mays</u> L. Planta, <u>59</u>, 535-562.
- Hertel, R., Thomson, K.S. and Russo, V.E.A. (1972). <u>In vitro</u> auxin binding to particulate cell fractions from corn coleoptiles. Planta, <u>107</u>, 325-340.

- Hewitt, S. (1980). Steroidal oestrogens and the regulation of growth and development in <u>Phaseolus vulgaris</u> L. Ph.D. Thesis, University of Glasgow.
- Hinman, R.L. and Bauman, C.P. (1964). Reaction of <u>N</u>-bromosuccinimide and indoles, a simple synthesis of 3-bromosuccinimide and indoles, a simple synthesis of 3-bromo-oxindoles. J. Org. Chem., <u>29</u>, 1206-1215.
- Hinman, R.L. and Lang, J. (1965). Peroxidase-catalysed oxidation of indole-3-acetic acid. Biochem., 4, 144-158.
- Hopping, M.E. and Bukovac, M.J. (1975). Endogenous plant growth substances in developing fruit of <u>Prunus cerasus</u> L. III. Isolation of indole-3acetic acid from the seed. J. Am. Soc. Hort. Sci., <u>100</u>, 384-386.
- Horng, A.J. and Yang, S.F. (1975). Aerobic oxidation of indole-3-acetic acid with bisulphite. Phytochem., 14, 1425-1428.
- Igoshi, M., Yamaguchi, I., Takahashi, N. and Hirose, K. (1971). Plant growth substances in the young fruit of <u>Citrus unshiu</u>. Agr. Biol. Chem., <u>35</u>, 629-631.
- Jacobs, W.P. (1952). The role of auxin in differentiation of xylem round a wound. Am. J. Bot., 39, 301-309.
- Jain, M.L., Kadkade, P.G. and Van Huysse, P. (1969). The effect of growth regulating chemicals on abscission and IAA-oxidising enzyme system of dwarf bean seedlings. Physiol. Plant., <u>22</u>, 1038-1042.
- Jones, R.L. (1973). Gibberellins their physiological role. Ann. Rev. Plant Physiol., <u>24</u>, 571-598.
- Kamerbeek, G.A. (1956). Peroxydase content of dwarf types and giant types of plants. Acta Bot. Neerl., 5, 257-263.
- Kang, B.G., Yokum, C.S., Burg, S.P. and Ray, P.M. (1967)*. Ethylene and carbon dioxide : mediation of hypocotyl hook opening response. Science, <u>156</u>, 958-959.
- Kenten, R.H. (1955). The oxidation of indoly1-3-acetic acid by waxpod bean root sap and peroxidase systems. Biochem. J., 59, 110-121.
- Kinashi, H., Suzuki, Y., Takeuchi, S. and Kawarada, A. (1976). Possible metabolic intermediates from IAA to β-acid in rice bran. Agr. Biol. Chem., <u>40</u>, 2465-2470.

- Kögl, F., Haagen Smit, A.J. and Erxleben, H. (1934)*. Heteroauxin abs stoffwechselprodukt niederer pflanzlichen organismen. Isolierung aus hefe. XIII Mitt. Zeit. physiol. Chem., <u>228</u>, 104-112.
- Kokkinakis, D.M. and Brooks, J.L. (1979). Hydrogen peroxide-mediated oxidation of indole-3-acetic acid by tomato peroxidase and molecular oxygen. Plant Physiol., 64, 220-223.
- Kopcewicz, J., Ehmann, A. and Bandurski, R.S. (1974). Enzymatic esterification of indole-3-acetic acid to <u>myo</u>-inositol and glucose. Plant Physiol., <u>54</u>, 846-851.
- Ku, H.S., Yang, S.F. and Pratt, H.K. (1970). Inactivity of apoperoxidase in indoleacetic acid oxidation and ethylene formation. Plant Physiol., <u>45</u>, 358-359.
- Kuiper, P.J.C. (1972)*. Water transport across membranes. Ann. Rev. Plant Physiol., <u>23</u>, 157-172.
- Lado, P., DeMichelis, M.I., Cerana, R. and Marré, E. (1976). Fusicoccin-induced K⁺-stimulated proton secretion and acid-induced growth of apical root segments. Plant Sci. Lett., <u>6</u>, 5-20.
- Lado, P., Rasi-Caldogno, F., Pennacchioni, A. and Marré, E. (1973). Mechanism of the growth-promoting action of fusicoccin : interaction with auxin and effects of inhibitors of respiration and protein synthesis. Planta, 110, 311-320.
- La Motte, C.E. and Jacobs, W.P. (1963). A role of auxin in phloem regeneration in coleus internodes. Dev. Biol., <u>8</u>, 80-98.
- Lang, A. (1956). Induction of flower formation in biennial Hyocyamus by treatment with gibberellin. Naturwiss., 43, 284-285.
- La Rue, C.D. (1936). The effect of auxin on the abscission of petioles. Proc. Nat. Acad. Sci., <u>22</u>, 254-259.
- Lau, O.L., John, W.W. and Yang, S.F. (1978). Inactivity of oxidation products of indole-3-acetic acid on ethylene production in mung bean hypocotyls. Plant Physiol., <u>61</u>, 68-71.

- Laurema, S. (1974). Indoleacetic acid oxidases in resting cereal grains. Physiol. Plant., 30, 301-306.
- Lavee, S. and Epstein, E. (1976). Metabolic differences in degradation of ¹⁴C-IAA in apple and olive tissue cultures. Abstracts 9th International Conference on Plant Growth Substances. pp 216-218.
- Lee, T.T. (1974). Cytokinin control in subcellular localisation of indoleacetic acid oxidase and peroxidase. Phytochem., 13, 2445-2453.
- Lee, T.T. (1980). Effects of phenolic substances on metabolism of exogenous indole-3-acetic acid in maize stems. Physiol. Plant., <u>50</u>, 107-112.
- Letham, D.S. (1963). Zeatin, a factor inducing cell division isolated from Zea mays. Life Sci., 8, 569-573.
- Libbert, E. and Rische, H. (1969). Interactions between plants and epiphytic bacteria regarding their auxin metabolism. V. Isolation and identification of IAA producing and destroying bacteria from pea plants. Physiol. Plant., 22, 51-58.
- Libbert, E., Wichner, S., Schiewer, U., Rische, H. and Kaiser, W. (1966). The influence of epiphytic bacteria on auxin metabolism. Planta, <u>68</u>, 327-334.
- Little, C.H.A., Heald, J.K. and Browning, G. (1978). Identification and measurement of indoleacetic and abscisic acids in the cambial region of <u>Picea sitchensis</u> (Bon) Carr. by combined gas chromatography-mass spectrometry. Planta, <u>139</u>, 133-138.
- Loveys, B.R. (1977). The intracellular location of abscisic acid in stressed and non-stressed leaf tissue. Physiol. Plant., <u>40</u>, 6-10.
- Lyons, J.M. and Pratt, H.K. (1964). Effect of stage of maturity and ethylene treatment on respiration and ripening of tomato fruits. Proc. Am. Soc. Hort. Sci., <u>84</u>, 491-500.
- McCune, D.C. (1961). Multiple peroxidases in corn. Ann. N.Y. Acad. Sci., <u>94</u>, 723-730.
- McDougall, J. (1978). Indole-3-acetic acid in <u>Phaseolus</u> <u>vulgaris</u> L. Ph.D. Thesis, University of Glasgow.

- McDougall, J. and Hillman, J.R. (1978). Purification of IAA from shoot tissues of <u>Phaseolus</u> <u>vulgaris</u> and its analysis by GCMS. J. Exp. Bot., <u>29</u>, 375-386.
- Macháčková, I., Gančeva, K. and Zmrhal, Z. (1975). The role of peroxidase in the metabolism of indole-3-acetic acid and phenols in wheat. Phytochem., 14, 1251-1254.
- MacLachlan, G.A. and Waygood, E.R. (1956a). Catalysis of indoleacetic acid oxidation by manganic ions. Physiol. Plant., 9, 321-330.
- MacLachlan, G.A. and Waygood, E.R. (1956b). Kinetics of the enzymicallycatalysed oxidation of indoleacetic acid. Can. J. Biochem. and Physiol., <u>34</u>, 1233-1250.
- Magnus, V., Iskrić, S. and Kveder, S. (1971). Indole-3-methanol, a metabolite of indole-3-acetic acid in pea seedlings. Planta, <u>97</u>, 116-125.
- Magnus, V., Šimaga, S., Iskrić, S. and Kveder, S. (1982). Metabolism of tryptophan, indole-3-acetic acid and related compounds in parasitic plants from the genus <u>Orobanche</u>. Plant Physiol., <u>69</u>, 853-858.
- Mann, J.D. and J**q**worski, E.G. (1970). Minimising loss of indole-acetic acid during purification of plant extracts. Planta, <u>92</u>, 285-291.
- Marré, E., Colombo, R., Lado, P. and Rasi-Caldogno, F. (1974). Correlation between proton extrusion and stimulation of cell enlargement : effects of fusicoccin and of cytokinins on leaf fragments and isolated cotyledons. Plant Sci. Lett., <u>2</u>, 139-150.
- Martin, H.V., Elliott, M.C., Wangermann, E. and Pilet, P.E. (1978). Auxin gradient along the root of the maize seedling. Planta, 141, 179-181.
- Menschick, R. and Hild, V. (1976). Decarboxylation of IAA in coleoptiles in relation to phototropism. Abstracts 9th International Conference on Plant Growth Substances. pp 252-254.
- Metzler, D.E. (1977). Biochemistry : The Chemical Reactions of Living Cells. Academic Press, London.
- Meudt, W.J. (1971). Interaction of sulfite and manganous ion with peroxidase oxidation products of indole-3-acetic acid. Phytochem., 10, 2103-2109.
- Michalczuk, L. and Bandurski, R.S. (1980). <u>In vitro</u> biosynthesis of esters of indole-3-acetic acid. Plant Physiol., 65, Suppl. 157.
- Miginiac, E., Sossountzov, L. and Dugue, N. (1978). Modes of transport and metabolism of ¹⁴C-IAA and ³H-IAA in relation to morphogenesis of axillary buds in <u>Scrophularia arguta</u>. Physiol. Plant., <u>44</u>, 335-344.
- Miller, C.O. (1956). Similarity of some kinetin and red light effects. Plant Physiol., 31, 318-319.
- Miller, C.O. and Skoog, F. (1953). Chemical control of bud formation in tobacco stem segments. Am. J. Bot., 40, 768-773.
- Mittelheuser, C.J. and Van Steveninck, R.F.M. (1969). Stomatal closure and inhibition of transpiration induced by (R,S)-abscisic acid. Nature, <u>221</u>, 281-282.
- Mollan, R.C., Donnelly, D.M.X. and Harmey, M.A. (1972). Synthesis of indole-3-acetyl aspartic acid. Phytochem., 11, 1485-1488.
- Moloney, M.M., Elliott, M.C. and Cleland, R.E. (1981). Acid growth effects in maize roots : Evidence for a link between auxin economy and proton extrusion in the control of root growth. Planta, <u>152</u>, 285-291.
- Moloney, M.M. and Pilet, P.E. (1981). Auxin binding in roots : A comparison between maize roots and coleoptiles. Planta, 153, 447-452.
- Mulkey, T.J., Evans, M.L. and Kuzmanoff, K.M. (1981). Effects of aminoethoxyvinylglycine on the auxin responsiveness of intact roots of <u>Zea mays</u>. Plant Physiol., <u>67</u>, Suppl. 2.
- Nakajima, R. and Yamazaki, I. (1979). The mechanism of indole-3-acetic acid oxidation by horseradish peroxidase. J. Biol. Chem. 254, 872-878.

Nowacki, J. and Bandurski, R.S. (1980). <u>Myo-inositol esters of indole-3-acetic</u> acid as seed auxin precursors of <u>Zea mays</u> L. Plant Physiol., <u>65</u>, 422-427.
Ockerse, R., Gustin, M.K. and Lambert, N.J. (1975). Indoleacetic acid oxidase in terminal pea buds : multiple molecular forms and response to gibberellic acid. Plant Physiol., <u>56</u>, Suppl. 30.

- Ogawa, Y. (1961)*. Uber die wirkung von kinetin auf die blütenbildung von Pharbitis nil. Pl. Cell Physiol., 2, 343-359.
- O'Malley, B.W., Towle, H.C. and Schwartz, R.J. (1977). Regulation of gene expression in eukaryotes. Ann. Rev. Genetics, 11, 239-275.
- Osborne, D.J. (1978). Target Cells New Concepts for plant regulation in horticulture. Scientific Horticulture, 30, 1-13.
- Páal, A. (1919)*. Ueber phototropische reizleitung. Jahrb. Wiss. Bot., 58, 406-458.
- Parups, E.V. (1969). Involvement of free radicals in the oxidative degradation of indole-3-acetic acid. Can. J. Biochem., 47, 220-224.
- Pharis, R.P. and Morf, W. (1968). Physiology of gibberellin-induced flowering in conifers. In: Biochemistry and Physiology of Plant Growth Substances. Proceedings 6th International Conference on Plant Growth Substances. Eds. F. Wightman and G. Setterfield. pp 1341-1356. Runge Press, Ottawa.
- Phinney, B.O. (1960)*. In: Plant Growth Regulation. Ed. R.M. Klein. pp 489-495. Iowa State Univ. Press, Ames.
- Pilet, P.E. (1976). Fusicoccin and Auxin effects on root growth. Plant Sci. Lett., 7, 81-84.
- Pilet, P.E. and Elliott, M.C. (1981). Some aspects of the control of root growth and georeaction : the involvement of indole-3-acetic acid and abscisic acid. Plant Physiol., <u>67</u>, 1047-1050.
- Pilet, P.E., Elliott, M.C. and Moloney, M.M. (1979). Endogenous and exogenous auxin in the control of root growth. Planta, <u>146</u>, 405-408.
- Pilet, P.E. and Senn, A. (1980). Root growth gradients : a critical analysis.
 Z. Pfl. physiol., <u>99</u>, 121-130.
- Piskornik, Z. and Bandurski, R.S. (1972). Purification and partial characterisation of a glucan containing indole-3-acetic acid. Plant Physiol., <u>50</u>, 176-182.
- Powell, L.E. (1967). Determination of infra-red spectra on microgram quantities of some plant growth regulating compounds. Plant Physiol., 42, 1460-1462.

- Pratt, H.K. and Goesch1, J.D. (1969). Physiological roles of ethylene in plants. Ann. Rev. Plant. Physiol., <u>20</u>, 541-584.
- Rao, N.R., Naithani, S.C. and Singh, Y.D. (1982). Physiological and biochemical changes associated with cotton fibre development. II. Auxin oxidising system. Physiol. Plant., 55, 204-208.
- Ray, P.M. (1958). Destruction of auxin. Ann. Rev. Plant Physiol., <u>9</u>, 81-118.
- Ray, P.M., Dohrmann, U. and Hertel, R. (1977). Specificity of auxin binding sites on maize coleoptile membranes as possible receptor sites for auxin action. Plant Physiol., <u>60</u>, 585-591.
- Ray, P.M. and Ruesink, A.W. (1962). Kinetic experiments on the nature of the growth mechanism in oat coleoptile cells. Dev. Biol., 4, 377-397.
- Rayle, D.L. and Cleland, R. (1970). Enhancement of wall loosening and elongation by acid solutions. Plant Physiol., 46, 250-253.
- Reeve, D.R. and Crozier, A. (1977). Radioactivity monitor for high-performance liquid chromatography. J. Chrom., 137, 271-282.
- Reeve, D.R. and Crozier, A. (1980). Quantitative analyses of plant hormones. In: Encyclopedia of Plant Physiology, New Series, vol. 9 : Hormonal Regulation of Development 1. Ed. J. MacMillan. pp 203-280. Springer.
- Reinecke, D.M. and Bandurski, R.S. (1981). Metabolic conversion of ¹⁴C-indole-3-acetic acid to ¹⁴C-oxindole-3-acetic acid. Biochem. Biophys. Res. Comm., <u>103</u>, 429-433.
- Ricard, J. and Job, D. (1974). Reaction mechanisms of indole-3-acetate degradation by peroxidases. A stopped-flow and low-temperature spectroscopic study. Europ. J. Biochem., 44, 359-374.
- Ricard, J., Mazza, G. and Williams, R.J.P. (1972). Oxidation-reduction potentials and ionisation states of 2 turnip peroxidases. Europ. J. Biochem., <u>28</u>, 566-578.
- Richmond, A.E. and Lang, A. (1957). Effect of kinetin on protein content and survival of detached Xanthium leaves. Science, 125, 650-651.

- Riov, J. and Gottlieb, H.E. (1980). Metabolism of auxin in pine tissues : Indole-3-acetic acid conjugation. Physiol. Plant., <u>50</u>, 347-352.
- Rivier, L. and Pilet, P.E. (1974). Indoly1-3-acetic acid in cap and apex of maize roots : Identification and quantification by mass fragmentography. Planta, 120, 107-112.
- Roberts, L.W. (1974). Does 3-methyleneoxindole possess auxin activity? J. Exp. Bot., <u>25</u>, 761-763.
- Rodriguez, A.B. (1932)*. Smoke and ethylene in fruiting of pineapple. J. Dept. Agric. P.R., <u>26</u>, 5-18.
- Ryugo, K. and Breen, P.J. (1974). Indoleacetic acid metabolism in cuttings of plum (<u>Prunus cerasifera x P. munsoniana</u> cv. Marianna 2624). J. Am. Soc. Hort. Sci., 99, 247-251.
- Sabater, F., Cuello, J. and Sanchez Bravo J. (1976). Absence of biological activity in oxidation products of indoleacetic acid. Biologia Pl., <u>18</u>, 460-463.
- Sacher, J.A. (1959). Studies on auxin-membrane permeability relations in fruit and leaf tissues. Plant Physiol., 34, 365-372.
- Sachs, T. and Thimann, K.V. (1964). Release of lateral buds from apical dominance. Nature, 201, 939-940.
- Saleh, A.N. (1981). The effect of kinetin on the indoleacetic acid level and indoleacetic acid oxidase activity in roots of young plants. Physiol. Plant., 51, 399-401.
- Sandberg, G., Andersson, B. and Dunberg, A. (1981). Identification of 3indoleacetic acid in <u>Pinus sylvestris</u> L. by gas chromatography-mass spectrometry and quantitative analysis by ion-pair reversed-phase liquid chromatography with spectrofluorimetric detection. J. Chrom., <u>205</u>, 125-137.
- Sandberg, G., Odén, P.C. and Dunberg, A. (1982). Population variation and diurnal changes in content of indole-3-acetic acid of pine seedlings (<u>Pinus sylvestris L.</u>) grown in a controlled environment. Physiol. Plant., <u>54</u>, 375-380.

205

- Scheel, D. and Sanderman, H. (1981). Metabolism of 2,4-dichlorophenoxyacetic acid in cell suspension cultures of Soybean (<u>Glycine max</u> L.) and wheat (Triticum aestivum L.). 1. General results. Planta, 152, 248-252.
- Schlenk, H. and Gellerman, J.L. (1960). Esterification of fatty acids by

diazomethane on a small scale. Analytical Chem., <u>32</u>, 1412-1414.

- Schneider, E.A. and Wightman, F. (1974). Metabolism of auxin in higher plants. Ann. Rev. Pl. Physiol., 25, 487-513.
- Schurzmann, M. and Hild, V. (1980). Effect of indoleacetic acid, abscisic acid, root tips and coleoptile tips on growth and curvature of maize roots. Planta, <u>150</u>, 32-36.
- Scott, T.K. and Wilkins, M.B. (1968). Auxin transport in roots. II. Polar flux of IAA in Zea roots. Planta, 83, 323-334.
- Seeley, S.D. and Powell, L.E. (1974). Gas chromatography and detection of microquantities of gibberellins and indoleacetic acid as their fluorinated derivatives. Anal. Biochem., <u>58</u>, 39-46.
- Sembdner, G., Gross, D., Liebisch, H.W. and Schneider, G. (1980). Biosynthesis and metabolism of plant hormones. In: Encyclopedia of plant physiology, New Series Vol. 9 : Hormonal Regulation of Development. I. Ed. J. MacMillan. pp 281-444. Springer.
- Sequeira, L. and Mineo, L. (1966). Partial purification and kinetics of indoleacetic acid oxidase from tobacco roots. Plant Physiol., <u>41</u>, 1200-1208.
- Shaw, S., Gardner, G.K. and Wilkins, M.B. (1973). The lateral transport of IAA in intact coleoptiles of <u>Avena sativa</u> L. and <u>Zea mays</u> L. during geotropic stimulation. Planta, <u>115</u>, 97-111.
- Shaw, S. and Wilkins, M.B. (1974). Auxin transport in roots. X. Relative movement of radioactivity from IAA in the stele and cortex of <u>Zea</u> root segments. J. Exp. Bot., 25, 199-207.
- Shindy, W.W. and Smith, O.E. (1975). Identification of plant hormones from cotton ovules. Plant Physiol., 55, 550-554.

- Siegel, B.Z. and Galston, A.W. (1967). Indole-acetic acid oxidase activity of apoperoxidase. Science, 157, 1557-1559.
- Sirois, J.C. and Miller, R.W. (1972). The mechanism of the scopoletin-induced inhibition of the peroxidase catalysed degradation of indole-3-acetate. Plant Physiol., 49, 1012-1018.
- Skoog, F. and Miller, C.O. (1957)*. Chemical regulation of growth and organ function in plant tissues cultured <u>in vitro</u>. In: The Biological Action of Growth Substances, Symp. Soc. Exp. Biol., <u>11</u>, 118-131. Ed. H.K. Porter.
- Skytt Anderssen, A., Møller, I.B. and Hansen, J. (1972). 3-methyleneoxindole and plant growth regulation. Physiol. Plant., <u>27</u>, 105-108.
- Snow, R. (1935). Activation of cambial growth by pure hormones. New Phytol., <u>34</u>, 347-360.
- Stahl, E. (1965). Thin layer chromatography. Academic Press, New York.

Steen, M. and Hild, V. (1980). Geotropic curvature of decapitated <u>Avena</u> coleoptiles after application of tips of maize roots, tips of <u>Avena</u> coleoptiles, indoleacetic acid or abscisic acid. Planta, <u>150</u>, 37-40.

- Stoess1, A. and Venis, M.A. (1970). Determination of submicrogram levels of indole-3-acetic acid : a new highly specific method. Anal. Biochem., <u>34</u>, 344-351.
- Stonier, T. (1970). The role of auxin protectors in autonomous growth. In: Colloques Internationaux C.N.R.S. No. 193. Les Cultures de Tissus de Plantes. pp 423-435.
- Stonier, T. (1976). The role of auxin protectors in cell division. In: Abstracts 9th International Conference on Plant Growth Substances. Ed. P.E. Pilet. pp 376.
- Stonier, T., Stasinos, S. and Murthy Reddy, K.B.S. (1979). The masking of peroxidase-catalysed oxidation of IAA in <u>Vigna</u>. Phytochem., <u>18</u>, 25-28.
- Stutz, R.E. (1957). The indole-3-acetic acid oxidase of <u>Lupinus albus</u> L. Plant Physiol., <u>32</u>, 31-39.

- Summons, R.E., Entsch, B., Letham, D.S., Gollnow, B.I. and MacLeod, J.K. (1980). Regulators of Cell division in plant tissues. XXVIII. Metabolites of zeatin in sweet corn kernels : purifications and identifications using high-performance liquid chromatography and chemical-ionisation mass spectrometry. Planta, 147, 422-434.
- Sussman, M.R. and Goldsmith, M.H.M. (1981). Auxin uptake and action of N-1-Naphthylphthalamic acid in corn coleoptiles. Planta, 151, 15-25.
- Suzuki, Y. and Kawarada, A. (1978). Products of peroxidase catalysed oxidation of indoly1-3-acetic acid. Agric. Biol. Chem., <u>42</u>, 1315-1321.
- Suzuki, Y. and Takahashi, N. (1968)*. Effects of after-ripening and gibberellic acid on the thermoinduction of seed germination in <u>Solanum</u> melongena. P1. Cell. Physiol., 9, 653-660.
- Swartz, H.J. and Powell, L.E. (1979). Determination of indoleacetic acid from plant samples by an alkali flame ionisation detector. Physiol. Plant., 47, 25-28.
- Sweetser, P.B. and Swartzfager, D.G. (1978). Indole-3-acetic acid levels of plant tissue as determined by a new high performance liquid chromatographic method. Plant Physiol., 61, 254-258.
- Tang, Y.W. and Bonner, J. (1947). The enzymatic inactivation of indoleacetic acid. I. Some characteristics of the enzyme contained in pea seedlings. Archs. Biochem., <u>13</u>, 11-25.
- Tillberg, E. (1974). Occurrence of indol-3yl-aspartic acid in light and dark-grown bean seedlings (<u>Phaseolus vulgaris</u>). Physiol. Plant., <u>31</u>, 271-274.
- Thimann, K.V. (1935). On the plant growth hormone produced by <u>Rhizopus</u> suinus. J. Biol. Chem., 109, 279-291.
- Thimann, K.V. (1948). Plant growth hormones. In: The Hormones, Vol. 1.
 Ed. G. Pincus and K.V. Thimann. pp 5-74. Academic Press, New York.
 Thimann, K.V. (1969). The auxins. In: Physiology of Plant Growth and Development. Ed. M.B. Wilkins. pp 2-45. McGraw-Hill.

- Thimann, K.V. and Skoog, F. (1933). Studies on the growth hormone of plants. III. The inhibiting action of the growth substance on bud development. Proc. Nat. Acad. Sci. U.S.A., 19, 714-716.
- Trewavas, A. (1981). How do plant growth substances work? Plant Cell Environ., <u>4</u>, 203-228.
- Tuli, V. and Moyed, H.S. (1967). Inhibitory oxidation products of indole-3acetic acid : 3-hydroxymethyloxindole and 3-methylenoxindole as plant metabolites. Plant Physiol., 42, 425-430.
- Tuli, V. and Moyed, H.S. (1969). The role of 3-methyleneoxindole in auxin action. J. Biol. Chem., 244, 4916-4920.
- Ueda, M. and Bandurski, R.S. (1969). A quantitative estimation of alkalilabile indole-3-acetic acid compounds in dormant and germinating maize kernels. Plant Physiol., <u>44</u>, 1175-1181.
- Ueda, M., Ehmann, A. and Bandurski, R.S. (1970). Gas-liquid chromatographic analysis of indole-3-acetic acid <u>myo-inositol</u> esters in maize kernels. Plant Physiol., <u>46</u>, 715-719.
- Vacha, G.A. and Harvey, R.B. (1927). The use of ethylene, propylene and similar compounds in breaking the rest period of tubers, bulbs, cuttings and seeds. Plant Physiol., <u>2</u>, 187-194.
- Walton, D.C., Galston, E. and Harrison, M.A. (1977). The relationship between stomatal resistance and abscisic acid levels in leaves of water-stressed bean plants. Planta, <u>133</u>, 145-148.
- Weiler, E.W., Jourdan, P.S. and Conrad, W. (1981). Levels of indole-3-acetic acid in intact and decapitated coleoptiles as determined by a specific and highly sensitive solid-phase enzyme immunoassay. Planta, <u>153</u>, 561-571.
- Went, F.W. (1928)*. Die erklärung des phototropischen krummurgsverlaufs. Rec. Trav. bot. Néerland., <u>25</u>, 1-116.

Went, F.W. and Thimann, K.V. (1937). Phytohormones. MacMillan, New York.
Wetmore, R.H.and Rier, J.P. (1963). Experimental induction of vascular tissues
in callus of angiosperms. Am. J. Bot., 50, 418-430.

Weyers, J.D.B. (1978). The hormonal control of stomatal aperture in <u>Commelina communis</u> L. by abscisic acid. Ph.D. Thesis. University of Glasgow.

- Wilkins, M.B., Cane, A.R. and McCorquodale, I. (1972a). Auxin transport in roots. VII. Uptake and movement of radioactivity from IAA-¹⁴C by <u>Zea</u> roots. Planta, 105, 93-113.
- Wilkins, M.B., Cane, A.R. and McCorquodale, I. (1972b). Auxin transport in roots. IX. Movement, export, resorption and loss of radioactivity from IAA by Zea root segments. Planta, 106, 291-310.
- Wilkins, M.B. and Scott, T.K. (1968). Auxin transport in roots. III. Dependence of the polar flux of IAA in Zea roots upon metabolism. Planta, <u>83</u>, 335-346.
- Wilkins, M.B. and Whyte, P. (1968). Relationship between metabolism and the lateral transport of IAA in corn coleoptiles. Plant Physiol., <u>43</u>, 1435-1442.
- Wright, S.T.C. (1961). Growth and cellular differentiation in the wheat coleoptile (<u>Triticum vulgare L.</u>). I. Estimation of cell number, cell volume and certain nitrogenous constituents. J. Exp. Bot., <u>12</u>, 303-318.
- Wright, S.T.C. (1966). Growth and cellular differentiation in the wheat coleoptile (<u>Triticum vulgare</u> L.). II. Factors influencing the growth response to gibberellic acid, kinetin and indole-3-acetic acid. J. Exp. Bot., <u>17</u>, 165-176.
- Yamazaki, H. and Yamazaki, I. (1973). The reaction between indole-3-acetic acid and horseradish peroxidase. Arch. Biochem. Biophys., <u>154</u>, 147-159.
- Zeevart, J.A.D. (1976). Physiology of flower formation. Ann. Rev. Plant Physiol., 27, 321-348.
- Zenk, M.H. (1961). 1-(Indole-3-acetyl)-β-D-glucose, a new compound in the metabolism of indole-3-acetic acid in plants. Nature, <u>191</u>, 493-494.
- Zimmerman, P.W., Hitchcock, A.E. and Crocker, W. (1931). The effect of ethylene and illuminating gas on roses. Contrib. Boyce Thomson Inst., <u>3</u>, 459-481.
 Zmrhal, Z. and Macháčková, I. (1978). Isolation and characterisation of wheat peroxidase isoenzyme B1. Phytochem., 17, 1517-1520.
- [References not actually consulted, but cited through another authority, have been denoted with an asterisk.]

210

ADDENDUM I

Subsequent to the compilation of this thesis a sample of metabolite peak 10 was purified from a methanolic extract of root segments which had been incubated in aqueous $IAA-2-{}^{14}C$ ($10^{-2}mol m^{-3}$) for 2h and derivatised using diazomethane. The resulting methyl ester co-chromatographed on HPLC with the methyl ester of standard oxindole-3-acetic acid, thus giving further evidence that peak 10 represents this compound.

ADDENDUM II

Recently a report has been published describing HPLC analysis of IAA metabolites from <u>Cucumis sativus L.</u> (Purves and Hollenberg, 1982). Hypocotyl segments were incubated with ³H-IAA or ¹⁴C-IAA for varying lengths of time (from 0.5 to 48h) and the products extracted with methanol. Reverse-phase HPLC analysis of metabolites revealed an unspecified number of products, most of which were more polar than IAA. Two products were identified by co-chromatography as IAA-aspartate and IAAglutamate. The identity of the other products was not investigated.

Reference: Purves, W.K. and Hollenberg, S.M. (1982) Metabolism of exogenous indole-3-acetic acid to its amide conjugates in <u>Cucumis</u> <u>sativus L.</u> Plant Physiol., 70, 283-286

GLASCOW	*****22	te p	ľ
UND		,	Ĭ
LIBKAN.			Second Br

211

212 . -ERRATA

.