



University  
of Glasgow

<https://theses.gla.ac.uk/>

Theses Digitisation:

<https://www.gla.ac.uk/myglasgow/research/enlighten/theses/digitisation/>

This is a digitised version of the original print thesis.

Copyright and moral rights for this work are retained by the author

A copy can be downloaded for personal non-commercial research or study,  
without prior permission or charge

This work cannot be reproduced or quoted extensively from without first  
obtaining permission in writing from the author

The content must not be changed in any way or sold commercially in any  
format or medium without the formal permission of the author

When referring to this work, full bibliographic details including the author,  
title, awarding institution and date of the thesis must be given

Enlighten: Theses

<https://theses.gla.ac.uk/>  
[research-enlighten@glasgow.ac.uk](mailto:research-enlighten@glasgow.ac.uk)

THE CARBOL DIOXYDE CONTENT AND THE pH OF THE PLASMA OF CATS  
AND THE CHANGES ASSOCIATED WITH DISTURBANCES OF INSPIRATION

by

R. M. FERDOR, B.Sc., M.R.C.V.S.

A Study Submitted for the Degree of Doctor of Philosophy  
in the Faculty of Medicine,  
The University of Glasgow.

1959

ProQuest Number: 10647543

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



ProQuest 10647543

Published by ProQuest LLC (2017). Copyright of the Dissertation is held by the Author.

All rights reserved.

This work is protected against unauthorized copying under Title 17, United States Code  
Microform Edition © ProQuest LLC.

ProQuest LLC.  
789 East Eisenhower Parkway  
P.O. Box 1346  
Ann Arbor, MI 48106 – 1346

## INDEX

	Page
General Introduction	1
 <b>Part X</b> 	
<b>The carbon dioxide content and the pH of the plasma of normal cattle and their physiological variations, comparative determinations in the cow and the ventilation rate and the respiratory rate of normal cattle</b>	
Review of literature	5
Methods	
(1) Blood sampling	12
(2) Plasma carbon dioxide content	22
(3) Plasma pH	45
(4) Ventilation rate	64
(5) Respiratory rate	73
Results	
(1) A comparison of the effects of errors of sampling on the carbon dioxide content and the pH of plasma.	75
(2) The carbon dioxide content of the plasma of	
(a) Normal Ayrshire cattle	82
(b) Normal dogs	88
(3) The pH of the plasma of	
(a) Normal Ayrshire cattle	89
(b) Normal dogs	90

Results (contd.)	Page
(4) The physiological variations of the carbon dioxide content and the pH of the plasma of normal cattle.	92
(5) The ventilation and respiratory rates of Ayreshire cattle.	102
Discussion	108
Conclusions	109

## Part II

### The change in the carbon dioxide content and the pH of plasma caused by disturbance of respiration

Introduction	160
<b>Section I.</b> The effects of some general anaesthetic agents, Fluothane in particular, on the plasma pH and the plasma carbon dioxide content of cattle.	
Review of the literature	163
Method	
(1) Fluothane	174
(2) Thiopentone sodium	179
(3) Chloral hydrate	180
Results	
(1) Fluothane	
(a) Cattle	182
(b) Sheep	189
(c) Dogs	196
(d) Sheep	203

<b>Section 2.</b>	<b>Results (cont'd.)</b>	<b>Page</b>
(2)	Thiopentone sodium in cattle	260
(3)	Chloral hydrate in cattle	267
	<b>Discussion</b>	269
	<b>Conclusion</b>	272
<b>Section 3.</b>	<b>The disturbance of respiration caused by pneumonia.</b>	
	<b>Review of the Literature</b>	273
	<b>Methods</b>	274
	<b>Results</b>	275
	<b>Discussion</b>	283
	<b>Conclusion</b>	290
<b>Section 3.</b>	<b>The effects of alterations in the atmosphere presented to calves to breaths on the ventilation rate, the plasma pH and the plasma carbon dioxide content of cattle.</b>	
	<b>Review of the Literature</b>	292
	<b>Methods</b>	293
	<b>Results</b>	
	(1) Hypoxia	296
	(2) Hypercapnia	299
	(3) Anoxia	306
	<b>Discussion</b>	309
	<b>Conclusion</b>	326

Discussion (Part II)	237
Acknowledgments	231
References	232
Appendix I	236
Appendix II	237
Appendix III	239

THE CARBON DIOXIDE CONTENT AND THE pH OF THE PLASMA OF CATS  
AND THE CHANGES ASSOCIATED WITH DISTURBANCES OF RESPIRATION

## GENERAL INTRODUCTION

The most important factors regulating respiration are, in all probability, the oxygen tension and the carbon dioxide tension in the blood, the hydrogen ion concentration of the blood, the nervous centres within the central nervous system and the array of impulses impinging on these centres. These numerous factors form also the substratum out of which respiration develops. Changes are continually taking place in these factors so that the respiratory pattern at any one time is the resultant of the interplay of all these factors.

The relationship between the partial pressure of carbon dioxide ( $pCO_2$ ), the carbon dioxide concentration and the plasma pH is expressed by one form of the Henderson-Hasselbalch equation:


$$pH = pK + \log \frac{\text{Total } CO_2}{\alpha \times pCO_2}$$

where  $pK$  = constant = 6.2


$\alpha$  = constant = 0.0301 at  $37^\circ C$ .

The effects of carbon dioxide were simply demonstrated many years ago by Haldane who showed that inhalation of carbon dioxide caused a marked hyperventilation. The effect of inhalation of carbon dioxide is to raise the alveolar  $pCO_2$  and hence the arterial  $pCO_2$ . The increase in the arterial partial pressure of carbon dioxide is stimulant to respiration both centrally and to a lesser extent at the chemoreceptors of the carotid and aortic glans. As the  $pCO_2$  within the

blood is increased, a greater hydration of  $\text{CO}_2$  within the plasma will occur forming more carbic acid.



The increased quantity of carbic acid will react according to the equation



The hydrogen ions produced by the hydration of carbonic acid are buffered by the buffering systems in the plasma with a slight fall in pH and the bicarbonate ions remain in the plasma. At the same time due to the increase in arterial  $\text{pCO}_2$  a greater quantity of carbon dioxide diffuses into the erythrocytes where buffering mechanisms are available to deal with it. The concentration of bicarbonate increases within the erythrocytes and because of the increase in concentration within the erythrocytes diffusion of bicarbonate ions takes place out into the plasma while chloride ions from the plasma replace the bicarbonate ions in the erythrocytes. Thus as a result of the increase in the  $\text{pCO}_2$  there is an increase in the bicarbonate ions in plasma and thus an increase in the total  $\text{CO}_2$  content of the plasma while the plasma pH is lowered.

It can also be shown that by decreasing arterial  $\text{pCO}_2$  below the normal value the total  $\text{CO}_2$  is lowered and the plasma pH is raised and ventilation is diminished.

It has been demonstrated above that changes in the  $\text{pCO}_2$  lead to changes in plasma pH but it has been shown by Guy (1950) that alterations in plasma pH by themselves affect respiration. A lowering of plasma pH increases ventilation while raising of the plasma pH decreases ventilation.

-3-

The site of action is uncertain, the predominant effect is thought to be central (Hoff and MacKenzie, 1955).

Changes in the partial pressure of oxygen within the blood have also been shown to affect respiration. A lowering of the oxygen tension stimulates respiration by increasing chemoreceptor activity (Nulop, Tillquist and Hobson, 1939; Whelchel, Bartels and Hochstetler, 1956) but hyperoxia depresses centrally (Webb, Durbin and Coombs, 1943). Breathing pure oxygen has been shown to decrease chemoreceptor activity by Nulop et al. (1939) and to cause a decrease in ventilation of treated dogs by Webb et al. (1943). This decrease in ventilation disappeared after de-afferentation of the chemoreceptors. In man an initial decrease in ventilation following inhalation of 100% oxygen has been observed by Dejours, Ichiroisse, Leyraud and Tellier (1957), Dupuis and Coombs (1947), and Amuson and McNamee (1956). Dupuis and Coombs (1947) noted also a secondary increase in ventilation following the initial decrease. They ascribed this increase to an indirect effect of pure oxygen.

By analysis of the blood it is possible to determine the concentration in the blood of some or all of the major components affected by or affecting respiration, e.g., carbon dioxide, hydrogen ions and oxygen. This type of analysis can also be used to study alterations caused by disturbances of respiration due to disease, drugs or the administration of gas mixtures different in composition from the air normally breathed. There are many advantages in such a study inasmuch as provided a method of obtaining blood samples can be employed which

- 30 -

will give a true picture of the concentrations of carbon dioxide and oxygen and hydrogen ion actually affecting the central nervous system and the chemo-sceptors. If arterial blood samples can be obtained, these will give a true indication of the plasma concentrations of carbon dioxide and hydrogen ion concentration of the blood impinging on the respiratory centre and chemo-sceptors. In accurate analysis of alveolar air and expired air may not immediately reflect changes in the blood chemistry, changes liable to cause disturbances of respiration particularly when a greater than normal barometric exists in the lung areas which diffusion takes place, e.g. when alveolar hydrolization and epithelialization have occurred. In addition although alveolar gas analysis and analysis of expired air are possible, the gas samples obtained may not be true samples due to distribution of respiration caused by fear, breathing and excitement when collecting blood samples and also in the stadium by the addition of crushed musk (gas).

In the present study suitable methods of obtaining blood samples have been evolved. Using these methods of sampling, the carbon dioxide concentration of the plasma and the pH of the plasma of healthy adults cattle and calves have been determined as have the variations in respiration taking place in healthy cattle when at rest during the day. Studies have been made also of the alterations in these two concentrations, as a result of lung damage due to pneumonia, as a result of the central depression of respiration caused by general anaesthetics and as a result of the inhalation of mixtures of nitrogen, oxygen and carbon dioxide different in amount to those present in the normal atmosphere breathed.

For comparative purposes a study has been made of the concentrations of carbon dioxide and the pH of the plasma of normal dogs and also of the effects of anaesthesia on the plasma pH and plasma carbon dioxide content of species other than cattle, e.g. horses, dogs and sheep.

In order to describe the disturbances of respiration observed with anaesthesia, anaesthetics and inhalation of foreign gas mixtures measurements have been made of the ventilation and respiratory rates of cows and calves.

PAGE X

THE CARBON DIOXIDE CONTENT AND THE pH OF THE PLASMA OF NORMAL CATS  
AND THEIR PHYSIOLOGICAL VARIATIONS. COMPARATIVE DETERMINATIONS IN THE DOG  
AND THE VENTILATION RATE AND THE RESPIRATORY RATE OF NORMAL CATS.

## REVIEW OF THE LITERATURE

The carbon dioxide content of the plasma or serum obtained from venous blood of cattle has been determined by a number of workers. Some of these workers have also determined the blood or plasma pH.

Anderson, Gayley and Pratt (1930) in an analysis of the blood of 53 cattle gave a mean value of 58.9 volumes per cent (26 m.moles/litre) as the plasma bicarbonate concentration. Of these 53 animals only five were over one year of age and for these five a mean value of 43.4 volumes per cent (39.2 m.moles/litre) was found, which value is low in comparison to other workers' findings. Anderson *et al.* stated that the samples were collected without stasis from the jugular vein in such a manner that the blood was protected from exposure to the atmosphere by a layer of liquid paraffin. They gave the method of determination of the carbon dioxide content but did not explain how the values of carbon dioxide content were converted to plasma bicarbonate since plasma pH was not determined at the same time.

Milne (1952) in a study of the effects of thermal stress on the venous plasma carbon dioxide content and the plasma pH of two Ayralike cattle found a total carbon dioxide concentration of 69 volumes per cent (30.6 m.moles/litre) and 61 volumes per cent (27 m.moles/litre) during

the control period. He also found a plasma pH of the two calves to be 7.51 and 7.52 during the control period. For each determination a sample was taken under oil into heparinized centrifuge tubes and centrifuged immediately and the determinations then carried out. Although only two animals were sampled the values for total carbon dioxide concentration found were within the range of other authors. The pH determinations were less valid since readings were taken at room temperature and corrected to the temperature of the animal using a temperature coefficient found by Weston (1953) whose paper on temperature coefficients was concerned with the determination of temperature coefficients for the pH of human plasma and urine.

Burge (1978) in another study of the effects of thermal stress on the blood composition of dairy cattle found a mean value of 55.3 volumes per cent (25 m.moles/litre) for the carbon dioxide capacity of the plasma of six bovine subjects when in an ambient temperature of 55-60° F. prior to exposure to thermal stress. The method of taking the blood samples was not mentioned but since carbon dioxide capacity and not content was measured sampling methods were less critical.

Greige (1977) hypothesised that the cause of hypocalcaemia in cattle was an alkaloosis. He made 16 determinations on the plasma of five cows and found a mean value for the carbon dioxide concentration of plasma of these cows to be 25 m.moles/litre. He did not state how the blood

samples were banded or from what blood vessels they were obtained. On another occasion Craig, Johnson, Blackburn and Coffin (1949) made a more complete study of three cows over a period of one month. Maximum and minimum values were given for the carbon dioxide content of each cow and these were between 23 and 22 m.moles/litre. Similarly maximum and minimum values of 7.45 and 7.29 were given for plasma pH. It was not stated from which vessel the blood samples were obtained but it was stated that the samples were withdrawn without exposure to the atmosphere.

Dale and Brody (1953) in studying the effects of thermal stress on the cold base balance of four cows gave mean control values of 54.6 volumes per cent (25 m.moles/litre) for the carbon dioxide concentration and 7.32 for the blood pH. They took the blood samples from the jugular vein with minimal stasis and minimal exposure to air into oiled heparinised syringes.

Dale, Gobindram and Brody (1954) carried out a similar study on the effects of starvation on cold base balance and gave a mean value for the carbon dioxide capacity of the plasma of four cows to be 25.4 meq./litre. This plasma was obtained from jugular venous blood. Blood pH was determined on the same four cows and a range was given of 7.29 - 7.40. They took blood samples from the jugular vein into oiled heparinised syringes with minimal stasis and minimal exposure to air.

Krapf (1943) found the carbon dioxide capacity of the venous plasma

of 119 cattle to be between 55 volumes per cent (23 m.moles/litre) and 75 volumes per cent (33 m.moles/litre) while the plasma pH was from 7.35 - 7.50. No breed, sex or age differences were noticed. The samples were withdrawn with anaerobic precautions from the jugular vein.

McSherry and Grinley (1954) measured the carbon dioxide content and the pH of bovine serum. The samples were collected with the minimal trauma of the blood from the jugular vein into test tubes in such a manner that collection was under a layer of liquid paraffin. The samples were then allowed to clot at room temperature for two hours but despite this pH values were found similar to the values found by other authors. This is not what one would expect to find according to Rosenblat (1948) who showed that the estolytic taking place in blood samples which were allowed to stand at room temperature for one hour would cause a lowering of pH by as much as 0.1 pH units. McSherry and Grinley found a mean pH value for 86 adult cattle of  $7.42 \pm 0.06$  and  $7.44 \pm 0.14$  for the pH value of 20 calves. A mean bicarbonate concentration for the adults of  $39.5 \pm 1.96$  meq./litre and for the calves of  $31.03 \pm 1.9$  was found. The bicarbonate concentration was calculated from the total carbon dioxide content and the serum pH.

Simpson and Haydon (1939) measured the carbon dioxide content of the plasma of 26 normal cows. A mean value was determined of 63.3

volumes per cent (23 n.moles/litre) with a maximum value of 79 volumes per cent (35 n.moles/litre) and a minimum value of 54 volumes per cent (25 n.moles/litre). No description was given of the methods of collection or handling of the blood sample.

The most extensive study so far of the respiratory performance of cattle was made by Brody (1949). He conducted numerous experiments and constructed many types of spirometer for respiratory measurements. From the hundreds of results obtained he plotted a graph of the logarithm of the ventilation rate in Litres per minute against the logarithm of the body weight in Kilograms. From this graph he obtained a regression for all the determinations. From this regression he predicted that a calf of 50 kg. body weight would have a ventilation rate of 25 Litres per minute while an adult cow of 500 kg. body weight would have a ventilation rate of 87 Litres per minute.

Doyle, Patterson, Warren, Dotweiller and Reynolds (1958) have published some observations on respiratory function of adult cattle but weights of animals were not given. A mean value of 82 litres per minute was found for the ventilation rate of six cows while the respiratory rate averaged 36 respirations per minute.

Many authors in studying the disturbances of respiration produced experimentally have included control measurements made prior to the experimental alteration. McDowell, Lee, Pohman and Anderson (1953)

when studying the effects of thermal stress on respiration gave a mean value of 83 litres per minute for the ventilation rate, 43 respirations per minute for the respiratory rate of 10 adult cows whose weights averaged 450 kg., the ambient temperature varying from 57-84° F.

Hindley (1955) carrying out similar studies on calves in an environmental temperature of 20° C. gave an initial ventilation rate of 30 litres per minute and a respiratory rate of 34 respirations per minute but did not state the weights of the calves. In a full paper Hindley (1956) stated that he studied 30 four month old calves which had a mean control ventilation rate of 44 litres per minute and a respiratory rate of 70 respirations per minute at an environmental temperature of 20° C. This respiratory rate is much above that found by other workers. Such a value for respiratory rate indicates that his calves were excited, under a heat stress or affected with pneumonia. Again the body weights of the animals were not given.

Hansson and Johannsson (1958) studying the effects of anaesthesia gave ventilation rates of 38, 56, 49, 78 and 51 litres per minute and respiratory rates of 14, 13, 17, 23 and 16 respirations per minute for the control values of five Swedish Red cattle prior to anaesthesia. These values were somewhat lower than values found by other authors which may indicate a breed variation of Swedish Red cattle.

Summarising the values of the pH and the carbon dioxide content of

the plasma of cattle obtained by other workers it appeared that the pH of the venous plasma of cattle was between 7.30 and 7.50 pH units and that the venous plasma carbon dioxide concentration was between 20 and 33 m.moles/litre. From the values obtained for the respiratory rate of normal cattle it appeared that the normal respiratory rate of cattle was between 20 and 45 respirations per minute, probably depending on the prevailing ambient temperature. It was observed that the ventilation rate varied with the size of the animal and that a 50 kg. bovine subject had a ventilation rate of about 27 litres per minute while a 500 kg. bovine subject had a ventilation rate of about 90 litres per minute.

## METHODS

- (1) Blood sampling
- (2) Plasma carbon dioxide content
- (3) Plasma pH
- (4) Ventilation rate
- (5) Respiratory rate

### (1) Blood sampling

Venous blood samples were obtained from both the jugular and mammary veins. The jugular veins of cattle were not obvious on inspection or palpation unless some cardiac defect led to a rise in venous pressure. It was necessary to raise the vein by digital compression below the point of puncture. With the vein raised it was entered by means of a rapid stick puncture using a one inch 16 British Wire Gauge needle. Once the vein was penetrated the digital compression was removed and the blood allowed to flow out of the needle for at least 30 seconds prior to taking a sample. The other vein from which venous samples were removed was the mammary or subcutaneous

abdominal vein. This vessel was not suitable in young animals but provided a readily accessible source of blood in adult cows. Compression of this vein was not necessary. The vein was entered at the so-called "milk wall" where the vein entered the abdomen one handbreadth lateral to the xiphoid cartilage.

Blood samples approximating mixed venous samples were obtained by catheterisation of the jugular vein. An area over the upper third of the jugular vein was clipped and washed with a solution of trimethyl allyl ammonium bromide<sup>2</sup>. Five millilitres of a stetilo 5% solution of procaine hydrochloride was infiltrated beneath the skin in this area. After raising the jugular vein by means of a rope around the neck the vein was entered with a two-inch number 9 British Wire Gauge needle and the rope was removed. A 40 cm. length of 2 mm. diameter sterile nylon tube was inserted into the jugular vein through the needle and the needle was removed. A small piece of rubber tubing was attached to the nylon catheter and the catheter was filled with heparinised saline to keep it patent. The rubber tube was then closed with a Kehr's clip. The catheter was palpable within the jugular vein to the root of the neck but it was not possible to ascertain the exact position of the tip of the catheter. Figure 1 illustrates the jugular venous catheter in situ.




Fig. 1

Jugular catheter in situ.

To obtain mixed venous samples from the right ventricle of the human subject Cournand (1941) determined the exact position of the catheter tip radiographically. While it may have been possible to determine the position of the catheter in this manner in adult cattle the high dosage of X-rays necessary to penetrate the bovine chest was considered to constitute too great a hazard to handlers of the animals to adopt the practice.

Arterial blood samples have been obtained from cattle in many ways. Von Ecorous (1923) described how in dogs it was possible, by surgical intervention, to bring the carotid arteries to a superficial position enclosed in a roll of skin. This procedure has been carried out in cattle (Jennings, 1956) but suffered the disadvantage that the suture broke down after three months. Moreover this could be carried out on experimental animals only.

Arterial blood has also been obtained from cattle by direct puncture of the carotid artery in the middle third of the neck (Sollers and Remingay, 1951). They infiltrated the subcutaneous tissue with procaine hydrochloride and inserted the needle above the jugular vein and into the artery. However, when this method was repeated it was found that considerable restraint was necessary, and that slight movement of the neck withdrew the needle from the artery.

Swedish workers Hildebrand and Obel (1958) have described how they affected entrance into the abdominal aorta of cattle. They used a special needle 30 cm. in length, the last 8 cm. of which was curved and of 1.8 mm. external diameter for actual entrance into the artery, while the remainder of the shaft was of 10 mm. diameter. With the tip of the needle protected by a rubber finger shield the needle was inserted into the rectus, through the rectal wall into the abdominal aorta. This method was not used in the present study.

In the present study it was found possible to obtain arterial blood samples from cattle of all ages by puncture of the brachial artery at the root of the neck (Fisher, 1956). The animal to be sampled was tethered by means of a halter so that its head was held slightly upward and to the right. The hand of the operator was placed under the point of the shoulder as shown in figure 2 and the brachial artery was palpated at the point where it crossed the first rib.



Fig. 2  
Palpation of the brachial artery

In thin cows the brachial artery was easily found and rolled between the fingers and the first rib but in fat, thick-necked animals it was not possible even to feel the pulse at this point.

The proposed puncture site was infiltrated with a sterile 5% solution of procaine hydrochloride. A four inch 15 British Wire Gauge, long pointed needle was directed into the cow at a point just medial to the second finger as shown in figure 2, parallel to the long axis of the cow and at about  $15^{\circ}$  to the horizontal. The needle was inserted to a depth of one to three inches in order to effect entrance to the artery. Figure 3 illustrates this.



Fig. 3

Needle inserted into brachial artery

No fatalities have occurred in healthy animals using this method. Post mortem examinations of animals slaughtered for other reasons and on which arterial punctures had been carried out revealed only slight haematomata at the site of puncture in a few cases. The wall of the artery showed small puncture wounds but no significant damage had occurred.

In obtaining blood samples from cattle the following measures were adopted to ensure that the plasma samples provided an accurate estimate of the plasma carbon dioxide content and that contamination of these samples did not occur.

All blood samples were collected into clean Pyrex centrifuge tubes (fig. 4).



Fig. 4

Collection of blood into Pyrex centrifuge tube

- 69 -

These centrifuge tubes were thoroughly washed with soap and water to remove any trace of former samples. They were then rinsed twice in distilled water, soaked in concentrated hydrochloric acid and then rinsed again with two separate quantities of distilled water before drying in a hot air oven. Duplicate blood samples were taken into two separate tubes and duplicate estimations were carried out on each.

Blood on which an analysis was to be made was allowed to fill the soft plastic connecting tube and was then collected <sup>into heparinised centrifuge tubes</sup> under a layer of liquid paraffin so that no exposure to the atmosphere took place. In order to prevent a loss of carbon dioxide from the blood to the liquid paraffin, the quantity of liquid paraffin allowed to remain in contact with the blood was kept to a minimum. During collection blood was allowed to overfill the centrifuge tube so that most of the liquid paraffin was displaced from above the blood and a small volume only was left over the blood. The tube was then sealed with a soft rubber bung so that no air was included between the blood and the bung and more paraffin was extruded out of the centrifuge tube (fig. 5). The centrifuge tube was kept sealed until analysis of the plasma was undertaken.

Von Slyko showed many years ago that if blood was allowed to stand at room temperature, changes of autolysis took place and the pH fell. This was confirmed by Rosenthal (1946) who also showed that 30 minutes




Fig. 5

Extrusion of liquid paraffin from above the blood

could elapse before changes took place and that these changes could be prevented by refrigeration. Blood samples in the present study were collected and the analysis for the carbon dioxide content of the plasma and the plasma pH was started within 30 minutes of collection. On occasion this was not possible and the samples were refrigerated for as long as necessary and before analysis were brought up to room temperature.

## (2) Plasma carbon dioxide content

Two methods were used to determine the carbon dioxide content of plasma. The Monometric Method of Van Slyke (1924) was used for most of the estimations on normal animals. In the second half of the study the method of Conway (1950) was used when it was necessary to carry out large numbers of estimations in a short space of time.

In the Monometric Method of Van Slyke (1924) the gases present in the plasma are released by means of lactic acid and exposure to a vacuum within the special apparatus. The pressure ( $p_1$ ) exerted by the gases when held at constant volume and constant temperature is measured. The carbon dioxide is absorbed by sodium hydroxide and the new pressure ( $p_2$ ) is measured. From these two values,  $p_1$  and  $p_2$ , it is possible, knowing the temperature, to calculate the carbon dioxide content of the plasma by reference to a special table produced by Van Slyke (Fig. 6). The method is described below.

### Reagents

- (a) 2.0 normal lactic acid.
- (b) 1.0 normal sodium hydroxide solution.
- (c) Distilled water.

These three reagents were made air free by extraction in the manometric apparatus. After they were made air free they were collected into special containers without contact with air.

- (d) Caprylic alcohol.

**FACTORS FOR CALCULATION OF CO<sub>2</sub> CONTENT OF BLOOD (FROM VAN SLYKE AND SENDROY (62))**

TEMPERA TURE °C.	FACTORS BY WHICH MILLIMETERS P <sub>CO<sub>2</sub></sub> ARE MULTIPLIED TO GIVE:									
	Millimoles CO <sub>2</sub> per liter of blood					Vol. per cent CO <sub>2</sub> in blood				
	Sample = 0.2 cc.		Sample = 1.0 cc.			Sample = 0.2 cc.		Sample = 1.0 cc.		
	S = 2.0 cc. a = 1.037	S = 0.5 cc. a = 1.037	S = 3.5 cc. a = 1.017	S = 1.0 cc. a = 1.017	S = 2.0 cc. a = 1.037	S = 0.5 cc. a = 1.037	S = 3.5 cc. a = 1.017	S = 1.0 cc. a = 1.017	S = 2.0 cc. a = 1.037	S = 0.5 cc. a = 1.037
15	0.1514	0.0313	0.1229	0.0341	0.1335	0.3370	0.0697	0.2735	0.0758	0.2974
16	.07	11	22	38	25	54	93	19	52	50
17	0.1499	10	15	35	15	38	89	04	46	28
18	.92	08	08	33	06	22	86	0.2690	41	06
19	86	06	02	31	0.1297	07	82	75	36	0.2886
20	79	05	0.1196	28	88	0.3292	78	62	31	66
21	72	03	90	26	79	78	75	48	26	48
22	66	02	83	24	70	63	71	34	21	28
23	59	00	77	22	62	48	68	20	16	08
24	53	0.0299	71	19	53	34	65	07	11	0.2790
25	46	97	65	17	45	20	61	0.2594	07	72
26	40	96	60	15	37	06	58	81	02	53
27	34	94	54	13	29	0.3193	55	69	0.0698	36
28	28	93	49	11	22	79	52	57	93	20
29	22	91	43	10	15	66	49	45	89	04
30	16	90	38	08	08	53	46	33	85	0.2688
31	11	89	33	06	01	40	43	22	82	74
32	05	88	28	05	0.1195	28	40	11	78	59
33	00	86	23	03	88	15	37	00	74	44
34	0.1394	85	18	01	82	03	34	0.2489	71	30

To obtain factors for a sample other than 1 cc., divide the above factors for 1 cc. by the cubic centimeters of sample analyzed: e.g., for a 2-cc. sample, S, A, and a being the same, the factors are one-half of those for a 1-cc. sample.

### Method

After cleaning the extraction chamber a drop of caprylic alcohol was drawn into the capillary above the cock at the top. A small amount of this caprylic alcohol was drawn into the extraction chamber. The stopcock was closed and any caprylic alcohol left in the cup was removed by suction. 2.5 ml. of carbon dioxide free water were placed in the cup. The sample of plasma in the special pipette with a rubber tip was run into the chamber under the water already in the cup. The stopcock was closed and the pipette removed. Any small amount of plasma left in the cup was washed in with some of the water. In a similar manner 0.2 ml. of 2 Normal lactate acid was run into the chamber and washed in with the remainder of the water until the mark on the lower end of the cup was reached. A small amount of mercury was placed in the cup and this was run into the stopcock to form the mercury seal.

The apparatus was evacuated by lowering the levelling bulb until the mercury in the chamber had fallen to the 50 ml. mark. The cock leading to the extraction chamber was closed and the reaction mixture was shaken for three minutes. When extraction was completed the mercury was re-admitted slowly until the gas volume in the chamber had been reduced to 2 ml. The reading on the manometer was then taken; this gives the value for  $p_2$  in millimetres of mercury. The mercury in the levelling bulb was lowered once more to increase the gas volume to

about 10 ml. 1.2 ml. of 1 Normal air-free sodium hydroxide were placed in the top cup and 1 ml. of this drawn slowly into the reaction chamber. About 30 seconds were taken for running in the sodium hydroxide and during this time absorption of the carbon dioxide was completed. As previously the stopcock was sealed with a mercury seal. The volume of gas was once more reduced to 2 ml. The manometer reading now gave a value for  $p_2$  in millimetres of mercury. The pressure of carbon dioxide ( $p_{CO_2}$ ) was obtained from the equation  $p_{CO_2} = p_1 - p_2 - c$ , 'c' being the correction previously determined. To obtain the carbon dioxide content of the plasma, the  $p_{CO_2}$  value was multiplied by the proper factor from the table given in Figure 6.

The 'c' correction was obtained by performing blank analysis, in the manner directed excluding the plasma. The  $p_1 - p_2$  value found in the blank constituted the 'c' correction.

The 'c' correction may be the sum of two components. One of these is a slight excess of gas which may be yielded by the reagents themselves. The other is the fall in manometer reading caused even when no gas is removed between the  $p_1$  and the  $p_2$  readings, by the introduction of a given volume of absorbent solution such as the 1.0 ml. of 1 Normal sodium hydroxide used to absorb carbon dioxide in blood gas determinations. The introduction of solution between the two manometer readings,  $p_1$  and  $p_2$ , lowers the  $p_2$  value merely by increasing the volume of fluid between

the mercury surface and the water meniscus at the moment of reading. thereby the level of the mercury surface in the chamber and hence also in the manometer is lowered. The extent of this effect can be determined by means of blank analyses in which, with the same amount of reagents but no gas in the chamber, the usual amount of absorbent solution is admitted. The manometer is read with water meniscus in the chamber at the same mark before and after the admission of the absorbent solution. The precise shape of the apparatus causes the area of the meniscus in the chamber to vary according to the volume of solution present and according to the location of the water meniscus, causes the mercury meniscus to be located at points of slightly different cross section in the conical upper portion of the chamber. Table I below gives the results of determination of the 'c' correction factor for the particular Van Slyke manometric apparatus used.

Table I

Determination of the 'c' Correction Factor

$P_1$	$P_2$	$P_1 - P_2 = 'c'$
90.8	89.4	1.4
90.0	88.8	1.2
91.5	90.1	1.4
91.6	90.2	1.4
92.4	91.0	1.4
93.1	91.9	1.2
93.9	92.6	1.3
93.0	91.7	1.3
94.8	93.5	1.3
93.5	92.0	1.5

Note: 1 c = 1.3 millimeters of mercury.

Determinations were carried out of the carbon dioxide content of a standard solution of sodium bicarbonate containing a calculated 20 n.moles of carbon dioxide per litre. The results are presented below.

Table 3

Recovery of Carbon Dioxide from Bicarbonate Standard

Temp. ° C.	P <sub>1</sub> mmHg	P <sub>2</sub> mmHg	P <sub>1</sub> -P <sub>2</sub> × 10 <sup>4</sup> mmHg	Factor	n.moles CO <sub>2</sub> per litre
19	263.2	101.8	161.4	0.1202	19.16
19	267.2	103.0	164.2	0.1202	19.73
20	268.9	96.5	170.7	0.1203	20.70
18	267.0	95.0	172.7	0.1203	20.70
18	266.0	95.5	171.2	0.1203	20.69
18	269.5	96.5	173.7	0.1203	20.69
19	264.5	104.0	160.9	0.1202	19.13

$$\bar{x}_{CO_2} = 20.2 \pm 0.6 \text{ n.moles/litre}$$

It was concluded that the method as being used was suitable for carrying out a number of determinations.

In the method of Conway, within a gas-tight Conway unit the carbon dioxide is released from the plasma by means of a strong mineral acid. The carbon dioxide diffuses into the centre well of the Conway unit

where it is fixed by means of a solution of barium hydroxide so that a precipitate of barium carbonate is formed. The excess barium hydroxide is determined by titration with hydrochloric acid. Since the strength and volume of barium hydroxide originally present in the unit are known, the amount of carbon dioxide released from a known volume of plasma can be determined.

Certain modifications were made to the method as published by Conway (1950). Micro-Conway units were used at first but it was not found possible to obtain either accuracy or repeatability with the volumes used in these small units. It was decided therefore to use large size Conway units since there was no problem of volumes of plasma available from cattle.

Vaseline was found unsatisfactory as a seal for the Conway units used in the determination of carbon dioxide since diffusion of carbon dioxide appeared to take place through the vaseline. The gel triglycerin-glycerin-acid mixture mentioned by Conway was found to be satisfactory in this respect but if used according to the directions of Conway was far too viscous to spread and form an efficient seal. Water was therefore added to the mixture.

Each batch of barium hydroxide solution used for determinations was adjusted in strength so that 1 mL (the volume placed in a Conway unit) required 4 to 4.5 mL of N/10 hydrochloric acid. In this manner

the maximum accuracy was obtained. It was found that 1 ml. of barium hydroxide solution gave the optimal area for absorption of carbon dioxide. It was also found that the total volume which could be contained in the centre well of the Conway unit without overflow was 5.5 ml. so that 4 to 4.5 ml. of acid could be added.

The barium hydroxide with the indicator added was kept in a carbon dioxide free atmosphere. In this manner large volumes (400 ml.) of working solution could be made up, standardised and stored. The time for diffusion of the carbon dioxide from plasma was found to be in excess of the three to four hours given by Conway. The best results were obtained if the units were left overnight. To overcome the buffering capacity of plasma, it was found necessary to use a strong mineral acid (0.5 ml. of N/2 sulphuric acid) to release the carbon dioxide.

Titration of the Conway units was not accurate when performed open to the atmosphere, but titration in a carbon dioxide free atmosphere gave good results. Atmospheric carbon dioxide did not appear to interfere with the results but a sudden increase resulting from the operator exhaling over the units led to errors. A shield was designed to eradicate this error.

#### Apparatus

- (a) A number of no. 1 size Conway units.


(b) Syringe pipettes of the "walking stick" pattern. Two sizes were used, 0.5 ml. and 1.0 ml.

(c) One 5.0 ml. Green Line<sup>®</sup> burette with automatic refill.

(d) A magnetic stirrer.

(e) A transparent shield behind which the Conway units were titrated.

The burette, magnetic stirrer and transparent shield were combined to form the titrating unit illustrated in figure 7.



## Reagents

- (a) A saturated solution of barium hydroxide which was kept in a sealed bottle.
- (b) From this stock solution, a solution of barium hydroxide was made up which was used in the determination. This solution consisted of a 1 in 4 dilution of the stock solution of barium hydroxide in distilled water together with about 5 ml. of the mixed indicator solution described below. The strength of the solution was adjusted so that 1 ml. required 4 to 4.5 ml. of N/30 hydrochloric acid for neutralisation. Once made up this solution was stored in a carbon dioxide free atmosphere. The carbon dioxide free storage system illustrated in figure 8 consisted of a storage bottle connected to a U tube full of soda lime so that air entering the bottle passed through the soda lime and became carbon dioxide free. The storage bottle fed by gravity the barium hydroxide solution to a two-way stopcock. This stopcock was normally kept closed but when estimations were carried out a quantity of the carbon dioxide free barium hydroxide could be fed into a narrow tube from which sized volumes could be removed to fill the centre wells of the Conway units.
- (c) A solution of N/2 sulphuric acid.
- (d) Indicator solution which consisted of a mixture of phenolphthalein

and thymolphthalein.

(e) A solution of hydrochloric acid made up accurately to be of N/80 strength.

(f) A seal for the lids of the Conway units which consisted of a mixture of tragacanth gum, glycerin, dilute sulphuric acid and water. The inclusion of acid in this mixture prevented a diffusion of carbon dioxide from the atmosphere into the sealed Conway units.



Fig. 8




Carbon dioxide free storage system.

## Method

The glass rings and lids of the Conway units were cleaned with the sealing compound. The 0.5 ml. N/2 sulphuric acid was then delivered into the outer wall of each Conway unit diametrically opposite to the point where the plasma was added later. The lids were then placed on the Conway units leaving about one-half of the centre wall uncovered. By means of a sprung pipette 1.0 ml. of the barium hydroxide Indicator solution was delivered to the centre well of each unit and the units were immediately sealed. Some of these units set up in this manner were left like this to act as blanks to allow for the slow deterioration of barium hydroxide which occurred in the carbon dioxide free storage system.

The plasma in 0.5 ml. volume was removed from the sealed centrifuge tubes so that minimal exposure to the atmosphere took place. The lid of a Conway unit was moved over slightly and the plasma was delivered into the outer well of a Conway unit after which the unit was immediately sealed. The Conway unit was then rotated slightly to bring the sulphuric acid into contact with the plasma. The units were then left on the bench until next morning when the barium hydroxide left was titrated against the N/50 hydrochloric acid in the titration stand illustrated in Figure 7. The colour changes of the indicator were blue to pink to colourless. Blank units were titrated first to determine

the amount of acid (B ml.) required to neutralise the barium hydroxide present in each unit before the addition of carbon dioxide. The units containing the plasma were then titrated to determine the amount of acid (U ml.) required to neutralise the barium hydroxide left after the absorption of the carbon dioxide released from 0.5 ml. of plasma. The calculations given below demonstrate how the concentration of carbon dioxide present in the plasma samples was obtained from the two titration values B and U.



51

Blank reading (B) gives amount  $\text{Ba(OH)}_2$  present in solution

Unknown reading (U) gives amount  $\text{Ba(OH)}_2$  left over after

absorbing the  $\text{CO}_2$ .

Therefore  $(B - U)$  gives the amount of  $\text{Ba(OH)}_2$  required by  $\text{CO}_2$ .

But from equation (a) 1 m.mole  $\text{Ba(OH)}_2$  requires 1 m.mole  $\text{CO}_2$

and from equation (b) 1 m.mole  $\text{Ba(OH)}_2$  requires 2 m.moles  $\text{HCl}$ .

Therefore 1 m.mole  $\text{CO}_2$  is 2 m.moles  $\text{HCl}$ .

The strength of acid used was N/60 and the volume of plasma used 0.5 ml.

Therefore the number of m.moles  $\text{CO}_2$  in 0.5 ml. plasma is


$$\frac{(B-U)}{2} \times \frac{1,000}{60} \times \frac{1}{1,000}$$

Therefore the number of moles  $\text{CO}_2$  per litre of plasma is

$$\frac{(1-0)}{2} = \frac{1,000}{30} = \frac{2,000}{1,000} =$$

(B-5) 100

The method of Conway was used excepted several innovations and therefore a more complete analysis was made than was made for the Van Slyte method. A series of standard solutions of sodium carbonate was made up from Analytical reagent grade anhydrous sodium carbonate. These solutions were fairly stable at room temperature but tended to pick up carbon dioxide from the atmosphere to form sodium bicarbonate as shown in the following equations.



The solutions were freshly prepared and used on the same day that they were made. Solutions were made up which contained 10, 15, 20, 25, 30 and 35 millimoles of carbon dioxide per liter. Table 3 below gives the weights of sodium carbonate which dissolved in 1 liter of distilled water gave the required concentration of carbon dioxide.

Table 3  
Spectral Solutions of Sodium Dithionite

10 m.moles/liter required	1.06 gm. Na <sub>2</sub> CO <sub>3</sub> per liter of water
35	n
30	n
25	n
30	n
35	n

-2-

Eight blank Geomax units were set up together with seven Geomax units for each standard solution. These were left overnight and titrated in the morning. Thus there were eight values determined for the blank and seven values determined for each standard. By considering each blank titration value against each standard titration value a total of 56 results were obtained for each standard solution. These results are shown in tables 4 to 9. A mean result has been calculated for each standard solution together with the standard deviation. Frequency distribution curves were plotted for each standard and these are reproduced in figures 9 and 10.

Table A  
Recovery of 10 micromoles/litre of Carbon Dioxide

Bubbles (U) (ml.)	Number of blanks with this reading	Standard (U) (ml.)	Number of standards with this reading	$U - V$ (ml.)	micromoles/litre
3.62	1	2.79	1	0.83	10.38
		2.81	2	0.81	10.23
		2.82	2	0.80	10.0
		2.83	3	0.79	9.88
3.63	3	2.79	1	0.84	10.5
		2.81	2	0.82	10.25
		2.82	2	0.81	10.13
		2.83	2	0.80	10.0
3.64	3	2.79	3	0.85	10.63
		2.81	2	0.83	10.38
		2.82	2	0.82	10.25
		2.83	3	0.81	10.13
3.65	2	2.79	1	0.86	10.75
		2.81	2	0.84	10.5
		2.82	2	0.83	10.38
		2.83	2	0.82	10.25
Mean value = 10.2 ± 0.39					

Table 5

## Recovery of 16 micrograms/Ultre of Carbon Dioxide

Blanks (U) (ml.)	Number of blanks with zero reading	Standards (U) (ml.)	Number of standards with zero reading	$U - U$ (ml.)	micrograms/Ultre
3.62	1	2.44	6	2.19	14.95
		2.45	2	2.17	14.69
3.63	3	2.44	6	2.19	14.87
		2.45	3	2.18	14.75
3.64	3	2.44	6	2.20	15
		2.45	2	2.21	15.105
3.65	1	2.44	6	2.21	15.19
		2.45	1	2.22	15.25
Mean value = 14.96 ± 0.79					

Table 6  
 Recovery of 20 n.moles/liter of Cadmium Monoxide

Dilution (D) (ml.)	Number of dilutes with tide reading	Standard (S) (ml.)	Number of standards with tide reading	D - S (ml.)	n.moles/liter
2.62	2	2.02	1	1.6	20.0
		2.04	1	1.68	19.75
		2.05	2	1.67	19.63
		2.07	1	1.59	19.36
2.63	3	2.02	1	1.61	20.13
		2.04	4	1.59	19.86
		2.05	1	1.58	19.75
		2.07	3	1.56	19.5
2.64	3	2.02	1	1.62	20.25
		2.04	4	1.60	20.0
		2.05	1	1.59	19.86
		2.07	1	1.57	19.63
2.65	2	2.02	1	1.63	20.36
		2.04	4	1.61	20.13
		2.05	2	1.60	20.0
		2.07	1	1.58	19.86

Mean value = 19.92 ± 0.27

## Table 2

## Recovery of 25 n.moles/25 ml. of Quinac Mordant

Blanks (B) (ml.)	Number of blanks with title reading	Standard (U) (ml.)	Number of standards with title reading	B - U (ml.)	n.moles/25 ml.
3.62	1	3.62	2	2.00	25.0
		3.64	2	1.98	24.75
		3.66	3	1.96	24.5
3.63	3	3.62	2	2.01	25.13
		3.64	2	1.99	24.88
		3.66	3	1.97	24.65
3.64	3	3.62	2	2.02	25.25
		3.64	2	2.00	25.00
		3.66	3	1.98	24.75
3.65	3	3.62	2	2.03	25.5
		3.64	2	2.02	25.13
		3.66	3	1.93	24.75
Mean value = 24.8 $\pm$ 0.27					

Table 8

## Recovery of 30 n.moles/115cc of Dextro-Diiodide

Blanks (B) (ml.)	Number of blanks with thin wedging	Standards (U) (ml.)	Number of standards with thin wedging	B - U (ml.)	n.moles/115cc
3.62	3	1.26	5	2.36	29.5
	.	1.27	2	2.35	29.35
	.	1.28	2	2.34	29.25
3.63	3	1.26	5	2.37	29.75
	.	1.27	2	2.36	29.5
	.	1.28	2	2.35	29.35
3.64	3	1.26	5	2.36	29.75
	.	1.27	2	2.37	29.65
	.	1.28	2	2.36	29.5
3.65	2	1.26	5	2.39	29.95
	.	1.27	2	2.38	29.75
	.	1.28	2	2.37	29.65
Mean value = 29.67 ± 0.25					

Table 9

## Recovery of 26 n.moles/116cc. of Carbon Dioxide

Blinks (D) (nl.)	Number of blanks with this reading	Standard (U) (nl.)	Number of standards with this reading	D - U (nl.)	n.moles/116cc
3.62	1	0.84	2	2.74	24.25
		0.86	4	2.76	24.5
		0.88	2	2.78	24.75
3.63	3	0.84	1	2.79	24.03
		0.86	4	2.77	24.63
		0.88	2	2.75	24.25
3.64	3	0.84	2	2.80	25.0
		0.86	4	2.78	24.70
		0.88	2	2.76	24.5
3.65	2	0.84	2	2.81	25.12
		0.86	4	2.79	25.0
		0.88	2	2.77	24.63
Mean value = 24.76 ± 0.21					

SODIUM CARBONATE STANDARDS  
frequency distribution




Fig. 2

Frequency distribution curves of sodium carbonate standards  
containing 10, 15 and 20 m.moles/litre of carbon dioxide

## SODIUM CARBONATE STANDARDS

frequency distribution

CALCULATED

25


DETERMINED

242 (mean of 56)  
± 0.27

30

2967  
(mean of 56)  
± 0.15

35

34.76 (mean of 56)  
± 0.21No.  
of  
DETS.Fig. 10

Frequency distribution curves of sodium carbonate standards containing 25, 30 and 35 m.moles/litre of carbon dioxide

A number of plasma samples were also set up to test repeatability and the results are given in table 10.

Table 10  
Receptability of Plasma Samples

Bloks (B) (ml.)	Number of bloks with this reading	Urinom (U) (ml.)	Number of unknowns with this reading	B - U (ml.)	CO <sub>2</sub> conc. in miller/3 liter
<b>Plasma 1</b>					
4.37	2	2.36 2.37	3 2	1.91 1.90	23.66 23.75
4.38	4	2.36 2.37	3 2	2.92 1.91	24.69 23.89
					Mean value = 23.92 ± 0.06
<b>Plasma 2</b>					
4.37	3	2.34 2.36	4 1	2.09 2.01	25.26 25.13
4.38	4	2.34 2.36	4 2	2.04 2.02	25.30 25.23
					Mean value = 25.22 ± 0.1
<b>Plasma 3</b>					
4.37	2	2.66 2.68	2 3	2.52 2.49	31.38 31.29
4.38	4	2.66 2.68	2 3	2.52 2.50	31.50 31.25
					Mean value = 31.25 ± 0.12
<b>Plasma 4</b>					
4.37	2	2.03 2.09 2.10	2 2 1	2.09 2.08 2.07	26.19 26.00 25.88
4.38	4	2.03 2.09 2.10	2 2 1	2.10 2.09 2.08	26.26 26.19 26.00
					Mean value = 26.12 ± 0.115

### (3) Plasma pH

Since pH varies inversely with temperature the plasma pH had to be measured either at body temperature or at a lower temperature and temperature coefficients applied to the values obtained. Several colorimetric methods of determination of plasma pH have been described, many using phenol red as the indicator (Hastings and Henderoy, 1924; Shock and Hastings, 1934; Rusbridge, 1948, and Holden, 1953). The following method described by Van Slyke, Veldeinger and Van Slyke (1949) also using phenol red has been adopted to the EELC colorimeter<sup>a</sup> and has been used in the present study.

#### Reagents

- (a) A stock solution of disodium hydrogen phosphate containing 142 gm. of the anhydrous salt per litre.
- (b) A stock solution of potassium dihydrogen phosphate containing 236.3 gm. of the anhydrous salt per litre.

These two solutions were kept in stoppered Pyrex flasks in the refrigerator at 4° C.

- (c) Phenol red. A stock solution of 0.15 phenol red was made up by dissolving 100 mg. of phenol red in 20.2 ml. of 0.01 N NaOH and diluting to 100 ml. This was prepared several days before required.
- (d) A 0.003% dilution of phenol red was made by dilution of 20 ml. of

<sup>a</sup> Evans Electrocolorimeter Ltd., Harlow, Essex.

the 0.3% stock to 250 ml. with water. This solution was also stored in a refrigerator at 4° C.

(e) A 22.5% solution of sodium chloride.

(f) A 0.9% solution of sodium chloride.

(g) Neutralized liquid paraffin. This was made by mixing about 200 ml. of liquid paraffin with 200 ml. of distilled water to which a few drops of 0.3% phenol red solution had been added. The mixture was shaken vigorously and 0.02 N NaOH was added drop by drop until the watery emulsion of Andover became permanently pink. The oil was separated, centrifuged, re-separated and stored in a stoppered flask.

(h) Neutralized saline-dye solution with 8 mg. of phenol red per litre.

This solution was made immediately before use since the pH fell if it stood for more than an hour or two. Into a 100 ml. volumetric flask were placed 4 ml. of the 22.5% solution of sodium chloride and about 80 ml. of water. 10 ml. of the 60 mg. per litre phenol red solution were measured accurately and added. Drop by drop 0.02 N NaOH was added, stirring by rotation after each addition until the pH was about 7.4 (a colorimeter reading of 24). The flask was then filled with water until about 0.2 ml. of the mark and more 0.02 N NaOH added a drop at a time until the pH was between 7.4 and 7.6 (colorimeter readings 24 - 26). This flask was kept stoppered to retard absorption of atmospheric carbon dioxide.

Preparation of standard phenolate-dye solutions of known pH

From the 0.5 N stock solutions previously prepared mixed solutions of disodium hydrogen phosphate and potassium dihydrogen phosphate were prepared. These solutions mixed in the proportions shown in Table II gave the pH values indicated when they were diluted to 1/25.

Table II  
Volumes of Phenolate Solutions  
To be Mixed to Produce the pH Values Required

0.5 N stock solution		pH obtained at temperatures indicated when diluted to 1/25	
Na <sub>2</sub> HPO <sub>4</sub> , ml.	KH <sub>2</sub> PO <sub>4</sub> , ml.	20° C.	35° C.
50	22.63	7.13	7.30
50	33.40	7.43	7.40
50	5.97	7.93	7.70

To obtain the correct 1/25 dilution and correct pH, three separate curvatures were measured 8.15 ml. of water, 1.35 ml. of the appropriate 0.5 N mixed phenolate buffer and 1.0 ml. of the 50 mg. per liter phenol red solution. After addition of the dye, the mixture was each stirred with a footed rod and immediately stoppered to protect from atmospheric carbon dioxide.

During the development of the method standard phosphate-dye solutions were prepared and readings of optical density were made in a Unicam S.B. 600 spectrophotometer<sup>5</sup>. A non-linear calibration curve was obtained as described by Van Slyke et al. (1949) with specially selected test tubes in the test tube carrier. This is illustrated in Figure 11.

It was possible to obtain repeatability at 20° C. but not at 30° C. unpredictable errors occurred and the source of these errors was found to lie in the heavy metal construction of the cuvette carriers. A cuvette containing a phosphate buffer solution warmed to 30° C. and placed in the spectrophotometer, rapidly lost heat to the metal carrier so that the optical density was read at an unknown and random temperature somewhere between 20° C. and room temperature. The rapidity of heat loss from such a solution when placed in a test tube cuvette carrier in the Unicam S.B. 600 spectrophotometer is shown in Table I2.

This source of error was overcome with the special carrier by heating the metal cuvette carrier prior to placing in the spectrophotometer and thus slowing the heat loss from the cuvettes enabling a reading to be taken.

A calibration curve was then attempted using an EEL<sup>6</sup> colorimeter fitted a no. 625 ELLIOTT goldline filter whose absorption peak lies in the region of 545 m. $\mu$ . The phosphate-phenol red buffer mixtures were made

<sup>5</sup> Unicam Instruments Co. Ltd., Cambridge.

Table 12  
Temp. Loss from Test Tubes in the  
Test Tube Guards Carrying in the Spectrophotometer

Time (sec.)	Sample 1 Temp., °C.	Sample 2 Temp., °C.	Sample 3 Temp., °C.
0	39	39	39
5	38.5	38	38.5
10	38	37.5	38
15	37.5	37	37.5
20	37	36	36.7
25	36	35.4	35.0
30	34.6	34	35
45	33.5	33	34

up in the colorimeter tubes. Prior to reading at 20° C. the colorimeter tubes were heated in a water-bath to 23° C. The calibration curve obtained at 20° C. using a number 625 filter is illustrated in figure 11 together with a calibration curve obtained on the spectrophotometer.

From the experience acquired in attempting to obtain a calibration curve of the phosphate buffer-dye solution in the spectrophotometer at 36° C., a check was made of the temperature loss from the colorimeter tubes containing 10.5 ml. of fluid when placed in the "ELV" colorimeter.




Fig. 11

Comparison of calibration curves obtained  
from the spectrophotometer and the colorimeter.

A slower loss of heat was observed than occurred in the spectrophotometer and this was thought to be due to the plastic holder for the colorimeter tubes within the colorimeter. The results obtained in the colorimeter compared with heat loss in the spectrophotometer are given

In table 23.

Table 23  
Comparison of Heat Loss from Specimen  
Within the Colorimeter and the Spectrophotometer

Blue (cces.)	Temperature in degrees centigrade G.E.C. colorimeter			Spectrophotometer		
	Sample 1	Sample 2	Sample 3	Sample 1	Sample 2	Sample 3
0	39	39	39	39	39	39
5				38.5	38	38.5
10	39	38	38	38	37.5	38
15				37.5	37	37.5
20	38	36	36	37	36	36.5
25				36	35.5	35.5
30	37	36	37.5	34.5	34	35
35	36.5	37.5	37.5			
40				33.5	33	34
45						
50	36	36.5	37			

Calibration curves were then carried out using an O.G.R.I. sample glass filter. A straight line calibration curve was obtained at 20° C. and at 30° C. Since reproducible, straight line calibration curves were obtained with the O.G.R.I. filter, the method was then used to

determine plasma pH and the temperature coefficients of blood and plasma.

These calibration curves are shown in figure 12.




Fig. 12

Calibration curves phosphate buffers

For each determination three colorimeter tubes were used. Into two of these were pipetted 10 ml. of the neutralised saline dye solution for duplicate determinations and these solutions were immediately covered with a layer of neutralised liquid paraffin. Into the third

colorimeter tube 10 ml. of 0.9% saline were pipetted to receive a plasma blank. Into each of the two colorimeter tubes containing the 10 ml. of saline-dye solution was placed 0.5 ml. of plasma.

To obtain the 0.5 ml. of plasma from the centrifuge tube in which it had been collected, the rubber bung was removed and the plasma was withdrawn using a long tipped 0.5 ml. syringe pipette. A third 0.5 ml. of plasma was run into the colorimeter tube containing the saline. The contents of each colorimeter tube were mixed with a clean footed rod and the two tubes were placed in a water-bath at 22° C. for five minutes.

The zero of the colorimeter was set with the plasma-saline blank and the readings of the colorimeter tubes taken. By reference to a previously determined calibration curve the plasma pH was found.

It was necessary to check first the repeatability of the method with reference to plasma pH determinations and to do this pH determinations were made on 10 plasma samples. The results of the pH determinations at 20° C. are given in Table II. The results are not corrected to the body temperature of the animal. From these results it appeared that this method could be used for the determination of plasma pH of cattle.

Table 14  
Effect of Consecutive Heterologous transfusions on the same Sample

Sample	a	b	c	d	e	f
1	7.695	7.695	7.695	7.695	7.695	7.695
2	7.670	7.670	7.655	7.655	7.655	7.655
3	7.680	7.675	7.680	7.675	7.680	7.680
4	7.490	7.485	7.485	7.485	7.485	7.485
5	7.625	7.625	7.625	7.625	7.620	7.620
6	7.420	7.420	7.420	7.420	7.425	7.425
7	7.780	7.770	7.755	7.770	7.770	
8	7.755	7.755	7.760	7.760	7.770	
9	7.740	7.725	7.740	7.740		
10	7.770	7.760	7.760	7.760		

Since it was possible neither to find the blood pH directly at 35° C., nor to separate the plasma at 35° C., it was necessary to determine temperature coefficients for both blood and plasma. Rosenthal (1946) found that in order to obtain pH values for plasma, exactly equal to those of plasma as drawn, it was necessary to centrifuge at 25° C. If blood were cooled to 20° C. and centrifuged at that temperature and the separated plasma was then warmed to 35° C.

For pH measurements, the pH obtained was about 0.03 units greater than that observed when the blood was centrifuged at 20° C. It is shown that the reason for this was the difference in temperature coefficients for blood pH and plasma pH, the blood temperature coefficient being greater. Van Slyke *et al.* (1949) considered that since practically all the data in the literature concerning the pH of human plasma were derived from blood centrifuged at room temperature, for comparison it was justifiable to continue to centrifuge at room temperature.

When blood samples were drawn from cattle they were not drawn in the relatively warm atmosphere of a hospital ward and immediately transported to a warm laboratory. Blood samples were withdrawn into a relatively cold atmosphere of a byre and as a consequence were subjected to considerable cooling prior to centrifugation. Although it was found that the temperature of the centrifuge used was in the region of 20° to 25° C. while in use, it was considered that the cooling, prior to centrifugation, may have affected the results obtained. A standard procedure of bringing blood samples to 20° C. prior to centrifugation was therefore adopted and temperature coefficients were found for both blood pH and plasma pH.

The temperature coefficient of the pH of bovine plasma was found by taking plasma samples and finding the pH difference when read on the "EEL" colorimeter at two known temperatures, 20° C. and 25° C., for which calibration curves had been prepared using the phosphate buffer-tyro solution. Results are given in table 26.

**Table 15**  
**Temperature Coefficient of Dovlop Plasma (33 samples)**

Sample	pH 20° C.	pH 30° C.	Difference pH units
1	7.695	7.465	.240
2	7.630	7.435	.195
3	7.665	7.510	.155
4	7.690	7.520	.160
5	7.730	7.570	.160
6	7.670	7.495	.175
7	7.695	7.490	.205
8	7.740	7.555	.185
9	7.745	7.560	.185
10	7.635	7.425	.240
11	7.645	7.510	.135
12	7.705	7.545	.160
13	7.680	7.525	.155
14	7.635	7.465	.160
15	7.645	7.490	.155
16	7.630	7.530	.150
17	7.630	7.495	.135
18	7.640	7.470	.170
19	7.630	7.495	.135
20	7.705	7.555	.150
21	7.655	7.570	.145
22	7.690	7.530	.160
23	7.655	7.490	.165
24	7.775	7.595	.180
25	7.740	7.580	.160
26	7.640	7.450	.190
27	7.660	7.575	.145
28	7.660	7.495	.165
29	7.655	7.525	.130
30	7.570	7.420	.150
31	7.460	7.290	.160
32	7.600	7.430	.170
33	7.650	7.405	.245

Mean  $\pm .165 \pm 0.018$

A mean difference in pH of 0.255 pH units was found for 20° C. A temperature coefficient for the pH of bovine plasma of 0.009 pH units per degree centigrade was calculated; in a similar manner a temperature coefficient was found for canine plasma. The results obtained are given in Table 36.

Table 36  
Temperature Coefficient of Canine Plasma (10 samples)

Sample	pH 20° C.	pH 30° C.	Difference pH units
1	7.66	7.43	.23
2	7.56	7.33	.23
3	7.69	7.32	.21
4	7.62	7.40	.21
5	7.67	7.46	.21
6	7.76	7.50	.26
7	7.69	7.42	.21
8	7.72	7.45	.26
9	7.70	7.47	.23
10	7.73	7.49	.24
Mean = .232 ± 0.013			

A mean difference of 0.222 pH units was found for 25° C. A temperature coefficient for the pH of canine plasma of .012 pH units per degree centigrade was calculated.

The temperature coefficient of bovine blood was determined in an indirect manner since a method for the determination of blood pH had not yet been developed. Rosenthal (1940) pointed out that blood and plasma had the same pH at the temperature of separation. Therefore if the plasma could be separated at two different temperatures and the plasma pH found at those different temperatures, then the blood pH could be determined at those temperatures; if the blood pH was known at two temperatures it was possible to find the blood pH temperature coefficient.


Two blood samples were taken from the same animal at the same time and separated at 0° C. and 25° C., respectively, and the plasma pH was found for each at 20° C. The plasma pH values were corrected to the respective temperatures of separation to obtain blood pH at 0° C. and 25° C. and from these two values the temperature coefficient of the pH of this bovine blood could have been determined. However, in order to deal with a number of blood samples from different cattle a slightly different procedure was adopted. Two samples were obtained from each animal and after separation at 0° C. and 25° C. the pH values were found for the plasma at 20° C. The results obtained are given in table XV.

Table XI  
The Difference in Plasma pH Caused by Centrifugation  
of Blood at 25° C. and 0° C.

Sample	Plasma pH at 25° C.		Difference pH units
	Separation at 25° C.	Separation at 0° C.	
1	7.510	7.569	.059
2	7.538	7.585	.047
3	7.595	7.670	.075
4	7.620	7.675	.055
5	7.555	7.630	.075
6	7.535	7.600	.065
7	7.560	7.630	.070
8	7.625	7.705	.080
9	7.535	7.585	.050
10	7.400	7.460	.060
11	7.545	7.620	.075
12	7.450	7.530	.080
13	7.550	7.595	.045
14	7.605	7.660	.055
15	7.625	7.720	.095
16	7.620	7.690	.070
17	7.610	7.685	.075
18	7.565	7.635	.070
19	7.625	7.720	.095
20	7.585	7.660	.075
			Mean = .070 ± 0.015

From this table the mean difference in plasma pH due to difference in temperature of centrifugation of the blood was found to be 0.07 pH

units. A graph was constructed, as illustrated in figure 13, of temperature against pH and on this graph two parallel lines were drawn, of slope, the temperature coefficient of plasma and these two lines were separated by the mean difference of pH due to difference of temperature of centrifugation.



Temperature coefficients bovine blood and plasma

The points marked on this graph where blood and plasma had the same pH. These were the points on each plasma line at the temperature of centrifugation. By joining these two points a line of slope, the temperature coefficient of bovine blood, was obtained. The temperature coefficient of bovine blood found in this way was 0.012 pH units per degree centigrade.

Similarly the temperature coefficient of canine blood was also found. Duplicate samples were taken, one at 0° C. and 25° C. and the plasma pH found at 20° C. The values obtained are given in table 13.

Table 13

The Difference in Canine Plasma pH Caused by Centrifugation

of Blood at 25° C. and 0° C.

Sample	Plasma pH at 20° C.		Difference pH units
	Separtion at 25° C.	Separtion at 0° C.	
1	7.66	7.74	.080
2	7.58	7.69	.030
3	7.61	7.70	.090
4	7.66	7.73	.070
5	7.49	7.51	.020
6	7.38	7.48	.100
7	7.60	7.77	.070
8	7.73	7.80	.070
9	7.73	7.77	.040
20	7.51	7.59	.080
			Mean = .060 ± 0.018

The mean difference due to difference in temperature of centrifugation was found to be 0.06 pH units. A graph was drawn as illustrated in figure 14 with the two plasma lines separated by this mean difference.




Fig. 14  
Temperature coefficients canine blood and plasma

The pH of plasma and hence of blood at the temperature of separation was found and the line joining these was drawn, the slope of which was the temperature coefficient of the pH of canine blood. The temperature coefficient of the pH of canine blood found in this way was 0.0255 pH units per degree centigrade.

When temperature coefficients had been found for bovine and canine blood and plasma, respectively, the following method was adopted to determine the pH of cattle and dog plasma at 30° C. Blood samples were taken with strict anaerobic precautions as previously described, brought to the laboratory and placed in a water-bath at 20° C. for fifteen minutes to ensure a constant temperature before being placed in the centrifuge. The blood samples were centrifuged at a known temperature (usually 20° C.) and then the plasma pH was found using the "EEL" colorimeter as previously described. To the plasma pH found at 20° C. it was necessary to apply temperature coefficients to obtain the plasma pH at 30° C. The plasma pH at 20° C. was first corrected to the plasma pH at the temperature of separation by application of the temperature coefficient of bovine plasma and then the blood temperature coefficient was applied to correct the pH to 30° C. in the following manner.

For bovine blood and plasma if X be the temperature of separation and Y the temperature of reading then the plasma pH at 30° C. =  
pH at Y° C. + (X - Y)(0.009) - (20 - X)(0.012) where 0.009 is the calculated temperature coefficient of plasma and 0.012 is the calculated temperature coefficient of blood.

#### (4) Ventilation rate

Ventilation rate may be defined as the volume of gas expired in one minute. It is affected by the rate and depth of respiration, both of which may undergo alteration.

The extensive study reported by Brady (1950) was conducted using a variety of spiroimeters. In the present study the ventilation rate was measured by means of a face mask connected to unidirectional valves in such a manner that the bovine subject inspired from the atmosphere and expired to the atmosphere through a gas meter.

The face masks (Gerr's mask and Burle's mask) used for the anaesthetization of adult cattle were examined for their suitability for the measurement of the ventilation rate of cattle and were rejected since they were unsightly, not gas tight and also the dead space within the masks was too large. A mask was therefore fabricated to overcome these disadvantages. For the main body of the mask a open aluminium cone as was used for shop lighting was required. This had a diameter of six inches at its base and was 10 inches long. A one and a half inch brass nipple was forced through the apex of the cone and secured in this position by means of jam-rings with rubber washers acting as gas seals. Attached to the open base of the cone, by means of adhesive Septaflex<sup>®</sup>, was a six inch width of flexible rubber tubing of five inches

diameter. The rubber tubing enabled a gas tight seal to be made over the animal's face and the spun aluminium cone over the animal's nostrils could be moulded to the particular animal's facial contour. This is illustrated in figure 15.



Fig. 15

Face mask in position on a cow.

For calves a similar mask was fabricated. A smaller spun aluminium cone of length five inches and diameter of base four inches

was selected. A three-quarter inch nipple was forced through the apex and secured with the appropriate size jam rings and rubber washers. The air seal of the base of the mask to the face of the calf was made in one of two ways. On some occasions a rubber ring obtained from the cuff of a surgical rubber glove was used to make the seal while on other occasions an inflatable cuff, as used for a human type face mask, was attached.

The mask for adult cattle is illustrated in figure 16.




Fig. 16

Mask for adult cattle.

The masks used for calves are illustrated in figures 17 and 18.



Fig. 17

Calf mask with inflatable cuff seal.



Fig. 18

Calf mask with rubber ring seal.

Each mask was connected by means of the brass nipple at the apex to a 'Y' piece of appropriate size. For the adult mask the nipple screwed into the one inch stem of a 'Y' piece made from one and a half inch diameter copper tube. The other two arms of the 'Y' piece were connected to an inspiratory and expiratory valve, respectively. These connections were made by means of 18 inch lengths of flexible plastic tube of one and a quarter inch internal diameter ("Hooverflex"<sup>®</sup>). For the calf a 'Y' piece made of three-quarter inch diameter copper tube was used and the connections to the unidirectional valves were of one inch long thin flexible rubber tubing of the kind used in an anaesthetic apparatus.

The unidirectional valves used were of the diaphragm type. For the adult cow all arms of the valves were of a diameter of one and a quarter inch internal diameter while the rest of the valve was of appropriate size. For the calf the arms were of three-quarter inch internal diameter. Drawings of the valves giving the dimensions are shown in figures 19 and 20.

The expiratory side of the unidirectional valve was connected directly to the inlet of the gas meter. The gas meter used was the Parkinson Comin no. 004 (1½" outlet)<sup>®</sup> which measured a gas flow of up to ~~700~~ 11 litres per minute with a minimal resistance. This gas meter is

<sup>®</sup> Hoover Ltd., Farnham, Middlesex.

<sup>®</sup> Parkinson Comin Ltd., City Road, London.

shown in figure 21.




Fig. 19  
Unidirectional valves for adult cattle.




Fig. 20  
Unidirectional valves for calves.

This particular meter was obtained after consultation with Dr. J. V. A. Dinsdale who was using two such meters in the Scottish Hepatitis Research Trust unit in Glasgow.

Measurements were made of the resistance to gas flow of the gas meter together with the additional valves and connecting tubes as used when pulmonary ventilations were recorded. The results are given below.

	Rate of flow in litres per min.	Resistance in centimetres of water
(a) Small valves and gas meter	50	2
	100	5
	150	6
	180	14

Suitable for the measurement of pulmonary ventilations up to 50 litres/minute.

(b) Large valves and gas meter	130	2.5
	240	2.5
	320	4.5
	400	7.0

Suitable for the measurement of pulmonary ventilations over 50 litres/minute and less than 130 litres/minute.

7-015

Ability of the Parkinson-Green Gas Meter to Record

Accurately Pulsatile Air Flow

The ability of the gas meter to record accurately pulsatile air flow was tested by the use of an apparatus which had been developed by Deunda and Huxley (1959) to calculate the Kestrel respirometer. This apparatus gave a pulsatile gas flow at physiological respiration frequencies (32 to 60 per minute) and covered a range of pulsatile flows up to 200 litres per minute.

The volume control on the apparatus was adjusted first to give a steady flow of 200 litres per minute and then the interrupter was started so that a pulsatile flow corresponding to a respiratory rate of 36 per minute was produced. Reduction of total flow took place to 60 litres per minute.

The output from the apparatus was put through the gas meter and the volume as recorded by the gas meter was noted. The volume was collected into a Douglas bag and afterwards measured.

The same experiment was repeated with a pulsatile flow corresponding to that obtained with a respiration rate of 60 per minute and a ventilation of 80 litres per minute.

The steady flow was then increased to 400 litres per minute and measurements were made with the interrupter adjusted first to correspond

to a respiratory rate of 36 per minute and a ventilation of about 146 litres per minute then measurements were again made and also after adjustments had been made to the interruptor to correspond to a respiratory rate of 60 per minute and a ventilation of 168 litres per minute. Table 18(a) gives the results obtained.

Table 18(a).

Accuracy of Gas Meter in Recording Pulsatile Flow

Experiment	Steady Flow litres/min.	Respiratory Rate	Ventilation litres/min.	
			Gas Meter	Douglas Bag
1	200	36	68	68
2	200	36	67	67
3	200	60	79	81
4	200	60	80	82
5	400	36	146	142
6	400	36	144	140
7	400	60	168	168
8	400	60	164	168

The average error was 1.6% which would not interfere with the results.




Fig. 21

Parkinson Cowen CO<sub>2</sub> gas meter  
with calf mask and unidirectional valves attached.

Having constructed the apparatus, the method of use was then developed. The respiratory efforts of a bovine subject were observed when the animal was at rest and not connected to the apparatus. Observation was made of both the rate and depth of respiration. The animal was then fitted with the face mask alone and observation was

again made of the depth and rate of respiration. It was noted that the majority of cattle resented the fitting of the face mask and required slight restraint to maintain the mask in place. Most animals, after the initial resistance to fitting the mask, required only their heads to be held gently with the mask in position so that within two minutes of fitting the mask the rate of respiration had returned to normal. With the face mask on the animal, connection was made to the unidirectional valves by means of the 'Y' piece and the flexible tubes. The ventilation rate was measured from the gas meter for periods of five or 10 minutes.

Calves were found to be more difficult subjects than adult cattle since on occasions, after they had apparently settled down, they suddenly decided to struggle violently, settle down and then struggled violently again. When this occurred studies had to be abandoned in some calves.

The measurement of the ventilation rate of a calf is illustrated in figure 22.



Fig. 22

Measurement of the ventilation rate of a calf.

### (5) Respiratory rate

The respiratory rate was counted while the ventilation rate was being measured. The clicks of the clippings of the expiratory valve were counted and the total for each minute recorded.

## RESULTS

- (1) A comparison of the effects of oxides of ammonia on the carbon dioxide content and the pH of the plasma.
- (2) The carbon dioxide content of the plasma of (a) normal Ayshire cattle  
(b) normal dogs.
- (3) The pH of the plasma of (a) normal Ayshire cattle  
(b) normal dogs.
- (4) The physiological variations of the carbon dioxide content and the pH of the plasma of normal Ayshire cattle.
- (5) The ventilation and respiratory rates of Ayshire cattle.

## Standardized Analysis

Throughout this thesis where mean values are shown they are followed by the standard deviation.

Differences stated as significant indicate that the probability of the given results having occurred by chance is less than 5%. Student's "t" test was used throughout.

(1) A comparison of the effects of storage of samples on the carbon dioxide content and the pH of the plasma.

(a) Contamination of tubes

The possibility of errors arising from the contamination of blood sampling tubes was considered as follows. Duplicate blood samples were taken and duplicate examinations were carried out on the plasma of each sample. The results are given in table 19.

Table 19  
Variations in the CO<sub>2</sub> Content and the pH of Plasma  
Due to Contamination of Sampling Tubes

Cow	CO <sub>2</sub> , m.moles/litre				pH			
	Tube 1	Tube 2	Tube 1	Tube 2	Tube 1	Tube 2	Tube 1	Tube 2
1	30.5	30.6	30.5	30.5	7.435	7.43	7.43	7.43
2	29.5	29.5	29.5	29.3	7.44	7.44	7.47	7.47
3	30.8	31	30.8	31	7.47	7.47	7.485	7.485
4	27.6	27.8	27.5	23	7.42	7.42	7.43	7.43

From these results it will be seen that with the precautions adopted for the cleaning of glassware, variations from this source were not more than 0.3 m.moles/litre in the carbon dioxide content and not

more than 0.03 pH units in the determination of the plasma pH. Nevertheless the variations in the determination of the plasma pH were of sufficient magnitude to justify the practice of taking duplicate samples and doing single pH determinations on them.

#### (b) Exposure to atmosphere

It has been emphasised repeatedly that blood or plasma samples examined for their carbon dioxide content should not be exposed to the air because of the loss of carbon dioxide to the atmosphere (Garroway, 1950). To study this source of error blood samples were obtained from the brachial arteries of five cows. The samples were taken at the same time from each cow and one sample was collected with exposure to air while the other was collected under liquid paraffin and sealed with a rubber bung. The time between taking the first sample from the first cow and analysis of the plasma was 25 minutes while the interval between taking the samples from the last cow and analysis of the plasma was approximately 15 minutes. The results are given in table 20.

Table 20  
 Effect of Exposure of Blood to the Atmosphere  
 on the  $\text{CO}_2$  Content and the pH of Plasma

Case	$\text{CO}_2$ content in microliters/liter		$\text{CO}_2$ loss in microliters /liter	Plasma pH		pH difference
	Aerobic	Anaerobic		Aerobic	Anaerobic	
1	35.9	37.1	1.2	7.50	7.44	.06
2	37	37.8	0.8	7.50	7.42	.08
3	39.9	39.4	0.5	7.35	7.30	.05
4	30	30.4	0.4	7.37	7.36	.01
5	35.6	37.5	0.7	7.37	7.37	.00

The results showed once more that blood samples must be removed without exposure to the atmosphere otherwise a loss of carbon dioxide of up to 1.2 microliter/liter might occur with a pH difference as great as 0.06 pH units.

### (e) Liquid paraffin seal

Van Slyke (1949) stated that absorption of carbon dioxide by a protective layer of liquid paraffin could lead to errors due to absorption of the carbon dioxide by the paraffin. This source of error was studied by collecting blood in 30 ml. centrifuge tubes on one occasion, and on another occasion by collecting in 25 ml. centrifuge

tubes. In the 50 ml. tubes the blood was covered by two inches, one inch, half an inch and minimal layers of liquid paraffin, while in the 25 ml. tubes the layers of liquid paraffin were half inch, quarter inch and minimal layers of liquid paraffin. The results are shown in table 23.

Table 23.

Effect of the Layer of Carbon Dioxide to the Protective Layer of Paraffin on the CO<sub>2</sub> Content and the pH of Plasma

	CO <sub>2</sub> content	Electro. pH
<u>50 ml. centrifuge tubes</u>		
2" Liquid paraffin above blood	30.25	7.56
1" Liquid paraffin above blood	30.9	7.54
½" Liquid paraffin above blood	31.25	7.53
Minimal liquid paraffin above blood	31.5	7.52
<u>25 ml. centrifuge tubes</u>		
½" Liquid paraffin above blood	23.25	7.48
¼" Liquid paraffin above blood	23.625	7.46
Minimal liquid paraffin above blood	23.75	7.44

It can be seen that as great as 1.2 milligrams/liter in the plasma carbon dioxide content and as great as 0.05 pH units can occur

due to absorption of carbon dioxide by liquid paraffin when the layer was greater than minimal. Accordingly minimal layers of liquid paraffin have been used in this study.

#### (d) Autolysis on plasma pH

Holloway and Baityor (1954) determined colorimetrically the pH of serum samples obtained by adding blood samples with anticoagulant and allowing them to remain on the bench for a period of two hours in order that the serum might separate out. It is difficult to see how they obtained serum samples from bovine blood after two hours when bovine serum usually takes much longer to separate out unless the samples were refrigerated. Using this method they obtained apparently normal values in spite of the fact that Rosenthal (1948) had demonstrated that the effects of autolysis in blood samples standing for this length of time at 20° C. would cause a very marked fall in plasma pH.

An experiment was designed to check the effects of autolysis on plasma pH and to find out if immediate centrifugation would limit autolysis. Eight blood samples each in a 50 ml. centrifuge tube were taken from the breechled uterine of a cow. All the tubes were placed in ice for 35 minutes to reduce to a constant temperature, after which they were exposed to room temperature (16° C.). Tubes 1 to 5 were immediately centrifuged for 15 minutes and plasma pH of tubes 1 and 5 determined. After one and a half hours exposure to 16° C., tube 6 was

centrifuged and the pH determined on tubes 2 and 6. This procedure was repeated at half hour intervals on tubes 3 and 7, and 4 and 8, respectively. The results obtained are given in table 22.

Table 22

Effect of Autolysis on Plasma pH

	Centrifugation at start			Centrifugation at intervals		
	Tube no.	pH	pH	Tube no.	pH	pH
Initial	1	7.90	7.90	5	7.45	7.475
½ hour	2	7.905	7.905	6	7.45	7.465
1 hour	3	7.465	7.465	7	7.36	7.39
1½ hours	4	7.42	7.42	8	7.33	7.39

The results confirmed Repenthal's findings that autolysis caused a considerable drop of plasma pH (0.3 pH units) in one hour but that separation of the cells to the bottom of the centrifuge delayed the autolytic effect. As a result of these observations all blood samples were centrifuged and the pH determined within half an hour of collection. Where this was not possible, samples were placed in ice until the determinations were carried out.

It is difficult to draw any conclusion as to how McSherry and Ordway (1954) obtained their normal results by determination of the

serum pH. The only possible explanation is that by reading at room temperature they obtained a pH higher than the true pH of the serum samples since the effect of not taking into account a temperature coefficient for plasma or serum would counteract the lowering of serum pH caused by autolysis.

- (2) The carbon dioxide content of the plasma of (a) normal Ayrshire cattle  
(b) normal dogs

(a) Normal Ayrshire cattle

Blood samples by puncture were taken from the jugular veins of 29 healthy Ayrshire cows and the carbon dioxide content of the plasma was determined. The results are given in Appendix I. A mean value of  $27.4 \pm 1.9$  m.moles/litre was obtained. Blood samples were also taken from the jugular veins of 10 cows by means of a jugular venous catheter as already described. It was assumed that these samples approximated to mixed venous samples of cattle. The mean value of  $29.2 \pm 2.7$  m.moles/litre obtained for the carbon dioxide content of the plasma derived from mixed venous samples was not significantly higher than the mean value obtained for the carbon dioxide content of venous plasma derived from jugular venous blood obtained by puncture. The individual results are given in Appendix II.

Over a period of three years 159 arterial blood samples were taken from healthy cows in the University herd. This herd consisted of non-pregnant Ayrshire cows and was run on a commercial basis so that unproductive cows were quickly removed from the herd. This factor, combined with culling because of age, meant that there was a constant turnover of animals in the herd over the three years.

These cows were in varying stages of pregnancy and lactation. During the winter they were housed for most of the day but allowed out for a short time for exercise. In spring and summer they were housed at night while in summer they were out day and night except for milking. They were fed a production mixture of cattle cake, beans and oats throughout the year and in addition, when grass was not available, they received hay, silage, roots and kale.

The arterial samples were all taken from three cows at approximately the same time each day (8 a.m.), one hour after morning milking. Notes were made of the condition of the animal; whether dry or lactating, pregnant or non-pregnant. Samples were taken on nine different occasions during those three years as shown in table 20. The number of cows sampled on each occasion varied from eight to 23 and duplicate samples were taken from each cow on each of which a single estimation was made. Detailed results are given in Appendix III, while the mean values together with the standard deviations of the carbon dioxide content of the plasma obtained on each occasion are given in table 20.

The mean value obtained for the carbon dioxide content of the arterial plasma of 159 dairy cows was not significantly different from the mean values obtained for the carbon dioxide content of jugular venous plasma **and** mixed venous plasma and was identical with that given by Davenport (1950) for the arterial plasma carbon dioxide content of the

Table 22

Mean Values of the CO<sub>2</sub> Content of the Plasma of Ayrshire Cows  
Together with the Dates of Sampling

Year	Month	No. of cows	Mean value CO <sub>2</sub> content m.moles/litre
1955	April	20	27.6 ± 2.2
1955	June	20	25.4 ± 1.9
1955	October	10	27.5 ± 2.2
1956	January	23	27.3 ± 2.5
1956	May	20	27.0 ± 2.1
1957	June	20	25.2 ± 2.3
1958	January	19	23.2 ± 1.4
1958	March	6	27.2 ± 2.7
Total 159			Mean 27.2 ± 2.6

lassen subject.

From this series of observations no significant difference was found between any of the mean values obtained so that it can be inferred that no significant variations occurred in the arterial plasma carbon dioxide content of these cows due to seasonal variation.

The results obtained from pregnant and non-pregnant cows are compared in table 24.

**Table 24**  
**Mean Values of the CO<sub>2</sub> Content of the Fleeces**  
**of Pregnant and Non-Pregnant Cows**

Condition of cows	No. of cows	Mean CO <sub>2</sub> content m.moles/litre
Pregnant	65	26.5 ± 1.73
Non-pregnant	76	27.3 ± 2.36

It will be observed that there was no significant difference.

Similarly the results obtained were analysed on the basis of the lactational status of the cows. The results are given in table 25 and again there was no significant difference.

**Table 25**  
**Mean Values of the CO<sub>2</sub> Content of the Fleeces**  
**of Dry and Lactating Cows**

Condition of Cows	No. of cows	Mean CO <sub>2</sub> content m.moles/litre
Dry cows	15	27.7 ± 1.71
Lactating	126	27.0 ± 2.12

Arterial blood samples were also collected from 83 Ayrshire calves and the carbon dioxide content of the plasma was determined. These calves were purchased in a local market and brought to the Veterinary Hospital where they were housed in individual huts. For the first month of their lives they were fed "Ostomilk" no. 2\*, a dried whole milk product, at a strength of 3 lb. to the gallon of water and they received six to eight pints daily in two feeds. At three weeks of age they were introduced to hay and calf weaner meal, and at four weeks the "Ostomilk" stopped. A variety of antibiotic supplements was given during the first two weeks of life to prevent diarrhoea.

Blood samples were removed from the bronchial arteries of the calves the ages of which varied from one to eight weeks of age. The results obtained are presented in Table 26.

The mean value obtained,  $29.8 \pm 2.5$  n.mole/litre was significantly higher than that found for adult cows.

\* Glaxo Laboratories, Ltd., Greenford, Middlesex.

Table 26

Mean Volume for the  $\text{CO}_2$  Content of the Arterial Plasma  
of Calves Divided on the Basis of Age

Age	Number	Mean $\text{CO}_2$ content $\text{ml. moles}/100\text{ ml.}$
7 days	18	$20.53 \pm 1.7$
14 days	20	$20.3 \pm 2.5$
4 weeks	8	$20.6 \pm 3.0$
6 weeks	21	$20.3 \pm 2.2$
8 weeks	14	$20.9 \pm 1.6$
Total 81		Mean $20.3 \pm 2.6$

(b) Normal dogs

The carbon dioxide content of the arterial plasma of dogs has been reported to be lower than in most species (Spector, 1956). The carbon dioxide content of the arterial plasma of 16 dogs was determined to confirm the reported results. The dogs from which the samples were taken were healthy normal dogs which were being maintained in the Veterinary Hospital. Blood samples were removed from the femoral artery. The mean value obtained,  $21.19 \pm 2.9$  m.moles/litre, was similar to the value of 21.4 m.moles/litre reported by Spector (1956). Those values for the carbon dioxide content of the arterial plasma of dogs are significantly lower than the values found for the arterial plasma of cattle.

(2) The pH of the plasma of (a) normal Ayrshire cattle

(b) normal dogs

(c) Normal Ayrshire cattle

As described in the section on methods, blood samples were brought to a constant temperature in a water-bath, centrifuged at a known temperature, 23-25° C., and the plasma pH was found at 20° C. Using the temperature coefficients previously determined the pH values were corrected to 38° C.

Duplicate determinations of the arterial plasma pH of 169 ml of the venous plasma pH of 55 healthy Ayrshire cows were carried out. Some of these animals were housed in the Veterinary Hospital while others were cows in the University herd. The mean value obtained for the arterial plasma pH at 38° C. was  $7.43 \pm 0.045$  pH units, and the mean value for the venous plasma pH at 38° C. was  $7.39 \pm 0.05$  pH units.

Duplicate determinations were also made on arterial plasma samples from 10 normal calves housed in the Veterinary Hospital and a mean value of  $7.42 \pm 0.036$  pH units was obtained. Venous samples were also obtained from 19 calves and the mean value found for the plasma pH was  $7.38 \pm 0.02$  pH units. These values for plasma pH were similar to those found in adult cattle.

(b) Normal dogs

Determinations were made of the plasma pH of 10 dogs.

Temperature coefficients had already been found for dog blood and plasma as described in the section on methods. Using these temperature coefficients the pH at 33° C. of the plasma obtained from femoral arterial blood of 10 dogs was determined. A mean value of 7.42 pH units with a standard deviation of  $\pm 0.06$  pH units was obtained.

These values are very similar to the values obtained for cattle in this study ( $\text{pH} \approx 7.43 \pm 0.045$ ) but higher than the value given by Spector (1958) for dogs ( $\text{pH} \approx 7.36$ ).

(4) The physiological variations of the carbon dioxide content and the pH of the plasma of normal Ayrshire cattle

Davenport (1950) gave 1.1 m.moles/litre as the difference in the carbon dioxide content of arterial and venous plasma obtained at the same time from the same human subject. He also stated that no variation existed between venous plasma samples taken from different vessels in the same subject as did Holdens and Peleshay (1926). In the literature there is no record of the content of the arterio-venous differences in the carbon dioxide content of the plasma of cattle or of differences in the carbon dioxide contents of the plasma obtained from blood taken from different vessels of cattle. The differences in the carbon dioxide content and pH of plasma obtained from different vessels were studied in five cows. Blood samples were obtained at the same time from the brachial artery, the jugular vein and the mammary vein of each cow. The results obtained for the plasma carbon dioxide contents are given in table 27.

The results demonstrate that an arterio-venous difference in the carbon dioxide content of plasma could be detected but that its magnitude depended on the vein selected. The mean arterio-venous difference between the carbon dioxide content of the plasma from the brachial artery and that from the jugular vein was 1.2 m.moles/litre while the mean arterio-venous difference between the carbon dioxide content of the plasma from the brachial artery and mammary vein was 2.9 m.moles/litre. In the five cows sampled, the mammary venous plasma carbon dioxide content was greater than the jugular venous carbon dioxide content, the mean jugular venous difference being 1.7 m.moles/litre.

Table 27

Arterio-venous Differences in the Plasma  $\text{CO}_2$  Content

Cow	Vessel	Plasma $\text{CO}_2$ m.moles/ litre	Brachial- jugular difference	Brachial- mammary difference	Mammary- jugular difference
32640	Brachial artery	33			
	Jugular vein	30.5	2.5	3.4	1.9
	Mammary vein	32.4			
32652	Brachial artery	33.1			
	Jugular vein	34.25	2.15	2.8	1.65
	Mammary vein	36.9			
32653	Brachial artery	33.75			
	Jugular vein	30.1	1.25	2.75	1.6
	Mammary vein	31.5			
32656	Brachial artery	31.0			
	Jugular vein	36.25	0.25	3.0	2.75
	Mammary vein	39.0			
32657	Brachial artery	25.0			
	Jugular vein	26.63	1.63	2.63	1.0
	Mammary vein	27.69			
		Mean	1.2	2.9	1.7

The results for the differences in plasma pH are given in Table 28.

Table 20  
Arterio-venous Difference in the Plasma pH

Cov	Vessel	Plasma pH	Brachial-Jugular difference	Brachial-Mammary difference
12640	Brachial artery	7.40		
	Jugular vein	7.45	0.05	0.06
	Mammary vein	7.44		
12652	Brachial artery	7.45		
	Jugular vein	7.35	0.1	0.05
	Mammary vein	7.43		
12655	Brachial artery	7.54		
	Jugular vein	7.47	0.07	0.1
	Mammary vein	7.44		
12666	Brachial artery	7.50		
	Jugular vein	7.44	0.06	0.06
	Mammary vein	7.44		
12667	Brachial artery	7.50		
	Jugular vein	7.44	0.06	0.02
	Mammary vein	7.43		
		Mean	0.07	0.06

It will be observed that there is an arterio-venous difference in plasma pH between the brachial artery and both the mammary and jugular veins. The pH of plasma from the mammary vein was consistently different from the pH of the plasma from the jugular vein.

Variations have also been observed in the plasma carbon dioxide

content and the plasma pH of the healthy human subject throughout the course of a day which were unconnected to any specific activity (Cullen and Earle, 1929). A study was made of the variations of plasma pH and plasma carbon dioxide occurring in seven healthy adult cattle during the course of a day or part of the day when they were standing in the byre. Those animals were given their morning feed of cattle coke at 7 a.m. and thereafter had hay and water ad libitum through the day. They were noted to eat, drink, ruminate, defaecate and urinate and behaved in every way as normal cattle, throughout the period of study.

Blood samples were withdrawn at intervals during the day from the same point in the earo vessel.

The results obtained for those cows which have been numbered 1 to 7 are given in tables 29 to 32. Graphic representation of the results obtained from cows 1 and 2 are given in figures 23 and 24.

Table 22

Variations in the CO<sub>2</sub> content and the pH of the plasma  
obtained from lymphokinetic arterial blood of monkey

Time of sampling	Plasma pH	Plasma CO <sub>2</sub> content u.moles/litre
9.15	7.40	27.6
10.26	7.36	27.6
11.15	7.42	27.0
12.25	7.36	27.2
13.35	7.42	27.6
14.45	7.41	27.6
15.35	7.41	27.1
16.36	7.39	27.0

The variations observed in the plasma carbon dioxide content were small, being less than 1 u.mole/litre. The variations in plasma pH were likewise small. No correlation was apparent over the day between changes in carbon dioxide content and pH.




Fig. 23

Graphic representation of the results in table 29

Table 30

Variations in the CO<sub>2</sub> Content and the pH of the Plasma Obtained from Jugular Venous Blood of Cow no. 2 During the Course of an Afternoon

Time of Sampling	Plasma pH	Plasma CO <sub>2</sub> content m.moles/litre
14.00	7.43	29.5
14.35	7.42	32.5
15.10	7.41	29.5
15.45	7.32	30.3
16.00	7.31	29.9
16.10	7.36	28.8




Fig. 24

Graphic representation of the results in table 30

This cow which was sampled six times in just over two hours showed much more marked changes in both the pH and the carbon dioxide content of the plasma.

Table 27  
 Variations in the pH and the  $\text{CO}_2$  Content of Plasma  
 Obtained from Bronchial Arterial Blood of Cow No. 3  
 During the Course of an Afternoon

Time of sampling	Plasma pH	Plasma $\text{CO}_2$ content m.mole/liter
13.00	7.42	26.6
14.00	7.53	26.3
15.00	7.53	27.9
16.00	7.51	26.5

The variations in the plasma carbon dioxide content were small but there was a greater change in plasma pH.

Table 32  
 Variations in the CO<sub>2</sub> Content and the pH of Plasma  
 Derived from Bovine Arterial Blood of Cow no. 4  
 Taken at Short Intervals Over 60 Minutes

Time of sampling	Plasma pH	Plasma CO <sub>2</sub> content m.moles/liter
13.35	7.42	27.3
13.32	7.42	27.3
13.35	7.42	27.9
13.37	7.42	27.9
13.40	7.42	27.9
14.45	7.42	30.0

Six arterial samples were taken from this cow in 60 minutes during which time no change in plasma pH and only very small changes in plasma carbon dioxide content occurred except at the last sample where a rise of 2 m.moles/liter took place unconnected with a change in plasma pH.

Table 22

Variations in the  $\text{CO}_2$  Content and the pH of Plasma

Obtained from Jugular Venous Blood of Seven Cows, 5, 6 and 7

Time of sampling	Cow 5		Cow 6		Cow 7	
	Plasma pH	Plasma $\text{CO}_2$ n.moles/ litre	Plasma pH	Plasma $\text{CO}_2$ n.moles/ litre	Plasma pH	Plasma $\text{CO}_2$ n.moles/ litre
21.35	7.40	23.6	7.39	23.3	7.38	23.5
12.30	7.40	23.6	7.41	23.3	7.40	23.6
13.45	7.39	23	7.43	23.3	7.40	23.0
15.35	7.37	23.6	7.42	23.5	7.40	23.6
16.30	7.34	23.0			7.34	23.1

These three cows which were studied at the same time showed slight variations in both the carbon dioxide content and the pH of their plasma.

Studies on these seven cows indicated that changes in the pH and carbon dioxide content of the plasma which occurred during the day interconnected with very activity, were small and within the normal values already determined for adult lymphatic cattle.

### (5) The ventilation and respiratory rates of Ayvalik cattle

The main purpose of this study was to determine the normal values for the plasma carbon dioxide and pleural pH of Ayvalik cattle and also to determine the changes associated with disturbances of respiration.

To study these disturbances of respiration it was necessary to measure both respiration rates and pulmonary ventilation. The methods of so doing have already been described and it is necessary only to state that each measurement in table 24 represents the mean of at least five and often 10 measurements taken during a period of 25 to 30 minutes.

The respiratory minute volumes in table 24 were corrected to B.T.P.S. It was found however that when this was done, due to the fact that the temperature of the two rooms in which these measurements were made was relatively constant between 13 and 15° C., although atmospheric pressure varied from 747 - 770 mm.Hg., the conversion factor to correct to B.T.P.S. volumes was found to be a constant 1.112. This factor can be used to convert any other measured ventilation in these cattle to B.T.P.S.

The results show that the respiratory rate does not increase with body weight but the pulmonary ventilation does and as a consequence so must the tidal volume. The relationship between body weight and pulmonary ventilation is not linear according to Brody (1950) but logarithmic. Our results are in agreement with those of Brody.

Table 24

Respiratory Rate, Ventilation Rate, and the Tidal Air

of Normal Cattle Assumed According to Body Weight

Body weight (kg.)	Ventilation rate lit/sec/min. (B.T.P.S.)	Respiratory rate /minute	Tidal air lit.
27	7	16	270
33	6	19	324
39	7	24	276
40	7	22	265
36	9	20	327
37	7	20	246
40	16	31	502
45	24	35	606
50	20	30	963
52	23	32	720
52	27	30	690
55	31	30	1097
123	46	22	2973
255	72	20	3550
364	79	25	3025
376	86	22	3092
386	67	32	3739
432	110	24	4692
445	122	30	5705
450	100	24	4290

## DISCUSSION

The development of a method of obtaining arterial blood from the brachial artery of unanaesthetised cattle of all ages has made it possible for the determinations to be made of the carbon dioxide content of arterial plasma of cattle. It is interesting to note that the mean value obtained for the arterial plasma of adult cattle was identical to that given by Davenport (1950) for the arterial plasma carbon dioxide of the human subject but both cattle and humans have a significantly higher arterial plasma carbon dioxide than does the dog.

The mean values obtained in this study for carbon dioxide content of venous plasma of both cows and calves are similar to the values given by the majority of other workers whose results are given in the review of the literature at the beginning of this section.

The mean value for the carbon dioxide content of the arterial plasma of dogs found in this study was almost identical to the value given by Spector (1953).

Similarly the mean plasma pH values found in this study were identical with the venous and arterial plasma pH values given by Davenport (1950) for the human subject and the venous plasma pH values were similar to those found by other workers.

Variations in the carbon dioxide content and the pH of the plasma of normal cattle occurring during the course of a day were similar to changes observed by Cullen and Basile (1929) in the human subject.

The observations made in this study of the respiratory rate of cattle of all ages agreed with those of all the other workers with the exception of Findley (1956) whose control values given for the respiratory rates of calves indicated the presence of excitement, thermal stress or painende in his calves.

The values found for the ventilation rate of cattle were in agreement with the far more extensive study of Brody (1950) and with results of others where a comparison on a body weight basis could be made.

## CONCLUSIONS

From the various estimations made in Part I of this study certain conclusions were drawn:

- (1) The following precautions were necessary in the collection and handling of blood samples.
  - (a) When duplicate blood samples were taken and single estimations were carried out on these samples, the variations obtained by this method were not more than 0.3 n.moles/litre in the carbon dioxide content and not more than 0.03 pH units in the plasma pH.
  - (b) Unless the collection of blood was made under a layer of liquid paraffin, carbon dioxide loss to the atmosphere caused errors no greater than 1.2 n.moles/litre in the carbon dioxide content and 0.05 pH units in the plasma pH.
  - (c) Unless this layer of liquid paraffin left in contact with the blood was kept to a minimum, errors arose due to the absorption of carbon dioxide by the liquid paraffin of as much as 1.25 n.moles/litre carbon dioxide content and 0.05 pH units when this layer in a 50 ml. centrifuge tube was two inches deep.
  - (d) Unless determinations were carried out on plasma samples within

half an hour of collection, autolysis caused a lowering of plasma pH of as much as 0.2 pH units in one hour.

- (2) Both the Conway method and the Manometric method of Van Slyke were found to be suitable for the determination of the carbon dioxide content of plasma, the Conway method being the more suitable for large numbers of determinations carried out in a short time. Using these methods the following results were obtained.
- (a) The mean value for the arterial plasma carbon dioxide content of 159 adult Ayrshire cows was found to be  $27.2 \pm 2.7$  m.moles/litre. No significant differences were found due to seasonal variations, lactational status or pregnancy.
  - (b) The mean values for the carbon dioxide content of venous and mixed venous blood were found to be  $27.4 \pm 2.9$  m.moles/litre and  $29.2 \pm 2.7$  m.moles/litre, respectively.
  - (c) It was found that a mean difference of  $1.7$  m.moles/litre existed in the carbon dioxide content of jugular and coronary venous plasma of five cows, the coronary venous plasma carbon dioxide contents being greater.
  - (d) It was found that the mean value of arterial plasma carbon dioxide in 81 calves was  $29.8 \pm 2.6$  m.moles/litre, a figure significantly lower than the mean for adult Ayrshire cows.
- (3) The colorimetric method of determination of plasma pH using phenol

and on the indicator was examined and shown to be reliable and repeatable.

- (a) Using this method, temperature coefficients were found for bovine plasma of 0.009 and for bovine blood of 0.012 pH units per degree centigrade.
- (b) Using these temperature coefficients and correcting all results to 30° C., mean values of  $7.43 \pm 0.045$  pH units for the arterial plasma pH of 269 Friesian cows, of  $7.39 \pm 0.05$  pH units for the venous plasma pH of 55 Ayrshire cows, and of  $7.42 \pm 0.036$  pH units for the arterial plasma pH of 10 calves were found.
- (c) Temperature coefficients of 0.0155 pH units and 0.012 pH units were found for cow's blood and plasma, respectively.
- (d) Using these temperature coefficients and correcting all results to 30° C., a mean value of  $7.43 \pm 0.045$  pH units was found for the arterial plasma pH of 10 dogs.
- (e) No significant variations in the values of plasma carbon dioxide content and plasma pH were observed when seven cows were sampled repeatedly during periods of up to six hours.
- (f) It was shown that cattle of all ages had respiratory rates between 30 and 35 per minute and that the ventilation rate varied with the body weight, not in a linear manner but logarithmically so that a calf of body weight of 90 kg. had a ventilation rate of **25** Litres per minute while a cow of 500 kg. body weight had a ventilation rate of about 100 Litres per minute.

PART II

THE CHANGES IN THE CARBON DIOXIDE CONTENT AND THE PH OF THE PLASMA

OF CATS CAUSED BY DISTURBANCES OF RESPIRATION

## INTRODUCTION

Disturbances of respiration may be induced in a number of ways. It is possible to alter the central respiratory drive by factors affecting the respiratory centre either directly or reflexly as the result of stimuli elsewhere. It is possible also to produce changes in respiration by quantitative alterations in the blood, carbon dioxide or H ion concentration.

Little is known of the central respiratory mechanism of cattle or of the factors affecting it. It is assumed that cattle are the same as all other species studied with a respiratory centre which may be affected by other centres in the brain, by changes in its blood chemistry, by impulses arising from peripheral stimuli such as pain or limb movement and by impulses from chemoreceptors which respond to changes in blood chemistry.

In the present study certain disturbances of respiration of diverse origin were investigated in order to ascertain the changes in plasma pH and plasma carbon dioxide content as a result of these disturbances. Ventilation and respiratory rates have also been measured to define these disturbances of respiration.

The disturbances of respiration on which studies were made were

chosen because of availability of material. An intensive investigation had been started on the suitability of volatile and gaseous anaesthetics for cattle (Richter and Jennings, 1956(a), 1956(b)). This investigation was extended to determine the effects of certain anaesthetic agents, in particular Fluothane®, on the plasma pH and the plasma carbon dioxide content of cattle together with the changes in the ventilation and respiratory rates.

The studies of Jarrett, Jennings, McEntee, Malligan and Dugdale (1957) on the parasite *Dityosomum viviparum* made available many calves in different stages of parasitic pneumonia. Investigations were therefore made of the changes in plasma pH and the plasma carbon dioxide content of calves affected with parasitic pneumonia. Measurements were made at the same time of the ventilation and respiratory rates of calves with this disease.

In an attempt to explain the response of pneumonia and anaesthetised cattle, studies were carried out of the effects on respiration of gas mixtures different in composition to the atmosphere.

Since these three types of disturbances of respiration are widely different they are each presented in a separate section in this part of the thesis.

## SECTION X.

### THE EFFECTS OF SOME GENERAL ANAESTHETIC AGENTS, XYLOTOLONE IN PARTICULAR, ON THE PLASMA pH AND THE PLASMA CARBON DIOXIDE CONTENT OF CATTLE.

#### REVIEW OF THE LITERATURE

General anaesthetic agents, because of their effects on the central nervous system have been shown to cause changes in respiration with changes in plasma pH and plasma carbon dioxide content of man and many other animals. A very extensive literature has been accumulated on the clinical effects of general anaesthetics operating in man, the advent of each new anaesthetic agent leading to the production of numerous articles. For example, between the introduction of Fluothane in 1956 and the middle of 1958 at least 30 articles appeared on its clinical use in man. Fewer reports have appeared on the biochemical effects of general anaesthetic agents on the human subject but the amount of published material is still extensive. Similarly a voluminous literature has been accumulated on the effects of general anaesthetic agents on dogs,

cats and monkeys when these animals were used as experimental subjects.

On the other hand the effects of general anaesthetics on the respiration of cattle have not been studied very fully. Until recently volatile and gaseous anaesthetics have been considered unsuitable for cattle because of the copious salivation and outpouring of bronchial secretions with consequent risk of post-anaesthetic pneumonia (Wright, 1952). Ether was stated to be ineffective as a general anaesthetic for large animals (B. Vet. Codex, 1953) although Hendrill (1955) described its use as a general anaesthetic for horses. However, the successful use of volatile and gaseous anaesthetics in cattle has been reported by Hall (1957), Harrison and Johansson (1958), and Fluhler and Jennings (1960).

Hall (1957) in a clinical evaluation of the effects of Fluothane on four cows stated that respiratory depression occurred together with a slowing of the pulse rate. Harrison and Johansson (1958) reported the successful use of a mixture of chloropentoxide sodium, muscimylecholine and nitrous oxide as a general anaesthetic for cattle. They observed slight changes in the partial pressure of carbon dioxide in the blood and slight changes in the blood pH. Some of the changes they reported may not have been of respiratory origin since on occasion a rise in the partial pressure of carbon dioxide occurred with a rise in the plasma pH. As they also assisted ventilation by means of a pump it was not possible to draw conclusions about the effects of the anaesthetic mixture used on

respiration. In another experiment they reported a slight decrease in pulmonary ventilation together with a decrease in carbon dioxide production and a slight decrease in oxygen consumption when thiopentone sodium alone was used as the anaesthetic agent. Height (1952) reported an increase of pulse rate and respiratory rate during maintenance of general anaesthesia in cattle with chloral hydrate. Henderson (1944) has reported that the use of thiopentone sodium in young cattle caused a decrease in respiratory rate and an acceleration of the pulse. Ruxy (1950) reported favourably on the use of Anervon<sup>16</sup> in cattle but Height (1957) reported tachypnoea and dyspnoea in similar circumstances.

## METHODS

### (1) Fluothane

The closed circuit method of administration of Fluothane was used. A circle absorber apparatus was constructed of a size considered suitable for adult cattle and horses. The airways were of one and a half inch internal diameter, and an eight gallon reservoir bag was used. The outer cylinder of the soda-line absorber was 10 inches in diameter and 10 inches in height while the inner cylinder holding the soda-line was eight inches in diameter and eight inches in height and had a capacity of six litres.

This apparatus is illustrated in figures 25 and 26.




Fig. 25

Photograph of circle absorber




Fig. 26  
Diagram of circle absorber

Animals were not starved before anaesthetisation and they were sometimes anaesthetised shortly after normal feeding. No pre-anaesthetic drugs were used. A catheter was introduced into the jugular vein about one hour before anaesthetisation to enable blood samples to be withdrawn easily from the site before, during and after an operation. Just before an operation, the ventilation rate was

was recorded for a period of 10 minutes using the face mask and gas meter with the unidirectional valves previously described. At the end of this period a blood sample was taken, the plasma was separated and the pH and carbon dioxide content determined. On some occasions the subject was made to breathe pure oxygen for a further period of 10 minutes during which time ventilation was recorded and afterwards blood samples were again taken, the plasma separated and the pH and carbon dioxide content of the plasma determined.

Induction of anaesthesia was commenced in every case with the animal in the standing position. As anaesthesia advanced the animals were either allowed to sink to the ground or, when a mechanically operated table was used, the table was brought from the vertical to the horizontal position so that the animal was pulled into lateral recumbency. Two methods of induction were used. In the first, anaesthesia was induced by means of a 10% solution of thiopentone sodium in normal saline. When thiopentone sodium was used for the induction of anaesthesia it was not usual to record the ventilation rate during induction. In the second method Fluothane was used as the inducing agent and in some of these cases ventilation was recorded during induction by including the gas meter in the anaesthetic circuit. When Fluothane was used for induction, the close fitting face mask was applied to the animal and the circle absorber was connected to the face

neck. Fluothane was administered to the animal from a Trilene<sup>®</sup> vapouriser bottle by passage of oxygen through the Fluothane. Once general anaesthesia had been induced a cuffed endotracheal tube was introduced into the trachea. The circle absorber was connected to the endotracheal tube and anaesthesia was continued. Ventilation tubes were measured during anaesthesia by enclosing the gas meter in the circuit between the respiratory valve and the circle absorber. In those experiments where a study was made of changes in plasma pH and plasma carbon dioxide, a blood sample was taken for analysis as soon as possible after induction.

Once surgical anaesthesia had been attained after connection to the circle absorber, the oxygen input was allowed to bypass the Fluothane in the Trilene vapouriser but increments of Fluothane were added at intervals to make up for losses due to leakage from the circuit. During the studies of Fluothane on ventilation and plasma carbon dioxide content and plasma pH, ventilations were recorded continuously and blood samples were taken at half-hourly intervals for analysis of the plasma pH and plasma carbon dioxide content.

After the period of anaesthesia the anaesthetic apparatus was disconnected but the endotracheal tube was left in situ until a stage of recovery was reached where the subject would no longer tolerate it.

n Superfine Chemical (Pharmaceutical) Ltd., Winslow, Cheshire.

Ventilation rates were recorded to this point and only occasionally afterwards. Blood samples were usually taken until the plasma pH had returned to the control value.

In the comparative studies on Fluothane anaesthesia carried out in dogs, sheep and horses, the same general procedure was adopted. For the studies in dogs and sheep the same anaesthetic apparatus was employed but the large unidirectional valves were replaced by the three-quarter inch diameter unidirectional valves described in the section on the measurement of the ventilation rate of calves, and the tubes connecting these valves to the animal were also of three-quarter inch internal diameter. Control ventilations were not usually attempted in sheep or horses as both species became excited when they were fitted with a mask. For this reason it was possible to carry out induction of the horse with Fluothane on only one occasion.

### (2) Thiopentone sodium

Thiopentone sodium was administered as a 10% solution in normal saline. On one occasion the calculated dose was injected rapidly into the jugular vein and changes in plasma pH and plasma carbon dioxide content followed. In two calves induction was carried out more slowly until the stage of surgical anaesthesia was reached. Further increments of thiopentone sodium were added as required to maintain the animal in this plane of anaesthesia.

### (3) Chloral hydrate

Chloral hydrate was administered to one cow and the changes in ventilation, plasma pH and plasma carbon dioxide content followed.

A catheter was introduced into the jugular vein and a 10% solution of chloral hydrate administered slowly. Pulmonary ventilation rate was followed before and during the administration of the chloral hydrate and during the recovery period. The cow was maintained at the stage of etyonal anesthesia for approximately one hour by the addition of small increments of chloral hydrate intravenously. Blood samples were taken at approximately half hour intervals and the plasma pH and plasma carbon dioxide content determined.

## RESULTS

The results of the studies in this section are arranged in the following order.

- (1) Fluothane      (a) Cattle
- (b) Horse
- (c) Dog
- (d) Sheep
- (2) Thiopentone sodium in cattle.
- (3) Ocular hydrate in ~~one~~ cow.

## (1) Electroshock

### (a) cattle

Experiments were carried out on eight cattle. In two of these experiments the effects of Electroshock on the ventilation rate, the plasma pH and the plasma carbon dioxide content were followed. In six experiments the effects of breathing pure oxygen for 10 minutes on the ventilation rate, the carbon dioxide content and the plasma pH before the animal was electroshocked were also examined. Each experiment is described separately and each set of results is illustrated by a graph and given in a table.

Experiment I. Cow 850. In this experiment pre-anaesthesia control values of ventilation rate, carbon dioxide content and pH of plasma were determined and similar determinations were made over the period of anaesthesia. An hour after the cow had been removed from the anaesthetic circuit, a final blood sample was taken. The changes taking place are demonstrated in figure 27, and summarised in table 35.




Fig. 27

Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during Fluothane anaesthesia.

Table 25  
 Changes in Ventilation Rate, Plasma Carbon Dioxide Content  
 and Plasma pH During Ether-Anesthesia

Time	Before anesthe- sia	During Anesthesia (minutes)					During Recovery (minutes)		
		5	30	60	90	120	5	30	60
pH <sup>23</sup>	7.35	7.19	7.39	7.38	7.35	7.35			7.30
CO <sub>2</sub>	27.5	27.9	20.7	29.5	29.9	30.5			20.5
Vent.	93	44	47	42	54	53	76		

From these results it will be observed that depression of pulmonary ventilation occurred and that at the same time there was a fall in plasma pH and a rise in plasma carbon dioxide. The changes in pH and carbon dioxide were greater than those observed when following hourly variations in the normal unanesthetized cow. There was evidence of some carbon dioxide retention and a lowering of plasma pH as long as one hour after removal from the anesthetic atmosphere.

2. Throughout ideals pH = Plasma pH

CO<sub>2</sub> = Plasma carbon dioxide content in micro/mole/100 ml

Vent. = Ventilation ratio in Litres/minute.

Experiment II. Bull 400. This experiment is similar to the previous one. The changes taking place during the whole period of anaesthesia and recovery are illustrated in figure 28, and summarised in table 36.




Fig. 28

Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during Fluothane anaesthesia

Table 26

Changes in Ventilation Rate, Plasma Carbon Dioxide Content

and Effects of Inhaler Thiodene Anæsthetic

Time	Before anæsthe- tic	During Anæsthetic				During Recovery				
		5	30	60	90	5	30	60	120	180
pH	7.47	7.27	7.31	7.29	7.31			7.42	7.43	7.45
CO <sub>2</sub>	39.2	32.4	32.3	32.1	32.7			39.3	39.1	38.1
Vent	78	34	45	48	48	61				

In this experiment a depression of ventilation again occurred with fall in plasma pH and rise in plasma carbon dioxide content. One hour after anaesthesia the carbon dioxide content had returned to near the pre-anaesthetic value but the plasma pH took a further two hours to return to pre-anaesthetic values.

—ACR—

Experiment III. Bull 400. The changes taking place during anaesthesia and recovery are illustrated in figure 29 and summarised in table 37.




Fig. 29

Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during Fluothane anaesthesia

Table 27

Changes in Ventilation Rate, Plasma Carbon Dioxide Content

and Recovery of Breathing Fluorophore Anesthesia

Time	Before Anesthesia		During Anesthesia (minutes)					During Recovery (minutes)		
	Air	Oxygen	5	30	60	90	120	5	30	120
pH	7.36	7.35	7.25	7.29	7.29	7.30	7.30	7.25	7.25	7.25
CO <sub>2</sub>	39.6	31.9	32.5	33.4	35.6	35.9	35.9	39.5	39.75	39.75
Vent.	76	55	40	52	53	54.5	57	56		

When the animal was unanesthetized and breathing pure oxygen there was some reduction of pulmonary ventilation and a rise in plasma carbon dioxide content but the plasma pH was not significantly different. During anesthesia a depression of ventilation occurred but throughout the main part of anesthesia ventilation was not markedly different from that observed when the animal breathed pure oxygen. There was the expected carbon dioxide retention with a fall in pH and a rise in plasma carbon dioxide content above values observed in normal animals. Two hours after recovery from anesthesia the plasma pH and plasma carbon dioxide content were at their pre-anesthetic control values.

Experiment IV. Cow 900. This experiment was similar to the previous one. The changes which took place during the period of anaesthesia and recovery are illustrated in figure 30 and summarised in table 38.




Fig. 30

Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during Fluothane anaesthesia

Table 21

Change in Ventilation Rate, Mean Arterial Blood Content

and Plasma pH During Anesthesia

Time	Before Anesthesia			During Anesthesia (minutes)				During Recovery (minutes)		
	Alt	Oxygen		5	30	60	90	220	5	30
pH	7.45	7.45	7.37	7.38	7.38	7.29	7.29	7.30	7.44	7.44
CO <sub>2</sub>	26.8	23	23.8	23.8	23.8	23.8	23.8	23	32.5	32.5
Vent	71	49	38	26	29	30	37	50	69	69

In this animal there was pre-anesthetic reduction of ventilation with pure oxygen and during anesthesia a depression of ventilation occurred. The plasma pH and plasma carbon dioxide content both rose during inhalation of pure oxygen. During the anesthetic period retention of carbon dioxide occurred with a fall in plasma pH. Plasma pH had returned to the pre-anesthetic control value but there was still some retention of carbon dioxide 30 minutes after the administration of anesthetic had ceased.

Experiment V. Heifer 822. In this experiment which was similar to experiments III and IV the endotracheal tube was ejected in the middle of the experiment. The changes taking place during the period of anaesthesia and recovery are illustrated in fig. 31 and summarised in table 39.




Fig. 31

Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during Fluothane anaesthesia.

Table 22

Changes in Ventilation Rate, Plasma Oxygen Dioxide Content

and Plasma pH During Fluothyme Anesthesia

Time	Before Anesthesia		During Anesthesia (minutes)				During Recovery (minutes)		
	Mr	Oxygen	5	30	60	90	5	15	60
pH	7.38	7.38	7.29	7.23	7.24	7.23	7.39	7.39	7.39
CO <sub>2</sub>	26.2	26.7	31.2	31.6	32.8	33.6	27.5	28.2	28.2
Vent.	70	69	30	59	40.6	52	91	90	60

This animal also showed some reduction of ventilation during the administration of pure oxygen but struggled violently so that the reduction was less marked. The plasma carbon dioxide content gave no evidence of rebreathing during the administration of pure oxygen. During anesthesia depression of ventilation occurred with carbon dioxide ventilation and a fall in plasma pH. During recovery some hyperventilation occurred. At the end of an hour the plasma pH and plasma carbon dioxide content had returned to their pre-anesthetic control values.

Experiment VI. Cow 940. This experiment was similar to the previous one. The changes which took place during the period of anaesthesia and recovery are illustrated in fig. 32 and summarised in table 40.




Fig. 32

Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during Fluothane anaesthesia.

Table 40

Changes in Ventilation Rate, Plasma Carbon Dioxide Content

and Plasma pH During Fluothane Anesthesia

Time	Before Anesthesia Air Oxygen	During Anesthesia (minutes)					During Recovery (minutes)		
		5	30	60	90	220	5	15	60
pH	7.38 7.36	7.20	7.11	7.10	7.09	7.19	7.25	7.28	
CO <sub>2</sub>	36.5 35.25	30.9	31.9	29	26	32.9	32.5	30.1	
Vent.	99 65	39	36	29	26	35.6	50		

Inhalation of pure oxygen caused a **decrease** in ventilation but no marked changes were detected in the plasma pH and the plasma carbon dioxide content. A **depression** of ventilation occurred during Fluothane anesthesia with carbon dioxide retention and a fall in plasma pH. The plasma pH returned to the control value one hour after the car had been removed from the Fluothane atmosphere but there was still some carbon dioxide retention.

Experiment VII. Calf 272. This, too, was similar to the previous experiments. The changes which took place during the period of anaesthesia and recovery are illustrated in figure 33 and summarised in table 41.




Fig. 33

Changes in ventilation rate, plasma carbon dioxide content,  
and plasma pH during Fluothane anaesthesia

Table A2

Changes in Ventilation Rate, Plasma Carbon Dioxide Content,

and Plasma pH During Fluothane Anesthesia

Time	Before Anesthesia		During Anesthesia (minute)				During Recovery (minute)	
	Air	Oxygen	5	30	60	90	5	30
pH	7.30	7.30	7.21	7.18	7.22	7.24	7.24	7.30
CO <sub>2</sub>	29	29	25	34	32	22	21.6	24
Vent	42.5	31.0	29	29	35	26	31	26

In this calf similar changes to those in the adult cows took place. There was pre-anesthetic **lowering** of pulmonary ventilation during inhalation of pure oxygen but changes were not detected in the plasma pH and the plasma carbon dioxide content. **An depression** of ventilation occurred during Fluothane anaesthesia and although this was marked immediately after intubation, over the whole period of anaesthesia the depression was not great. Changes of plasma carbon dioxide and the changes of plasma pH reflected the ventilation changes. During recovery the plasma pH returned to the control value after half an hour although ventilation was still depressed and there was still some carbon dioxide retention.

Experiment VIII. Calf 270. This experiment was similar to the previous experiments. The changes which took place during the period of anaesthesia and recovery are illustrated in figure 34 and summarised in table 42.




Fig. 34

Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during Fluothane anaesthesia.

Table A2

Changes in Ventilation Rate, Plasma Carbon Dioxide Content  
and Plasma pH During Fluothane Anesthesia

Time	Before Anesthesia			During Anesthesia (minutes)					During Recovery (minutes)		
	Air	Oxygen		5	30	60	90	220	5	30	45
pH	7.30	7.31		7.32	7.35	7.34	7.34	7.35	7.36	7.39	7.32
CO <sub>2</sub>	33	33.0		32.0	32.2	32.6	32.7		32.2	32.4	32.2
Vent	35	34		39	16	18	27	17	22		

This calf showed only slight pre-anesthetic hypventilation, implying pure oxygen and a slight rise in plasma carbon dioxide content. During anesthesia a marked depression of ventilation occurred with carbon dioxide retention and a fall in plasma pH. During recovery, after 30 minutes, pH had returned to near the control value but after 45 minutes there was still evidence of carbon dioxide retention.

-2,027-

(b) Normal

Experiments were carried out on three horses. In the first experiment induction was by means of Fluothane while in the other two experiments thiopentone sodium was the inducing agent. In one experiment only was it possible to follow ventilation throughout. In all three experiments blood samples were taken and checked in the carbon dioxide content and the pH of the plasma followed.

Experiment I.II. Hunter 1006. Induction of anaesthesia in this animal was by means of Fluothane. The results obtained during the period of anaesthesia and recovery are illustrated in figure 35 and summarised in table 43.




Fig. 35  
Changes in plasma carbon dioxide content and plasma pH  
during Fluothane anaesthesia

Table A3

Changes in Plasma carbon Dioxide Content and Plasma pH

During Anesthesia

Time	Hofers Intraop- ticain	During Anesthesia (minutes)				During Recovery (minutes)		
		5	30	50	70	5	30	60
pH	7.38		7.33	7.33	7.35			7.38
CO <sub>2</sub>	39.5		30.75	32.5	30.75			39.5

During anesthesia the rise in plasma carbon dioxide and fall in plasma pH indicated a depression of ventilation. One hour after the animal had been removed from the anesthetic atmosphere, plasma pH and plasma carbon dioxide had returned to the control values.

Experiment II H. Horse 1071. Induction and intubation were carried out using thiopentone sodium as the anaesthetic agent. Ventilation was recorded after induction. The results obtained during the period of anaesthesia and recovery are illustrated in figure 36 and summarised in table 44.




Fig. 36

Changes in ventilation rate, plasma carbon dioxide content and plasma pH during Fluothane anaesthesia.

Table 14

Changes in Ventilation Rate, Plasma Carbon Dioxide Content

and Plasma pH During Nitrous Oxide Anesthesia

Time	Before Anesthesia	During Anesthesia (minutes)				During Recovery (minutes)		
		5	30	50	70	5	30	60
pH	7.40	7.31	7.30			7.40		
CO <sub>2</sub>	23.26	24.9	24.9			23.12		
Vent.		30	27			57		

Depression of ventilation occurred but was not so marked as it appears from the graph since under thiopental and after intubation no respiratory stimulation occurred. The changes in plasma carbon dioxide content and plasma pH were small since the period of anesthesia was short. Five minutes after removal from the anesthetic atmosphere the plasma pH and plasma carbon dioxide content had returned to the pre-anesthetic control values.

-144-

Experiment III H. Pony 792. It was possible in this animal to follow ventilation rate in addition. The results obtained during the period of anaesthesia and recovery are illustrated in figure 37 and summarised in table 45.




Fig. 37  
Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during Fluothane anaesthesia

**Table A5**  
**Changes in Ventilation Rate, Plasma Carbon Dioxide Content  
 and Plasma pH During Fluothane Anesthesia**

Time	Before Anes- thesia	During Anesthesia (minutes)				During Recovery (minutes)		
		5	30	60	70	5	30	60
pH	7.37	7.19	7.23			7.36		7.37
CO <sub>2</sub>	39.23	25	28.5			28.5		29.23
Vent.	40.5	40	39			45		74

Following induction and intubation with thiopentone sodium some respiratory stimulation occurred. The plasma carbon dioxide at this point indicated hyperventilation but the plasma pH had fallen and may have indicated some degree of a metabolic acidosis as reported by Banbox, Beechey, Briggs, Baumann and Barnes (1951) during ether anesthesia in dogs. Beyond this point some respiratory depression occurred with a rise in plasma carbon dioxide content. During the recovery period, after five minutes the pH and carbon dioxide contents were nearly back at the control values and after one hour they had returned to the control value. One hour after anesthesia when the ventilation rate was measured, some hyperventilation was observed.

~~10/17/68~~  
**(c) 1923**

Experiments were conducted on three greyhounds. In each experiment it was possible to follow ventilation before, during and after anaesthesia. Blood samples were taken and the changes in the carbon dioxide content and the pH of plasma followed throughout each experiment.

Experiment I.D. Greyhound 71. Induction of this animal was by means of Fluothane. The results obtained during the period of anaesthesia and recovery are illustrated in figure 38 and summarised in table 46.




Fig. 38

Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during Fluothane anaesthesia

Table 46

Changes in Ventilation Rate, Plasma Carbon Dioxide Content

and Plasma pH During Fluothane Anesthesia

Time	Before Anes-	thetic	During Anesthesia				During Recovery		
			5	20	60	90	5	25	55
PaCO <sub>2</sub>	7.40	7.36	7.36		7.38		7.39		
O <sub>2</sub>	29	30.3	30.5		30.3		30.3		
Vent.	7.6	4.6	5.0		5.0		12.5		

The passage of the endotracheal tube after Fluothane induction caused a marked diminution of respiration which was not recorded. Once the respiratory stimulation had ceased the ventilation rate was again recorded. A depression of ventilation occurred during Fluothane anesthesia but marked changes were not detected in plasma pH and plasma carbon dioxide content. During the recovery from Fluothane concentrations considerable hyperventilation occurred.

Experiment II D. Greyhound 59. This dog also received Fluothane for induction. The results obtained during the period of anaesthesia and recovery are illustrated in figure 39 and summarised in table 47.




Fig. 39

Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during Fluothane anaesthesia

Table A2

Changes in Ventilation Rate, Plasma Carbon Dioxide Content  
and Plasma pH During Fluothane Anesthesia

Time	Before Anesthe- ticide	During Anesthesia (minutes)				During Recovery (minutes)		
		5	30	60	90	5	25	53
pH	7.40	7.40	7.39	7.38	7.41	7.40	7.40	7.40
CO <sub>2</sub>	33	27	27.2	28	28.2	26.5	26.2	
Vent	7.7	12	6	5	6	11	27	

As in guinea-pig 71, intubation caused stimulation of respiration and hyperventilation. Following this hyperventilatory respiratory depression occurred. There were very small changes of plasma pH during the experiment. The plasma carbon dioxide content fell as a consequence of the hyperventilation and then gradually rose again. During the recovery period the plasma pH reached normality but the plasma carbon dioxide content fell as the animal was hyperventilating above the determined pre-anesthetic control ventilation.

-175-

Experiment III D. Greyhound 70. Induction was by means of Fluothane. The results obtained during the period of anaesthesia and recovery are illustrated in figure 40 and summarised in table 48.




Fig. 40  
Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during Fluothane anaesthesia

Table A3

Changes in Ventilation Rate, Plasma Carbon Dioxide Content  
and Plasma pH During Fluothane Anesthesia

Time	Before Anes- thesia	During Anesthesia (minutes)					During Recovery (minutes)		
		5	20	60	90	5	25	95	
pH	7.45	7.51	7.41	7.39	7.42		7.40		
CO <sub>2</sub>	26.7	25.5	27.5	26.8	27.5		26.5		
Ven's	30.0	22.6	5.0	5.3	10.0		10.5		

Inhalation caused a marked hyperventilation which lowered the plasma carbon dioxide content and caused the plasma pH to rise. Following the hyperventilation respiratory depression took place causing slight carbon dioxide retention and a fall in plasma pH. Twenty-five minutes after the animal had been removed from the Fluothane atmosphere the pulmonary ventilation was similar to the control value but the plasma pH was below the control value while the plasma carbon dioxide content was above the plasma control value.

(d) Sheep

Experiments were carried out on three sheep. Changes in the carbon dioxide content and the pH of plasma were followed in all three experiments but ventilation rate was measured in the first experiment only. Control measurements of ventilation rate in this experiment indicated the effect of exhalation on ventilation rate of the anaesthetized sheep and thereafter this measurement was abandoned in sheep.

Experiment I.S. Sheep 805. Induction of anaesthesia was by means of Fluothane. The results obtained during the period of anaesthesia and recovery are illustrated in figure 41 and summarised in table 49.




Fig. 41.

Changes in ventilation rate, plasma carbon dioxide content and plasma pH during Fluothane anaesthesia.

Table A9

Changes In Ventilation Rate, Plasma Carbon Dioxide Content

and Plasma pH During Fluothane Anesthesia

Time	Before Anes-	During Anesthesia				During Recovery			
		(minutes)	5	30	60	80	20	30	40
pH	7.50	7.00	7.02	7.05			7.42		
CO <sub>2</sub>	25	24	33	34.5			27.5		
Vent	35	1.0	1.5	2.2			4.0		

Marked respiratory depression occurred with striking changes in both the plasma pH and the plasma carbon dioxide content. Once the animal was removed from the Fluothane atmosphere both the plasma pH and the plasma carbon dioxide returned to near the control values.

Experiment II S. Sheep 813. In this animal ventilation was not recorded. Induction was by means of Fluothane. The results obtained during the period of anaesthesia and recovery are illustrated in figure 42 and summarised in table 50.




Fig. 42

Changes in plasma carbon dioxide content and plasma pH  
during Fluothane anaesthesia

Table 50

Changes in Plasma Carbon Dioxide Content and Plasma pH

During Fluothane Anesthesia.

Time	Before Anes- thesia	During Anesthesia (minutes)				During Recovery (minutes)		
		5	30	60	60	20	30	40
pH	7.47	7.10	7.11	7.10	7.10		7.46	
CO <sub>2</sub>	24	26.9	26	23	27		27	

Stabilizing changes both in plasma pH and plasma carbon dioxide content were observed during Fluothane anesthesia. During the recovery the plasma pH returned to the control pre-anesthetic value before the plasma carbon dioxide content.

Experiment III S. Sheep 823. In this animal ventilation was not recorded. Induction was by means of Fluothane. The results obtained during the period of anaesthesia and recovery are illustrated in figure 43 and summarised in table 51.




Fig. 43

Changes in plasma carbon dioxide content and plasma pH  
during Fluothane anaesthesia.

Table 52

## Changes in Plasma Carbon Dioxide Content and Plasma pH

## During Fluothane Anesthesia

Time	Before Anes- thesia	During Anesthesia (minutes)				During Recovery (minutes)		
		5	30	60	30	20	30	60
pH	7.34	7.13	6.93	7.14				7.35
CO <sub>2</sub>	36	33.4	34.3	35.4				33

Marked changes in both plasma pH and plasma carbon dioxide occurred during Fluothane anesthesia but both plasma pH and plasma carbon dioxide returned to the control values 40 minutes after removal from the Fluothane atmosphere.

(2) Thloperitone sodium in catfish

Three experiments were carried out. In the first experiment thloperitone sodium was used as a single dose of 6 gm, while in the other two experiments thloperitone sodium was used not only to induce but also to maintain anaesthesia by the addition of increments of thloperitone sodium intraperitoneally throughout the experiments.

Experiment TB I. This adult cow received an inducing dose of 6 gm. of thiopentone sodium. It collapsed in an anaesthetised state 17 seconds later. The results obtained during the period of anaesthesia and recovery are illustrated in figure 44 and summarised in table 52.




Fig. 44

Changes in plasma carbon dioxide content and plasma pH  
during thiopentone sodium anaesthesia

Table 52

Changes In Plasma Carbon Dioxide Content and Plasma pH  
During Thiomalate Sodium Anesthesia

Time	Anes- thesia	During Anesthesia		During Recovery		
		10	30	15	40	70
pH	7.38	7.30	7.27	7.26	7.20	7.22
CO <sub>2</sub>	33.5	20.5	20.45	23.0	27.5	29.25

The changes in plasma pH and plasma carbon dioxide contents were not marked. It was noted that the period of surgical anesthesia was about 30 minutes, and was followed by a period when the animal was at a higher level of anesthesia when no operative interference would have been possible. After 50 minutes the cow could not be considered anesthetized in any degree but she remained on her side and was not able to get up for a further 40 minutes. The cow was then able to maintain breast recombancy but did not get on to her feet for a further hour.

Experiment 1B II. Four week old calf. Blood samples were removed from the brachial artery and changes in blood carbon dioxide content were followed. Anaesthesia was maintained with increments of thiopentone sodium. The results obtained during the initial period of anaesthesia are illustrated in figure 45 and given in table 53.




Fig. 45  
Changes in blood carbon dioxide and pH  
during thiopentone anaesthesia

Table 52

## Changes in Blood Carbon Dioxide and pH

## During Nitroprusside Anesthesia

Time	Before Anesthe- sia	During Anesthesia (minutes)			
		30	60	90	120
pH	7.32	7.19	7.21	7.21	7.26
Blood CO <sub>2</sub>	23	23	23	23	23

Retention of carbon dioxide occurred and there was the usual fall in blood pH. The cat did not recover from anesthesia for a period of 15 hours.

Experiment TB III. Six week old calf. Blood samples were removed from the jugular vein by means of a catheter. The results obtained during anaesthesia are given in figure 46 and summarised in table 54.




Fig. 46  
Changes in ventilation and respiratory rates,  
plasma carbon dioxide content and plasma pH  
during thiopentone anaesthesia.

Table 5A

Changes in Ventilation and Respiratory Rate,Plasma Carbon Dioxide Content, and Plasma pHDuring Nitrogenous Anesthesia

Time	Before Anes-	During Anesthesia (minutes)				
		10	40	50	60	80
Roop	32	23	25	43	23	30
Vent	11	4.6	6	7.2	6.4	5.8
pH	7.33	7.26	7.23	7.26	7.23	7.26
$\text{CO}_2$	39.125	39.25	39.375	39.625	39.375	39.125

A marked depression of ventilation occurred but the changes in the plasma carbon dioxide content were not great. The calf was still anesthetized at the end of the experiment and remained so for a further 12 hours.

\* Throughout this Roop = Respirations per minute.

### (3) Chloral hydrate in cattle

One experiment only was carried out the results of which are given in figure 47 and table 55.




Fig. 47

Changes in ventilation rate, plasma carbon dioxide content  
and plasma pH during chloral hydrate anaesthesia

Table 56

Changes in Ventilation Rate, Plasma Carbon Dioxide Content  
and Plasma pH During Chloral Hydrate Anesthesia

Time	Before Anes- thesia	During Anesthesia (minutes)			
		5	30	50	60
Vent.	200	30	21	26	27
pH	7.36	7.23	7.19	7.14	7.16
CO <sub>2</sub>	36	28	23.26	23.25	23.5

Respiratory depression occurred and as a consequence carbon dioxide retention and a fall in plasma pH. After chloral hydrate administration had been discontinued respiration still remained depressed but the plasma carbon dioxide content returned to the control pre-anesthetic level, although the pH was still below the control value.

DISCUSSION  
(Part II section 1)

The changes in pulmonary ventilation, plasma pH and plasma carbon dioxide concentrations of cattle during Fluothane anaesthesia were compared with repeated oxygen in general anaesthesia. There was a greater depression of ventilation than with the other anaesthetics, isoflupredone sodium and chloral hydrate, which were studied. Fluothane was administered together with pure oxygen and it is possible that the increased oxygen tension eliminated oxy-heme-sulfur stimulation of ventilation that might have arisen due to hypoxia caused by central depression of ventilation due to anaesthesia (Cawse, Forster, Dubois, Deluca and Morrison, 1955). It is possible that the cardiovascular effects of Fluothane in producing a fall in cardiac output and a fall in blood pressure (Burn, Nyström, Folger and Patch, 1957) may have produced interference with gaseous exchange by alterations of pulmonary blood flow. It has been shown that if Fluothane is administered with air to the lamb subject cyanosis tends to occur (Burn, Huskin, Oggins and Robertson, 1957). The rapid excretion of Fluothane through the lung once the animal has been removed from the anaesthetic atmosphere ensured a rapid decrease in pulmonary ventilation during recovery. The changes observed in cattle were similar to those reported by

Bovine, Hamilton and Pittenger (1958) in the human subject and qualitatively similar to the effects on the ventilation of horses, sheep and dogs. Accompanying the depression of ventilation there was a rise in venous plasma carbon dioxide content during Fluothane anaesthesia to a value greater than that observed in the venous plasma of normal cattle. In prolonged periods of anaesthesia there was usually stabilisation at a higher concentration of  $\text{CO}_2$  than normal and towards the end of experiments a tendency for the carbon dioxide concentration of the plasma to fall. These observations of stabilisation and tendency for the carbon dioxide content of the plasma to fall may have been explicable in the light of Quantel's (1952) work when he showed that oxidative metabolism was decreased during general anaesthesia in which case there would be a decreased carbon dioxide production. This decreased carbon dioxide production could possibly be handled by the depressed ventilation.

The changes of carbon dioxide content of the plasma during general anaesthesia were qualitatively similar in all species studied but quantitatively much greater in the sheep. Although ventilation was not generally recorded it was observed that many sheep became apnoea during Fluothane anaesthesia which might explain the high values obtained. The increases in the carbon dioxide content of cattle plasma during Fluothane anaesthesia were more marked than those observed with chloral hydrate or thiopentone sodium.

Plasma pH fell quite sharply after induction and remained at a low level during isoflurane. The changes observed were correlated with carbon dioxide retention and a respiratory acidosis. The changes observed in all species were qualitatively similar but the sheep showed greater changes. The decreases which took place in plasma pH during Fluothane anaesthesia were greater than those observed with chloral hydrate or thiopentone sodium. It was noted that towards the end of isoflurane anaesthesia and during recovery that the plasma pH returned to the pre-anaesthetic control value before the plasma carbon dioxide content returned to the control value. This may have been due to some carbon buffering capacity or buffering regulating mechanisms coming into play during recovery. This could indicate interference with renal regulation during Fluothane anaesthesia as has been shown to occur with barbiturate anaesthesia in dogs (Bilke, 1967).

CONCLUSIONS

From the results obtained in these experiments the following conclusions were drawn.

- (1) A depression of pulmonary ventilation occurred during the administration of Fluothane to cattle, horses, sheep and dogs.
- (2) Of the species studied, the greatest respiratory depression occurred in sheep, these animals often becoming cyanotic during the administration of Fluothane.
- (3) Accompanying the depression of ventilation there was a rise in the plasma carbon dioxide content, the extent of which depended on the depression of ventilation.
- (4) The retention of carbon dioxide caused a lowering of plasma pH and hence the production of a respiratory acidosis.
- (5) Recovery from the anaesthetic effects of Fluothane was rapid, the plasma pH in most cases returning to the pre-anaesthetic value one hour after removal from the Fluothane atmosphere, while at this point the carbon dioxide content was still elevated.

## SECTION 2

### THE DISTURBANCE OF INSPIRATION CAUSED BY PNEUMONIA.

#### REVIEW OF THE LITERATURE

The pathological changes taking place in the lungs of cattle as a result of pneumonia have been extensively studied by Jarrett (1936) and Jarrett, Jennings, McIntyre, Mulligan and Urquhart (1957). The disturbance of the respiratory rate in pneumonia caused by *Methyococcus glycogenii* has been followed by Jarrett, McIntyre and Urquhart (1954) and has been used by them to assess the severity of the pneumonia process. Scarpini and Haydon (1925) reported no change had taken place in the carbon dioxide capacity of the plasma of cattle with pneumonia but did not indicate the severity of the disease in the animals studied.

## METHOD

The calves used in this series of experiments were treated in a similar manner to that described in Part I.

The pneumonia was produced by giving 4,000 *Mitchondria vivipara* larvae to each calf when it was about eight weeks of age. Three weeks after infection a severe clinical pneumonia was invariably present in these calves.

The calves described as normal in these experiments were not infected with *Mitchondria vivipara* larvae but sometimes had a slight disturbance of respiratory rate due to residual lesions of cuffing pneumonia, a ubiquitous disease of calves. The ventilation and respiratory rates were measured as described in Part I. Blood samples were taken and estimations carried out on the plasma as described in Part I.

## RESULTS

The results of five experiments are given and the variations in design are described separately.

### Experiment I

The plasma carbon dioxide content of arterial blood was determined in two groups of calves of 12 weeks of age. In this experiment one group of 10 pneumonic calves was studied together with another group 10 normal calves. Over a period of nine days 29 blood samples were taken from the pneumonic group, all but one of the calves being sampled on two occasions. From the normal group of 10 calves 36 blood samples were taken, only six of the calves being sampled on two occasions. The results of the determinations are given in table 56.

The mean value for the determinations made on the pneumonic calves was significantly higher than the mean value obtained for the normal calves indicating that some carbon dioxide retention had occurred in these pneumonic calves.

Table 16  
Carbon Dioxide Content of the Arterial Plasma  
of Normal and Pneumonic Calves

Date	Normal Calves		Pneumonic Calves	
	Calf no.	Total CO <sub>2</sub> m.moles/litre	Calf no.	Total CO <sub>2</sub> m.moles/litre
4/6	28	30.2	39	32.9
	29	33.4	34	32.6
			35	32.0
			36	32.4
			37	34.6
7/6	35	31.1	30	33.7
	36	31.9	31	33.7
	37	31.0	32	30.5
	38	31.4	33	35.6
	39	32.0	34	32.5
11/6	40	30.6	35	32.0
	41	30.6	36	31.7
	42	31.4	37	31.4
	43	30.8	38	33.7
	44	33.2	39	31.0
12/6	26	33.6	30	31.9
	27	35.6	31	33.7
	28	33.8	32	30.5
	29	34.3	34	32.2
Mean values		33.9 ± 2.3	32.6 ± 1.6	

### Experiment III.

In this experiment the arterial plasma pH values of a group of 13 panniculate calves were compared with the plasma pH values of 13 normal calves. Both groups of calves were about 12 weeks of age. The panniculate group had been suffering from panniculitis for about three weeks. The results obtained are given in table 37.

There were no significant differences for the values for the plasma pH obtained in the two groups of calves and all values found were within the range of values determined for calves in Part I of this chapter.

Table 57

## The Arterial Plasma pH of Normal and Pneumonia Calves

Normal Calves Calf no.	Plasma pH	Pneumonia Calves Calf no.	Plasma pH
9722	7.47	9836	7.38
9723	7.44	9831	7.44
9724	7.42	9832	7.42
9725	7.44	9834	7.44
9726	7.45	9833	7.40
9727	7.47	9828	7.39
9728	7.44	9823	7.39
9729	7.40	9821	7.44
9730	7.47	9819	7.38
9802	7.42	9816	7.42
9732	7.38	9914	7.36
9733	7.42	9817	7.40
9512	7.39	9127	7.39
Mean values		$7.40 \pm 0.04$	

### Experiment LIX

In this experiment determinations of the carbon dioxide content and the pH of plasma were made simultaneously in two groups of calves, one consisting of five normal calves and the other of 19 pneumoniae calves. The results obtained are given in Table 50.

These results confirm those of the previous two experiments that parasitic pneumoniae in calves caused a carbon dioxide retention with a significantly higher plasma carbon dioxide content and a significantly lower plasma pH than normal calves exposed to similar conditions.

Table II  
The Arterial Plasma pH and Carbon Dioxide Content  
of the Plasma of Normal and Diuretic Calves

Normal Calves			Diuretic Calves		
No.	pH	$\text{CO}_2$ m.mole/liter	No.	pH	$\text{CO}_2$ m.mole/liter
31	7.33	27.4	4	7.42	27.8
32	7.47	27.8	5	7.44	29.1
33	7.46	27.9	617	7.42	31.2
34	7.36	26.8	618	7.38	26.4
931	7.42	27.6	62	7.40	30.4
Mean values			61	7.42	30.1
$7.43 \pm 0.04$			55	7.39	32.0
$27.9 \pm 0.54$			56	7.34	32.6
			57	7.36	32.4
			58	7.40	29.1
			59	7.39	33.5
			60	7.36	30.5
			37	7.33	29.8
			38	7.40	28.9
			39	7.36	29.8
			49	7.36	26.8
			932	7.33	29.0
			933	7.46	29.9
			763	7.30	28.3
			Mean values		
			$7.38 \pm 0.04$		
			$29.8 \pm 2.0$		

#### Experiment IV

The procedure except in this experiment was different from that in all other experiments in that the calves had picked up their parasitic infection on a heavily contaminated pasture. These calves were therefore much more severely infected than the calves in the other parasitic groups and in fact all the calves died a few days after the measurements were made. The results obtained are given in table 59.

There was a considerable difference in the plasma carbon dioxide contents of the two groups of calves but still no significant difference between the plasma pH values. The severity of the pneumonia is indicated by the great increase in respiratory rate. Although the ventilation rates in the parasitic group had increased by nearly 50%, the tidal air had decreased by about one half.

Table 39

The Plasma oil and the Carbon Dioxide Content Compared with  
the respiratory and ventilation rates of normal and pregnant calves

Calf	CO <sub>2</sub> m.moles/33.620	pH	Ventilation litres/min.	Respiratory rate/min.	Plasma oil
<b>Normal calves</b>					
a	23.6	7.49	21	42	630
b	21.3	7.38	22	35	620
c	22.4	7.38	23	30	660
d	22.1	7.38	24	30	800
Mean	22.6	7.38	23	34	740
<b>Pregnant calves</b>					
e	27.6	7.26	39	75	400
f	21.3	7.30	30	62	400
g	26.9	7.35	40	80	500
h	26.3	7.40	23	57	430
i	22.3	7.35	49	62	540
Mean	24.3	7.33	37	77	470

## Experiment V

In this experiment a group of five calves was studied over a period of 40 days during which time six examinations were carried out approximately at weekly intervals. The first examination was carried out 10 days after an infecting dose of 4,000 Mycobacterium tuberculosis bacilli had been given, i.e., at the early pneumonia stage.

The results for each calf are given in tables 60 (a) - (f). In table 60 (f) the mean values for each measurement are given.

In this experiment it was possible to follow the changes from the early to the severe pneumonia state. The plasma carbon dioxide content increased until the fourth week from infection after which they tended to stabilize at a high **value**. This was also true of respiratory rate. On the other hand the pH did not alter significantly.

Table 60

Changes in Plasma pH, Plasma Carbon Monoxide Content,  
 Ventilation and Respiratory Rates of 3 Calves  
 Following *P. mytilinus* Infection

Table 60 (a)

Table 55

Date	$\text{CO}_2$ m.moles/litre	pH	Ventilation litres/min.	Respiratory rate/min.	Ridol etc
12/9	29.1	7.38	24	20	800
19/9	27	7.36	25	47	930
26/9	36.7	7.44	30	55	945
3/10	31	7.39	32	92	625
10/10	34.3	7.38	32	50	640
23/10	36.3	7.28	31	55	560

Table 60 (b)

G.R.C. 97

Date	$\text{CO}_2$ m.mole/litre/min	pI	Ventilation litres/min.	Respiratory rate/min.	Tidal litre
12/9	29.4	7.35	26	30	670
19/9	26.9	7.39	26	32	560
26/9	26.0	7.44	26	32	540
3/10	29.4	7.40	26	37	460
10/10	26.6	7.38	26	43	420
21/10	26.3	7.36	26	73	490

Table 60 (c)

G.R.C. 98

Date	$\text{CO}_2$ m.mole/litre/min	pI	Ventilation litres/min.	Respiratory rate/min.	Tidal litre
12/9	30.6	7.35	22	44	372
19/9	28.4	7.39	25	60	250
26/9	30.5	7.41	26	69	260
3/10	28.5	7.46	26	66	272
10/10	28.5	7.39	26	78	203
21/10	28.6	7.36	26	126	215

Table 60 (a)

Case 52

Date	$\text{CO}_2$ m.moles/litre	pH	Ventilation litres/min.	Respiratory rate/min.	Tidal air
12/9	23.8	7.40	23	61	630
19/9	23.6	7.38	24	50	480
26/9	21.5	7.44	27	75	360
3/10	30.6	7.33	33	57	560
10/10	31.6	7.37	39	50	520
23/10	31.6	7.33	37	69	540

Table 60 (b)

Case 200

Date	$\text{CO}_2$ m.moles/litre	pH	Ventilation litres/min.	Respiratory rate/min.	Tidal air
12/9	21.3	7.38	22	35	630
19/9	21.3	7.36	31	93	330
26/9	36.2	7.30	24	78	208
3/10	25.8	7.36	32	97	300
10/10	37.9	7.34	30	69	337
23/10	35.1	7.32	37	80	462

Table 60 (3)

## Mean Values of 5 Calves

Date	$\text{CO}_2$ m.mol/100/12620	pH	Ventilation 12 breaths/min.	Respiratory rate/min.	Tidal litres
22/9	29.8	7.37	20	36	555
23/9	33.4	7.38	23	36	410
25/9	33.7	7.42	25	66	378
3/10	32.6	7.39	26	66	393
10/10	34.9	7.37	25	62	403
20/10	22.9	7.33	23	79	426

DISCUSSION  
(Part II section 2)

Calves affected with a parasitic pneumonia showed an increase in ventilation and respiratory rates, the increase in the respiratory rate being the greater so that the tidal air fell indicating shallower and less efficient respiration. The carbon dioxide content of the plasma of these calves was significantly higher than that of normal calves. It was noteworthy that, with the exception of one anoxaemic severely ill calf which died the next day, the plasma pH of these pneumoniae calves was never below normal. A compensated respiratory acidosis therefore existed in these calves.

The pneumoniae animals showed a continuing increased respiratory response to pneumonia so that there must have been a continuous stimulation to respiration. The increased carbon dioxide content of the plasma would also act as a stimulus, but since the plasma pH was within the normal range, the pH would not act as a stimulus.

It was observed that the arterial blood of these pneumoniae calves was darker in colour than arterial blood from normal calves. The lowered oxygen content which this indicated could also have acted as a stimulus to respiration by its effect at the chemoreceptors although this might be counteracted by the fact that carbon dioxide has a

dopamine effect on respiration.

## CONCLUSIONS

- (1) Calves affected with a parasitic pneumonia showed an increase in both the respiratory and ventilation rates compared to normal calves.
- (2) The increase in the respiratory rate was greater than the increase in ventilation rate so that respiration became shallower and less efficient.
- (3) These pneumoniae calves showed an increase in the plasma carbon dioxide content but no marked change in the plasma pH.

SECTION 3

THE EFFECTS OF ALTERATIONS IN THE ATMOSPHERE  
ASSOCIATED TO CHANGES IN THE VENTILATION RATE.  
THE PLASMA pH AND PLASMA CARBON DIOXIDE CONTENT OR OXYGEN

REGULATION OF THE RESPIRATION

Alterations in the atmosphere can affect breathing effect respiration and the changes in plasma carbon dioxide tensions, plasma carbon dioxide content and plasma pH have been extensively studied in man and experimental animals.

The predominant role that carbon dioxide plays in the day by day regulation of respiration emerged from the studies of Haldane many years ago whose results are still quoted in most textbooks of physiology.

A good summary of the effects on respiration of the tensions of oxygen and carbon dioxide in the blood and the plasma pH was given by Guy (1950). He stated that an increase in the arterial carbon dioxide tension ( $\text{PaCO}_2$ ) caused hyperventilation and a lowering caused hypoventilation. He described the stimulant effect of hypoxia on respiration

as slight, no appreciable change in respiration taking place until the content of oxygen in inspired air fell below 14%. Gray discussed the reasons for this and suggested that the hyperventilation caused by the lowered arterial  $pO_2$  resulted in acapnia the depressant effect of which counteracted the stimulation of hypoxia so that the resultant respiratory stimulation was slight. He also stated that the effects of increasing the oxygen content of inspired air were negligible in the normal human subject. However, there is some evidence for the normal arterial  $pO_2$  of the unanesthetized human subject acting as a tonic stimulus to respiration. The measurements of arterial  $pO_2$  in normal human subjects breathing room air found by Amusco and Molton (1956) indicated values below the threshold found in animal experiments by Macleod et al. (1935). Maffinay, Holt et al. (1943) found evidence for the presence of tonic chemoreceptor sensitivity at the normal arterial  $pO_2$  of dogs. They produced experimentally a decrease in ventilation if dogs were given pure oxygen to breathe. This decrease no longer took place when the chemoreceptors were denervated.

Examination of the literature revealed no evidence of studies of the respiratory behavior of cattle when exposed to gas mixtures of different composition to atmospheric air.

1000000

Studies were carried out on Ayrshire calves between six and 32 weeks of age. The calves were lightly restrained in a dog stand and the close fitting neck previously described was fitted. This procedure seldom disturbed the calf provided their heads were held gently. The neck was connected by means of a brass 'Y' tube and non-kink rubber tubing to a pair of unidirectional valves. The expiratory portion of these valves led to the gas motor so that the ventilation could be recorded. The inspiratory side of the unidirectional valves was connected to a length of flexible tube (HooverFlex). At the end of this flexible tube was a 'T' junction to the end of which was attached a one and a half inch excess release valve which could be opened to the atmosphere. To the third arm of the 'T' an eight gallon reservoir bag was fitted. The other end of the reservoir bag had a small diameter rubber connection through which measured gas mixtures were fed into the reservoir bag.

The calves were allowed to inspire room air through the open excess valve for a period of 30 minutes during which time ventilation rates and respiratory rates were recorded. At the end of this control period, if blood changes were being followed, control blood samples were taken from

-1-

the jugular vein or brachial artery. The sphenic valve was then eluted and the gas mixture, the effects of which were being studied, was fed into the reservoir bag. Gas admixtures were fed in at such a rate as to keep the reservoir bag about half full, the animal's inspiratory efforts being plainly visible in the extraction of the gas mixture from the bag. Gas mixtures were either made up as required or purchased ready made from British Oxygen Co., Ltd., Rosehill Works, Polmadie, Glasgow. The following mixtures were purchased: (a) 90% nitrogen and 10% oxygen, (b) 95% oxygen and 5% carbon dioxide, (c) 200% carbon dioxide, and (d) 300% oxygen. Accurate analysis of the gas mixture presented to the animal was not attempted.

The calves were allowed to breathe the gas mixtures for varying periods of time during which respiratory and ventilation rates were recorded. Blood samples were taken at intervals, centrifuged and the plasma pH and plasma carbon dioxide content were determined. In a number of cases, at the end of an experiment, the calf was allowed to inspire pure air for a period while ventilation rates and respiratory rates were again recorded and blood samples taken for the determination of plasma pH and plasma carbon dioxide content.

RESULTS

The results of the studies in this section are arranged in the following order.

- (1) Hyperoxigenation
- (2) Hypoxospina
- (3) Anoxia

## (2) Hyperoxigenation

In this section a variety of experiments have been carried out. In experiments I and II the effects of hyperoxigenation on the ventilation rates of three calves and three dogs, respectively, were studied.

In experiments III and IV more extensive studies were carried out in that the effects of the gradual increase in the concentration of oxygen inspired on the carbon dioxide content and the pH of plasma, the ventilation and respiratory rates were studied.

In experiment V, similar determinations were made on a normal calf breathing pure oxygen for 35 minutes.

In experiments VI and VII the studies on the effects of pure oxygen were continued in two photogenic calves.

Experiment I

Each calf was allowed to breathe room air for a period of 10 minutes during which time the ventilation rate was recorded. At the end of this time the release valve was closed and each calf was made to inspire pure oxygen from the reservoir bag for a further 10 minutes and the ventilation rate was again recorded. The results obtained are given in figures 48 and 49.




Fig. 48

Depression of ventilation rate caused by inhalation of pure oxygen




Fig. 49

Reduction of ventilation rate caused by inhalation of pure oxygen

A reduction of ventilation of varying degree occurred in each calf.

## Experiment II

Each of three greyhounds was allowed to inspire room air for a period of 10 minutes during which time its ventilation rate was recorded. Each greyhound was then made to inspire pure oxygen for 10 minutes when the ventilation rate was again recorded. The results obtained are given in figures 50 and 51.




Fig. 50

Reduction of ventilation rate caused by inhalation of pure oxygen




Fig. 51

Reduction of ventilation rate caused by inhalation of pure oxygen

As in the calves in experiment I these dogs showed a varying reduction of ventilation rate.

### Experiment III


In this experiment one calf was submitted to alternate periods of 10 minutes of air and air/oxygen mixtures. On each of the three periods of oxygenation the concentration of oxygen was increased, terminating in pure oxygen on the third occasion. Determinations carried out are given in table 62 and illustrated graphically in figure 52.

**Table 62**  
**Effects of Inhalation of Air Enriched with**  
**Different Proportions of Oxygen**

Mixture Inhaled	Air + 1 Micro O <sub>2</sub> per min.		Air + 4 Micro O <sub>2</sub> per min.		Air + Pure O <sub>2</sub>		Air
	Air	Pure O <sub>2</sub>	Air	Pure O <sub>2</sub>	Air	Pure O <sub>2</sub>	
gH	7.40	7.32	7.41	7.36	7.98	7.23	
CO <sub>2</sub>	31	33	32.9	33	32.9	33.3	
Roop	57	62		63		65	59
Vet	43	39		35		34	39

11417 19.9.58

AIR	O <sub>2</sub> 1 LITRE/MIN.	AIR	O <sub>2</sub> 4 LITRE/MIN.	AIR	O <sub>2</sub> 100%	AIR
-----	--------------------------------	-----	--------------------------------	-----	------------------------	-----

Fig. 52

Effects of inhalation of air enriched with  
different proportions of oxygen

It was noted that a slight increase in the respiratory rate and a decrease in the ventilation rate occurred, the changes being greater the higher the concentration of oxygen in the inspired air. The changes in plasma pH and plasma carbon dioxide content were less obvious but each time enriched air was given the pH fell and the plasma carbon dioxide content rose but in the control periods between enriched mixtures neither pH nor carbon dioxide content returned to the initial values.

## EXPERIMENT XI

A similar experiment was conducted on a calf with a more severe pulmonary pneumonia. The periods between the administration of different levels of oxygen were lengthened to half an hour. The results obtained are given in table 62.

Table 62  
Effects of Inhalation of Air Enriched with  
Different Proportions of Oxygen

Mixture Administered	Air	Air + 1 liter O <sub>2</sub> per min.	Air	Air + 4 liters O <sub>2</sub> per min.	Air	Pure O <sub>2</sub>
Al	7.39	7.35	7.40	7.35	7.39	7.27
O <sub>2</sub>	39	38.4	38.1	38.5	39.1	39.4
Resp	94	95		87		92
Vent	36	36		32		31

It will be seen that a **reduction** of ventilation occurred with the high concentrations of oxygen but that no **reduction** occurred when 1 liter of oxygen per minute was added at a ventilation rate of 36 liters per minute. The respiratory rate was lowered slightly by high concentration of oxygen. The plasma pH showed the expected trends but apart from the marked fall during the administration of pure oxygen the

changes were not significant. The plasma carbon dioxide content, like the plasma pH, showed the expected trend in that it rose when the ventilation rate fell but the changes were again not significant.

## Experiment V

In this experiment the effects of more prolonged administrations of pure oxygen were studied in a two week old calf. The determinations made are given in table 62 and illustrated in figure 59.

Table 62  
Effects of Inhalation of Pure Oxygen

	Air 5 min.		100% O <sub>2</sub>		Air 7 min.
	5 min.	15 min.	35 min.		7 min.
Resp	12	37.6	18.5	39	22
Vent	6	6.9	4.7	5.9	6.0
CO <sub>2</sub>	35.5	26	35.4	36.6	35.6
pH	7.40	7.36	7.35	7.33	7.43

It was observed that a slight increase in the respiratory rate occurred during the administration but a noticeable decrease occurred afterwards. Ventilation fell during the administration of pure oxygen but rose again before the cessation of the administration. This initial rise not pneumonic but initial high values of arterial oxygen were observed for the plasma carbon dioxide content. Concomitant with the changes in ventilation the plasma carbon dioxide content rose and the plasma pH fell but the changes were small. There was a rapid return

to normal values once the calf again breathed air.




Fig. 53  
Effects of inhalation of pure oxygen

## Experiment VI

In this experiment the prolonged administration of pure oxygen to a severely premature calf was studied. The results of the determinations are given in table 64.

Table 64

### Effect of Inhalation of Pure Oxygen

Parameter	Air	Inhalation of pure O <sub>2</sub> (minutes)				Recovery (minutes)	
		10	20	30	45	5	10
pH	7.30	7.29	7.22	7.21	7.20		7.33
CO <sub>2</sub> , plasma	37.5	26	33.8	39.4	38.9		38.6
Pcap	96	94	97	95	94	95	94
Vout	24	29	33	30	32	30	27

It was observed that the respiratory rate was little affected by inhalation of pure oxygen but that the ventilation rate fell. After the calf had ceased breathing pure oxygen the ventilation rate rose above the initial level probably because of the higher than normal plasma carbon dioxide content. The plasma pH fell quite sharply and remained low. The plasma carbon dioxide content which was initially high rose to even higher levels during the administration of pure oxygen and had not returned to the initial level after 30 minutes.

### Experiment VII

The effects of the inhalation of pure oxygen on a pretermic calf were again studied. This Ayamhise calf was as severely affected as the previous calf. The results obtained are given in table 65 and illustrated in figure 56.

Table 65  
Effects of Inhalation of Pure Oxygen

Air		Inhalation of pure O <sub>2</sub> (minutes)					Recovery (min.)		
		5	10	25	35	45	5	10	20
pH	7.42	7.33	7.27	7.25	7.23	7.22	7.33	7.35	
CO <sub>2</sub>	39.4	39.6	41.3	42.1	43.0	43.1	42.3	42.3	
Resp	69	76	60	75	69	70	64	66	73
Vent	32	36	26	36	38	29	35	34	32

The ventilation rate was observed to fall as did the respiratory rate during the administration of pure oxygen. When the administration of pure oxygen had ceased respiratory rate rose as did the ventilation rate to values above the initial values. There was in this calf a marked rise in the plasma carbon dioxide content and a fall in plasma pH and even 10 minutes after the administration of pure oxygen had

ceased the pH and plasma carbon dioxide content had not returned to the initial values.




Fig. 54  
Effects of inhalation of pure oxygen

## (2) Hypoxia

In this section a variety of experiments have been carried out on the effects of hypoxia. In experiments I to IV the plasma pH and plasma carbon dioxide changes due to inhalation of 95% oxygen and 5% carbon dioxide were followed in normal two-month old calves. In experiments II, III and IV the ventilation rates were also measured and in experiments XII and XIV respiratory rates were measured.

The effects of hypoxia on calves with mild and severe pneumonia were studied in experiments V and VI, respectively.

Experiments I, II, III and IV

The results obtained from experiments I to IV are given in tables 66, 67, 68 and 69, respectively, and in addition the results from experiment IV are illustrated graphically in figure 56.

Table 66

Experiment I  
Venous plasma

Effects of Inhalation of 5% Carbon Dioxide

Parameter	Mean	5% CO <sub>2</sub> Inhalation (minutes)			Recovery (minutes)	
		10	20	30	5	35
pH	7.35	7.24	7.24	7.23	7.37	7.32
CO <sub>2</sub>	30.1	31.5	31.5	32.5	29.6	30.1

Table 67

Experiment II  
Venous plasma

Effects of Inhalation of 5% Carbon Dioxide

Parameter	Mean	5% CO <sub>2</sub> Inhalation (minutes)		Recovery (minutes)
		10	20	
pH	7.37	7.25	7.24	7.37
CO <sub>2</sub>	29.4	31.4	33.1	32.5
Vent	29	30	33	25

Parameter	AIF	% CO <sub>2</sub> Saturation (Minutes)			Recovery (minutes)		
		5	10	30	5	10	30
Pt	7.37	7.34	7.29	7.21	7.26	7.32	7.37
CO <sub>2</sub>	32.0	32.76	34.4	36.5	33.75	33.0	32.0
Bsyp	32	40	56	50	40	36	32

Parameter	Altitude	5% CO <sub>2</sub> Inhalation (minutes)			Recovery (minutes)	
		5	15	30	5	15
pH	7.35	7.37	7.37	7.35	7.35	7.35
CO <sub>2</sub>	27.9	29.5	29.6	30.5	29	27.5
Po <sub>2</sub> mm Hg	25	29	36	29	27	26
Vent	26	33	37	33	20	19




Fig. 55  
Effects of inhalation of 5% carbon dioxide

These four calves showed an increased plasma carbon dioxide content and a lower plasma pH. Where ventilation rate was measured this showed an increase as did respiratory rate when this was recorded. In all cases there was a rapid return to the initial values when carbon dioxide administration ceased.

### Experiment V

The results obtained from studying the effects of hypercapnia in a mildly pneumonic calf are given in table 70 and figure 56.

Table 70  
Effects of Inhalation of 5% Carbon Dioxide

	Air	5% CO <sub>2</sub> inhalation (minutes)		Recovery (minutes)		
		5	15	1	15	30
pH	7.37	7.23	7.18	7.26	7.29	7.35
O <sub>2</sub>	33.9	35.8	35.0	34.3	33.0	33.5
Resp	44	56	54	65	75	60
Vent	25	38	45	40	35	23

The respiratory rate of this calf continued to increase throughout the period of administration of the 95% oxygen and 5% carbon dioxide and this increase was maintained through about half of the recovery period. The ventilation rate rose and started to fall while the calf continued to breathe the gas mixture. The plasma carbon dioxide content rose upon administration of the carbon dioxide but started to fall before this administration had ceased. The plasma pH fell quite sharply and continued to fall throughout the administration of carbon dioxide.

During recovery which was slower than in the non-pneumonic calf, return to the initial values of carbon dioxide and pH took 30 minutes but at this point both respiratory rate and ventilation rate were still above the initial values.




Fig. 56  
Effects of inhalation of 5% carbon dioxide

## Experiment VI

The results obtained from studying the effects of hypoxia in a severely pneumonic calf are given in table 71 and figure 57.

Table 71

### Effects of Inhalation of 5% Carbon Dioxide

Time	Inhalation of 5% CO <sub>2</sub> (minutes)							Recovery (minutes)		
		14	20	25	32	37	46	5	10	20
pH	7.34	7.24	7.26	7.37	7.17	7.17	7.18	7.23	7.31	
CO <sub>2</sub>	34.9	38.6	40.3	41.9	41.8	40.4	40.5	32.4	37	
Vent	30	27	30	26	30	32	33	35	33	32
Room	96	70	63	62	66	66	72	61	66	65

This calf differed from the other calves in that during the administration of carbon dioxide the respiratory rate underwent a marked decrease but began to also again towards the end of the administration. The ventilation rate showed some decrease initially but a rise occurred towards the end of the administration of the gas mixture to above the initial value. The plasma carbon dioxide content rose and the plasma pH fell initially and then the carbon dioxide content fell and the plasma pH started to also when the ventilation increased during the administration.

of the gas mixture. Changes in plasma pH and carbon dioxide content were quite marked. During the first 20 minutes of the recovery period there was a slight return towards the initial values of plasma pH and plasma carbon dioxide content.




Fig. 57  
Effects of inhalation of 5% carbon dioxide

### (3) *Anoxia*

The experiments were carried out to study the effects of anoxia in calves by the administration of a gas mixture deficient in oxygen. These calves were made to inspire a gas mixture containing 90% nitrogen and 10% oxygen.

## Experiments I and II

The results obtained in experiments I and II are given in tables 72 and 73 and figures 58 and 59.

Table 72

### Effects of Acetyl on the Ventilation and Respiratory Rates,

### The Carbon Dioxide Content and the pH of the Plasma of a Cat.

AlP	Aer	Inhalation of N <sub>2</sub> mixture (minutes)					Recovery (minutes)		
		10	22	36	47	5	23	27	
		pH	7.40	7.47	7.48	7.45	7.35	7.32	
		CO <sub>2</sub>	26.6	25.6	25.3	25.8	25.6	25.5	25.2
		Resp	29	30	24	33	34	33	32
		Vent	36.4	21	20	21	21	26	24




Fig. 58

Effects of anoxia on the ventilation and respiratory rates,  
the carbon dioxide content and the pH of the plasma of a calf

Table 73

Effects of Anoxia on the Ventilation and Respiratory Rates,  
the Carbon Dioxide Content and the pH of the Plasma of a Calf

	Air	Inhalation of N <sub>2</sub> mixture (minutes)					Recovery (minutes)	
		12	30	45	55	10	20	
pH	7.43	7.52	7.53	7.58	7.57	7.44	7.44	
CO <sub>2</sub>	30	30.8	30	29.1	28.6	29.1	29.8	
Resp	18	24	25	27	27	20	20	
Vent	4	7	5	9	10	5	5	




Fig. 29

Effects of anoxia on the ventilation and respiratory rates,

the carbon dioxide content and the pH of the plasma of a calf

Both calves responded to the anoxic stimulus with an increase in the respiratory and the ventilation rates. As a result of the increased ventilation additional carbon dioxide was excreted so that the plasma carbon dioxide content fell and the plasma pH rose. It seemed that the lowest plasma carbon dioxide content and the high pH had a less stimulating effect on respiration because the ventilation rate then fell while the

-20-

respiratory rate showed greater variation although the anoxic mixture was still being administered. Plasma pH then fell and plasma carbon dioxide content rose. After the administration of the anoxic mixture had ceased the ventilation rate and the respiratory rate continued to fall while the plasma pH also fell and the plasma carbon dioxide content rose.

DISCUSSION  
(Part II section 3)

The administration of oxygen at one atmosphere caused a reduction of the ventilation of calves and the glycerinoids studied had little effect on the respiratory rate of the calves. The effect of the hyperventilation of the calves was to cause a rise in the carbon dioxide content and a fall in the pH of the plasma. In normal calves and calves not severely affected with pneumonia the changes produced by oxygen administration were small and when the subject was again made to breathe atmospheric air, a rapid return to the initial values took place.

Experiments by Buler et al. (1939) on cats showed that the chemoreceptors of the carotid glomeruli of this species stimulated at tensions of oxygen found in normal circumstances and that as the oxygen content was increased the frequency of the impulses decreased. These results suggested that in cats normal breathing reflects a certain amount of chemoreceptor drive due to oxygen lack. Using trained dogs Wark et al. (1943) found that a reduction of ventilation occurred immediately following the inhalation of pure oxygen. This effect was abolished by denervation of the carotid and aortic bodies.

Experiments conducted in man by Delphys and Gorwood (1947), DeJarnett et al. (1957) and by Amussen and Nelson (1958) have shown a decrease in

In ventilation immediately following the inhalation of pure oxygen. Ballico and Comroe (1937), however, noted a secondary rise in ventilation after the first one or two inhalations. They attributed this secondary rise to the excitant effect of oxygen. From their determinations of arterial  $pO_2$  in man, at rest, breathing room air, and a comparison with the threshold value for chemoreceptor drive found by Mitchell et al. (1935), Armstrong and Nielson (1956) concluded that with oxygen at the tensions present in blood in man breathing normal air there was stimulation of the chemoreceptors.

Findings in the present series of experiments on cattle of a decrease in ventilation with a fall in plasma pH and a rise in the plasma carbon dioxide content during inhalation of pure oxygen require more extensive investigation. It would not be surprising if cattle showed species differences in their respiratory behaviour compared to man since in addition to those results obtained with hyperoxigenation it has been shown that at a weight basis their pulmonary ventilation is high (Reedy, 1950), the packed cell volume of bovine blood is low as is the haemoglobin concentration compared to the human subject (Nielson, 1956).

When 100% oxygen was administered to severely pneumoniae calves a reduction of ventilation again occurred with greater changes in plasma pH and plasma carbon dioxide content. These calves took longer to return to the initial values when they were again breathing room air. However, the pneumoniae calves studied increased their respiratory

offered in order to remove the increased plasma carbon dioxide content produced by the hypoventilation due to the administration of pure oxygen. It is possible that the administration of pure oxygen for prolonged periods to a severely pneumonic calf making its normal respiratory efforts might prove deleterious. It has been shown that continual inhalation of pure oxygen at one atmosphere may produce pulmonary edema (Bain, 1953). The reduction of ventilation might reduce the plasma carbon dioxide content to below levels which the calf could not then get rid of.

The administration of a 95% oxygen and 5% carbon dioxide mixture to normal calves caused an immediate increase in the respiratory rate and pulmonary ventilation. At the same time there was a marked fall in plasma pH and a rise in the plasma carbon dioxide content. The calf mildly affected with pneumonia responded slightly differently in that the increase in ventilation was continued for a longer period during recovery from the administration of 95% oxygen and 5% carbon dioxide. The severely pneumonic calf responded to 95% oxygen and 5% carbon dioxide with a reduction of both pulmonary ventilation and respiratory rate although the plasma pH fell to very low values and the carbon dioxide content rose to very high values. This calf apparently responded to the 95% oxygen content of the gas mixture and was not responsive to the increased carbon dioxide intake until concentrations within the plasma above those normally found in pneumonia had arisen.

The findings in this calf are in agreement with the findings by Boutourline-Young and Whittenberger (1951) that the respiratory centre may in certain circumstances become adapted to chronically high carbon dioxide by losing its responsiveness.

The respiratory rates of normal calves when made hyperventilating were not as high as those observed in pneumonic calves although the plasma carbon dioxide content rose to values observed in pneumonic animals and the plasma pH fell below the values observed in pneumonic animals. An increased carbon dioxide content of the plasma and the rise in  $pCO_2$  could not therefore be the only stimulus to the increased respiratory rate of pneumonic calves.

When calves were exposed to hypoxia by the administration of gas mixtures containing less than 20% oxygen there was at first an increase in both respiratory rates and pulmonary ventilation while the plasma carbon dioxide content fell and the plasma pH rose. Although the hypoxic gas mixture was still being administered the pulmonary ventilation fell. These observations might be explained on the basis of Gray's hypothesis (1950) that the final drive to respiration due to changes in the blood gases is the resultant of a number of factors some stimulant, some depressant. In these calves the fall in ventilation which occurred while the animals were still being rendered anoxic was no doubt due to the resultant depression from the sum of central anoxic depression, central hypoventilatory depression, central alkalotic depression, reduction in stimulation of the chemo-receptors from hypoventilation and alkalosis and chemo-receptor stimulation due to hypoxia.

CONCLUSIONS

From the results obtained in these experiments the following conclusions may draw.

- (1) Hypoxycapnia of calves by the administration of 100% oxygen at one atmosphere caused a depression of pulmonary ventilation leading to an elevation of plasma carbon dioxide content and a fall in plasma pH.
- (2) These changes were much greater in pneumonic calves than in normal calves.
- (3) Hypoxycapnia caused an immediate increase in respiratory and ventilation rates, a rise in the carbon dioxide content and a fall in the pH of plasma.
- (4) Pneumonic animals took longer to recover from the effects of hypoxycapnia than normal animals.
- (5) Anæsthesia caused stimulation of respiratory and ventilation rates of calves which led to a respiratory alkalosis.

DISCUSSION  
(Part II)

The study of the disturbance of respiration of cattle caused by general anaesthesia with Fluothane, thiopentone sodium and chloral hydrate showed that there was a depression of pulmonary ventilation which was greatest when Fluothane was the anaesthetic agent. A part of the marked hypoventilation may be due to the concomitant administration of oxygen, which was shown to be responsible for some reduction of ventilation when administered to unanaesthetised cattle, but there is no proof of this. Horses, dogs and sheep also showed a depression of ventilation when anaesthetised with Fluothane, sheep often becoming apneic.

Because of the depression of ventilation, carbon dioxide retention took place so that the plasma carbon dioxide content rose to higher concentrations than those found in normal unanaesthetised animals. As a consequence of the carbon dioxide retention, the increased  $pCO_2$  caused an increase in the H ion concentration, a fall in plasma pH, and a respiratory acidosis was produced. The changes in plasma pH which occurred in general anaesthesia were greater than the variations observed in normal unanaesthetised cattle when the latter were studied over periods from one to six hours. It was noticed that in prolonged general anaesthesia of two to three hours duration the plasma carbon dioxide content did not always continue to rise but stabilised at a high

concentration or fall while the animal was still being maintained in a state of surgical anaesthesia. It was considered that this may have been as a result of a general depression of metabolism due to anaesthesia and hence a decreased carbon dioxide production.

During the anaesthetization of animals with Fluothane and oxygen it was noticed that venous blood became arterial in colour so that pulse was not a feature of this method of anaesthesia provided the circulation was maintained. The rapid recovery from Fluothane anaesthesia meant that the ventilation returned to normal or sometimes above normal quite rapidly and the carbon dioxide content of the plasma fell while the plasma pH rose. The return of plasma pH to the pre-anaesthetic control value before the return of the plasma carbon dioxide content to the pre-anaesthetic value may have been due to renal compensation which was previously inhibited by the anaesthetic agent. It has been shown that general anaesthesia can interfere with renal regulation (Blelo, 1957).

Uloperitone sodium and chloral hydrate anaesthesia in cattle produced changes that resembled those produced by Fluothane anaesthesia.

Bronomale calves showed an increase in respiratory rate and pulmonary ventilation, the greater increase occurring in the respiratory ratio, so that these calves had a smaller tidal volume and shallower respiration than normal calves. The plasma carbon dioxide concentrations of prewean calves were higher than those observed in normal calves but lower than the concentrations found in anaesthetized and the concentrations

obtained by the administration of 95% oxygen and 5% carbon dioxide to calves. It was noteworthy that with the exception of one animal, scarcely all calf the plasma pH was never below normal. This contrasted with anesthetized animals. A compensated respiratory acidosis therefore existed in these pneumoniae calves.

Hypoxia produced by the inhalation of pure oxygen at one atmosphere caused a reduction of the ventilation of calves and dogs. In calves it was shown that this reduction of ventilation caused a rise in the plasma carbon dioxide concentration and a fall in the plasma pH. The changes were small in normal calves and more marked in pneumoniae calves. There was little change in the respiratory rate of these calves. After removal from the pure oxygen normal calves quickly returned to the pre-administration values of plasma carbon dioxide content and plasma pH, while the pneumoniae calves studied were able to increase their respiratory efforts to get rid of the extra carbon dioxide which had built up in their circulation as a result of the hyperventilation caused by pure oxygen inhalation. It was thought that some pneumoniae animals may not have been able to do this so that the administration of pure oxygen to them might prove deleterious.

When the effects of the administration of 5% carbon dioxide and 95% oxygen were studied the results obtained in normal animals were as expected in that there was an increase in respiratory rate and pulmonary ventilation and plasma carbon dioxide content, and a fall in plasma pH.

Although the plasma carbon dioxide content rose to values observed in pneumonia, the respiratory rate did not increase to the values observed by Jarrett *et al.* (1957) in pneumonia.

Anoxic anaemia in calves produced by the inhalation of 90% nitrogen and 10% oxygen caused initially an increased respiratory rate and pulmonary ventilation so that a respiratory alkalosis was produced. Ventilation then decreased while this gas mixture was still being administered and as a consequence plasma pH fell and plasma carbon dioxide content rose again. Hypoxia by itself did not act as a marked stimulus to respiration.

The results of these studies on respiratory disturbances in cattle have solved some interesting problems on certain aspects and shed light on other aspects of clinical work. The fact that calves suffering from a severe parasitic pneumonia did not become acidotic is probably the most surprising result obtained. In this connection, too, it is interesting to note that it was not possible to produce as great a degree of tachypnoea by means of artificial hypoxaemia or hypoxia as did pneumonia. Further investigation is necessary into the locus for the receptors for the tachypnoea of parasitic pneumonia.

ACKNOWLEDGEMENTS

I wish to record my indebtedness to  
Professor W. E. Wolpaw and Dr. W. E. McNaughton for advice, guidance  
and criticism throughout this work.  
The staff of the Surgery Department, particularly Mr. S. Jennings, M.R.C.V.S.  
for enabling studies to be made on unanaesthetized animals.  
Dr. J. V. A. Duskin and Mr. J. Brockway of the Scottish Hospitals  
Infectious Research Trust Ltd., Glasgow for advice and the use of  
calibrating apparatus.  
Dr. W. P. H. Jarrett and Mr. H. G. C. Sharp, R.V.M.S., M.R.C.V.S. for  
the provision of mounted animals and the production of the  
experimental pneumonia.  
Misses Joyce Kerr, Gertrude Colquhoun and Helen Petrie, Mrs. Mary Dow and  
Mr. Bryan Reddick for their help in much of the technical work.  
Mr. A. Minale for taking the photographs.  
Miss Catalonia Cowan for being extremely patient and painstaking in the  
preparation and typing of this thesis.

REFERENCES

- Anderson, A.K., Gayley, H., and Pratt, L. (1930) J. Dairy Sci. 13, 335.
- Anderson, E., and Nicolson, H. (1950) Acta Physiol. Scand. 43, 355.
- Baker, W. (1955) J. agric. Sci. 45, 433.
- Baker, W.D. (1957) Am. J. Physiol. 191, 393.
- Bonhoure-Messing, H., and Huismanborger, G.L. (1951) J. Clin. Endocrin. 20, 626.
- Brit. Vet. Codex. London Pharmaceutical Press. 1950.
- Drody, S. Biomechanics and Growth. New York: Methuen Publishing Corporation. 1950.
- Bunker, J.P., Decker, H.H., Buzigge, D.D., Baerwalter, W.R., and Bussey, R.A. (1951) J. Pharmacol. & Paper, Theory. 103, 62.
- Burgo, Gloria (1949) Rec. Bull. No. agric. Eng. Sta. No. 433.
- Congdon, J.H., Pownall, R.H., Rubino, A.J., Bridges, H.A., and Guileen, B.Macabelli. The Lung (Clinical Physiology and Pulmonary Function Tests). Chicago: The Year Book Publishers Inc. 1955.
- Conney, H.J. Microdilution Analysis and Volumetric Factors. London: Crosby Lockwood. 1950.
- Couzannet, A., and Pongos, H.A. (1942) Proc. Soc. Exper. Biol. & Med. 46, 462.
- Craigie, A.H. (1947) Am. J. vet. Res. 8, 247.
- Craigie, A.H., Johnson, R.B., Blackman, B.G., and Coffield, F.M. (1949) J. Amer. vet. med. Ass. 104, 135.
- Gallen, C.E., and Kunkle, E.P. (1929) J. Mol. Chem. 53, 545.
- Dale, H.S., and Drody, S. Pyeocoding Book, A.V.M.A. (1953) 196.

- Dale, H.B., Gobaudson, O.K., and Brody, S. (1954) *Am. J. vet. Res.* 15, 297.
- Davenport, H.M. The ABC of Gold-Berg Gastroscopy. Chicago, Illinois University of Chicago Press. 1950.
- Dojoves, P., Labrousse, Y., Reynaud, F., and Tellion, A. (1957) *J. Physiol. (Paris)* 49, 215.
- Dovine, J.O., Kunkel, H.E., and Plotzinger, D.J. (1958) *Anesthetology*. 19, 12.
- Boyle, J.T., Robertson, J.L., Warren, J.H., Dohmeder, D.K., and Reynolds, M. (1953) *Vet. Rec.* 17, 23.
- Delpy, R.D., and Conaco, J.H. (1947) *Proc. J. Physiol.* 140, 277.
- Dawson, J.V.C.A., and Brookway, J.H. (1959) *Brit. J. Nutr.* 13, 43.
- Buller, U.S.V., Lijstewerk, G., and Rotteman, V. (1939) *Skand. Arch. Physiol.* 59, 132.
- Randall, J.B. (1955) *J. Physiol.* 130, 162.
- Randall, J.B. (1956) *J. Physiol.* 136, 300.
- Robert, R.H. (1956) *Vet. Rec.* 68, 692.
- Grey, J.S. Pulmonary Ventilation and its Physiological Control. Springfield, Illinois: Charles C. Thomas. 1950.
- Maldone, J.S., and Randell, J.B. Respiratory. Oxford: The Clarendon Press. 1935.
- Hall, L.W. (1957) *Vet. Rec.* 69, 762.
- Hansson, O.H., and Stohmann, H. (1958) *Nord. Vet.-Med.* 10, 469.
- Hansson, O.H., and Obel, H. (1958) *Nord. Vet.-Med.* 10, 457.
- Hornbake, A.S., and Sandoy, J. (1924) *J. Biol. Chem.* 61, 695.
- Hudson, H.H. (1954) *J. Comp. Path. & Ther.* 54, 213.
- Nonkols, P. (1938) *Deutsch. Tierarztl. Woch.* 46, 83.

- Raffaele, H.J. (1949) *Balt. Vet. J.* 211, 440.
- Schoenith, W.F.H. (1956) *Balt. Vet. J.* 212, 431.
- Schoenith, W.F.H., Sonnenburg, P.H., McNulty, W.X.H., McAllister, H., and Uegelman, G.H. (1957) *Vet. Rec.* 116, 1329.
- Schoenith, W.F.H., McNulty, W.X.H., and Grayburn, G.H. (1954) *Vet. Rec.* 66, 665.
- Sonnenburg, H. (1953) Personal communication.
- Uenzel, W. (1943) *Swiss. Vet. Med.* 67, 365.
- Kerfootson, C.J. "Gases and Vapors" in Dill, V.A. ed. *Physiology in Medicine*. New York: McGraw Hill Book Co. Inc. 1954.
- Wellenreiter, R.E., Rao, D.H.K., Polozan, M.H., and Anderson, R.S. (1953) *J. Anim. Sci.* 22, 573.
- McSherry, B.S., and Ordway, I. (1954) *Anat. J. vet. Res.* 25, 409.
- Meldeau, G. (1955) *Anat. J. Clin. Path.* 29, 710.
- Marie, L.P. "Respiration" in Fulton, J.V. ed. *A Text-book of Physiology*. Philadelphia & London: W.B. Saunders & Co. 1955.
- Perry, H.B. (1950) *Vet. Rec.* 62, 409.
- Quinnell, J.H. (1952) *Anesth. & Analg.* 31, 132.
- Rosenthal, T.B. (1948) *J. Biol. Chem.* 173, 25.
- Rutherford, R.C. (1948) *J. Lab. & Clin. Med.* 33, 601.
- Sampson, J., and Hayden, C.R. (1935) *J. Am. vet. med. Ass.* 86, 39.
- Sollesse, A.F., and Hendry, A. (1951) *Am. J. vet. Res.* 12, 99.
- Shock, N.W., and Hastings, A.B. (1926) *J. Biol. Chem.* 104, 565.
- Spector, W.S. ed. *A Handbook of Biological Data*. Philadelphia: W.B. Saunders & Co. 1950.

- Von Seeger (1911) *Fphys. Archiv. ges. Physiol.* 342, 577.
- Von Slyko, D.O., and Hoddle, J.H. (1926) *J. Biol. Chem.* 63, 503.
- Von Slyko, D.O., Hoddle, J.H., and Van Slyko, R.K. (1949)  
*J. Biol. Chem.* 176, 743.
- Wetts, J.G., Dunko, P.R., and Currie, J.H. (1953) *Anat. J. Physiol.* 135, 630.
- Wetts, J.G. (1953) *J. Applied Physiol.* 5, 619.
- Wiggle, R., Borbolla, H., Budde, H., and KochLaudel, H. (1955) *Fphysik.  
Arch. ges. Physiol.* 351, 211.
- Wrigth, J.G. Veterinary Anaesthesia. London: Ballilliere, Tindall &  
Cox. 1952.

THE CARBON DIOXIDE CONTENT AND pH OF THE PLASMA  
OF CALF AND THE QUANTITIES ASSOCIATED  
WITH DIFFERENCES OF RESPIRATION

By E. W. Fawcett

SUMMARY

A study was made of various methods of obtaining blood samples from cattle and a technique evolved whereby arterial blood samples could be obtained from the brachial arteries of cattle of all ages.

Examination was made of the errors liable to arise when determinations were made of the plasma carbon dioxide content and the plasma pH if certain precautions were not taken in the handling of blood samples. It was shown that it was necessary to collect blood samples under a layer of liquid paraffin but this layer must be kept to a minimum thickness to prevent the absorption by it of the carbon dioxide from plasma. Determinations had to be carried out within half an hour of collection of samples in order to prevent autolytic lowering of plasma pH.

Both the Gomory method and the nonconcentric method of Von Slyde were

found suitable for the determination of the carbon dioxide content of the plasma of cattle. The mean value for the arterial carbon dioxide content of 159 adult Ayrshire cows was found to be  $27.2 \pm 2.7$  m.moles/litre. No significant differences were found due to seasonal variations, lactational status or pregnancy. The mean values found for the carbon dioxide content of venous and mixed venous plasma were found to be  $27.4 \pm 1.9$  m.moles/litre and  $29.2 \pm 2.7$  m.moles/litre, respectively, and the mean value of the carbon dioxide content of the arterial plasma of 61 calves was found to be  $29.8 \pm 2.5$  m.moles/litre which value is significantly higher than similar determinations in adult cattle. The mean value of  $31.12 \pm 2.9$  m.moles/litre found for the arterial plasma carbon dioxide content of 15 dogs was significantly lower than the mean value found for the arterial plasma carbon dioxide content of cattle.

A colorimetric method for the determination of plasma pH was evolved and with this method, temperature coefficients for pH of blood and plasma of cattle were found to be 0.012 pH units per degree centigrade and 0.009 pH units per degree centigrade. Using these temperature coefficients a mean value of  $7.43 \pm 0.045$  pH units was found for the arterial plasma pH at  $35^{\circ}$  C. of 159 Ayrshire cows, and  $7.39 \pm 0.05$  pH units for the venous plasma pH of 55 Ayrshire cows. Temperature coefficients were also found of 0.0155 pH units and 0.012 pH units for equine blood and plasma, respectively, and with those temperature coefficients the mean value of  $7.43 \pm 0.045$  pH units was found for the arterial plasma pH at  $35^{\circ}$  C. of 30 dogs.

Studies were made of diurnal variations in the carbon dioxide content and the pH of the plasma of five cows. No significant variations were found over periods of up to six hours.

Respiratory and ventilation rates were measured in cattle of all ages. It was found that all cattle had respiratory rates between 26 and 35 per minute but the ventilation rate varied with body weight in a logarithmic manner so that a calf of 50 kg. body weight had a ventilation rate of about 25 liters/minute while a cow of 500 kg. body weight had a ventilation rate of 100 liters/minute.

With methods developed and normality defined, studies were made on three types of disturbance of respiration of cattle.

General anaesthesia caused a depression of the pulmonary ventilation of cattle which was greatest when Fluothane was the anaesthetic agent. Horses, dogs and sheep also showed a depression of ventilation when anaesthetised with Fluothane, sheep remaining unconscious. The plasma carbon dioxide content of anaesthetised animals rose above the levels found in unanaesthetised controls and the plasma pH fell below the normal values so that a respiratory acidosis was produced. Recovery from Fluothane anaesthesia was rapid; the ventilation rate quickly returned to the pre-anaesthetic control values, the plasma carbon dioxide content fell and the plasma pH rose. Plasma pH usually returned to the pre-anaesthetic control value before the plasma carbon dioxide content.

Promocaine calves showed an increase in both respiratory and ventilation rates, the decrease in the respiratory rate being greater so that the tidal air fell. The plasma carbon dioxide content of those calves was greater than the plasma carbon dioxide content of normal calves but generally lower than the levels found in anaesthetized. It was noteworthy that with the exception of one anaesthetic, severely ill calf the plasma pH was never below normal which contrasted with the effects of general anaesthesia. A compensated respiratory acidosis therefore existed in these calves.

Hypoxia caused a reduction of ventilation of calves and dogs. Calves showed an increase in the plasma carbon dioxide content and a fall in plasma pH but these changes rapidly reversed when the calves were again breathing room air. Promocaine calves showed greater changes and took longer to recover.

Hypoxia by the administration of 95% oxygen and 5% carbon dioxide caused an increase in respiratory and ventilation rates and in plasma carbon dioxide content, and a fall in plasma pH. The plasma carbon dioxide content rose to the values observed in anaesthetized but the respiratory rate did not decrease to that of preanaesthetic animals.

Hypoxia in calves produced by the administration of 90% nitrogen and 10% oxygen caused initially an increase in respiratory and ventilation rates as a consequence of hyperventilation, hypocapnia and alkalosis. Ventilation decreased while the hypoxic gas mixture was still being administered so that the plasma pH fell and the plasma carbon dioxide content rose.

Appendix I

Garden City Hospital Certificate of Venous Plasma Samples  
Obtained by Puncture of the Jugular Vein  
of NonVital Arteries from the Uninervous Hand

Date of Sampling April 1, 1954

Sample No.	$\text{O}_2$ content m.moles/liter	Sample No.	$\text{O}_2$ content m.moles/liter
1	26.94	35	23.75
2	27.40	36	23.82
3	27.50	37	26.75
4	26.09	38	25.81
5	26.86	39	26.93
6	29.24	40	26.72
7	26.34	41	25.72
8	26.34	42	27.45
9	21.08	43	26.20
10	23.52	44	23.95
11	27.72	45	23.45
12	27.92	46	21.92
13	23.23	47	29.19
14	25.09	48	26.36
		49	23.29

Mean = 27.42 ± 1.9 m.moles/liter

*Appendix II*

Gaseous Metabolite Content of Mixed Venous Plasma

Sample No.	$\text{CO}_2$ content n.molee/216m
1	24.5
2	29.1
3	29.6
4	26.8
5	20.1
6	26.0
7	22.4
8	22.0
9	26.0
10	23.5

Mean = 23.2  $\pm$  2.7 n.molee/216m

Appendix D

Autosomal Recessive Canyon Migraine Syndrome  
of Healthy Argentine Cons in the University Herd

Sampling Date: Aug 21, 1965

Flor	$\text{SO}_2$ content n.moles/litre
Bentan	22.3
Brownie	27.3
Poppy	20.1
Holl	26.8
Elsie	27.9
Averil	27.0
Mo	25.3
Gloria	25.4
Sylvia	27.6
Mirado	20.3
Eva	20.9
Quack	27.7
Dinah	20.6
Peggy	26.6
Crumpie	20.9
Ruby	20.6
Juno	25.7
Stratford	27.6
Some	20.8
Georgy	20.3

Mean = 27.6  $\pm$  2.195 n.moles/litre

Sampling Date June 1963

Day	<u>90<sub>a</sub> content n.moles/litre</u>
Eve	26.4
Keto	26.2
Elsie	26.1
Ivy	24.5
Gweny	26.0
Diego	26.9
Violet	25.1
Betty	25.6
Ruby	27.0
Anne	23.7
Juno	26.9
Blush	23.6
Quiff	26.7
Quags	26.7
Rebecca	23.6
Aster	24.3
Ivy	24.7
Poppy	24.6
Seato	29.1

Mean = 26.4 ± 1.9 n.moles/litre

Sampling Date October 1966

Dog	<u>CO<sub>2</sub> content n.moles/litre</u>
Dawson	20.3
Astor	26.4
Granny	28.5
Poppy	26.5
Audon	22.6
Eugie	25.3
Bra	20.6
Blade	27.4
Koko	28.5
Archie	25.0

Mean = 27.5  $\pm$  2.244 n.moles/litre

Scoring Date January 1956

Name	222.00000 B.Miles/13.500
Betty	26.2
Arlon	26.3
Sunie	27.1
Avell	26.8
Graney	29.8
Glenda	29.0
Poppy	27.9
Mitch	29.3
Pearl	29.0
Iva	29.7
Actor	26.6
Peggy	26.2
Ivy	29.9
Sylvia	29.0
Rita	22.6
Eve	23.8
Alla	20.2
Jack	26.9
Strawber	26.8

Scanning Date: January 1976  
(contd.)

Color	$\text{CO}_2$ content μmoles/liter
June	24.6
Pony	24.7
Steering	25.6
Violet	29.0

Mean = 27.5 ± 2.53 μmoles/liter

Sample Date: May 1996

Cor	$\text{O}_2 \text{ content}$ m.moles/litre
Melissa	27.3
Quinn	31.2
Hannah	28.5
Astor	27.7
Ivy	28.6
Poppy	30.3
Audon	27.6
Eliza	24.3
Kate	27.8
Sylvia	30.1
Rosy	27.0
Allie	24.0
Stormy	24.2
Dash	25.6
Violet	25.4
Jane	24.6
Ruby	29.1
Peggy	25.6
Dinah	25.8
Quint	23.6

$\text{Mean} = 27.0 \pm 2.365 \text{ m.moles/litre}$

Breeding Date, May, 1957

NAME	$\frac{\text{H}_2\text{O}_2 \text{ reaction}}{\text{m. nitroa/146m}}$
Denton	27.0
Arter	27.0
Ivy	26.7
Sarah	26.3
Aiden	26.2
Rita	26.0
Diana	24.9
Sybil	24.4
Avril	23.1
Glorie	24.0
Stephens	23.0
Eva	23.6
Suzanne	24.5
Sodie	23.5
Ruelind	25.7
Reva	26.0
Scout	25.1
Pony	25.6
Sandra	27.6
Shoppy	25.9

Mean = 26.2  $\pm$  1.237 m. nitroa/146m

Reproductive Data January 1958

Name	<u><math>\frac{99.2 \text{ mm long}}{n=2300/14.5\text{cc}}</math></u>
Hagan	27.6
Holona	29.0
Bentoni	26.3
Astor	28.2
Suzhi	27.2
Ardon	26.8
Suzie	26.0
Rita	26.8
Ema	27.1
Gladys	26.9
Theresa	29.2
Tina	30.2
Shirley	28.2
Sally	30.7
Jack	29.6
Ruth	30.0
Strong	26.2
Alla	27.7
Sophia	30.6

Mean = 28.2  $\pm$  1.47 mm long/14.5cc

Scanning Data March 1978

<u>Name</u>	<u>CO<sub>2</sub> content m.moles/litre</u>
Bonita	25.0
Gerry	29.1
Alice	26.7
Susie	27.0
Terry	31.2
Clotue	27.1
Juno	24.0
Sally	26.5

Mean = 27.2  $\pm$  2.725 m.moles/litre

Sampling Date May 1988

Cow	<u><math>\text{CO}_2</math> content nmole/liter</u>
Bonita	23.62
Gremio	27.50
Astora	23.75
Sorah	23.62
Audra	24.50
Rico	25.87
Rosie	20.12
Ira	23.00
Eva	25.87
Bonnie	23.75
Reindeer	23.75
Sondra	26.25
Shoppy	27.50
Jane	27.00
Bally	27.75
Jack	26.25
Violet	23.75
Dutch	25.50
Suzanne	26.00
Allie	25.90

$\text{Mean} = 26.9 \pm 1.8 \text{ nmole/liter}$