

https://theses.gla.ac.uk/

Theses Digitisation:

https://www.gla.ac.uk/myglasgow/research/enlighten/theses/digitisation/

This is a digitised version of the original print thesis.

Copyright and moral rights for this work are retained by the author

A copy can be downloaded for personal non-commercial research or study, without prior permission or charge

This work cannot be reproduced or quoted extensively from without first obtaining permission in writing from the author

The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the author

When referring to this work, full bibliographic details including the author, title, awarding institution and date of the thesis must be given

Enlighten: Theses <u>https://theses.gla.ac.uk/</u> research-enlighten@glasgow.ac.uk Thesis submitted to the University of Glasgow towards the degree of Doctor of Hedicine

4 .

1

# ON DECOMPRESSION SICKNESS IN DIVERS: SOME OBSERVATIONS MADE DURING TRIALS FOR

### THE ROYAL NAVY

By

Donald E. Mackay, M.B., Ch.B., Surgeon Commander, Royal Navy

Modical Department of the Admiralty,

Ministry of Defence,

Empress State Building,

London, S.W.6.

ProQuest Number: 10647904

All rights reserved

INFORMATION TO ALL USERS The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



ProQuest 10647904

Published by ProQuest LLC (2017). Copyright of the Dissertation is held by the Author.

All rights reserved. This work is protected against unauthorized copying under Title 17, United States Code Microform Edition © ProQuest LLC.

> ProQuest LLC. 789 East Eisenhower Parkway P.O. Box 1346 Ann Arbor, MI 48106 – 1346

#### SULMARY

Diving is an ancient activity and its mein complication. decompression sickness, has been described for at least a century. Theoretically treatment of this injury should present no problems other than these of technique. In practice there are many unexplained points which east doubt on the efficiency of current methods of treatment. Diving trials using air, or oxygen and helium breathing mixtures are described as well as submarine escape training without the use of self-contained breathing apparatus. Cases of decompression sickness occurring during the trials are described and clossified as major or minor depending on the therapy used. It is suggested that this classification is less confusing than others as there is no implication of actiological feators other than decompression. "Pulmonary Barotrauma" develops by a different mechanism and is considered separately from "Decompression Sickness". The progression from dissetisfaction with recommended therapeutic routines to the development of complementary ones is shown. The application of basic principles to cases arising under conditions not proviously encountered, is described. Case notes of 136 cases treated by recompression are given as well as comments on 264 other cases. Proposed therapeutic routines and illustrated records of some of the very deep therapies are included in the Appendices.

# CONTRAVIS

Title page

Certificate

Summary

List of contents

Chapter	I	Introduction	1
		Scope of Diving	1
		Effects of Pressure	2
		Decompression Injury	3
		Pattern of Trials Work	4
		Types of Diving Considered	Ą
Chepter	II	Common Terms used in Diving	б
Chapter	III	Historical Review	13
Chapter	IV	Principles of a Diving Trial	32
		Ethics	32
		Selection	35
		Number of Dives and Divers	36
		Assessment of Cases of Decompression Sickness	37
		Problems of Treatment	39
		Classification of Cases	40
		Assessment of Trials	41

# COMPENTS

Chapter V	Personnel and Materials in the Diving Priels	43
	and all a markety and a second the	15
	Baokground	43
	Personnel Qualifications	43
	Personnel: Physical Factors	44
	Equipment	44
	The Dase	52
Chaptor VI	Important Fectors	55
	Pressure Control	55
	Time Control	57
	Depth Keeping	58
	Sea Temperaturo	59
	General Health	60
Chapter VII	Trials of Standard Air Tables	62
	Baoleground	62
	Basic Decompression Theories	63
	Selection of Tables	64
	Triel Orders Requirements	64
	Period of Trial	66
	The Divers	66
	The Dives	68

# CONTEMPS

		Effect of Duration and Environment of the Dive	75
Х.		The Tables	78
		The Cases of Decompression Sickness	81
		Symptoms and Signs in Minor and Major Cases	84
		Notes on Major Cases	87
		Therapeutlo regimes	99
		Comment	102
Chapter	VIII I	hultiple Dives	105
		Problems	<b>1</b> 05
		Notes on Tochniques	106
		Results	108
		Case Notes	109
		Therapeutic mutine	111
		Comment	112
Chapter	IX S	jurface Decompression Divos	113
		Problem	113
		Notes on Technique	114
		Results	115

# CONTEMPS

	The Dives	116
	Case Notes	118
	Comment	124
Chapter X	Submarine Escape Training	125
	Beoleground	125
	Routine	125
	Therapeutic approach	126
	Case Notes	128
	Discussion	146
Chepter XI	Deep Diving Experiments	151
	Background.	151
	Notes on Techniques	151
	Approach to the Trials	153
	The Dives	153
	a) First period	153
	b) Second period	157
	c) Third period	160
	d) Fourth poriod	165
	Decompression Sickness	169
	Case Notes	178
	Comments	220

# CONTRATES

.

Chapter XII	Discussion	222
	Onitorion for Diagnosis	<b>S</b> 55
	Differential Diagnosis	223
- *85_ **	Problems of Treatment	224
••	Alternative Treatment	<u>ss1</u>
	Pressure required for measuresion	228
	<i>Invation of Stay at Pressure</i>	229
	Breathing Gas	229
	Therepeutic adjuvents	232
	Therapeutic Decompression Tables	235
	Signs and Symptoms	238
	Double-blind trial	239
	Acclimatisation	240
Chapter XIII	Conclusions and Recommendations	242
Chapter XIV	References	247
Acknowledgements		265
List of Figures		267
Figures		269
Appendix I : 1962	Statement of Decompression Sickness	286
	Signs, Symptoms and Treatment	

# CONTENTS

Appendix	11	Section I. U.S.N. Therapeutic Routine 1958	292
		Section II. R.N. Therapoutic Routine 1956	298
		Section III. R.M. Therapeutic Noutine 1964	302
		Section IV. Trials Therapeutic Routines	308
Appendix	III	Therapeutic routine for Submarine escape training accidents	314
Appendix	IV	Some interesting Therapeutic Routines	321

~ **ì** 

### CHAPTER T

#### THTRODUCTION

# Scope of Diving

For more than two centuries diving has been carried on mainly for the purpose of salvage. For two thousand years there has been a military fringe benefit in damaging shipe of the enemy. This military application was widened considerably in scope during the 1939 - 1945 war to include such activities as pre-invasion beach reconnaisance end attack on land-based installations, as well as being developed in its original role against shipping. The effect of this expansion was a great increase in the demand for knowledge of the effect on man of this underwater environment and in the number of trained men who were thrilled by the adventure. Technical advances led to the establishment, in the last two decades, of a new sport which now numbers its participants in hundreds of thousands. Thus an activity which 25 years ago affected perhaps 500 people in Great Britain, now involves over 1,000 in the Royal Navy alone and probably one quarter of a million in this country. At the present time breathing equipment has such a short endurance that the risk of decompression injury is slight for sportsmon, but with continuing advances in material, engineering, and design problems, breathing sets are becoming available that will place the divers using them at greater rick of specific injury due to diving. These cases may occur near coastal towns or inland villages, where water more than 20 feet deep to available for diving. This work will cover some

aspects of cases at decompression sickness found in divers doing experimental work and in men undergoing submarine escape training, and more commonly known as "bends".

### Effects of Pressure

1

Any reduction in the ambient pressure of man is a decompression. By physical laws the gases that such a man had been breathing must have been in solution in his tissue fluids as well as filling his aircontaining cavities. The main physical laws observed are those concerning the solubility of a gas in a fluid; the concept of partial pressures; the relationship of volume and pressure; and the process of diffusion. A man at 33 feet in water breathing air from an adequate supply, is breathing at a pressure equal to the weight of 33 feet of water plus the whole height of the atmosphere which is equivalent to another 33 feet of water - in this case therefore the pressure is equivalent to 2 atmospheres absolute or twice the pressure exerted on a man at sca level. In response to leve of physics as the environmental pressure round the subject is reduced, i.e. as the non returns to the surface, the excess gases absorbed must either expand or come out of solution. This problem is manified with increasing depth as the total pressure is increased by 1 atmosphere absolute for every 33 feet of water. If these expanding or evolving gases cannot reach the external environment, an injurious situation may arise and any detectable ill-offects are called decompression mickness.

ō.

### Decompression Injury

Decompression Sidmess is a term which ameans self-explanatory and by implication suggests a condition which should respond to the straightforward application of the physical principles involved. Through the years and with the use of several therapeutic or preventive measures for ailments involving the teeth the paramasal sinuses, and the middle ear space to name a few. the term decompression sickness new tends to be limited to conditions where gas bubbles are presumed to be present in the tissues of the body rather than in the body air spaces. Such bubbles may arise in situ or they may be introduced into the body accidentally. One important effect arising from the actiology of the bubble is that the cas in the bubble introduced accidentally into the tissues is more likely to dissolve in the tissue fluids as these fluids usually do not contain as much due in colution as those tissues in which the bubbles develop from superscturation. In diving most bubbles introduced into the body crise from demage to the lung and es such the conditions thus produced are probably better grouped under the title "mulmonary barotrauma". The term "decompression sickness" would then be kept for those conditions presumed to be due to the evolution of gas from tissue fluids. The treatment is based on similar principles for both conditions. There have been very many communications on the ailmont and related matters - more than enough to fill three large volumes of a bibliography. (85. 86. 87) Nowover, surprisingly few accounts of the

treatment of disease as seen in divers have been published compared with the numerous papers published dealing with the conditions as it occurs in tunnel workers, caisson workers, and in eviation.

### Pattern of Trials Work

The majority of the cases described in this paper occurred in the course of work associated with the development of safe diving routines for mon. The trials followed experimental work on animals, which was designed to test the theoretical calculations, and consisted of simulated dives in compression chambers followed by actual dives in water. The success or failure of the routino depended on the number of cases of decompression sidmess that developed; too many cases meant that the routine was too dangerous while no cases meant that too many safety factors had been included. Once a routine had an incidence of 1 - 2% cases, standard safety fectors would be added and the routine premulgated for general use. The trials medical officer was responsible for assessing each dive and for carrying out any examination or treatment. He was concorned in all matters that affected the divers and gave his advice in the planning stages of each trial as well as giving his opinion on progress.

### Types of Diving Considered

Five types of diving will be discussed. Firstly in the normal dive the diver descends breathing the appropriate mixture to his depth, carries out his task, and ascends carrying out a decompression routine

-4-

at verious stages in the shallower part of his ascent. In a second type, after carrying out his task the diver ascends directly to the surface and enters a compression chamber where the pressure is increased and then released in a pre-determined pattern - this routine is called "surfece decompression". Combined diving is a third variant and consists of a dive followed after a short interval at atmospheric pressure by another dive but the ascent from this second (or any subsequent dive) is so made that any remaining excess gas still procent in the diver's tissues from the preceding dive is taken into consideration. The fourth type of diving is known as "deep diving" and, while the exposure to pressure and time follows the normal descent, work, and then ascent pattern, the depth of such dives means that air cannot be used as the breathing medium by the divers. The fifth type. "submarine escape" dives are those in which the escaper or "diver" is esposed to rapid increases in pressure and after a minimal time at maximum depth. he carries out his ascent without breathing equipment. Circumstances have led to the problem being studied by the general practitioner away from skilled anoillary help, and in the nature of the activity producing the injury. this will almost invariably be the situation in which the doctor has to treat his patient. Fol and Watelle (124) in their paper which is comonly considered the first serious study of calcoon workers opologised for their work as it had been done "sans plen, sans programme. sans ordre" as there had been no intention of publishing it. Over a contury later, their continents are cohoed.

-

#### OHAPTER II

### COMMON TERMS USED IN DIVING

In very general terms a "diver" carries out a "dive" when a man. suitchly equipped, goes under pressure greater than atmosphere for a time. This dive has three phases, one of increase of pressure. or "compression", one of duration at maximum pressure or "bottom time" and one of return to atmospheric pressure. Usually the first two are together called. "the dive" and the last is the "decompression". From the point of view of the diver, the most important pressure is that of the depth at which he is working. This pressure is related to atmospheric pressure at sea level which is treated as unity or "1 atmosphere absolute". Atmospheric pressure at sea level varies with veather conditions and latitude but these fluctuations are small relative to the large pressures involved in diving and may be ignored. It is convenient to express this unit in round figures as 15 pounds per square inch, 1 Kilogram per square centimetre or 33 feet of water and such values can be referred to as "an atmosphere". It follows that a diver at 33 foot in the sea is subjected to an ambient pressure of 2 atmospheres absolute. and at 100 feet. to an ambient pressure of 4 atmospheres absolute. In most prossure work, gauges read zero at sea level so that gauge readings are always 1 atmosphere or its equivalent less than the absolute pressure, thus the above diver is subject to a prossure of one atmosphere and 3 atmospheres gauge respectively. For convenience, the pressures in diving tend to be expressed as x feet of

water (gauge understood) or y stmospheres (absolute understood) whomeas compressed air working pressures tend to be described in pounds per square inch (gauge understood).

The situation is rapidly reached where it is injurious to return directly to atmospheric pressure because of the onset of symptoms end signs of "decompression sickness". To evoid this the diver follows a "decompression mutine" described in "decompression tables". (Table II.1). Decompression tables are lists arranged in increasing order of depth in 10 feet intervals; in each interval, the duration of dives are arranged in lengthening order of time and usually in multiples of 5 minutes. Opposite a depth-time combination a row of figures gives the shallowest depth the diver may reach after a dive of this combination and the minimum time he must spend there to prevent the onset of decompression sickness before proceeding to his next depth/time newse; such depth/time pauses are "decompression stops" or "stages". The details for this "stage" decompression zoutine for a dive of a particular depth and a particular duration is referred to as a "schedule". If the decompression is carried out continuously the decompression routine is known as a "bleed" of so many units of pressure per minute. Decompression sidmess has various synonyms such as "caleson disease", "bends" or "compressed air illness" in such industries as tunnel building and marine selvage and such as "aero-embolism" and "barotrauma" in aviation.

-----?m

Depth not	Duration Time leaving surface		Stoppages at different (minutes)		depthe	Total Time for		
(ft.)	ascent not exceeding (minutes)		501	40*	301	201	101	pression (minutes)
80	Limiting	30 40 50 55 60 70 75	1 * 4 * 4 * 4 * 4 * 4	<b>₽</b> <b>₽</b> <b>₽</b> <b>₽</b> <b>₽</b> <b>₽</b> <b>₽</b> <b>₽</b>	音音音音音音	1100000	- 5 5 10 15 20 25	- 5 10 15 20 25 30
¢.	Lino	80 90 105 120 140 160	<b>₹ * * *</b>	111550	500000	500050 100050	30 35 4 50 50 50	40 50 65 85 100 <b>13</b> 0
90	Limiting	2532595565	<b>\$ \$ \$ \$ \$ \$ </b>	<b>₽ ₹ \$ \$ \$ \$ \$ \$</b>	11111100	1 1555555	5 5 10 15 20 25	- 5 10 15 25 30 35
	51no	70 75 80 90 100 100 120 135 150	11111100	11110000	5555555 <b>5</b> 2255	10 15 25 30 35 35 40	30 30 35 45 45 45 50 50 50	45 50 60 70 80 100 110 125 140
Table II.1								

AIR DECOMPRESSION

Decompression Table

Extract from R.N. Diving Manual (153) showing schedules for dives to 80 and 90 feet. Dives below the limiting line should only be done in exceptional circumstances. In a previous edition (132) all dives above the limiting line were grouped in Diving Table I and those below the limiting line in Diving Table II.

When a dive is carried out in the sea it is known as a "sea dive" or "wet dive" and the expression of the pressure as depth in feet of water is logical. Certain pressure vessels are made in which the diver may carry out a dive where the increase in pressure is obtained by supplying gas, usually air, from a system of compressors and storage cylinders. For convenience in diving research, the pressure is often also expressed as depth in feet of vater. Such vessels are known as "compression chambers" or "pots" and the divor carries out a "chamber" or "pot" or "dry dive". In commercial work the terms "caisson" or "tank" my be used for such vessels. The primery function of these chambers in neval work is the treatment of decompression sickness for which the patient is placed inside the "pot" and "recompressed". After relief of his injury the patient returns to the surface in a decompression routing intended to prevent reopportance of his signs and symptoms and generally referred to as a "theraveutic decompression".

When the decompression tables in general use are those for diving no deeper than 200 feet, they are called the "standard air tables" as air will be the breathing mixture used by the diver to sustain his life while underwater. If a diver carries out first one dive and later more dives, he is said to carry out "combined diving" or "repeated diving" or "multiple diving". Various instructions exist for discovering the decompression routine for such dives and an important point is the "surface interval" or time spent at atmospheric pressure between two dives. If for some reason the diver cannot carry out the appropriate

<u>.</u>

decompression for his dive and is forced to return to atmospheric pressure he must be recompressed in a chamber within a very short time to carny out an appropriate decompression routine and this mancouvre is referred to as "surface decompression".

"Deep diving" generally pefers to all dives deeper than 200 feet and is so distinguished because air is unsatisfactory as the breathing medium for the diver. This is due to the amount of nitrogen present which causes "nancosis" and the danger from the amount of exygen present of "organ poisoning". As a result the propertion of oxygen is reduced and the required bulk of the breathing gas usually then consists of helium. Such diving in the see may involve the use of a chember which can be lovered into the vater and which can be kept empty of water but. because it can also be closed with the diver inside and he can carry out his decompression routine while so confined, it is called a "submersible decompression chember". In two ships, one British and one Swedich. this submersible decompression chamber can be placed on top of a compression chamber and, once the pressure is equalized, the diver may "transfer under preseure" from relatively oranged to more comfortable surroundings to continue his decompression. If a diver is required at any time to very the gas mixture that he breathes, for example if he breathes 100% oxygen during the final stages, he is said to be carrying out "oxygen decompression". In experimental work "oxy-helius" is the colloquial torm for respirable oxygen and helium mixtures and is only qualified by the percentage of exygen e.g. 5% exy-helium. In such work

-10-

also the depths may be as shallow as 35 feet but as the aim is to dive to great depths, all ony-holium dives are considered "deep dives".

Once a technique of diving has been tried under very close supervision and in controlled conditions, it is taught in the training schools and can be used by men on their tasks. Shus it is changed from en "experimental technique" to an "operational routine". Gertain modifications may be made to decrease risks. For example in looking up decompression tables for a dive. if the man has reached an exact doubh or an exact duration. it is customany to use the decompression achedule for the next greater depth or longer duration. The operational teams consist of "Clearence Divers" who are highly trained volunteers. They have been taught to use self-contained breathing apparatus which is complete with its own gas supply whether it be air. or exygen. or oxygen and mitrogen mixtures and which may be open circuit (i.e. exhaling into the vator) or closed circuit or semi closed circuit. In addition they can use equipment, usually open elrouit, where the gas, usually sir. is supplied from the surface thus avoiding the problems of ondurence. These men are popularly known as "frogmon" and are fulltime divers. Other neval ratings carry out their own various duties such as mechanics, storemen, communicators and cooks and dive as a parttime estivity. Such divers are usually limited to self-contained or surface-supplied air equipment and are "Ship's Divers". The classical or"Stenderd Diver" with helmet and heavy boote has become redundant in the Royal Navy but he was used in most of the trials described here

because of the case with which his depth could be controlled.

The case of a diver's progress under water is affected by his buoyancy which can be increased as "positive" buoyancy by his equipment and gas contained deliberately or accidentally in his clothing. It is customary to have the diver and his gear slightly "negative" or "neutrally" buoyant by making him initially "positively" buoyant and then adding lead weights to over-compensate or balance this. These weights can be "ditched" or rapidly released as a safety measure so that the restoration of his positive buoyancy will bring the diver back to the surface. The importance of buoyancy in this work is the effect on the offert required by the diver to carry out his descent and ascent as well as on his ability to keep a constant depth during his dive.

-12-

#### CHAPTER III

#### HISTORICAL REVIEW

It is a metter for surprise that a subject which is normally dismissed in most modical test books in a peragraph should have an extensive bibliography. Most of this literature is contained in the three volumes of the "Bibliographical Sourcebook of Compressed Air, Diving, and Submarine Medicine" which lists more than 10,000 references up to 1955 (85.85.87). Even at that, it is stated by the compiler (85) that the list is not comprehensive as papers of the last century and the early part of this one are inserted as much for their own lists of references as for their intrinsic merits.

Han has long wished to study the response of his kin to exposure to altered onvironment - particularly when the reaction was not a simple reversible one as superficial study suggested. When the problems started to involve thousands of people where previously the victims had been numbered in tens, as happened when civil engineers started tunnelling and when aviation seared towards space, the stimulus to study was great. This meant that the physiology, toxicology and traumatic possibilities of atmospheric pressure, exygen, nitrogen and carbon diexide when altered from normal values, came under the scrutiny of hen in several disciplines of research and clinical work both for the study of avoidance of the problems as well as for the potential benefits to the individual, his well-being and his occupation. Behnke (18) considered that the mascent bubble had replaced the spirochaste as the great impersonator

and the bulk of the literature describes the effects of the presumed attacker on the various systems of the body, rather than on the whole individual.

Very zerely is there one single paper which is a milestone; 111 this field more often a series of articles is collected in one volume either by one author or on one topic. Pride of place must go to Boyle (23), who in the course of studies on respirable atmospheres in evecuated chambers, recorded, in passing, the appearance of a gan hubble in the eye of a viper. While it may be true that this bubble was due to vaporisation (2), rather than to superscturation of the aqueous humour, it is the first recorded observation of a bubble caused by reducing the emblent pressure. Injuries, including decompression sickness. must have occurred in sponge and year! divers throughout the ages but the records are vague and more suggestive of the effects of unbalanced excess external ambient pressure at dopth crushing the air-containing body cavities, especially the thorar and commonly known as "squeeze", than of decompression sickness. Triger (144.), who developed the first practical calsson, is credited in an ennotation of an address he gave in 1845, with the first description of a case of causen disease. Pol and Watelle (124) in 1854 described their experience with conversial calsson wonkers in which they took advantage of the vorkers' own connents, dolaberately to re-introduce a patient into compressed air as treatment for his condition - the first therapeutic compression - as vell as recommending a period of half-en-hour for return to surface for

-12-

all workers the first decompression routino. Many observers in this field produced concepts of the causation of decompression sideness which varied from the mechanical effects of high pressure and dense gas on circulation and respiration to more basic physical effects due to solution and evolution of gas in fluids. Hoppe-Soyler (88) in 1857 is credited by Hill (84) with being the first to point out that death occurred because of distruction of the circulation by bubbles and to explain that this sort of condition was due to incleanate time for gas to escape from venous blood during its passage through the lungs. It is difficult to accertain the order of priority of the many porkers, mainly French and Contral Buropeans, in the various steps taken as often the vital point is obscured by the literary style as well as by the vast amount of eneculation on the toplo. However Paul Bert's work culminating in the publication of his book "La Pression Barometrique" in 1878 (18) stands out like a lighthouse in thet his experiments established beyond argument that decompression sidcaess was due to gas absorbed during exposure to raised pressures and then released during the return to atmospheric pressure. His most prectical point was his suggestion for a steady rate of decompression from the working pressure so that the decompression time was related to the working pressure instead of being a fixed duration. These principles were gradually accepted and work by various groups has been directed to establishing the various durations of exposure and total decompression and the rates of pressure change as modified by energies, by temperature, by ventilation, by the respired

-15-

gases (from pure oxygen to oxygen and holium mixtures) and by environment from see as in diving, to air, as in comporcial works such as tunnelling.

The actiology of decompression sickness is the formation of intra and entravascular bubbles as described by Gorsh and Marvey and their co-versers (30, 60, 61, 62, 65, 68, 69, 90). The only recent query of the bubble actiology has been raised by End (58) who considered that high pressures led to intravascular sludging of the red corpuscies which blooked the circulation and led to the secondary appearance of bubbles as suggested by Swindle (143) but this work has not been confirmed nor does it appear to have been followed up.

On decompression theory, several groups have made major contributions the carliest being Heller, Mager and Von Schotter (71) who as a result of experiments and a review of the past in their exhaustive volumes recommended a decompression time of 20 minutes for exposure to each atmosphere greater than atmospheric pressure. Heldano with his coworkers, especially Boycott and Damant (22), reconsidered the stage method already touched on and discerded by Bort (48) and produced the first really practical diving tables based on work which showed that it was always safe to bring a man to half his absolute pressure if he had been saturated at that pressure - the 2 : 1 ratio concept. This meant that a diver could be raised to a depth for a period during which the excess gas would be evolved from his body thus enabling him to proceed to a shallower depth. It was for more preciseal to raise a diver through

-16-

the water in staps then to provide a constant rate of travel. Hill carried out a considerable amount of work on decompression especially with Lacleod which he gathered together in his book "Ceisson Sickness" (8) but his work was more valuable for his configmatory experiments then for any original concepts, he being a proponent of the blood rathor than the stage method. As a result of his work with Heldene. Demant (39) produced therapeutic decompression tables for compressed air workers suffering from celsson discess which introduced a decreasing rate of bleed. Smith (139) had previously suggested that a special treatment chamber be provided at compressed air vorkings but it seems that Moir (120) of Great Britain as a member of the consulting ongineoring 21m supervising the work on the Rudson tunnel in New York, was actually the first man to have such a chamber provided and he claimed a dramatic reduction in fatal cases of decompression cickness once its use was established. That there was a need for such measures is shown by the many sories including both fatel and non-fatel cases reported which ranged from the 63 cases by Pol and Watelle (124), in 1854, 25 cases by Littleton (99), 119 by Woodward (151), 3,692 by Keeye (90) to 689 by Cempbell Golding ot al (20) in 1960. Steadily since 1854, prodded perhaps by compensation claims for doath and disability, industry has gradually increased the decomprossion time from a fixed 30 minutes irrespective of pressure to the varying scale of the current Compressed Air Special Regulations (119) with its varying rates to a slowest of 9% minutes per pound per square inch (usually regarded as 10 minutes per pound in practice)

-17-

In the Haldonian theory it was postulated that there were hypothetical tissues of various half-times to seturation from which it ves possible to calculate diving tables on the assumption that each tissue could tolerate a supersaturation equivalent to double that of Saturation at ambient pressure. Through the years this concept has been eleborated by many workers who, as a result of experience, have increased the number of tissues, and varied the ratio for each tissue; novadays some workers have a multi-tissue model each tissue having a different half-time with its orn ratio which varies with the absolute pressure and with the inert gas being broothed. Hempleman and Rashbass and their colleagues (34. 35. 36. 37. 38. 54. 73. 74. 75. 76. 78. 126. 127) showed that Haldane's assumption (22) did not hold at greater pressures than 2 atmospheres absolute and they tried to simplify the procedure by treating the diver os if he was composed of one homogenous fluid which could safely tolerate a constant excess ros tissue tension. These two approaches led to the introduction of diving tables based on the ratio concept by Van der Aue et al (147) and by Dwyer (52) in 1957 into the United States Navy (145) and thence in most navies, and tables based on the single tissue idea by Grocker (35) into the Royal Nevy in 1950 (132). (A comparison of these tables is part of this work). A Swiss methematicion and diver, Hennes Keller, startled the diving world in the early 1960's with his spectroular dives to various depths to 1.000 feet (91); his calculations seemed to be based on a 3 tissue model instead of the multiple tissue idea and on the use of maximal

-18-

quantities of exygen in his breathing mixtures to produce maximal gradients for elimination of absorbed inert gas. Regrettably he has endeavoured to keep the details of his ideas secret for commercial reasons but it is generally felt that there is no real now principle involved but rather a difference in technique and acceptable safety standards.

The treatment of decompression sickness in divers continued along the lines suggested by Damant (39). Shilling (137) collected a large number of reports of cases occurring in divers and compressed air workers and he produced a classification, based on the system of the body apparently affected, which has stood the test of time. Erdman (57) had preceded him with a similar but less comprehensive classification limited to tunnel workers. Behnke joined the team working on these problems in the United States Navy and has taken part in very many studies over the past 30 years. On the problem of decompression sickness his most lasting work may well be his share of the project that led to the production of a new therapeutic routine for divers (51), later evaluated by treatment of 113 cases: - this work was so impressive it led to the adoption of the recommended tables with few modifications by almost every Navy in the world including the Royal Navy. (See Appendix II) (Bohnke incidentally in the 1940's visualised the use some day of a compression chamber for the treatment of eviators' decompression sickness (6) but this was not done till reports of dramatic successes in the late 1950's led to a drastic re-assessment of this form of therapy by

-19-

eviation medicine experts). The subject was surveyed and brought up-todate in 1954 in the book "Decompression Sickness" (59) where the emphasis was more on the condition as met at low pressures by eviation personnel but Behnke (5) contributed his experiences with high pressure work.

Indley (46) had summarised the basic principles of therapeutic recompression as follows: "The treatment of compressed air illness is inmediate recompression. Becompression is preferably done in a chamber especially designed for the purpose into which air can be pumped aspidly up to any required pressure and in which the rate of decompression is under exact control". In the 30 years since that passage was written. the main changes are that one would now talk of "decompression sickness" to include those cases arising while diving with breathing gases other then eir and one would add to the requirements of the chember the facility of access by a medical attendant as needed. 1.e. the chamber must have an access lock large enough to take a man. That recompression. when adequate, is the right treatment when treating cases of recent onset, is dramableally obvious to anyone who has worked in this field; as exemplified by the late Sir Robert Davis (42) who wrote "No one who has seen a victim of compressed air illness, gravely ill or unconscious. put into a chamber and brought back to life in a few minutes by the application of air pressure will forget the extreordinary officiency of recompression or be backward in applying it to a subsequent case of illness". This can be contrasted with the inexperience shown in a recent report by Emmary et al. (24) who treated cases of neuroeirculatory

-20-

collepse in decompression sickness by infusions and pressor drugs although a compression chamber was evallable in the same building.

The four variables of treatment are the pressure required, the stay at that pressure, the memor of the subsequent decompression, and the gas mixture breathed during the decompression. Any other treatment given in addition to recompression is encillery and is more akin to good nursing with its benefits then to specific therapy. Keavs for exemple, in his large series (90) followed the practice of some predocessors and recommended that the pressure be raised to that at which the tunnel worker had been exposed during his shift - 1.e. the working pressure. Damant (39. 40) recommended that the pressure be raised to that level at which the signs and symptoms disappeared - i.e. minimum effective pressure or depth of relief. The subsequent decompression was a steady bleed in Keay's portine and an increasingly slow "bleed-off" of pressure in Damant's method. In both methods air was breathed during the entire procedure. Gulffiths (28) used to recompress his pain cases which he called Type I Decompression Sickness - to 2 pounds per square inch above working pressure which he maintained for a period of 15 minutes after the disappearance of all signs and symptoms. If there was no relief, he then increased the pressure by 2 pounds per square inch every 15 minutes till relief occurred. He then decompressed fairly quickly to half the absolute pressure or 12 pounds per square inch (gauge) which ever was the greater and then continued decompression by reducing the pressure by 1 p.s.i. during 15 minutes. However he now uses the minimum

-21-

effective pressure for relief waiting there for half-an-hour after relief of symptoms before decompressing in the same manner (67); this routine has led to fewer recurrences. In more gravely ill patients - Type II he still returns to working pressure or higher if needed, and again after the disappearance of signs and symptoms (which may be several hours in clearing), decompression is started at 15 minutes per pound pressure to 15 p.s.i.g. and then 30 minutes per pound pressure to atmospheric pressure with a 4-hour pause at 12 p.s.i.g. and 2-hour pauses at 8, 4, and 2 p.s.i.g. In essence these routines started as "working pressure" methods but have been convorted to "minimum effective pressure" methods.

In contrast the U.S. Mavy Therapeutic tables (145) use the "minimum effective pressure" approach but once this has been found this is used to select an increment of pressure where the man is kept for half-an-hour before commencing decompression in stages gradually increasing in duration towards the surface. In addition, the use of 100% oxygen by the patient is optional although strongly recommended. These tables were based on decompression schedules devised to prevent the appearance of decompression schedules devised to prevent the appearance of decompression schedules and symptoms did not develop, then the table would probably be effective as a therapeutic regime in actual cases. The maximum pressure required was fixed at 165 feet (6 Ats.Abs.) as it was calculated that a further increase of 4 atmosphere or so would not have an appreciable effect and higher pressures introduced other complications as well as delaying the time

-22-

when 100% oxygen could be used (7). (An experienced medical officer could vary these limits if he felt the case needed such increases). During the treatment of the more seriously ill cases, periods of continued exposure to certain pressures for 12 hours and 6 hours were introduced to try and saturate the patient at that pressure and reduce recurrence by allowing time for elimination of the bubble. (The Haldanian concept of ratio was modified in this work). Henderson (82) had foreseen the possibility of decompression sickness in eviators in 1917 but Boothby and Lovelace in 1938 (21) were the first to attribute a case to decompression sidmess rather than to anoxia or some other diagnosis. Armstrong (2) described the similarity of the condition in aviators and divers but he was satisfied that the return to lower heights or ground level was adequate therapy. Due to his work attention was drawn to the problem during the war when bombers could not descend while over hostile territory to ease efflicted aircrew. Amongst the possible solutions Lawrence et al (97) reported on a "bends bag" into which the injured man could be placed and the pressure raised to 3.5 pound per square inch (equivalent to an altitude of 22,000 feet). This latter approach is an example of the use of the minimum effective pressure technique while the former methods can be considered as examples of the maximum evailable pressure routines. Behnke's suggestion (6) for compression of certain cases of aviation decompression sickness to pressures greater than atmospheric was a reasonable extension of the minimum offective pressure technique.

-23-

Several other matters affect the therapeutic decompression. The use of 100% oxygen has been investigated by several people (89. 157. 100) the most comprehensive work being by Behnke who believed it encouraged elimination of the inert gas in the noxious bubble (9). Donald (46) among others (122. 130. 158) believed in the occurrence of decompression sickness under certain circumstances when the diver was breathing only 100% oxygen. Oxygen and helium mixtures have been recommended (134. 135. 145) but the main use is for the comfort of the patient as difficulties in pulmonary ventilation are thought to be eased and the abolition of inert gas narcosis leads to a better attitude by the victim. Exercise during decompression has been the subject of controversy, being recommended by the Heldene group (22) and the Hill group (84) emong others (10). Van der Aue with his group (146) conclusively showed that exercise increased the incidence of decompression sickness. It would appear that quiet novements to keep the circulation moving in all parts of the body is preferable to stagnation but that it is casy to overdo it and make an attack of decompression sickness more probable. Thus exercise during a treatment routine would also seem to be contra-indicated.

Eventually, the collected experience of several years of treating decompression sickness in divers was analysed by Slark (138) and by Mackey (106) in Britain and by Rivera (131) in America. In essence these papers showed how errors in application of the tables could be made as detailed by Behnke (5) and the opinions on need for future work

-24-

vore similar. The most startling finding was that the treatment routino for a gravely ill person could on occasion produce decompression cickness in an attendant who had been with the patient throughout but who had not had a prossure exposure before treatment started. i.c. the routine not only did not prevent the occurrence of decompression sickness in a man who had not dived but an overt attack could be produced. In the meantime experimental work on deep diving was restarting in several countries and other ideas on therapy were developing. For example Erde (56) has investigated the use of hypothermia combined with high prossure for neurological cases of decompression sickness with delayed treatment with inconclusive results and Barthelmy (3) in France has investigated the use of Heparin which he has shown has a protective offect on onset in dogs but no real effect on treatment in man. The use of glucose coline infusion or plasma was recommended by Behnke (6) and used in a very difficult case by Cockett and Nakamura (31). Nicotinic acid has been tried with doubtful offect (153) Beton (53) has published his experience on the treatment of goats with decompression sickness in experimental work. His method depends on the minimum effective pressure which he meintains for a few minutes followed by decompression by a plow bleed until the animal becomes restless. The decompression is then stopped till the goat settles down when the bleed restarts; eccasionally animals with spinal cord lesions do not respond and have to be destroyed. Although the indications would be loss definite as divers tend to concerl the restlessness due to recurrence of symptoms till they are unbearable,

-25+

his approach is worthy of attention. During current work in the U.S.N. Workman and Goodman (196) have produced a routine using minimal effective pressures, work matched by Barnard and Mackey (108) in the N.N. Both groups use much lower pressures for selected cases than the official tables and take much shorter times to carry out the subsequent decompression.

There are two main problems of decompression sickness in experimental work; first the possibility of the injury arising at greater prossures then published routines envisage; and second, the pulmonary irritant (45, 140) effect of dry gas containing high partial pressures of oxygen on the divers in the consequent prolonged decompression. The section on deep diving will exemplify the trial and error routines developed along the lines of minimum effective pressure, high exygen levels, bleed-offs and prolonged pauses based on the experience of the past.

At this point it is worth considering the special problems that exise in diving to depths greater than 300 feat as far as physiology is concorned. The fact that exygen at a partial pressure of 2 atmospheres in the sea could be sentely toxic was described during the Second World War in the work of Donald (44.45). At 300 feet, or 10 Ats.Abs. the exygen, in the 20% exygen, 80% nitrogen mixture that is air, reaches toxic levels so that it is necessary to reduce the exygen content of respired gases at this and greater depths. In addition, the nitrogen content has an effect; whether direct, or indirect this effect is

-26-
narcotic in nature and has been a highly controversial subject emong research workers over the past 30 years. The current main protagonists are Bennett (13) in Great Britain and Carpentor (29) in the U.S.A. on the side of a direct effect on corebral colls and Sousing (136) in Germany and Duhlmann (25) in Switzerland who prefor the theory that the offect is due to disturbed elimination of carbon dioxide. As result of recent work by Bennett (14. 15. 16) and further consideration by Buhlmann (26) it seems that there are effects not entirely attributable to high cerbon dioxide levels and that nitrogen may exert its effect by impeding carbon diexide elimination in brain stem cells. Riessling and Mag (93) showed that this effect was detectable in impaired performance at 100 feet while Bennett and Glass (17) found E.E.G. changes at very challow dopths. Miles and Mackay (117) showed that the impairment was not so great as to endangor the life of the self-contained diver at 180 feat. These findings had confirmed the experience of deep divers of the past who had limited the normal maximum diving depth on air to 250 feet. When Bhd (55) introduced helium as the gas to supply the volume required for respiration on the suggestions of Hildebrand ot al (83. 134. 135) mental clarity and manual destority returned to the deep diver and it was reasonably straightforward to produce oxygen and helium mixtures which vould not be toxic. Molium had different solubility and diffusion coefficients to nitrogen (12) which meant that values already found for nitrogen decompression had to be the subject of much experimental work but the principles remained. Reasons of supply of the quantities needed

-27--

meant that the field was investigated mainly by groups in the United States of America. The other noble gases, xonon, krypton, argon and neon (14, 15, 16) have also been investigated and have been found to be more narcotic than nitrogen and as such impracticable. Hydrogen has many attractive qualities and Zottorstrom (19. 159) in Sweden carried out several successful dives before he lost his life through the inexperience in diving techniques of his surface attendants. Tho explosive properties of hydrogen entail certain precautions, and tedious procedures during the dive so that little interest has persisted in this cheaper gas. The use of mixtures containing oxygen. nitrogen. and helium has been suggested (149. 155) but so far no reports of practical experience with such multiple gas mixtures are available. The effect of the use of helium on decompression time has generally been, that co saturation occurs more quickly with helium than nitrogen, short duration dives have a longer decompression time then an equivalent dive on air but that longer dives reach a stage where an increase in duration does not lead to much increase in decompression time (121. 132). This concept was challenged (49. 50) and the results were available at the start of the deep diving trials. The trials described in this theois in the section on deep diving show that helium dives tend to have more severe symptoms then air dives (79. 80. 81. 103. 104. 105). It is difficult to be precise on the cause of this because of the other fectors of total duration of the dive and the high pressures of exposure of the diver.

-28-

So far the term decompression sideness has covered cases arising during a reduction in pressure when it is presumed gas appears in various tissues by evolving from supersaturated solutions although the reasons and mechanism for this process are not precisely known. Ges may also be introduced to the tissues directly without the intermediate stage of solution and evolution. as happens occasionally in clinical manosuvres at atmosphoric pressure. For many years it has been generally accepted that restriction of free expansion of gas in the lungs may lead to general or local high pressures with consequent rupture of lung tiesue and aproad of gas intravaccularly or interstitially (4. 28. 33. 47. 95. 98. 125. 125). The practical importance of this point may be that as the tissues are relatively unseturated in most cases compared. with a diver it should be possible to offect the disappearance of the bubbles more rapidly by recompression and so reduce the secondary damage due to hyporie or oedena. Behnke's first publication was in this field and many authors such as Alvis (123), Grocker (32), Burnett (27) and Scheoffer (98) have described cases sometimes demonstrating the lesion which led to the pocketing or trapping of gas at higher pressure. Tho signs and symptoms are similar to decompression sideness and usually. develop rapidly. Of course, such cases may occur after any period of compression so that the diagnosis of decompression sickness may not be differentiated from milmonary barotrauma. The question is raised whether cases of decompression slokness in which lesions such as lung cysts are described are due to an inadequate decompression routine or to inadequate

-29-

pulmonany vontilation (28). Wright (109. 110. 118) has cerried out many experiments on cadevers and animals which show the range of pressures required to lead to pulmonary damage as well as the fact that the damage starts on release of pressure. He has not yet succeeded in demonstrating histologically the actual passage of the gas through the pulmonary lining but this fits the observed cases in which the gas entry is not pervisiont so presumably the interruption of tissue continuity disappears on equalisation of pressures. Welder (148) has reported enimal experiments with drugs causing brenchospasm leading to pulmonary barotraina which may be interproted as showing that a clinically deconstrable losion is not always present or paraistent in man which again fits observations in such cases. The treatment in any event is recompression as often recommended. It is of interest that Lundin and Iundaron (100) have described the use of recompression therapy after several deve delay in the case of a woman who had had a large air embolism during a thoracic operation and that while there was no dramatic response. all attendants were convinced that her eventual complete recovery started while at high pressure in spite of the delay.

Muchly some montion must be made of the bone lesions soon in compressed air vorkers. The causation mechanism is still uncertain. The best curvey is that of Davidson (4.1) in tunnel vorkers and he found several reported cases of the asoptic necrotic lesions in divers. Preliminary studies have been carried out in the Royal (112 . 115) and United States Navies (20) suggesting that diving under their respective

-30-

regulations does not load to this sequel but that commercial undertakings are somehow responsible. Whether this is due to the longer time at pressure, the frequency of shifts, the abuse of decompression routines, the ignoring of minor signs and symptoms or in the inadequacy of recompression therapy, all these points have been discussed in extense wherever men concerned with this field have met. Due to the time lag of development of the radiological abnormalities and the recent awareness of the problem, it has not been possible to make any deductions on this matter in the various trials recorded here. However all men now taking part in experimental work have a survey carried out as recommended by Davidson (41) so that in the future some information may be gained, especially as the deep dives are now longer in duration and at higher pressures than commercial work.

-31-

# CHAPTER IV PRINCIPLES OF A DIVING TRIAL

#### Thios

In a disciplined service, there must be constant avarances of the possibility of a clash on some aspects of human experimentation. Where such experimentation is intended to lead to greater safety for or higher efficiency by the combatent. It is easy to gloss over the problem of the means to the end. The general guide used in this field is embodied in the Declaration of Helpinki (72). In a diving trial, the man is exposed to a probability of injury where the severity cannot be Ideally such trials divers would be volunteers with the best predicted. of motives and fully informed on ell the risks - en impossible situation where the risks are not fully known. In practice incentives were provided; formerly these had been of a financial nature then purely one of prestige but latterly they were both financial and endowing prestige. Opportunities vere regularly evailable for men to volunteer for such Any men who is sent to any diving teen hes the right to refuse work. to dive - at the risk of losing his qualifications if his refusal is considered unreasonable with the financial and posting consequences. The problem of "free choice" lies not only in such direct after offects but also with the indirect pressures that are brought to bear on a man by his colleagues as members of a select group, by the attraction of a shore posting in the base area, or by the possible comments in his record sheet. The statement by the Medical Research Council (111)

olearly delineated the problem in the case of procedures not of direct benefit to the individual.

There is no doubt that the discovery of a diving routine which will reduce the toll of injury and permanent disablement due to an activity which is carried on for military, connercial, and leisure purposes is a laudable end. With the co-operation of the responsible officers the othical difficulties of the participation of the men were safeguarded by the adoption of the following routine. When the individual is nominated for a trials team or he volunteers for such a posting, he joins the team temporarily to learn about the work involved. After an interview he may decide against such work and an alternative draft to a similar billet for sea or shore service is found; if he vishes to continue and he is considered suitable. he carries out a short probation period and only after mutual agreement is his posting confirmed. At a later date he may change his mind - or he may become unsuitable for reasons other than his own fault - and in that case. Stremuous efforts are made to transfer him to a suitable posting. This possibility is explained to him initially but no guarantee is given other than that attempts to find such a position will be made. (No may also Leave the teem for disciplinary or other administrative reasons). When a man leaves the team his record is marked to the effect that he took part in such work - and especially in any notable event - but no mention is made of any reasons other than disciplinary for his departure mless there is an enforced limitation on his future diving employment. In this vay

-33-

an acceptable method is provided where the diver may "contract out" with minimal indirect pressure whether he was a volunteer or selected instead of the ideal routine of "contracting in" with full knowledge.

In reaching his decision the prospective subject must have confidence first that the trial is worth doing; second that adequate proliminary work has been done: third that the control is adequate: fourth that he will be given the best available treatment if he is afflicted; and fifth that in the worst possible outcome of serious disability he and his dependents will be treated humanely. The first three points are essential to the performence of the work in any case and the fourth is patisfied by the services of experienced medical officers and attendants backed up by consultants. The traditional outlook on the fifth point is changing. In his training the characteristics of initiative, a sense of responsibility, and self-reliance are developed in divers and these attributes cannot be ignored in experimental work however less troublesome blind obedience might be. In this day and age these qualities have led to domestic arrangements such that the previous compensation of appropriate pensions is usually inadequate. While death of the diver may lead to his debts being paid, a disability pension is often insufficient to maintain hire-purchase payments. Some form of insurence to cover the outstanding debts either as a lump sum or as a regular supplement of income must be considered as part of the expense of an experiment where reduced earning capacity may occur however unlikely such an event may be. It is worthy of note that there

-34-

has been no leck of volunteers although it has not always been possible to arrange their velcase from their current tasks and that a few men who were drafted exercised their right not to take part.

## Selection

Having laid down the conditions on which men may be found for trials work, the next problem is how to discover their suitability. As knowledge of diving physiology is still relatively limited, the problems of selection are relatively simple. The mon obviously must be competent at their job and this means that they must be fully trained and experienced. Miles (114) has carried out a survey of some physical characteristics in divers and found no point in which they differed from national service men after initial entry training. Wise (150), in the light of work done with aircrew on susceptibility to decompression sickness (58. 66), found no predisposing factor in 1,003 divers examined for age, height, weight, and body type. A general clinical interview and examination and an X-ray examination of the chest are required in eech man to confirm his state of health and his general attitude. (In deep diving work N.N.G. and skeletal X-rays are now being taken as control examinations for future checking). The testing time is the first dive when the new man has to entrust himself to routines that have not been proved sele. For this reason, it is essential that one at least of those responsible for the trial should be in the sotual diving team and in particular it is of value if the doctor can be this man; another good choice is the senior diving officer responsible for the carrying

-35-

out of the programme; the designer of the diving equipment should also dive. No doctor can be as well trained as the divers but as long as he is known to be willing and actually does dive at some opportunity the divers will be satisfied. In deep diving he may be permitted to dive under vary closely supervised conditions at a later date - even if it is only the very last dive on a routine already accepted on the basis that they want to make sure he is always fit enough to treat them. Once a trial has been under way for some time, less experienced divers may take part if it is clear that a lower state of training is acceptable, and some such divers did take part in two of the trials discussed in this work.

As each diving trial is intended eventually as a vorking routine, the divers in the experimental team should be representative of an operational unit. In fact a case might be made for selecting divers known to be susceptible to the risks involved, so that the worst possible results are obtained.

#### Number of Dives and Divers

It has long been known that divers vary both between themselves and, for any one person, from day to day  $(h_{1})$ , in their susceptibility to diving injuries of various kinds. There is at present, no exiterion for the safety of a routine other than the incidence of decompression sickness, in spite of attempts to prepare scales based on severity of signs and symptoms (141. 142) or special post-dive tests (92) to assess dives in a quantitative manner. Superficially there is a 50/50 chance

-36-

because of the case/no case outcome of each dive but experience has modified this situation by excluding the extreme situations of all injured or none injured without however defining the variables concerned. This makes any attempt at applying sophisticated statistical rules a pointless pastime. Some decision had to be made on the number of dives that must be done before a perticular routine can be considered as tested, as neither unlimited time or unlimited numbers of men. - or unlimited money were available. In the past up to 5 dives was considered sufficient to check a routine for any one combination of depth and duration. Most of the trials here reported, were carried out with a target of 10 dives by 10 different individuals for each stop; this was not always achieved. This number was selected on an arbitrary and intuitive basis and was accepted by various authorities with little discussion (101). Because more dives were done by a greater number of individuals, these trials are more acceptable than previous work of a similar nature, nut no pretence is made that a satisfactory solution has been found to the question of what is a well-plenned decompression trial. A trials team has varied from 8 to 22 individuals at any one time but the inevitable spread of each trial over a period of time has meant that many more men have usually taken part in any one trial.

# Assessment of Gases of Recompression Sidness

The actual dive routine and various matters of equipment and decompression tables vary with each trial but a constant need is the assessment of the condition of each man during, and after, the decompression

-37+

of his dive. Almost every diver at the start of his experimental experience finds that questions about his bodily reactions result in sensations he had not previously noted; this response is complicated es he does not know what sensations he would have if he had developed decompression elekness and he has heard many bar room tales about the disabilities in undiagnosed or mistracted cases. Where a physical sign is present, there is little difficulty; usually the complaints are subjective and experience is required both by the diver and the dector in accertaining the cause and the severity before appropriate treatment may be given. At one time some neval divers suggested that all senior divers undergo on attack of decompression sickness to gain experience, so that trainee divors would learn directly from one with experience but this proposition was rejected out of hand by those divers and doctors with experience of the variability of individual susceptibility and case severity.

Once symptoms and signs have been accepted as due to decompression, the dive in question can be considered risky. However individual variability is such that an adequate routine may still be possible and the diagnosis could be wrong. Although all such cases should be diagnosed as decompression sickness, which is a serious condition, a further empirical classification is carried out into major and minor attacks. It is realised that this may be interpreted, as an indication of a disease which has a steady progression in severity proportional to the imadequecy of decompression, but this attitude yould be as wrong as a

₩<u>7</u>8~

claim that there was no such progression. Experience shows that if decompreion sickness occurs after a dive it may be of any severity; it is more likely to cccur in minor forms rather then major if the decompression is barely sufficient and conversely if the decompression is grossly inadequate.

# Problems of Treatment

There are several practical reasons for using severity as a suide but the main one is the therapy required. Once a man has parried out a standard therapeutic recompression regime, he is reluctant ever to carry out enother unless he feels. in himself. that it is essential. If the medical advisor treats all cases by recompression, the diver tries to essess his own condition: if he considers it trivial he will not report his symptoms and signs and he will try his own remedies. Decompression sickness is an injury that may become symptomless fairly quickly without specific therepy; or it may be eased by such home remedies as a bath, analgosics and rest; or it may be quiescent for some hours and then return with increased severity. In the first two cituations. no harm may be done to the viotim but assessment of the decompression routine may be erroneous, while in the third situation, the condition may become very difficult to treat. If the doctor treats his patient as well as the injury and uses symptomatic pelliatives when indicated as well as specific treatment, he has the advantages of cooperation, surveillance, and information to easist him in his task of assessment and therapy. This point may seem to have been laboured but

from personal experience men will forego financial recompense of sums of 620 and over for a treatment of 10 hours duration or so rather than report sick if a therapeutic recompression was probable. Men have also been known to tell several days later of symptoms that had occurred but which had since disappeared, this inspite of a flexible approach to treatment by the attending doctor. (The problem of bone damage is very worrying when dealing with men who have concealed their injury.) Classification of Gases

A preference for dividing cases into major and minor categories rather than by system involved has already been mentioned. Such a division has two practical advantages. First any illness not due to the decompression, such as a simple bruise can be considered as a minor case and there is no indication to underestimate signs and symptoms. Secondly, the classification helps in determining the rest period of a diver after an incident as in a minor case a "lay-off" of 24 - 48 hours is adequate but a period of 5 - 7 days is prudent after recovery from a major incident. The difference between the two grades rests on one factor - whether the diver required to be recompressed for therapy or not. Thus a major case is one where the diver had to be recompressed for the relief of his signs and symptoms. All other cases are minor ones.

Major cases have signs and symptoms which range from aching pain to cardiovascular and neurological involvement of various types. The amount of compression required depends more on the speed of development

-<u>l</u>.()---

of the condition and rapidity of therapy then on system or systems involved so that further sub-divisions of essessment are not worthwhile: eny recompression is enough to classify the attack as major. A minor case may present, as localised prantius, or environe or mild aches wound joints which respond to mild enalgesies or to heat therapy in the form of deep hot weter boths or to simple rest. The possibility of recompression may arise if signs and symptoms progress and these indicate that this would be better thorapy - in fact a hot bath may sometimes bo provocative in cases of doubt as to the best treatment for a particular case - and such cases would be reclassified as major ones. Often in cases where aching pains are the only symptoms, errors can be made by underestimating the severity first by the divor end then by his nedical attendants. In such cases only knowledge of the individual will reduce the possibility of error. A period of several hours of persistent mild aches is an indication of error in assessment and such cases should then be considered as major even if recompression therapy has not been used.

#### Assessment of Trials

It is admitted that the standards of assessment are arbitrary but successful operational routines have been evolved. It is considered that in trial dives an overall incidence of major cases of 2% is acceptable (101) as the addition of safety factors for operational use would reduce this considerably. No comparable standard is available for an emergency or an exceptional routine but adequate treatment facilities any negatized (132.133) wherever and whenever such a situation might axise - and the risk of decompression sickness as found at the trials

-41-

assessment is one of the factors taken into account before an exceptional dive may be permitted.

Where decompression theories are being tested the assessments are easier. One major case means the schedule is unacceptably dangerous as does the occurrence of more than two minor cases. The occurrence of one or two minor cases out of the group of ten dives on a particular routine would mean that the next step in the programme would be carried out very cautiously but not necessarily after any alterations in the plan.

#### CHAPTER V

#### PRISONALLE AND MATERIALS IN THE DIVING TRIALS

## Background

Diving qualifications in the Royal Navy are usually awanded after appropriate courses have been successfully completed. Maval equipment is designed for specific tasks such as salvage and mine clearance. Very few divers have experience of all types of diving equipment although the great majority have had an occasional dive in equipment other than that of their speciality. For example a Glearance Diver, normally a self-contained swimmer, is not always able to move with comparable speed and economy of effort to that of his Standard Diver colleague when wearing the latters' traditional surface supplied helmet and other equipment.

A change has been taking place in neval diving however and although three of the trials described here were carried out by men in standard equipment - although the run down in the Standard Diving branch led to some of the trials diversi being Glearance Diversible by basic training - the fourth trial was carried out by swimment in a role that had been exclusively the preserve of the helmeted man. The fifth trial described deals with submarine escape training in which equipment was used to provide buoyancy only.

## Personnel Qualifications

It is much to the credit of Diving Officers and men that the rivelry of those trained differently elways led to friendly beater and and competition mather than to biokering and lack of teanwork. This was helped by those few divers who had transformed from Standard to Oberrance Diving. The experience in diving varied from men with 20 years service in standard equipment, - that is before self-contained diving had been instituted in the Navy, - to men with a 3 weeks course completed some months before. The qualifications ranged from officers, qualified as Clearance Diving Officers or as Qualified Deep Divers, and matings, of the three grades in each of the Clearance Diving and Standard Diving branches, who were all full time divers, to men who had qualified in one of the two Ships' Diver's grades and who were part-time divers as they had other qualifications such as Mechanical Engineer, Stores Assistant, Sick Berth Attendant, or Seaman.

#### Personnel: Physical Factors

The non varied considerably in respect of such characteristics as age, height and build but all vere still apparently healthy enough to remain on active duty in Fulheems category P2. Age varied from 18 to 48; weight from 9 stone to over 15 stone; height from the minimum standard of 62 inches to 74 inches; and body build from extremely thin to obese (i.e. waist measurement much greater than chest measurement) Equipment

The equipment that was used varied with each trial and with the phase of the trial. These phases fall naturally into those where simulated dives were carried out in compression chambers and those where the dives were carried out in the ses.

-14-

# a) Compression Chambers

During a compression chamber dive the man is exposed to an ambient air pressure of his simulated dive but does not have the hydrostatic effect of pressure difference between head and feet. If he is carrying out a dive with air or some mixture of oxygen and helium as his breathing medium. the dynamics of his respiration is unencumbered by breathing equipment as he normally breathes chamber atmosphere. The energy he expende is less then on a comparable sea dive unless he deliberately exercises on a rowing machine or other work device. The diver is under constant observation through the port holes and he is in communication both by voice and coded signals so that a feeling of isolation is absent and rarely does he have any apprehensive feeling of failure of equipment. If he is one of a group diving, he may be particularly affected by the feeling of general hilarity and irresponsibility engendered by inert gas narcosis. He can wrap up in blankets to be warm and comfortable during decompression (Fig.15) and he may even have a map.

Altogether 7 different compression chambers were used in two establishments and one ship (Fig. 1 and Fig. 2). There were no differences in principle in the operation of the chambers but there were many minor differences of construction all of which had some bearing on the trials; all but two of these differences affected the comfort of the diver (Fig. 3). When it is realised that these chambers were also used for therapeutic compressions which sometimes took up to 5 days, it can be appreciated that the comfort of the available chambers was relevant to

-45-

the disclosure of signs and symptoms of decompression sickness and to the decision as to the best treatment in a particular case. Any dognatic instructions for therapy under all circumstances could be cruel to men liable to be under therapeutic conditions frequently either as a patient or as attendant to a patient, however valuable such instructions might be for the rare occurrence in an operational team.

The two differences of operation of the chambers concerned the accessibility of the patient and the equipment for administering alternative breathing mixtures to that of the enamber atmosphere. ALL ohembers have small compartments, commonly referred to as medical hand locks, through which small items such as food and drink. recreational matter, daugs, and for hygienic needs, could be passed. However if skilled medical attention was required for the petient - for examination. passive physiotherapy, attention to pressure points, catheterisation and blood sempling to mention a few actual requirements of the past, unless a suitably trained person had accompanied the patient into the chembor at the onset. a very difficult situation could arise. For this reason chambons should have at least two large communicating compartments one of which could be used as a man-look. Once such a look is evailable, opportunity can be taken to relieve the attendants, or to introduce another patient if such a need arises. One of the chambers on this trial was of single compariment construction with small hand two chembers had a large look on each side of the main compartlooks menta and the remaining four had the common large look and main

-46-

compartment construction.

Broathing the chamber atmosphere is the simplest erronament for the occupants. The earbou disxide concentration in the closed compariment will give but frequent ventilation by flushing ges through the chember will evold env danger from this cause. At times it may be desirable for the men to breathe a different orygen and helium mixture or pure oxygen for part of a similated dive, or of a recompression (Problems of chronic exposure risks and the fire hezend put therepy. the possibility of making pure oxygen the chamber etmosphere out of practical consideration). The solution must be some form of breaching equipment. In these triels several types were used either fitted to the chambers or portable to be telsen into a chamber. The most porular consisted of storage cylinders of the appropriate gas outside the chamber fod through a control panel to a main pipe inside the chamber. the men being eble to connect simple demand valves with monthpieces to this pipe (Fig. 4). Such a system kept bulky conference out of the chamber leaving a maximum amount of room for the perpents but it wested and caused a steady pressure rise in the chamber. Alternatively the control panel could be installed inside the chamber and the demend velves connected directly to it thus saving structurel modifications but decreasing the evailable space and still leading to a prossure rise from exhaled gas. The popular system was permanently fitted in one chamber and the internal panel was used in two others most of the time. All chambers were fitted with a system which had the supply cylinders

-17-

externally connected to volucers inside the chamber and the gas was then led to closed circuit breathing apparatus. This system had the advantages of room and gas exponent but the breathing equipment was heavy, and was only tolerable for short spells mainly because the gases became uncomfortably warm from the absorbent; generally it was unpopular and had a dotrimental effect on the morale of any invalid. In fact a fourth method was preferable and this was to nove cylinders, control panel and demand valves as one unit into the chamber in spite of the disedvantages of bulk and weight.

Other differences that existed between chambers were not important in the routine but wope of importance to the divers. The length and diameter were the most obvious as the dimensions determined such points as ability to sleep stretched out, ability to sit upright, and ability to stand. Small chambers had to be ventilated much more frequently to remove carbon dioxide and replenich oxygen when the chamber atmosphere was boing breathed.

During this process, noise was troublesome, especially if treatment was being undertaken; the noise of compression and decompression from stage to stage might be acceptable but the unevoidable insistent cound of release of compressed gas during ventilation provented normal sleep and caused fatigue. (Silencers were tried but were relatively ineffective).

Wide bench sents and folding chairs were desirable where they could be fitted as some dives took up to 22 hours in the chambers and

-48-

aons therapies were much longer. Eventually fean rubber seat cushions and floor and bench mattresses were provided.

Communications were either by use of coded knocks with a mallett or by means of a sound-powered telephone initially. Later four chambers were fitted internally with two-way microphone and loudspeaker fitting so that the occupants were always eventheard by the outer attendants. Later still the system was modified to receive music and the occupants once more had to knock to draw extention to their wich to transmit a message and the music was then turned off!

The final significant difference between the chambers concerned the thermal insulation. It was found to be nearly impossible to keep the inhabitants were in a steel chamber in the open air during a British Winter while a light alloy chamber in a warm foreign summer needed ice blocks on top to maintain a comfortable internal temperature. Swings of up to 30 degrees Centignade (55 degrees Fehrenheit) were recorded on several occasions during such similated dives. Hore stable conditions were obtained where the chamber was kept under cover in an even temperature and such conditions applied to 5 of the chambers, 2 of which were on board ship.

# b) Submarsible Chambers

As a link between equipment on lend and equipment in the sea, a brief description of submersible chambers is relevant (Fig. 5). Two such chambers were used. The principle is that of a diving bell which, with its occupants, is raised and lowered in the sea rather like a lift.

-49-

The veter level is kept down by gas supplied from the floating base or ship so that the occupatts are exposed to the ambient pressure in air. The chamber is lowered cither to a point where the diver then callies forth and returns or else to a point where the diver. already started his easent meets and enters it to complete his decompression. The chamber is then hoisted as required and when instructed the occupants can close a hatch to seal the mouth of the bell thus converting it into a pressure retaining vessel. The divers may either breathe the atmosphere of the bell or they may use breathing equipment. This chamber is then hoisted inboard where the pressure is released as plenned while it resto on the deck; in the case of the never bell it is hoisted on top of a compression chamber into which, once the pressure is related to match that in the upper chamber, the divors transfor. (43. Fig. 6a and Fig. 6b). Once the hatch is closed on the receiving chamber the pressure is released so that the bell can be used again while the first divers continue their decompression in greater comfort. The refinements of the second bell, compared with the first, other than the tremendous value of the possibility of transfer of the divers under prousure from sea to shipboard, are commutations, closed circult television conttoring, and gas supplies carried in cylinders round the upper part so thet during very deep dives the submersible chember becomes a base for operating swimmers without too many lines and hoses in the water and yet the surface controllers are kept fully informed about the sectiv of the men. As well as operational advantages, the transfer

-50-

system allows rapid medical essistance in case of injury to the diver during the decompression.

# c) Breathing Muipment

In the see, 42 dives were carried out using entirely selfcontained equipment; 664 dives were done using standard equipment

(Fig. 7). This is the familiar idea of a diver who has a surface supply (Fig. 7). This is the familiar idea of a diver who has a surface supply to a holmot and corrected attached to a subbarized total sufface supply to a holmot and corrected attached to a subbarized total sufface supply to a holmot and corrected by heavy boots and cheat and back weights (naturally the process of entering or leaving the water with 160 lbs. of equipment is energetic). A transistorised communication system was used and the wire was included in a braided rope which acted as lifeline. Enough air was provided to ventilate the belief to prevent carbon dioxide accumulation and it was supplied at a slightly higher pressure than that to which the diver was exposed; the diver regulated his buoyancy to just negative by controlling the escape of gas by the valve in his helmot. The self-contained equipment used was similar to the standard gear except that the gas supply was contained in four cylinders on the diver's back and breathing was carried out in a closed circuit system this equipment did not come into general use.

The remaining sea dives - all deep dives, 206 in number - were carried out by men, dressed in rubber suits entered through the neek and intended to keep the swimmers dry and with fins on their feet, and commonly described as fragmen. They were not entirely self-contained as the breathing equipment consisted of an open circuit system where gas from the surface - or from the cylinders of the submersible chamber - was supplied by aixpipe to a reducing system and demand and enhaust valve on the diver's back and then by breathing tubes to and from a monthpiece set in a wide vision mark (Fig. 6). The main supply was so enranged that an emergency supply of gas carried in two small cylinders on the diver's back would automatically come into use 1f the main pressure dropped. This "get-you-home" supply is the modern counterpart of the volume of gas available in the helmet and upper part of the suit of the standard diver and lasts just long enough for the diver to reach his base without delay.

#### The Base

Every dive is controlled from a base where all administrative details can be dealt with. N.H.S. Vernon, a training establishment carried out this function for the divers while they were carrying out the initial compression chamber dives. The Adminalty Experimental Diving Unit (A.E.D.U.) situated within the boundaries of H.M.S. Vernon, is a small unit of engineers, divers and draughtsmen responsible for devising and developing diving equipment. It is a mixed givil and uniform establishment of the Royal Neval Scientific Service. Both have their own compression chamber facilities and both have their own medical organisation. A.E.D.U. sponsored the trials on the air dives, combined dives, and surface decompression dives and the Unit medical officer had the major responsibility in seeing the work through.

H.H.S. Dolphin is elso a training establishment but only for

-52-

submarine energy. As part of the training, submarine escape is carried on in a special department which has a full time staff including a doctor, who has some advisors to call on if required. The advisors are doctors who have had experience in this field and are available either at A.E.D.U. or the R.M. Physiological Laboratory (R.N.P.L.). Compression chamber facilities are available at the training tonk.

R.N.P.L. 1s a research ostablishment of the Royal Naval Scientific Service and is staffed by scientists and technicians of the civil side of the Adminalty, with the assistance of two Medical Officers from the uniformed side. Much of the work concerns basic research into diving end enimal work is on important part of this. Co-operation between N.N.P.L. and A.E.D.V. is important and one of the doctors is appointed to both units. Thus the deep diving programs cane from discussions with both units, animal work by R.N.P.L. with the scientists having the major role. Then A.H.D.V. arranged for enough divers from H.M.S. Vomon to travel dealy to R.N.P.L. (geographically quite close) end carry out dives agreed between scientists, doctors, and divers of the two units in the most convenient compression chembers (sometimes the scientists travelled to N.M.S. Vernon). The teenwork was such that R.N.P.L. staff would operate the compression chembers, under the care of melified neval diving officer, from A.E.D.U. with the medical officer oremining, advising, essessing and if need be instructing on treatment measures from his dual post. The next stage was to plan the sea trip where the A.E.D.U. took the major responsibility to carry out the dives

-53-

required by R.N.P.h., the doctor continuing his role.

This type of shore base organisation led to considerable flexibility in arrangements in that roles were fairly closely defined but allowed considerable co-operation between the research side, the user side, and those who were responsible for the victims if all did not go well. Thus some of the trials were more in the nature of experiments carried out for Scientific Officers of R.N.P.L. (37. 39. 80. 81. 127) while others were really evaluations of routines devised on their information by Naval Officers of A.E.D.U. and Medical Officers of both units (34. 36. 38. 101. 102. 107. 117).

The see dives were all carried out from H.H.S. Backain which had been completed as a deep diving training and research ship but during the period covered in this work she was given an alternative role as a support ship (Fig. 9 and 10). Originally she carried a team of 2 Diving Officers and 12 diving ratings and 1 diving Medical Officer with adequate stonage, working and maintenance space. Latterly the needs of her other role had reduced the storage and maintenance space and her team was reduced to 1 Diving Officer and 6 diving matings and her Medical Officer was less familiar with diving problems. This deprevation meant that for the diving trials her team had to be augmented by up to 2 Diving Officers, 1 diving Medical Officer, 6 experienced divers and 2 engineers to cope with problems of gas mixing and supply and equipment generally. However the ship was still able to find the appropriate water conditions for trials whether in West Scotland see looks or Canaxy Teland harbours. Norwegian fjords or Therden bays.

-54-

# <u>CHAPTER VI</u> IMPORTATE FACTORS

-55-

Many factors are relevant to the problem of decompression slokness in divers and in view of the obscurity of such aspects as mechanism and pathology, doubtloss there are many still unappreciated. The pressures to which the diver is exposed and the duration of exposure are closely linked and fundemental espects. In this chapter some material points are discussed in more detail.

## Pressure Control

Pressure in a compression chamber is relatively easy to measure, the main problem being the desirable accuracy. Simple Budenberg gauges with dial display are common but exact accuracy on such a gauge is only possible at a particular point although the error at either end of the scale can be reduced by various means. These gauges need to be regularly calibrated by hydraulic dead load testers. All the sea dives and the majority of chamber dives were carried out using this type of gauge. Latterly a range of Bristol recording gauges were used on the main chamber and by selecting the appropriate instrument, very great accuracy in chamber diving was achieved (Fig. 2). In sea dives the gauges measure the pressure in the air-pipe carrying the gas supply to the diver or submersible chamber and this pressure must be the same as that of the depth of water at the diver's chest in standard gear if he is breathing confortably or at the water level when the submersible chamber is used. In standard dives this value was further excess checked by the use of a measured load-line. by echo-sounder. by the measured marks on the diver's life-line, or by the reading of the gauge on a weighted open ended air nive when gas escaped at depth. Such readings as were taken had to agree within 5 feet for satisfactory control but the master reading was the one on the gas sumly to the The depth of the submersible chamber was more difficult to diver. assess of the start of a dive os great volumes of gas. delivered at high pressures through a small pipe, were required to keep the water level down and the total reservoir pressure was needed to enable the diver to control this level. When the chember reached the prodetermined depth the main supply line was shut, the diver's control velve inside the chamber was opened wide, and the residual pressure on the line gave the depth of the chamber in feet of water. This figure ves the internal pressure of the chember so if the bottom hatch was shut. It did not necessarily record the actual donth of water. This method was of little value where it was desired to take a diver to a specific depth rather than to discover what depth he had reached. For the former purpose the only satisfactory means was to measure directly and mark appropriately the special non-spin, non-stretch hoisting wire of the chamber and to repeat this at wookly intervals when errors of up to 2 feet in 500 feet might be found. The depth of the chember could be cross-checked by echo-sounder and by recording the required number (previoualy counted) of revolutions of the winch drums.

-56-

# Time Control

All dives were recorded to the nearest helf-minute using various types of wrist-watches and clocks. In long dives discrepancies arose and in ship or establishment the most consistent buildheed clock was found and was used as the standard but cheeks your kent by stop wetches or by the individual recorder's wrist wotch. Diving practice hes been - and is - to use intervals of 5 minutes or multiples of 5 minutes for the duration of dives and the duration of decompression stops; this custom cannot be continued where changes of pressure or depth occupy an appreciable time. Thus in the air dives the compression or descent of the diver at 100 feet per minute was included as part of his stay at maximum pressure as such a pariod could not exceed two minutes; his escent to his first stop, at 60 feet per minute was included in the time spent at that stop which was adjusted to take this period into account; the time of eacent between stops - normally a distance of 10 feet at a similar speed - was included in the time of the shallower stop. In deep diving the increase in pressure had to take longer as the sene compression and descent rate was maintained and the time was then considered separately from the donation of the time et maximum pressure or bottom time; the escents were also considered es separate activities and after the initial rate of 50 feet per minute to the first few stops, the ascent was controlled to take a full or a half-minute between stops. It was fairly rapidly found that 5 minute intervals were not helpful in experimental deep diving and the duration

-57-

of bottom time and each decompression stop time was given in single minute values. Whatever the noutine, it was strongly emphasized that the rate of change of pressure in compression chambers or of descent and escent in sea dives, should be as smooth as possible; in fact accuracy of timing was, within limits, less important than regularity of rate. Smooth accurate movement was caster in a chamber then in sea dives; chamber compression could slow up towards depth if the cylinder supply pressure was low and the diameter of the piping system was inadequate, to pass the volumes required to raise the pressure inside the chamber.

In offect, the probable operational routines had a slight safety margin when considering the descent time as time spent at full pressure and a greater margin when rounding the ascent and stop duration to the next 5 minute interval; while calculations had to be made, in the experimental procedures, for the uptake and release of gas during the time the diver was undergoing pressure changes.

#### Donth Keeping

In the sea dives an important matter was the ability of the diver to maintain his depth. In tidal waters, especially in long dives, his actual depth could change with the state of the tide. Whatever the waters, his ability to maintain his depth during decompression could be affected by the strength of a current or by the rise and fall of swell. This latter factor could be exaggerated by the movement of the ship caused by weather conditions and result not only in a wide range of

-58-

movement by the diver but also in a snatching type of motion. This effect, most noticeable during the shallower decompression stops, could be avoided by the use of submersible chambors.

The decision on what constituted marginal weather conditions first in respect of the diver and second for the use of submarsible chambers - with the problems of control during transfer from sea to ship board - was difficult and always liable to appear wrong by sudden changes in the weather.

## Sea Tennomatium

A factor of note though of debatable importance was that of sea temperature. There is little argument on the preferable confort of a short dive in the warm waters of a Gauary Telends apring to that of a long dive in the cold vaters of a Hebridean winter. The morale amongst the surface attendants tended to vary directly with the temperature and inversely with the precipitation. The Standard Diver could dness with many layers of clothes - long cotton and heavy wool underware, flemel body belt, long stockings and woollen cap and even electrically heated combinations on occasion. Such clothing left only his heads exposed to the environmental temperature and to some ertent gloves could case this problem. An important item in very long dives was a man sized negkin to absorb unine if micturition because necessary. On the other head the swimmer could not wear so many undersuits because either he could not then don his neck entry rubber suit or he would become so buoyent he would require unacceptable amounts of lead to counteract this. His

-59-

normal underwear would consist of a combination suit of cotton and a similar thin anit of wool over a pair of bathing trunks or underrants. It was essential that he excreise during his stay in the water to keep van and the potential ill-effects of such activity during decompression were evolded by the use of the transfer under pressure facility. A few divors used "wet suits" - these one suits made of foamed synthetic mbber open at enklos, wrists and neck which permit a static thin film of water between suit and the bare skin of the diver and which repidly reaches body temperature. However the construction of the material is such that at depth the foan flattens and the diver might as well wear a suit of tissue paper. All in all, the divers kept reasonably warm so that no conclusion could be drawn on the effect of sea temperature on decompression sidness, although there was no doubt about the more pleasent diving conditions such as increased visibility in warmer waters. General Health

Finally the general health of the divers was important. While they were all generally fit men, opisodes of illness did occur, sometimes of infectious origin. As post-dive complaints were the main sources of information in assessing the dive, it followed that divers were expected and asked to declare any signs and symptoms present before a dive. These were not always reported and men dived with boils of the perincum; they dived when they were suffering from hangevers, from muscle stiffness after gardening; and one man dived in standard equipment while wearing his conset during a mild encombation of symptoms of a slipped disc.

-60-

(The conset was spotted while the diver was undressing in an obscure commer after his dive). At least two such man developed decompression sickness but the true incidence could not be assessed as undoubtedly ell the cases were not discovered. A frequent condition in British divers is a sub-acute minitis and it seen became apparent that men were diving without ill-effect who had nesal mucosel swelling and whose tympenic membranes could not be seen to move during a Velsalva manoenyze. Eventually it was found that those divers who compleined of a fullness of the head or heaviness of the face or inability to clear their cars vere those who should not dive end, notwithstending the clinical findings, the remainder usually succeeded in corrying out their dive. It was also noticed that men who used considerable effort during a Valsalya manoeuvre or who repeated the manoeuvre often - a common fault in the relatively inexperienced men - recohed a point where the manocurre became ineffective and the dive had to be abandoned. Inis presumably was a tissue reaction to unaccustomed violant movement as instruction and prestice residly produced a situation where the minimal rise in pressure obtained by closing off the mouth with the tongue and barely occluding the nostrils was adequate. No case occurred of timitus and deafness in these trials but it has occurred in divers who have to uso a little more force then usual to clear their cars because of sild congestion; this injury is more common at shellow depths but it emphasizes that divers must be able to equalize their middle cars nomally before diving.

-61-

#### -62-

#### CHAPTER VII

#### TRIALS OF STANDARD ATR TABLES

## Background

In point of time, these trials (107) followed the assessment of the two air diving routines discussed later but as the pattern is basic to all triels for diving experimental work, this account is placed first. New diving decompression schedulos vere introduced into Royal Naval service in 1958 in the form of two tables (35. 132). Table I was the routine working table designed to give a total decompression time of about 35 minutes or less for a dive and any douth-time combination that required a longer decompression was described in Table II. This timing was chosen as it scened a reasonable period during which a man would stay fairly inactive in the cold seas round Britain. The latter table was provided in case the limits of Table I were exceeded for accidental reasons or deliberately for operational reasons of such importance that a risk of decompression sickness was acceptable. Table I had been fully tested and experience confirmed the opinion in the report that it offered a reasonable economy in decompression time with a negligible risk of bends (35). Teble II was not tested as it was an emergency table but the new theories had after calculation resulted in schedules which were in general longer than these in the table it was replacing and hence it was presumably as sefe. However the proposed routines (36) of surface decompression and combined dives involved an increased use of Table II
and the results described later cast doubts on the merits of this table and thus on the assumptions on which the table had been based. The first step was to test the calculated table uncomplicated by the proposed routines and the second step would depend entirely on the results of this approach.

## Rasic Decompression Theories

Due to world-wide dissatisfaction with diving tables which were felt to be too safe in some parts and too dangerous in others. other countries had also been revising their routines and the United States Navy issued their latest tables (145) as work was about to start on testing the British tebles. The Royal Naval tables had been devised on a concept of a single homogeneous tissue which absorbed gas in an exponential manner on exposure to pressure and on subsequent exposure to a reduced pressure the release of gas followed a reflection of the curve so that the amount of gas in solution varied with the absolute pressure and the square root of the time; when this emount expressed in term of total absorbed cas in feet of sea water exceeded a value of 30 feet above ambient pressure, then the ascent of the diver was controlled to maintain this excess pressure at or below this value so that he could ascend to a shallower depth or surface without the risk of decompression slokness. The U.S.N. tebles were a sophisticated development of the Heldanian concept that the absolute gas tension in a tissue at pressure could be halved safely; the sophistication lay in the use of several hypothetical tiesues of different helf-times to saturation each with its

-63-

own ratio for reduction of pressure based not on the 2:1 cir ratio of Heldano but on a nitrogen content ratio which also varied for each tissue with different abient absolute pressures.

#### Selection of Tebles

It could that it would serve a useful comparative purpose to adapt the U.S.N. tables to R.N. diving techniques and to carry out the trials with the same men at the same time with both tables. The modification entailed the conversion of dive and decompression stop durations to 5 minute groups and inclusion in the stops of the time taken for the ascent. The R.N. tables were forwarded on a pecket card version (13.116) but the compiler did not inform these carrying out the trials that, in anticipation, he had already modified the tables arbitrarily and it was much later before a cross-check of the tables was made and the card version abandoned. Accordingly the adapted United States Navy tables were dived initially at the same time as the Gard version and latterly concurrently with the published Royal Navel tables. For convenience the several versions of the tables as finally used will be referred to as the American (A), Card (C), and British (B) tables respectively. Wriel Orders Requirements

The trial orders called for 10 dives to be done on selected schedules on each diving table. The depths of the dives selected were at each 10 feet interval from 70 feet to 180 feet. The durations of the dives alternated with successive depth intervals from the minimum time on Table II at a given depth with a duration 20 - 30 minutes longer at

-64×



-65-

the next greater depth (See Graph VII.1). Later the longer dives were abandoned and the shorter duration introduced for the majority of the intervals. Each selected schedule had to be dived in a compression chamber before being dived in the open water but this later proved to be of doubtful value in enticipating risks and was suspended. Period of Trial

# Over the period of five years that this work was carried on intermittently, various changes in technique and interpretation took place so that a careful review of the reports had to be made to decide which dives were within normal operating limits and which had gross deviations. Nost of these errors usually arose from the training received by diving personnel to use the next deepest or the next longer schedule whenever a dive nears the limit for depth or duration. For assessment, the trial dives were required to be as near these limits as possible. However the dives excluded from comparison were of value in demonstrating the incidence of cases of decompression sickness when the routines are abused and what variations there were both between men and in the same man from time to time.

#### The Divors

Altogethor 96 men took part in the trials but the diving team rarely exceeded 15 in number at any one time. Seven men carried out 20 or more dives each; twenty-six men carried out 10 to 19 dives each; twenty-one men 5 to 9 dives each; twenty-coven men 2 to 4 dives each; end twelve men carried out one dive each mainly for experience in the

-66-

techniques involved. Apart from the basic diving qualifications and the fitness resulting from being a member of an armed Service, there was no obvious common factor of experience or build. No detailed alinical examination records were made in keeping with the deliberate policy of the minimum departure from normal working routines. No selection was made for any man to dive on a particular table so that it was hoped there would be a random selection of divers on the schedules under test at any one time depending on the composition of the diving team at that time.

Twenty-three men dived only on the British tables with no major incidents and eight men had equally uneventful dives only on the American tables. Eighteen men dived on both British and American tables and 12 major cases of decompression sickness occurred in 8 men. Twelve men dived only on the Card tables and 6 men had 7 major attacks between them. Further sub-division into the effect of chamber and sea environment on the separate schedules is unprofitable because of the small numbers involved. Table VII. 4 shows the combination of diving tables dived by the numbers of men involved and the distribution of major eases of decompression sickness.

Tables used	A only	B only	C only	Λ & Β	A & 0	9 & C	A&B&C
Number of men diving	8	23	12	18	24	1	7
Number of major cases	0	0	7	12	6	1	Ļ.
Number of men affected in major cases	0	0	6	8	5	4	lş.
Number of dives	12	94	Ğl <sub>ŀ</sub>	165	256	5	98

# Table VII. 1

# Distribution of Decompression Tables and Divers

A . American tebles B = British tables C = Card tebles

#### The Dives

Altogether 764 dives were carried out but 89 (12%) had such departures from the desired southine that they were excluded and 19 dives which used the correct schedule but in which decompression was carried out in the submersible chamber were considered separately. In the remaining 656 dives there were 30 major cases of decompression sickness and 112 less severe cases an incidence of 4.6% major and 17.1% minor coses.

Twenty-one schedules were tested by dives but 4 had a total of only 6 dives between them; one other schedule was dived 10 times on only one set of tables; the remaining sixteen schedules were dived on at least two tables - some in fact were dived on two tables in the chember and three tables in the sea.

Envir	hvironment:		Dry			Wet	
Tablos		٨	P	Ç	A	Ð	G
Selie	dulos			1997 - 2097 - 2097 - 2097 - 2097 - 2097 - 2097 - 2097 - 2097 - 2097 - 2097 - 2097 - 2097 - 2097 - 2097 - 2097 -	des endered medalenses de for		
Feet	llinutes		an dan kata panén di katang kata	-	hing a life of the low symptotic rugs family	anda manjiki katala ji katala manji katala kata	
70 80 90 100 120 120 120 120 120 120 120 120 12	100 120 800 700 650 450 750 450 750 500 500 500 500 500 500 500 500 5	1 1 1 166 151 151 12 10 1 17 1 1	李 章 章 章 章 章 章 章 章 章 章 章 章 章 章 章 章 章 章	10 10 12 25 10 14 30 10 12 18 18 18	-910 10910 10910 10010 10010 10010	10 15 10 14 11 9 10 10 10 10 10 10 10 10 10 10 10	11111010010-1101111
<b>180</b> Ta	25 stel	12 112	\$	10 152	10 160	10 169	11 61

# Table VII. 2

Dives Carried out in en Acceptable Manner

-69-

# a) Chamber Dives

Only 2 dives out of 266 acceptable chamber dives were carried out on the British tables, the remainder being dived on the American and Card tables. From Table VII. 3 it might seem that if there was an attack of decompression sickness on the Card tables it tended to be more semious than an attack after a dive on the American version. 5 of the major cases on the Card tables occurred after 2 group dives of 4 men each on one particular depth-time combination, so that the overall differences may be due more to the chance of selection of men then to differences in the safety of the decompression routine.

Dives		liinon	Cases	Najor Cases	
Table Used	Moniper	Mmber	<i>%</i>	Number	%
A	112	31	27.7	3	2.7
U C	2 152	6S 1	13.2	т 8	(50) 5-3
Total	266	52	e zh <u>a jin eli ni</u> en energan ee	12	

### Table VII. 3

#### Decompression Sickness in Chember Dives

# b) Sea Dives

390 acceptable sea dives were carried out on the three tables under trial and are tabulated in Table VII. 4. It would appear that the relative severities of the American and Card tables are reversed when compared with the chamber dives. Again there was one particular schedule that upset the record - this time on the American schedules when the weather deteriorated rapidly during the dive - the only occasion this happened - and all 4 men in the sea suffered major attacks. If this point is taken into account there is apparently little difference between the British and American tables when the dives are carried out in a similar manner.

Dive	Minox	Ceses	liejor Ceses		
Table Used	Number	Number	the second se	Number	%
Δ	160	25	15.6	10	6.3
Э	169	23	13.6	7	4-1
C	6.1	12	19-7	1	1.6
Total	390	60	-	18	- -

# Table VII. 4

#### Decompression Sickness in See Dives

### o) Evoluded Dives

If the decompression schedule used was not the correct one for the depth-time combination, whether the dive resulted in a form of decompression sickness or not, it could not be used as a test of adequacy of the schedule or used in comparison with other routines. Such dives were still of value in suggesting where extra safety already existed or was needed; in showing by the variability of response in the men how reliable this method of assessment might be; and in providing evidence on the value of the routine therapeutic measures. (Table VII. 5). (1) Wrong schedule used: excess decompression

In 30 dives the decompression was prolonged for periods varying from 5 to 40 minutes from that intended. A major incident occurred in one dive that had 15 minutes extra decompression and 4 minor incidents occurred once each after 5 minutes; 5 minutes; 15 minutes; and 40 minutes extra. In detail, 15 dives were decompressed on the schedule for a dive 10 feet deeper than that actually carried out (2 minor cases); 11 dives on the schedules for a dive 5 minutes longer than the one performed (1 major and 2 minor cases); and 4 dives on a schedule both 10 feet deeper and 5 minutes longer than that needed for the actual dives (no cases).

(11) Wrong technique used: excess decompression

In addition 36 dives (including 5 dives using the submersible decompression chamber) had extra decompression added to the schedules through a misunderstanding. Previous tables specified a time for ascent to the first stop from the maximum depth and then a period at that stop. This meant that divers paid too much attention to the overall time and not enough to the manner of the ascent \* / R.N. divers carry out their own ascents under signal control from the surface instead of being hoisted on a stage as happens elsewhere. 7 It is strongly suspected that dives with irregular ascents alternating fast and slow periods are more productive of cases of decompression sickness. In this work, trial orders included instructions to ascend at a steady rate, 60 feet per minute being the target rate and the divers were not to leave the first

-72-

atop till a specified period after leaving the bottom had elapsed. The trial tables thus had the time for ascent included in the time of the first stop. Initially this was not appreciated by older divers and the whole of the combined period was spent at the first stop. It has always been thought that the depth and duration of this stop was more critical than the other stops and hence the extra periods which varied from 1 to 3 minutes disaqualified the schedule for comparative analysis and assessment. One major and seven minor cases occurred in this group.

# (iii) Miscellaneous: excess decompression

12 other dives - with one minor incident in the group - were excluded for various errors which also increased the total decompression time. These slips included errors in arithmetic by the recorders; accidents involving delay in the diver changing his depth on time, such as getting the lines fouled round the guide ropes; and taking too long to leave the water.

### (iv) Insufficient decompression

11 dives altogether were discounted because of inadequate decompression. The commonest fault (6 dives) was the use of a schedule for a shallower dive than the one done so that the reduction in time veried from 15 to 30 minutes - 7 major cases of degeompression sickness resulted. One dive had 15 minutes less decompression as a schedule for a dive of shorter duration than the one intended was used without illeffect on the diver. The remaining two dives, also without ill-effects

»73»

on the divers, had 5 minutes deducted because of arithmetical error, the slip being made at the shallowest stop.

(v) Submersible decompression chamber dives

In 19 dives the decompression schedules used wore the ones intended, 10 on the British table and 9 on the American table. However to evoid prolonged stops in cold water the submersible decompression chamber was used in such a way that the diver could climb into it at his first stop. The greater confort was desirable but the exercise involved in climbing into the chamber during decompression and removing parts of the equipment was potentially a cause of increased incidence of decompression sickness. The results when compared with the same dives causied out in the sea without the use of the submersible chamber were thought to show an increase in "bends". The use of the chamber was discontinued and the dives were not included in the final analysis. (All these dives were causied out at the same dopth for the same time interval).

-74-

Type	s of Faults	Minor Cases	Major Cases	
CLOSS	Comment	Number	Number	Number
an a	Actual dive shallower	15	2	0
	Actual dive shorter	11	ŝ	1
1kriza	Actual dive Shorter and shallower	Ą.	0	0
Decompression	Ascent edded	31	7	0
	Miscellaneous	12	2	0
este suetas un directo de tracito de actual de como	Sub-total	73	13	1
	Actual dive deeper	8	0	7
Insufficient	Actual dive longer	1	0	0
Decompression	Stop cut	2	0	0
nen ne mel mel keye tan ter piko Angana yakt keya tang angan yakt keya tang	Sub-total	11	0	7
Submorathle	Planned	19	4	1
Gramber used	Ascent added	5	0	1
nygesne o navyt zen nygynytten tieken nyt ynten hei innen ytt	Potel	109	17	10

# Table VII. 5

Summary of Dives Encluded from Analysia

# Effect of Duration and Duvironment of the Dive

When considering the project initially, it was thought that if the tendency of the decompression was to be inedequate, the longer the duration of a dive the more the deficiences would become apparent. It was also commonly accepted that a sea dive would disclose deficiences in a schedule not necessarily apparent after a chamber dive. It was decided that a chamber dive should be done first because a case of decompression sickness anising during decompression could be looked after better if the dive took place in the means of treatment, and because too great an incidence of cases would indicate that that schedule should not be used in the sea.

Table VII. 6 compares the relative durations of dives on each table. Table VII. 7 the environment of the dives on the tables and Table VII. 8 the effect of duration and environment of the dives.

				leses	Ma <b>jor C</b> asos	
Relative Duration	Table	Number	Number?	¥0	Munder	%
Short	л В <sup>.</sup> С	172 98 17:4	31 8 22	16.0 8,2 12.6	5 1 2	2.9 1.0 1.1
	Total	lj44	61	13.8	8	1.8
long	A B C	<b>10</b> 0 73 39	25 16 10	25.0 21.9 25.6	8 7 7	0.0 9.6 17.9
	Totel.	212	51	24.1	22	10.4

### Table VII. 6

### Duration of Dives and Tables

Note: "Short" and "Long" : See Graph VII. 1

-76-

D:				Minor Cases		Major Cases	
Environmenti	Table	Number	Number	9%	Mmber	<i>%</i>	
Dry	A B C	112 2 152	31 1 20	27.7 (50) 13.1	3 1 8	2.7 (50) 5.3	
	Total	266	52	19.5	12	4.5	
Wet	A B G	160 169 67	25 23 12	15.6 1346 19.7	10 7 1	6.3 4.1 1.6	
	Total	390	60	15.4	18	4.6	

# Table VII. 7

# Environment of Dives and Tables

Note: "Dry" - dive in a compression chamber

"Wet" - dive in the sea

Divos			Minor (	20869	Najor Cases	
Environment	Relative Duration	ilumbor	Number	%	Mumber,	%
Dry	Short Long	193 73	24 28	12.4 38.4	3 9	1.5 12.3
Werb	Shor <b>t</b> Long	251 139	37 23	14.7 16.5	5	2.0 9.4

# Peble VII. 8

# Environment and Duration of Dives

It will be seen that dives of short duration were less likely to lead to cased of decompression sickness than longer dives and that this tendency persisted whether the dives were done in the see or in a compression chamber. It was remarkable in view of the widely held opinion that see dives were always more dangerous than chamber dives that the results were comparable. The figures for each set of tables show dissourcically opposite tendencies between the American tables wish increasing - and the Card tables - wish decreasing. As the American tables generally had shorter decompression times than the British tables which were in turn shorter than the Card tables for the same dives, one gains an impression that for short dives, the longer the decompression the safer the dive. It is difficult to explain at present the reasons for the apparent increase in risk in longer dives as all decompressions become longer.

#### The Tables

There are two points of comparison for any given depth and duration of dive as far as decompression tables are concerned. These are the total decompression time and the distribution of the stops, - in particular the depth of the first stop. Both these points are shown in Table VII. 9.

-78-

Ĺ.	ivo	Decompression Tebles					83 546 546 646 646 647 547 547 547 547 547 547 547 547 547 5
Dopth	Depth Dunation		Total Time (min.) Depth First S				
feet	min.	A	В	C	А	1)	C
80	69	36	41	51	20	30	30
90	70	41	46	56	80	30	30
100	<b>60</b> <sup>-</sup>	44	46	51	20	30	30
110	50	36	41	46	20	30	30
120	45	46	41	46	80	30	- 30
130	40	41	41	46	20	30	.30
150	30	36	41	46	20	30	40
160	30	41	46	**	30	40	-
180	25	41	46	51	30	40	40
130	45	56		61	20	and the state of t	40
70	120	96	56	**	20	30	-
60	180	76	86	-	20	40	atia
120	75	101	111	116	30	<del>5</del> 0	50
140	60	101	111	126	30	50	50
150	60	116	131	and the second s	40	50	
160	60	136	146	156	40	60	60
170	45	101	106	121	40	50	60

# Table VII. 9

Comparison of Total Decompression Times and Depth of the First Stop for the Dives Done on the Various Tables

#### a) Total decompression time

As is elveys the case in diving triels, there is no consistent difference. In the shorter dives carried out. the total decompression times of the American tables were the same in 1 schedule. 5 minutes shorter in 7 schedules, and 5 minutes longer in 1 schedule than those in the British tables for the seme dives; those in the Card tables vero 5 minutes longer in 6, and 10 minutes longer in 2 schedules then those in the British tebles, and 15 minutes longer in 2, 10 minutes longer in 4, 5 minutes longer in 2 and of the same duration in 1 schodule when compared with the American tables. In the longer dives the American total decompression times were the same as those in the British table onco, 5 minutes shorter once, 10 minutes shorter 4 times and 15 minutes shorter once; the Card tables were caly dived on 4 schedulos for long dives and these were longer then those in the British tables by 5 minutes once, 10 minutes once and 15 minutes twice and longer then those in the Amenican tables 15 minutes once, 20 minutes twice and 25 minutes once.

b) Dopth of first stop

The first stop is considered to be especially important as this marks the maximum difference for elimination of absorbed gas between dissolved tonsion and alveolar pressures i.e. the extraction gradient. In this respect the American tables have a first stop 10 feet shallower then the British tables in the 9 shorter dives done on both tables and 10 feet shallower in 3 schedules and 20 feet shallower in 4 schedules for the longer dives. The Card tables have the same depth of first stop

-80-

in 7 schedules and 10 feet deeper in one compared with the short Dritish tables and 10 feet deeper in 7 and 20 feet deeper in two schedules compared with the short American tables. In the longer dives the Gand tables have the same depth at first stop in 3 and start 10 feet deeper than the British in one schedule and are 20 feet deeper than the American tables in all four schedules. The importance of the depth of the first stop is thus less critical then previously essuned when the difference is only a few feet between possible choices. The Gazet of Decompression Sickness

One would expect that the more often a man dived on an experimental table the more likely he was to suffer an attack of decompression sickness in some form and probably both minor and major in sevenity. This expectation is based on the chance that he will dive scener or later on an inadequate schedule. This is borne out by the details in Table VII. 10.

Group	Mumber of Divers	Numbor of Dives	Number of Minor Cases	Number of Major Cases
No trouble Minor trouble only Major trouble only	33 34 13	137 312 87	75 •	- 14
Minor and Major invable	16	.229.	54	26
Total	96	764	129	40

### Rable VII. 10

#### Divers Grouned According to Attacks of Decompression Sickness

The surprising groups are those who dived fairly frequently but suffered only one form, either minor or major. Consealed in the data are the records of individuals who were exceptional in their performance. For example 3 men carried out 52 dives between them without any incidents and 6 men who carried out 108 dives between them, had an average of 4 minor incidents per man. A simple analysis of the four groups is given in Table VII. 11. Study of the men in these groups did not show any obvious difference in experience, age or body build, nor was there any difference in the particular diving trip and season. Some of the more notable records in each group are given in Table VII. 12.

Group	Detei 1	Rengo	Avorage	Nedlan
No trouble	Dives	1 - 23	4.1	3
Minor troublo only	Dives Cases	1 - 22 1 - 7	9 <b>.</b> 2 2.8	8 2
Major trouble only	Divob Cosos	1 - 14 1 - 2	6.7 1.1	9 1
Ninor and Major trouble	Dives Minor Cases Major Cases	1 - 33 1 - 8 1 - 4	14.3 3.4 1.6	11.5 3 1.5

### Table VII. 11

Some Details of Groups of Divers

-82-

Diver	Minor Attacks	Major Attacks	Number of Dives	% Incidence of trouble
DA		***	23	
1(1)	****	-	13	
RUB		***	16	-
127	7		21	33
<b>TEA</b>	S	, min	4	50
ETI	2	-	4	50
RO	5	-	4	50
TH	4		55	18
BG		2		4,0
INJ	,3	1	. 7	57
OA	3	2	12	42
GN	l.	1	10	50
FIA	3	5	5	100
TIO	. 8	2	13	77
RA	3	<b>4</b> .	55	33
145	2	2	. 7	57
m	8	8	33	30
SØ	2	1	50	12
110	5	. 1	24	25

Table VII. 12

Some Notable Records

ş

.

# Symptoms and Signs in Minor and Mador Cases

(1) Pain

The commonest symptom was pain which varied in severity from such as to cause extreme restlemeness to a persistent ache which could lead to "syncopal attacke" and, in the mildest cases to an ache which was a vague disconfort constantly obtruding on the attention and difficult to locate accurately. In diving jargon these were differentiated as "pein", "ache" and "niggle" respectively. Very rarely could the location be given by pointing with one finger; more often the whole hand was placed over the area involved. The pain often radiated proximally or distally or in both directions. This symptom was recorded after 102 dives and in over 30 of these dives the pain was in more than one cite.

-04-

Eight dives resulted in the men having miscellaneous types of pein such as generalised aches, headaches, colicky abdominal pain and aching in jaws. The only type that would seem to have an obvious explanation are the abdominal cases in that gentric and intestinal distension from air swallowed while under pressure is well known but pain erising from other abdominal sites had to be excluded.

The joint and its surrounding area most frequently involved in this trial was the right shoulder which was specifically mentioned on 23 occasions and included in "both shoulders" on 4 more. By contrast the left shoulder was only specified on 9 occasions. The shoulder girdle and various parts of the upper limbs were reported as the site of pain on 78 occasions. The polvic girdle and lower limbs were recorded as the sites of pain on 31 occasions (left knee 9 times, right knee 7 times, both knees once). The predominance of the upper limbs in this trial is probably due to the need to exercise the arms to maintain position at a stop during decompression; to change depth with stops; and to operate the control valve high on the right side of the helmst... Often the man preferred to wait to see the progress of his pain before reporting and with experienced man, it was a useful guide to ask them if they wished to be recompressed. A great many cases responded to local heat and rest. Analgesies were rarely given lest masking of developing severity of pain occurred.

(11) Itoh and rash

Itching was noted after 68 dives and its distribution varied from head, face, and nocky limbs; to torse but nover on hands or feet even in chamber dives. The true incidence was difficult to assess in that if a diver was asked if he had an itch, he developed one. Some man stated that they always itched after a dive but as it was nover troublesome they did not worny about it and rarely reported it. No diver was ever seen to have exceriminant of his skin due to scratching and in any case a hot shower or bath cleared the sympton from all cases. Little importance was attached to this symptom as experience was compatible with some reported lineratory investigations (129).

Rash was reported after 19 dives. Typically it was irregular blotoby with no malsed edges and varied in colour from dusky blue to bright red even in a single patch and described as mottling or marbling (Fig. 11). The lesions were confined to the shoulders and trunk and were normally noticed during the undressing of the diver or in his customary post-dive shower or bath when his skin usually returned to normal. If the diver delayed his ablutions, the blotchy lesions could persist for some hours. Recompression cleared the discolourations.

-86-

No particular significance is attached to itching or to rashes when no other symptoms were present but they were considered, when volunteered or observed amongst the other factors in assessing the adequacy of the decompression schedule. Heny men are not apparently affected and therefore such ailments could be skin signs of decompression sickness. The effect of local heat suggests that both itches and rashes might be due to bubbles present in the tissues of the skin. This is not necessarily a guide to the degree of tissue saturation elecenters in the body as these cases usually arise where the divers clothing permits adequate air space round his body with the possible effects of percutaneous diffusion of gas adding to the local tissue saturation levels.

### (iii) Control Norvous System

Thirteen times a diver became pale, light-headed, and looked about to faint. The whole picture suggested shock but the pulse was usually about normal in rate for the individual and of adequate volume on palpation. This picture was sometimes complicated by low chest pain, nausea and vomiting, dizziness and exhaustion. Rapid recompression was the treatment of choice and was rapidly effective.

Disturbed sensations also occurred; misty vision of the left eye once; diminished sensation of a thigh, or of a hand, or of a foreann, after other dives; paraesthesia in the distribution of an ulnar nerve twice; and weakness of a limb three times. Four times there was a sensation of stiffness, once in the neck and three times in the right knee, without associated pain, swelling or limitation of movement.

(iv) Oodema

The most baffling sign was the painless, non-pitting, ordena which occurred in 4 men, once in the right ankle, once in the left forearm, once over the dorsum of the right hand, and twice in the same man on the face where it started symmetrically over the malar areas. /Figure 12 illustrates the facial lesions/ The fluid gradually dispersed, being aided by gravity especially noticeable in the facial cases. Notes on major cases of Decompression Stolmess

After each case a note is made on whether the case was treated according to the instructions recommended in the N.N. Diving Menual (Appendix II); if not it is noted whether the therapy was longer or shorter than the recommended procedure.

a) <u>Chember dives</u>

<u>Diver PF</u> reported helf-an-hour after surfacing from a dive to 150 feet for 30 minutes on the Card table. He had fairly severe aches in his right arm and right hip which cleared at 22 feet on recompression.

-87-

He surfaced after treatment on Therepeutic Table I with no ill effect. (Recommended)

<u>Diver KA</u> reported two and a helf hours after surfacing from a dive to 140 feet for 75 minutes on the British table. His main completint was mild pain in his left knee of doubtful causation so he was treated by a hot bath which relieved the pain but a sensation of weakness slowly developed. The pain recurred and one hour after reporting, he was recompressed. His symptoms cleared at 120 feet. Therapy on Table II was completed successfully. (Recommended)

<u>Diver MS</u> reported four and a half hours after surfacing from a dive to 120 feet for 75 minutes on the Card table. An ache in his right knee was relieved at 55 feet. He was successfully treated on Therapeutic Table I. (Recommended)

<u>Diver Hit</u> reported three hours after surfacing from a dive to 120 feet for 75 minutes on the Card table. An ache in his back was relieved at 22 feet. There was no recurrence after treatment on Therapeutic Table I. (Recommended)

<u>Diver PP and Diven MS</u> reported on surfacing from a dive to 160 feet for 60 minutes, on the Gard table. Diver FP had pain in both forearms and upper arms while Diver MS had pain in his left knee. Both men were recompressed and they were relieved by 105 feet. Therapeutic decompression Table II was used. (Recommended). <u>Diver FU</u> carried out the dive with these two men but he did not report till the next day with general aches and a headache. He was recompressed and all his symptoms

-88-

cleared at 55 feet. In view of the delay and site of symptoms, it was decided to use Therapeutic Table II. (Longer). He surfaced feeling well and there was no recurrence of symptoms. / If the headache had been considered to be due to contral nervous system injury, then Table III should have been used and his treatment was shorter than recommended/

<u>Diver SL and Diver SC</u> carried out a dive to 160 feet for 60 minutes on the Card table. Diver SL reported a severe pain in his right shoulder on surfacing and he was immediately recompressed, relief being obtained at 45 feet. Therapeutic Table I decompression was carried out uneventfully. (Recommended). Diver SC did not report till the following day when he complained of persistent mild aches in both knees which were not entirely relieved in spite of therapy on Table III. (Longer)

<u>Diver NP</u> reported with severe pain in his right shoulder after a dive to 150 feet for 30 minutes on the American tables. He was recompressed and after his symptoms were relieved at 49 feet, he was decompressed on Therapeutic Table I. (Recommended)

<u>Diver WS</u> carried out a dive to 100 feet for 60 minutes on the American tables one afternoon. Next day, he reported with severe pain in an ankle and he was recompressed. The pain cleared at 44 feet and treatment was then carried out on Therapeutic Table I. (Recommended)

<u>Diver IN</u> carried out a dive to 140 feet for 60 minutes on the American tables and reported with pain in his right shoulder and arm. He was treated by recompression but the remaining details have been lost. (Assumed recommended as he himself was the doctor in attendance)

-<u>89</u>-

## b) Sea dives

<u>Diver HO</u> reported severe pain in his right shoulder on surfacing after a dive to 130 feet for 40 minutes on the Card table. He was recompressed and, after he had obtained relief at 42 feet, Therapeutic decompression Table I was used. (Recommended). Shortly after surfacing, for the second time, a residual bruised sensation in his shoulder became more painful. He was again treated by recompression and decompression was carried out from 41 feet, the depth of relief, on Eherapeutic Table IV. (Recommended for a recurrence)

<u>Diver HO</u> reported on another occasion with an ache in his right shoulder after a dive to 120 feet for 45 minutes on the American tables. On recompression his symptoms were relieved at 49 feet; he was treated with Therapeutic Table I. (Recommended)

<u>Diver MO</u> developed an ache in his right shoulder after a dive to 120 feet for 45 minutes, on the American tables. His symptoms cleared at 60 feet on recompression. Treatment started on Therapeutic Table I (Recommended) but after surfacing, his symptoms recurred. He was again compressed and this time his symptoms were relieved at 40 feet from where he was decompressed on Therapeutic Table IV. (Recommended for a recurrence)

<u>Diver GR</u> carried out a dive to 150 feet for 30 minutes on the American tables, using the submensible decompression chamber in the Letter part of the escent. 3 hours after surfacing, he reported complaining of severe pain in his right upper arm and forearm. He was

~90~

recompressed, his symptoms being completely relieved by the time the pressure was raised to 64 feet. Treatment on Therepeutic Table I was successful. (Recommended)

<u>Diver NM</u> also cerried out a dive to 150 feet for 30 minutes using the submersible decompression chamber but on the British tables. During the escent he developed a pain in his night shoulder which eased and then increased in severity on arrival at the surface. He was recompressed and his symptoms cleaned by the time he reached 66 feet. Therapeutic Table I was used successfully. (Recommended)

<u>Niver FR</u> reported 20 minutes after reaching surface from a dive to 140 feet for 60 minutes on the British tables. His main complaint was itching of both foreanes with other troubles in the form of a mild ache in his right elbow and a faint mottled rash over his foreares. After a hot bath, the itch and the rash disappeared and the elbow ache was improved. However 5 hours after surfacing, the ache radiated into the upper and and he developed somewakeess of his grip. He was recompressed, his symptoms being malieved at 100 feet, and Therapeutic Eable II was used. (Recommended unless weakness not secondary to pain, in which case routing shorter than recommended)

<u>Diver ST</u> dived to 90 feet for 70 minutes on the British tables. Soon after surfacing he reported with a pain in his left shoulder which after a hot shower, radiated up the left side of his neck. On recompression the pain became more severe and was eased a little by tablets of aspirin. During the decompression, at the 120 feet stop, the pain disappeared and

#91#

decompression on Therapeutic Table II was completed. (Recommended)

<u>Diver KA</u> cerried out a dive to 80 feet for 80 minutes on the British tables. After surfacing, he had some itching of his arms and back and 5 hours later he reported with some pain in his left knee and misty vision in his left eye. He was kept under observation for another two hours when although his vision was back to normal, his knee pain was unchanged; he was then recompressed. His symptoms cleared by 66 feet so Therapeutic Table I was used. (Shorter. His visual signs, although clearing quickly, indicated his contral nervous system had been affected). 2 hours after surfacing from this compression, his symptom of pain recurred so he was again compressed. His symptoms cleared at 65 feet and Therapeutic Table III was used for the subsequent treatment. (Shorter than recommended for a recurrence)

<u>Diver Wil</u> developed pain in his right shoulder about 2 hours after surfacing from a dive to 120 feet for 75 minutes on the British tables. On examination there was some minimal loss of power of his right hand which was unaltered after a hot bath. Half-an-hour later there was a definite weakness in flexion of the right elbow, so he was recompressed. As all signs and symptoms had cleared by 66 feet, he was treated on Therepeutic Table T. (Shorter)

<u>Diver 20</u> dived to 120 feet for 75 minutes on the British tables. He reported 2 hours after reaching surface with pain in his left shoulder and upper arm. He looked unwell and he had weakness of his left arm and hand. On recompression, he made a complete recovery at

-92-

25 feet and he was treated offectively on Therepeutle Table I. (Shorter)

<u>Diver WL</u> complained of pain in his left shoulder after reaching surface from a dive to 80 feet for 80 minutes on the American tables. The pain radiated in a mild stabbing fashion to his left wrist and into his neek. The whole incident was not severe in character and was improved by local heat and codein compound tablets. However there was no further improvement in the next few hours so he was recompressed. Initially the pain was aggravated and he developed weakness of his left hand but he became symptom free at 100 feet. He was treated on Therepeutic Table II and surfaced with a very mild residual bruised sensation which were off steadily. (Shorter)

<u>Diver AS</u> reported about one hour after surfacing from a dive to 120 feet for 75 minutes on the American tables with general malaise which became more severe over the next hour during which he also developed aching in all his limbs. He was recompressed with complete relief of his completints at 22 feet. After surfacing from decompression on Sherapeutic Table I, he had a slight residual ache in his left knoe which gradually cleared. (Shorter)

<u>Divor KA</u> dived to 120 feet for 75 minutes on the American tables. One and a half hours after surfacing he reported with pain in his left knee, temporarily relieved by a hot bath and aspirin tablets. He was compressed with relief of his symptoms at 100 feet. He was treated on Therapeutic Table II (Recommended) but on reaching surface his symptoms recurred. He was again compressed; his symptoms were not fully cleared till 165 feet vill 105 feet was reached and he was then decompressed on Therapeutic Table III successfully. (Shorter than recommended for a recurrence)

<u>Diver CR</u> dived on the same day as the two preceeding cases to 120 feet for 75 minutes but he was decompressed on the British tables. He reported with mild aches in his left elbow and both legs after surfacing but he preferred to carry out some social duties than be treated. 4 hours later he felt very unroll and complained of severe pain in both hips, across his abdomen and in both legs. He was palo and cold with a pulse rate of 64 per minute and he suddenly fainted while being taken to the compression chamber. He regained consciousness at 20 feet on compression and there was only a slight residual ache in his left elbow at 465 feet. However on beginning decompression he developed a slight ache in his right shoulder which persisted unchanged during the decompression on Therapeutic Table III and gradually cleared at the surface. (Shorter)

<u>Diver BG</u> dived to 140 feet for 60 minutes on the British tables. He folt generally unvell on surfacing and he reported 1% hours later when he was seen to be unsteady in his gait, pale and cold, with symmetrical malär symbling and general weakness. He complained of a pain in the front of his chest, his pulse was 80 per minute and his blood pressure 95/65 mm.Hg. A short period of recumbency did not lead to improvement but on recompression his condition improved steadily and he had recovered- except for the facial swelling - within 10 minutes at 165 feet. After surfacing from a decompression on Therapeutic Table III

-91.-

he felt tired and bruised. (Recommended). The facial swelling decreased over the next two days; his gait was stiff legged for about a day.

<u>Diver KA</u> dived to 150 feet for 60 minutes on the American tables. 1 hour efter surfacing he reported with aching in his calves and his left knee, slight pain in his chest and general malaise. He was kept under observation and in another hour as his general condition had deteriorated he was recompressed with relief of his signs and symptoms at 150 feet. Decompression on Therapeutic Table III was successful. (Recommended)

<u>Diver 36</u> surfaced from a dive to 160 feet for 60 minutes on the British tables apparently well but 2 hours later he complained of general malaise, abdominal pain, cold and sweating, and feeling faint. In spite of lying flat, his condition was unchanged so he was recompressed and steadily he improved till after 15 minutes at 165 feet he was completely recovered. He was decompressed on Therapeutic Table III uneventfully. (Recommended)

<u>Diver Mi</u> dived to 160 feet for 60 minutes on the British tables. Soon after surfacing he reported an ache in his left shoulder which gradually vorsened. Aspirin tablets and local heat had no effect. He gradually developed loss of power of his left arm and hand due to pain and he was recompressed. His troubles cleared at 64 feet and Therapeutic Fable I recompression was carried out successfully. (Shorter)

-95-

Divor MU. Diver OR. Diver HA and Diver HI all dived one afternoon to 170 feet for 45 minutes on the American tables. The vector suddonly deteriorated, while the men were at various stages of their decompression routines. It was decided that the men would complete the dives in the sea instead of surface decompressing but in the final stages they had great difficulty in maintaining their depth on the stops because of the disturbance of the water. Diver CR started his symptoms with itching and a mild ache in his left upper arm at his 20 feet stop and which were not too severe on reaching surface. His condition deteriorated very clowly and 2 hours later, he was recompressed with relief of his symptoms at less than 60 feet. Diver NA had only mild niggles at his 20 feet stop but they became aches after surfacing and 3/4 of an hour later he was recompressed with relief at less than 60 feet. Moor M developed pain in his right knee at the 20 feet stop which was severe on reaching surface so he was recompressed with relief of his symptoms at less than 60 feet. These three men were compressed simultaneously in one chamber and were all satisfectorily decompressed on Therapeutic Table I. (Recommended for all three). Diver MI was the last to reach surface (within 5 minutes of Diver MJ) and as he reported well except for exhaustion, the therepeutic routine started without him. Shortly afterwards he developed a severe ache in his upper arms and suddonly fainted. He was immediately recompressed in a second chamber and completely recovered by 60 feet and decompression started on Therapeutic Table I. (Shorter). He developed nausea and he had an attack of

-96-

vomiting during the decompression; on surfacing he had a mild pain in his right knee and a bilateral facial swelling of non-pitting orders over the malar areas. As sheep was impossible because of restlessness due to the irritating ache, he was again compressed with marked improvement in his confort. Therapeutic decompression Table III was used (Shorter) but he still had a marked residual bruised feeling in his knee and the facial availing which had gravitated to the areas over the jaws. Over the next two days the bruised censation cleared and the swelling dispersed over his neck. / This case was complicated as it transpired during therapy that he had concealed a "hangover" and a very late night in the hope that by his turn to dive in the late afternoon, he would have recovered.

<u>Diver HA and Diver HE</u> dived on the same day to 460 feet for 60 minutes on the American tables. At the 10 foot stop Diver HA developed an ache over his right elbow which became more severe at the surface although he did not report for 1% hours. He was obviously in pain, and there was general swelling of the dorsum of his Right hand with patches of paracethesis of his fingers. He was recompressed with relief of all his troubles, except the swelling, by 30 feet. Diver HS also was relieved at 30 feet of his symptoms which had been aches in his Right shoulder and Right elbow on surfacing slowly becoming more severe till there was limitation of power of all muscle groups at the shoulder and distally. Therepeutic decompression on Table I was used uneventfully. (Shorter for Diver HA. Shorter for Diver DE if weakness was not secondary to pain)

Diver DA. Diver MJ. Diver MI and Diver MD dived on the same day in a flat calm sea using the British tables, to a depth of 130 feet for 40 minutes. Diver DA reported 24 hours after reaching surface that he felt faint and giddy; that niggles in his right shoulder on surfacing had gradually been developing into severe sches; and he was found to have percepthesis of the almar distribution of his left hand. He was recompressed with relief of his symptoms at 165 feet and Therapeutic decompression Table II started. (Shorter). At 10 feet he had a mild recurrence of his symptoms and on reaching surface. he developed pain in his left shoulders, left hip and left thigh so he was again compressed. His symptoms cleared at 25 feet and decompression was carried out from there on Therapeutic Cable IV. (Recommended). Diver MI reported 1 hour after his dive and at the same time as Diver DA. but his compleint was headache and mild aches in his right shoulder unrelieved by a hot bath. Re was compressed with his colleague and he also did not get relief till 165 feet. Therepeutic decompression Table II was the one used. (Recommended) and he reached surface without compleint; on arrival he reported continuing malaise and pain in his left shoulder so he also was recompressed, again with relief at 165 feet. Decompression on Thorapeutic Table III was used (Shorter) but 4 hours after surfacing he again had symptoms and recompression was again required - this time therapy was on Therepoutto Table TV. (Recommended). Diver KI reported 40 minutes after reaching surface that he had discovered a general mottled body rash

-98
while he was having a hot bath for some aches in his left upper arm and across his back. As he reported while the other cases were being considered, he was recompressed with them and his symptoms cleared quickly. After the completion of the decompression on Therapeutic Table II (Longer) he had a residual bruised sensation which cleared gradually over 24 hours without further therapy. <u>Diver NB</u> reached surface after the initial therapy of his three colleagues had started and, after a medi and shower, he had a map from which just more than 2 hours after surfacing, he wakened to find he had difficulty in standing, and in moving his limbs because of severe pain. He was recompressed and all his symptoms cleared at 40 feet. He was decompressed on Therapeutic Table I with no ill-effects remaining. (Shorter)

-99-

## Therapeutic Regimes

169 cases of decompression sickness occurred but only 38 were recompressed. It is difficult to say whether this proportion is small or large but under the conditions of close observation in such work it is probable that some cases that were treated by local heat and rest might, elsewhere, have been recompressed. Part of the reason for not recompressing marginal cases immediately was the desire to build up mutual confidence so that men would report or admit their symptoms however trivial as soon as possible. It is obvious from the case historics that the time to reporting did decrease as they are arranged in chronological order (except for the last 3 chamber cases which occurred after the first few see cases). Residual symptoms were not always entered in the records as a sensation of stiffness and bruising was accepted as a common temporary terminal phase of treatment in confined conditions.

On the whole, the therapoutic tables in the N.N. Diving Hanual (132) were used as recommended; this particularly applied to cases of uncomplicated pain, especially those where examination disclosed no abnormality and the diagnosis was perforce made on the history alone. Sometimes when the pain was accompanied by a history of long delay in reporting, or when effects such as limitations of movement or syncope were thought to be secondary to pain, the basic tables for pain were still used. Other cases which had neurological signs and symptome should always have been treated on the longer routines recommended. The longest routine was also the one supposed to be used for recurrences arising during a therapeutic decompression with the decompression restarting from the depth of relief.

16 cases were not treated as recommended, 3 of these using a longer routine. 6 of them were treated on a shorter routine with acceptable results. In the tenth case, his pain changed to "weakness" during decompression which cleared before maximum pressure was used and he was treated as if he had pain only but there was a residual bruised sensation on surfacing. The eleventh case had had malaise with his pain but he was treated on a pain routine which left him with a residual mild ache. The last five cases recurred; two after following the recommended tables and three after following shorter therapeutic routines.

-100-

The first pair were treated the second time on a shorter table than the one recommended; the second group were also recompressed with the diver having the recommended routine responding as well as one of his colleagues and a little better than the third in his group both of whom had a shorter therapy. Of the 22 cases treated as instructed, 2 men had a recurrence of signs or symptoms which resulted in further recompression once each along recommended lines.

The deterioration in the symptoms of two men during compression is of interest. The correct action was clear in the one whose symptoms cleared while the compression continued but the explanation for the man who suddenly improved during the course of subsequent decompression was not clear. While the relief of the patient is often so dramatic that one can find a precise offective pressure. it is perturbing that sometipes after a further increase in pressure. Often considerable. the patient may report further symptoms. The other point that raises questions about the assumptions on which the therapoutic tables are based was the appearance of pain during or after the use of the tables longer than those intended for pain cases on two out of three occasions. An explanation of these unexpected events might 11e along the following lines. Dramatic relief of signs and symptoms occurs where the size of the traumatic bubble is rapidly reduced. If the reduced bubble does not redissolve but persists, then, if in an intravascular site, it could be driven further into a consistive area. It it persists, whether intravascular or extravascular in site, then reactive oedema might occur

-101-

with imperiment of see exchange so that the hubble does not dissolve but may even grow, both by bubble physics leve and by any subsequent decompression. On this explanation, signs and symptoms should then be treated as evidence of a inibile and the nature - whether pair or collepse or some other - is of interest only as site. A loose englogy is the freetune of a bone which is treated primarily by the basic principle of immobilization and secondary by firstion, traction, or splinting. A separation of decompression sidess cases into "pain cases" and "others" suggests that symptom is more important than site whereas the injury has to be established and then appropriate treatment given. While generally cases with a particular symptom might be treated in a similar memory it does not follow that the injury is similar in size, site, or reaction. There is no clear evidence as to whether such inibles are intravascular (8) or extravascular (65) although animal work and post-mortems have suggested extrevescular sites, clinical progress of a case as often is indicative of intravescular. The impression is geined that once bubbles are presumed to appear then all theoretical curves of ges elimination are at best very poor approximations. Under these circumstances, the administration of pure oxygen was not really indicated and any beneficial offects were probably countered by the disconfort of the equipment for its administration evailable during this trial.

## Comment

These results show the crudity of this approach to the evaluation

-102-

of decompression tables and the futility of theories and experiments designed to add a minute here and to subtract a minute there when planning new tables. It seems that the larger increments edded operationally by using the decompression schedule for a deeper and/or a longer dive, than the one actually dived, may reduce but not abolish the risk of decompression sickness. A radical view might be that the assumptions on gas elimination by maximum extraction gradients are wrong; that all tables have the first stop too shallow; and that there is a possibility of giving too short a decompression time at deeper stops and too long at shallower depths. An alternative view is that the lesion of decompression sickness, the hypothetical bubble, may arise or indeed always arises - during the initial decompression and thereafter any distribution of decompression time is almost immeterial; signs and symptoms would then appear in a proportion of cases varying with the actual site of the lesion, the size and numbers of such bubbles, and the susceptibility of the individual on that day.

There was little to choose between the American and British tables on such crude testing and no real assessment of the theories behind each table could be made. It was found that environment did not have much effect, provided the depth could be accurately controlled. It appeared that increasing the total decompression time did not always lead to an increase in safety suggesting that there is an optimum time - perhaps for the individual rather than the dive. No explanation was obvious for different susceptibilities to decompression sickness

-103-

among the divers. The treatment nontines had some surprising immediate feilunes in view of the apparent specific corrective procedure applied to a traumatic physical agent. Cortain departures from recommended mutilities demonstrate an approach which might be of value in modifying initial therapy.

## GHAPTER VIII

## MULPIPLE DIVIS

## Problem

All methods of celculating decompression schedules assume that the diver will reach atmospheric pressure with excess ges dissolved in his tissues which will. however. cause him no harm. While he remains at atmospheric pressure, he will gradually come into equilibrium with his environment. If he dives before he has climinated all the excess gas previously absorbed, he will start taking up gas on top of a higher tissue level then if he had started at coullibrium. Diving tables are devised primarily for the diver in equilibrium and they may entail too great a risk of decompression sickness for routine application to the man carrying out more than one dive in reasonable succession. Some formula is required before the tables can be applied to the dive to discover the decompression needed. Various mathematical manipulations are available to find a solution. The principle is similar in all. that is to attempt to convert into a measure of time the excess gas remaining in the diver's tissues at the start of a subsequent dive. This time may be considered either as duration to be edded to the time on the bottom of the subsequent dive or as decompression time to be added to the total decompression time of such a dive. This increment is added as appropriate to the duration or decompression of the current dive and the actual decompression schedule to be used is found in the In offect an equivalent fictional dive is obtained and combined tables.

with the actual dive to produce a schedule that will pommit the diver to surface with an acceptable level of tissue gas saturation. In these trials the extra time was added to the initial total decompression time and the distribution of stops found by looking in the tables for the dive at the same depth with the same or next greatest total decompression time as the combined figure. (102)

## Notes on Technique

The diving method was routine as far as depth assessment and the actual performance of each dive was concerned; standard helmot-diving practice as previously described was the method used. It was intended to camp out 10 pairs of dives in each series so that a dive would be followed after a specific interval at atmospheric pressure by a repotition of his first dive by each man. In practice because of tides, a dive was paired with enother depth and time combination which was considered to be equivalent in gas uptake and elimination. The "double dive" plan was prepared with different surface intervals as well as several groups of paired dives and the envisaged programe was rather daunting in the total number of immersions anticipated. To make the maximum use of time and personnel a complicated time table was evolved. which, in any one day, mixed different combinations of dives with different surface intervals and which, it was honed would even out the effect of variables such as weather and individual responses. The dives carried out before the programe was abandoned are summarized in Table VIII. 1.

-106-

Table VIII. 1 Record of lives

÷

			E	IS	88	a ə	Diver		
ಸಕಾರಿ	<b>≅</b> ⊐ ನ∰ನ ⊐	€~∞ võ	J.	4	or W	יס גע Dive		Chronological Order of	-
140	140	14.0	180	190	180	160	Depth Ft.		
15	5	25	20	20	20	20	Duration Min.		First Dive
10	<b>t</b>	8	30	35	30	23	Total Decompression Time Min.		
65	125	9577	125	70	5	5	Interval Min.		Q11110000
041	140	140	180	180	180	180	Depth Ft.		
15	Ĵ	25	20	20	20	20	Duration Nin.		- Second D
25	20	55	55	65	65	65	Calculated	Decompre	ive
ଝ	3	65	65	33	85 75	ઙૢ	ssion Mins. Actual		
		SH Bend 11 hours	HU Bend 2 hours		D) Bend 14 hours		Notes with time to reporting bend		

-107-

. . . . .

ł

1

•

It will be seen from Table VIII. 1 that 7 men dived on only one combination each and that 6 mon did the remaining 13 pairs of dives. What is not apparent is, that, as a result of a misunderstanding, only dives 5, 13, 15, 19 and 20 were carried out correctly. When the tables were consulted using the calculated total decompression time. if there was a schedule with the some figure then that schedule gave the distribution of the stops. If there was no such schedule then the next longer schedulo was taken. The latter procedure was done correctly for the five dives mentioned - but the next schedule was also emoneously used for the remaining dives. In spite of this mistake which one would expect would have decreased the risk of decompression sickness, all 4 ceses which needed recompression developed in this group of divors. Some minor cases also occurred but, due to inexperience, any complaints such as rash or itch were regarded as trivial and no record. was made. It will also be noticed that the cases occurred after diver which themselves were routine but where the decompression schedule for the second dive was to be found in Table II in the 1950 edition (132) or below the limiting line in the 1964 edition of the Royal Navy Menual (133). Unusual points are the very long period to reporting signs and symptoms in the case of two of the divers as experimental work often leads to complaints within minutes of reaching surface; and the fact that the extra 45 minutes on the surface interval for Diver PE did not put him less at misk then his colleagues FR and ME, or even better off

## Case Notes Diver HU

This man was very stout being a heavy beer drinker but he could carry out a tolerance (chair-step) test within the 45 seconds allowed for recovery, and his very active organising job with no apparent difficulty. Two hours after reaching the surface, he developed newses and abdominal pain of a colloky nature but examination was negative. Meny possible diagnoses were considered but recompression had to be the first treatment and all his symptoms cleared at 60 feet. It was decided to decompress him on Therapeutic Table I (Shorter). Four hours after he had completed this therapy apparently successfully, his symptoms recurred, again with negative findings on examination. Again on recompression. his symptoms cleared at about 60 feet but this time he was decompressed on Therapeutic Table III (Shorter than recommended for a recurrence). 3 hours later, he again reported feeling sick but it was decided that further recompression would be of little avail as he was thought to be suffering mainly from exhaustion. He was put to bed but 2 hours later still he complained of pain in both knees and it was decided to recompress him for the third time. As soon as he was on his feet he felt better and after 12 minutes walking up and down, his only complaint was extreme exhaustion. He returned to his bed and after an extremely sound night's sleep, he had fully recovered in the morning. It is relevant that he had been instructed to reduce weight the previous week - which he had decided to do with his own diet of cabbage. wun and beer and had lost

3 1bs. It is a matter for discussion whether he ever had decompression sickness but he had to be treated as if he had the injury. It is also debatable whether the first table should have been Table III and the recurrence should have been treated on Table IV from 60 feet. He had had a recurring bend in his right shoulder 3 months before.

## Diver DO

This young man was lean and athletic and carried out his dive satisfactorily, reaching surface at 1700. Next morning he reported that on waking, he had no power in his right hand and fingers and examination confirmed that only very weak movements of the hand were present. Although his lesion was neurological, when power returned completely at 60 feet on recompression, it was decided to decompress on Therapeutic Table I (Shorter); this regime was successful. He had had a bend in his right wrist and forearm 4 months before.

## Diver SM

A new in his middle twentics, he was very stocky and heavy. About 11 hours after sumfacing he complained of pain in both his knees but no abnormality was found on examination. His symptoms were relieved on recompression at 80 feet; he was brought back to atmospheric pressure on Therapeutic Table II (Recommended). (Subsequently, in other work, he developed similar knee bends but never in a shorter period than 8 hours after his dive).

#### Diver PE

This joing man was lean in build. Two hours after reaching the

-110-

surface, he developed pain in his left shoulder, with no abnormal findings on examination. His symptoms cleared at 25 feet on recompression and the use of Therepeutic Table I (Recommended) was successful. He had not had decompression sickness before.

#### Thorapeutio Routine

The initial high incldence of cases of decompression sidness badly affected morale, and it was obvious that no matter how successful the subsequent diving might be, either the basic tables had to be changed or a new mutine devised. Other points to be noticed were that two of the deses were not treated according to instructions, one completely successfully and one with recurring trouble; that the case of Diver HU had to be recompressed whatever the possible diagnosis as decompression sidness was more likely then e.g. gestric distension or eppendicitie; the great variability in response between men which makes assessment difficult; and the possibility of excessive decompression causing trouble rether than preventing it. This trial was carried out at a time when the divers, instead of receiving a daily retaining fee irrespective of work, were receiving a monetary amand based on a rate per minute: this rate varied with the maximum depth of dive and also applied during the recompression treatment for that diver the attendants received a proportion of the total sum avanded to the diver. This encouraged the development of symptoms without signs and only close knowledge and observation enabled the distinction to be made between a gomine and a "financial" bend. In this particular trial, there were

-111-

no such problems; the four cases were genuine and none of the others had any symptoms which might have been misdiagnosed.

## Comments

The decompression theory may possibly be inadequate and when manipulated for this routine of repetitive diving the risk of decompression sickness is too high. The Therapeutic Tables may be used deliberately as four separate routines, each one being used as appropriate irrespective of differentiation by signs and symptoms.

#### CHAPTER IX

#### SURPACE DECOMPRESSION DIVES

#### Problem

Sometimes weather conditions are such that a diver cannot carry out his decompression in the water because strong currents or heavy sees prevent reasonable depth keeping on decompression stops by excessive movement - which may in turn lead to see sickness. His ship might even have to mun for shelter. On other occasions, in self-contained equipment, it may be that the task is of such a nature that it is preferable for the diver to use all his limited gas supply in performing his task rather than be forced to abandon it unfinished. In such cases the advantages resulting from the ability to bring the diver directly to the surface safely and then to recompress him in a chamber are obvious. Such advantages include operations in worse weather, greater confort for the diver, more efficient use of limited gas supplies, a continuous effort on the task, direct observation of the diver, direct reporting of his findings, and less problem in keeping depth during decompression stops. This type of routine had been available with previous tables for emergency use, especially in the case of the helmeted diver who became over-inflated and "blew-up" to the surface. It was necessary to check that the new tables could also be used for such a manoeuvre and some changes in the drill were made in keeping with the principles of celculation of the tables (36).

Notes on Technique

Whereas the divers described earlier were docompressed to stnospheric pressure so that the excess gas saturation did not load to obvious bubble formation, in surface decompression the diver returned to the surface without stopping and was thus in a potentially dengerous situation. Previous routines had forbidden direct surface decompression if the depth of the diver was greater than 200 feet unless one or two stops were made in the water. The drill that was to be superseded was that the diver ascended as quickly as possible without losing control and, at the surface, attendants divested him of his surface lines, weights and helmet before pushing him into a compression chamber where he was recompressed to his original depth. This whole menoeuvre had to be completed in 5 minutes or less. The decompression schedule used to bring him book to atmospheric pressure was that for a dive of his actual depth and of a duration which was a total of actual dive time plus time of drill plus 5 minutes that he spent at depth in the chamber. If the dive was near permissible limits of duration, this addition of 10 minutes could meen that a dive on Diving Table I was decompressed on a Diving Table II schedule (or on a dive below the limiting line - see Table II. 1). The new drill aimed at providing less stimulus to bubble formation by insisting on a steady rate of ascent of 60 feet per minute; and at reducing the amount of excess gas absorbed in the final compression, by returning the diver for a period of 5 minutes, to a pressure only one atmosphere greater

-114-

then the depth his first stop would have been 11 he had been carrying out his decompression in the water. The unknown factor was the presence of decompression bubbles developing during the initial ascent, would the subsequent compression be adequate to return them to solution and only then could one assess how effective was the subsequent decompression. The general timing remained the same except that the actual time of drill was treated as a constant of 5 minutes so that the schedule found in the tables would always be the one for a dive 10 minutes longer then for the dive actually done.

## Results

210 acceptable dives were carried out by 39 individuals over the period September 1957 to March 1958. 12 men carried out one trial dive each while of 18 men who did 5 dives or more each, the maximum number by 1 men was 15. The vest majority of dives were carried out by men dressed in standard diving equipment but 42 dives were done by men wearing the cumbersome self-contained Mine Recovery Outfit which eventually did not reach operational Service.

26 cases of decompression sidness that responded to local therapy and/or time were noted. It was felt that some divers considered the minor symptoms as unavoidable complications of any dive and that they did not report them unless they became troublesome - an impression perhaps borne out by the fact that 21 of the reported cases occurred in 7 men. 1 of these minor cases was unusual and it is described in more detail in the notes on Diver MA.

-115-

9 cases of decompression sidness requiring treatment by recompression occurred in the dives properly carried out. A 10th case - again in Diver MA - developed in one of two dives which were grossly under decompressed, when by error the recorder omitted to add the 10 minute increment when looking up the correct schedule in the manual. This meant that the divers were surface decompressed for the dive actually done. These two dives are excluded from the analysis. The Dives

It had been intended that dives would be cannied out in paired depths c.g. 120 and 130 feet and that they would be of such durations that one of the pair would be well short of the limit in Table I and the other would be on the border-line between Table I and Table II with 10 dives at each of the chosen points. (Well above the limiting line and on the limiting line in the latest R.N. Diving Manual (133)). On studying the records, the dives were found to be widely scattered over the depth range (from 60 feet to 160 feet) and in duration (from very short to over the limit). The dives were thus better classified according to whether the final decompression schedule remained on Diving Table I or whether it was to be found on Diving Table II (132).

-116-

Decompression	Number of Dives	Cases of Decompression Sickness					
Table Used		IJ	nor	Najor			
ana an farman air bha ann an ann ann ann ann ann ann ann an		No.	· 95	No.	ø		
Teble I	105	9	8.6	5	1.9		
Teble II	105	17	16.2	7	6.7		
Totel	210	26	12.4	9	4.3		

#### Teble IX. 1

#### Results arrenard by Decompression Table used

Concealed in the table lie minor departures from the instructions to the attendants conducting the dives. On six occasions between 7 and 9 minutes were added to the actual time of dive, when finding the correct decompression schedule as had been customary on the previous routine, so that the divers underwent a shorter decompression than was intended. On 19 occasions there was no dive schedule for a dive exactly 10 minutes longer than the dive actually done; a schedule for a dive 15 minutes longer was used and the divers underwent a longer decompression than was intended - and one man had to be uncompressed for an attack of decompression sickness. On 9 occasions, the recorders used the schedules for the next deeper interval i.e. 10 feet deeper again leading to extra decompression. Adjusting the figures by excluding these dives led to an overall incidence of major cases of 4.5% and the difference between the incidence of minor and major cases on Diving Table I and Diving Table II remained similar. Case Motes

Diver MA had two serious episodes in his 4 dives on this trial. On the first occasion he was decompressed for the dive he actually did instead of for a dive 10 minutes longer. This meent that he had 30 minutes less decompression time and his first stop was 10 feet shallower then it should have been. Because the error lay in the schedule. the maximum presence of recompression in the chamber was also 10 feet shallower than it should have been. He reported while he was on his 10 foot stop but his symptoms in fact had appeared at his 20 foot stop. He reported nausea and further questioning disolosed paraesthesia of limbs and forchoad, low backache, and dyspnoce; on attempting to stand his less gave way. Examination disclosed some loss of power of his left tendon and plantar reflexes were present, equal, and normal; and legi he had full voluntary movement. He was recompressed and all signs and symptoms cleared by 50 feet. He was decompressed successfully on Therapentic Table I (Shorter). In view of his history in other diving work with other medical advisers, he was considered unfit to dive in experimental work for 3 months. Shortly after he resumed, he carried out another surface decompression trial dive. This time there were no errors in the drill end he reported he was fit after it. He was due to dive eqain the next day but just before his dive he stated that the previous day, about 1/2 an hour after his dive, he had developed extreme weakness of his body from the weist down which had cleared over a period of 15 hours with the aid of a hot bath and he had not informed

-118-

and person. Examination at the time of his reporting disclosed no abnormality but he was forthwith removed from all diving in the Royal Navy. (Note: 3 years later in another part of the world he persuaded the local naval diving team to let him accompany them on a routine practice dive in a compression chamber with a well tried schedule and a similar lower limb paralysis developed, treated successfully by therapeutic recompression).

Diver PI carried out 3 dives at widely spaced intervals and needed recompression after 2 of them. On the first occesion he developed en ache in his left upper an between 10 and 15 minutes after surfacing. A therapeutic test of a hot bath was tried; his symptoms increased in severity and spread to his right ann and involving both elbows and both shoulders. He also developed an aching sensation in both thighs, the enterior abdominal well, and scalp. No was recompressed 50 minutes after surfacing and all his symptoms cleared at 35 feet. He was treated on Thereneutic decompression Table I (Recommended) returning to atmospheric pressure at 2211. At 0315 he reported evain. In obvious pain which he stated was sited in his upper left leg and which had wakened him 22 hours earlier when it was a mild pain in his left knee and left hip. He was again recompressed but this time his symptoms did not clear till the pressure reached 155 feet. He surfaced after a therapeutic decompression on the Table III routine (Shorter than recommended for a recurrence). with stiffness and some residual bruised sensation in the left thigh which eased with exorcise.

-119-

h months later, without a dive in the intervening period, he again developed trouble after a trials dive. His ascent had been very irregular in rate. On leaving the compression chamber, he complained of a constition of lumps in his scalp, loss of appetito, stiffness and pain in the right olde of his chest causing difficulty in breathing, and disturbed censation of his right leg from buttook to toes. Examination could not confirm any of his completants but he was recompressed and reported all exaptons relieved at 35 feet. In spite of the neurological complaints. he was treated on Therapentic Table I (Shorter). When he left the 10 feet stop he developed a slight nagging pain in his left shoulder. Regrettably his personality had raised strong suspicions that he was - and had been malingering to earn the fee paid per minute under pressure to divers in those days. He was put to bed with placebos initially but such later it was obvious that his case had genuinely recurred and he was then treated with strong analgesics and sedatives. These were used as 1t was considered that in the natural course of the injury, recompression therapy would not be so effective after the delay before reconsideration and that the duration of recompression therapy would probably be longer than effective bed treatment. Eventually 48 hours after he had last reached surface he had recovered although for a short time he was conscious of a sharp pain in his left shoulder tip on coughing. (Eventually for executive reasons he was considered generally unsuitable for diving and he was relieved of his diving qualifications.)

-120-

<u>Myer MH</u> Scatures in these accounts, mainly because of the amount of trouble he developed in each trial. He dived very often in this surface decompression trial his 15 dives were 3 more than any other diver and he had two opisodes treated by recompression. For his first incident 3 hours after surfacing, he complained of an ache in his right shoulder more severe than a dull ache in his right ellow; he had had a dulled feeling immediately after the dive in the same areas. No abnormality was found on examination. He volunteered the information that one week before he had had a pain in his right shoulder after he had acted as attendent to Diver SH (g.v.) during a therapeutic recompression but which he had not hitherto reported. (Since then he had dived every day without trouble). He was recompressed and his symptoms closered when a depth of 150 feet was reached. The use of Therapeutic Table II was completely successful (Recommended).

His other main opisode occurred about 3 weeks later when he reported on reaching surface that he had pain in his left shoulder and upper arm which had been felt for a short time on arrival at 20 feet, and again at 10 feet, and yet again at the surface. Examination revealed tenderness over the shoulder with slight limitation of movements and it transpired that he had had a knock to his shoulder the previous day. The clinical picture was skin to decompression sickness so he was recompressed with relief at 78 feet and he responded to the use of Therapeutic Table II (Recommended). Bruise discoloration appeared on the medial aspect of his left upper arm below the axilla during the

-121-

therapy, but bubbles may have formed in the damaged ticsue so they had to be treated whatever the final diagnosis.

<u>Diver LE</u> carried out a routine dive using the self-contained equipment. He reached surface and complained of abdominal distension and numbross of his right leg. He rapidly deteriorated with numbress developing in his left leg and lack of co-ordination apparent in both legs. The obdominal pain was severe. He was rapidly recompressed without an examination and his condition observed from outside the chamber as the medical officer was temporarily unfit to be compressed. His pain cleared at 10 feet and his neurological signs and symptoms disappeared at 50 feet. He was treated on Therapeutic Table I with complete success (Shorter).

<u>Diver WH</u> also used the self-contained equipment but his initial ascent was fast and uneven - in fast he was told three times to slow down and steady up. He appeared to be well till 2 hours after the end of his dive when he developed - but did not immediately report - a throb under both knee caps which became more a sensation of warmth. Examined 3 hours later there was no abnormality and a hot bath was prescribed. 5 hours later still a sharp pain developed in the right knee and a dull ache in the left knee. He was recompressed with relief at 55 feet and a routine Therapeutic Table I decompression was uneventful (Recommended).

<u>Diver SM</u> finished his dive at 1018 with no fil-effects. It was his first dive after an opicode of decompression sickness 8 days earlier during multiple dive trials (q.v.). About 2145 as he had some discomfort

-122-

in both his knees, he had a hot shower whereupon his sensation changed to dull ache and stiffness. Although physical examination was negative he was recompressed and all his symptoms cleared by 73 feet. Routine therapeutic decompression using Table II from 165 feet was successful (Recommended).

<u>Diver FR</u> was an experienced diver who developed signs of peptie ulceration during these trials. He was allowed to continue diving and carried out 12 dives in this trial. On one dive he developed a nagging pain in his left upper arm while at the 10 foot stop which persisted on reaching surface; in view of its reported mildness and in the absence of physical findings it was decided to await further developments. These occurred about  $t_{\rm H}^4$  hours later when the pain increased and he lost fine sensation of the fingers of his left hand. After relief of signs and symptoms at 65 feet on recompression, he was decompressed on Wherapoutic Table I with no further trouble (Recommended if neurological signe secondary to pain). (He was referred for routine annual chest K-ray on return to port two months later when a shadow of early adult tuberculosis was seen on the X-ray plate).

<u>Diver DR</u> surfaced from a routine dive at 1030 and, shortly afterwards he noticed some discomfort in his left fore-arm which he reported at 1230 - more to obey instructions than for any other reason. He refused any treatment at that time. During the afternoon the pain spread to involve his whole arm and "was becoming a muisance". A hot both produced some waxing and waning in his symptoms and when he was soon again at 1630 loss of gulp of his left hand was found. He was recompressed with restoration of power at 140 feet and he was decompressed on Therapeutic Table II (Recommended if weakness secondary to pain). His ache was cased but didnot clear during treatment and it lested with some stiffness in the erm for about three days after his therapy. Germenta

# The most noticeable point is the frequent lack of physical findings in many cases which must be due to decompression sickness as evidenced by the response to recompression. The spread to involve nerves, presumably by local ordema in cone cases is an indication of delay which leads to greater pressures being required for improvement and consequently longer periods spent in the recompression chambers. Frequently this results in a prolonged post-therapy constition of bruising. In spite of a financial benefit to petiont, and attendents under conditions then prevailing, the chorter treatments used on two ecceptors without illeffects - and without the problems of personality clashes - were popular.

It was felt that the manipulations of the theories had led to a satisfactory routine which was adopted later as a routine technique. There was grave doubt about the adequacy of the decompression for longer dives. This led to the extension of the investigation into the basic tables described in Chapter VI. When the cases were properly assessed, the depth of rolief of the signs and symptoms was thought to be a better guide to the subsequent therapeutic decompression than to use their actual nature as on aid in selecting the routine to be used.

ee 24er

#### CHAPPER X

#### SURMARINE ISCAPE TRAINING

#### Beeleground

This training is not experimental in nature but it does involve a risk of injury due to the rapid changes of pressure undergone by the traince. The signs and symptoms often resemble serious cases of diver's decompression sickness. The patients are treated in a similar manner by recompression. The training is carried out in a warm well-lit cylindrical tank of 20 feet diameter and 100 feet depth (Fig. 13). Access looks are situated at 30 feet, 60 feet and the bottom. In addition there is a diving bell that can travel the depth of the tower. Routing

The trainee usually wore light goggles, swimming trunks and an inflatable lifejacket, with a selief value, to provide buoyency; or else he dressed in the Submarine Escape Immersion Suit (Fig. 14). Each class of 9 men and an instructor was compressed in the appropriate look by flooding up with water and when the pressure equalised, the communicating door into the tank opened. In turn, each trainee took a breath from the appropriate air supply and was pulled backwards through the door by two instructors who held him until a senior instructor was satisfied that he was exhaling steadily. The trainee was then released; his natural buoyancy plus the buoyancy of the lifejacket took him rapidly to the surface under constant surveillance by instructors attationed in the water and at the surface. On arrival at the surface he left the vater unassisted and he was kept standing near by under observation for a short period. Any abnormality of behaviour, or physical condition could be immediately noticed. Unless there was an immediate alternative explanation, the trainee was assumed to be suffering from some form of pulmonary barotrauma and he was recompressed rapidly - 30 seconds from suspicion of injury to start of compression being considered slow. While there have been no fatalities in training in the Royal Navy, they have occurred in other circumstances (98) and then most commonly in less then 5 minutes from reaching surface. Each trainee, after a practice dive to 100 feet in a compression chamber, carried out two ascents from 30 feet, one from 60 feet and one from the bottom of the tank in the course of 2 days.

The incidence of cases that needed recompression, including those in which an alternative explanation was later found, was between 1 in 0,000 and 1 in 10,000, escents by trainees (the escents by instructors were done without lifejackets and there were only 2 incidents in 8 years). The signs and symptoms depended on the site reached by the bubbles of air which enter the tissues from the alveoli.  $\int Complete$  responsibility for treatment was limited to a few cases ab initio but the remainder of the cases described became my responsibility when I was called in as adviser on the subsequent decompression.

## Therementic Approach

The therapy used was recompression on the assumption that the ges pookets or bubbles must be reduced in volume before ischeemic or pressure

-126-

damage became irreversible. Such treatment should lead firstly to the bubbles in the circulation passing on to be filtered out in a relatively hamless area or in the lungs or to reduce the area of effective damage; or to gas in the pockets or bubbles going into solution in the tissues; or finally, in certain circumstances, to allowing time for other therapeutic measures to be taken. It wes noteworthy how frequently the presumed initial breach of tissue continuity in the lung appeared to be immediately scaled as evidenced by the disappearance and failure to recur of signs and symptoms. The patient had then to be returned to atmospheric pressure and it was customary . but not invariable . to use the therapeutic tables that wore used by divers for decompression sickness. In the light of experience in divers, it was felt that the situation of the escape training casualty was different. In decompression sickness the hypothesis was that bubbles tended to form de nove in tissues supersaturated with gas. In pulmonary barotrauma, especially the air embolism forms, the bubbles were thought to be introduced into tissues where the atmospheric gases in solution were still essentially at normal etmospheric tension as there had been insufficient time at high pressures to alter the levels to any great degree. This led to the attitude that the therapeutic recompression of a submarine training accident could be regarded as a double chamber dive carried out by the patient at rest. A routine was evolved on these lines for cases which had apparently responded completely to the initial compression. It became apparent

-127-

that 1f the bubble had not completely redissolved or been filtered out so that signs and/or symptoms persisted then a prolonged stay under pressure was required till the effects of the bubble disappeared as pressure could not be further increased, because of equipment limitations. The suggested decompression routine (Appendix 3) was based on standard air Diving Table II (132) and this sometimes caused confusion with Therapeutic Table II (132) (new editions of the Diving Manual (133) now refer to these tables as Table I below the limiting line and Table VB respectively). Important subsidiary factors in investigating the value of a shorter routine were an ambient temperature of 90°F, high humidity, and close confinement in the small compression chamber.

#### Case Notes

Induce CK carried out an ascent from 100 feet while on requalifying course. About 30 feet he thought he was running out of air because of tightness in his chest as if it was about to burst. While climbing out of the tank he appeared to have some difficulty and then his less gave way. He was immediately rushed to the chamber (about 15 feet away) and though his arms seemed to become flaceid, he did not appear worried and his speech was clear and rational. He was compressed to 100 feet but as he was not completely recovered, pressurisation continued till he reached 165 feet. Physical examination at that depth disclosed no abnormality. He had remained conscious with unimpaired speech, sight and hearing and he had deliberately not tried to move till

+128+

he was asked to do so during the compression. Decompressed on standard air diving tables as if he had dived to 165 feet for 35 minutes i.e. maximum depth for total time from initial pressure exposure that day and he regained the surface 2 hours from the start of treatment. Exequent examinations during this period were negative and subsequent chest X-rays were normal. The clinical picture was that of cerebrospinal air embolism but in the absence of positive findings the site and diagnosis are not confirmed (Shorter).

Trainee 05 courled out a 30 feet ascent on this his third mutine course (second regualifying course). After climbing out of the water. an instructor questioned him because of suspicion that he was puffing and blowing excessively and that he was not moving vory easily. The petient denied any difficulties; on questioning a little later he edultted feeling giddy; a check on his vision by asking him to follow the exeminer's fingers was normal but an attempt to make him walk disclosed on inco-ordinated spartic gait. As he was being lifted to the chamber, he developed a flexor spass first in his right arm and then in his left arm which caused some difficulty in passing him through the chember hatch. He was compressed to 165 feet and examination was negative: except for emnesia from the time of request to follow the finger - that is he did not remember seeing the finger or responding till the chember had nearly reached maximum pressure. He was decompressed on Therapeutic Table III (Recommended). While at the 50 feet stage, he complained of retrosternal disconfort, suspected to be due to mediestinal emphysema;

-129-

it was decided to continue the decompression and at the 40 feet stage, sungical emphysicms was detected in the right supraclavicular triangle. A careful watch was kept on the patient's condition as decompression continued but as his disconfort was unaltered and the only positive finding was a slight increase in the emphysicms in the neck, no other action was needed. On examination after transfer to hospital 4 hours after reaching surface, E.E.G. showed no focal disturbance but chest X-ray confirmed the mediastinal emphysicms. On repeat X-ray one week later no abnormality was seen. The diagnosis was pulmonary barotreana presenting as corebral air embolism and complicated by interstitial emphysicms.

<u>Indiaco (M</u> This man on his first course, appeared to be apprehensive as shown by excessive frequency in carrying out the Valselva manocuvue during his practice charber dive. We carried out his water drill satisfactorily. On elimbing out of the tank after his 60 feet ascent, he seemed to waver and to have difficulty in finding the rang of the ladder. When he did not reply when he was asked if he was alright, he was muched to the chamber and he reached 112 feet within 90 seconds of reaching surface. This unusual depth was not deliberately selected but happened to be the level where the attending doctor stopped compression having decided that the incident was not due to barotrauma but possibly due to hysteria and abdominal distension as a result of excessive air swallowing during car clearing. The patient reported that he had had a sudden pain in his epigzstrium on the ladder and that

-130+

he had not lost consciousness but the attendant reported that he had to easist the patient to clear his ears during the therepeutic compression by closing the patient's nostalls. Decompression on standard air tables for a dive to 120 feet for 40 minutes was carried out (Shorter). He was examined on surfacing, by the author, who could find no abnormality and who recommended observation for 4 hours in the training tank building (standard practice in case of recurrence). Halfen-hour later he was seen by the original medical officer because of developing restlessness and lack of co-operation. Physical examination was negative and again a presumptive diagnosis of hysteria was made. 1 hour later still he was seen by a third medical officer who diamosed acute confusional state, and after recalling the author who confirmed the disturbed montal state; the patient was recompressed to 165 feet. As a result of the long period to the start of adequate therapy there was no dramatic improvement but experience had shown that Therapoutic Table III usually resulted in necovery in this type of case so it was used instead of the longer Therapeutic Table IV. At 50 feet his conversation became rational and when he was re-examined at 10 feet by the author, the only abnormality found was some interstitial emphysema in the right enterior triangle of the neck. Chest X-ray in hospital showed this emphysema but no mediestimel ges shedow was detected and the E.E.C. was a normal record. This man suffered from pulmonary barotrama presenting as cerebral air embolism and later interstitial emphysema but an error in diagnosis by an inexperienced doctor led to

-131-

difficulties in subsequent therapy. When seen after the first compression, there did not appear to be any indications for recompression; presumably this was in the latent period of a partially treated lesion before reactions such as local ordema develop; the wisdom of a period of close observation is apparent.

Instructor HG This man. surfaced from a 100 foot unassisted ascent efter the last man of his class. He complained after about 10 minutes of pain in his left shoulder redisting to his elbow and wrist and which then developed into paralysis of his left and and forearm with paraesthesia of the two lateral fingers of his left hand. He was recommessed and his signs and symptoms cleared at 110 feet. Ac examination showed no abnormality he was then further compressed to 165 feet before decompression on standard air tables for a dive to that depth for 50 minutes (Shorter). Progress was uneventful. It later transpired that this man had been treated while on leave one month before for bronchitis (with a runour of pneumonia). Full plate X-ray, taken with knowledge of his task but ignorance of his illness 5 days before this incident, was reported normal. Hospital investigations including chest X-reys and E.E.G. disclosed no abnormality. The most probable diagnosis was cerebral air embolism; decompression sickness could not be excluded, as it has been known to occur exceptionally after a dive to 100 feet for 10 minutes as this man had done. (Diving tables give a minimum duration of 20 minutes at 100 feet before decompression is required).

-132-

<u>Trainee MM</u> This was surfaced from 60 feet during his initial training and lay on the surface of the vator with his oyus closed. While his deed weight was being pushed into the chamber, he began to protect but he was compressed to 165 feet. General examination showed no abnormality and he seemed fully aware of his surroundings and events. He was decompressed on standard air diving tables for a dive to 165 feet for 35 minutes with no apparent ill-offects (Shorker). Re-examination disclosed a small posterior traumatic perforation of the left car drum. No chest 2-ray was done in hospital; his E.E.G. was normal. The diagnosis is obscure, but observers were convinced he was unconscious and hence cerebral air embolism was a possibility and treatment had to be given.

<u>Traince ND</u> loft the vator after his first ascent from 30 feet apparently well but within a minute complained of disziness and inability to use his right hand. (Later it transpired that he also noticed numbress round his mouth, numbress on the medial aspect of his right abov, and in the approximate distribution of the ulnar nerve in the right hand). He was immediately recompressed to 165 feet. As on arrival at this pressure some numbress was present round his mouth and over the medial three fingers of his right hand and there was inequality of lower limb reflexes with right brisker than left, he was decompressed on Therapeutic Table III (Recommended). During this period full sensation returned and his reflexes became equal. Chect X-rays and E.E.C. on admission to hospital disclosed no abnormalities but the clinical picture was air emboliem.

Trainco MI This can had ettended the course twice before. On the first occusion he had carried out the full course but shortly after surfacing from 100 feet, he had been found apparently unaware of his surroundings; he had not been recompressed and observation in hospital had not disclosed any abnormality after he had quickly regained his conse of his surrounding. On the second occasion, he did not carry out his 100 foot accent because he developed a couch on the morning of the second day. On this third course he surfaced from 60 feet and complained of a twitching in his hunstrings on climbing out of the tank although his valking seemed normal. While he was being examined he developed paraesthesia of his knees and then loss of active movement of his knees and recompression was started. These complaints steadily improved during compression but he then found difficulty in raising his arm to clear his cars. By the time 165 feet was reached, all signs and symptoms had disappeared. He was kept thore for 11 minutes and decompressed on standard air diving tables for 1 hour at that depth - i.e. a total decompression time of 22 hours - and no signs or symptoms recurred over 24 hours observation (Shorter). This can showed neurological abnormalities possibly due to air ombolism.

-134-

Traince EU was carrying out his second escape course. On the first occasion, he had developed numbress of his 4th and 5th fingers left hand and he had been treated by recompression; initially it seems his decompression had lasted 1 hour (no details of depth known) but when he
had then developed weakness in his legs. Therapeutic Table III had been used. This time, again after his 100 foot run, he developed numbress in his right little finger and he was compressed till this cleared. This depth was 420 feet where he was kept for 11 minutes and then decompression was carried out over 2 hours (Shorter). While this man's complaints were minimal on both cocasions, the initial failure of a very short routine on the first occasion and the success of a slightly longer procedure (but still chorter than the routine therapeutic tables) on the second are of interest though the diagnosis of air embolism is only presumed.

Trainee 12 was on his fifth encape course and had just completed the last accent from 100 feet, when he began to feel unwell as he climbed out of the tank. He then fell to the floor, overcome by weakness and dissiness but he later multiplied he did not lose consciousness. He was immediately recompressed with such marked improvement to his general condition that numbress and weakness of his right arm became noticeable. At 165 feet, the refignes of his right arm became noticeable. At 165 feet, the refignes of his right arm vere increased and there was loss of fine finger movement so a therapeutic table was indicated and Fable II was the one selected (Shorter). During the 10 hours decompression, the signs improved although on reaching surface the right bloops reflex was still brisker than the left. Observation in hespitel showed continued progress in a petient who had fairly classical clinical cerebral air embolism.

-135-

<u>Grainee 103</u> carried out a most unsatisfectory accent from 60 feet with no apparent exhalation for the first 25 feet from release. (No had exhaled satisfactory before he was released so his chest was not fully expanded at the start of this period). 2 minutes after surfacing he compleaned of heaviness in his right arm and he suddonly lost consciousness as he was entering the compression chamber. During recompression, he steried to recover, initially being confused and disorientated but within one minute at 165 feet he became rational and co-operative. No ebnormality was found on examination and he complained only of very transient pine and needles over his left antecubital fossa in addition to a headache over the vertex (where it was thought his head had hit the chamber wall during entry). He was decompressed on standard air diving tables, the whole procedure taking 3 hours (Shorter). Observation and investigations in hospital disclosed no abnormality. This case clinically resombled those with corebral air embolism and responded to a short routing, in spite of the persistence of some signs on arrival at maximum pressure, usually considered an indication for one of the standard therapeutic decompression routines.

Trained Mi on his first course seemed to be unable to leave the water after his 100 foot escent. He was promptly transferred to the compression chamber and compressed to 165 feet where he was found to be dazed and frightened. He had some bleeding from his nose but was otherwise well. He was then rapidly decompressed to 100 feet and examined again. Standard air decompression was then started but at 50 feet, he

-136-

compleined of right inframemary pain so pressure was increased to 60 feet and in a few minutes the first signs of a right pneumothorax were found. Pressure was maintained and interstitiel emphyseme tracked behind the right clevicle after about 20 minutes. The patient was extremely envious and complained of difficulty in vision and insbility to breather the former appeared to be due to a wendering scotome and the latter was not confirmed by examination. After various pressure fluctuations, he was given chloropomazine and his visual defect, inbuler in oheractor by this stage, gradually cleared. The decompression thereafter was a modified Therapeutic Table I (Shorter). Observation and investigations in hespital confirmed the pneumotherar and he was found to have an unusual H.B.G. depression reaction to chlorenomezine. The initial rapid visit to 165 feet and back to 100 feet was unusual and in the author's view led to subsequent complications in the treatment of undoubted pulmonary barotrauma presenting as cerebral air embolism. pnounothozez, and intenstitial emphyseme.

<u>Insince M</u> completed his second course but 6 minutes after his final ascent - from 100 foet - he was seen to be clumsy and inefficient with his suit fastenings and as his replies to questions were confused he was recompressed to 100 feet before any medical officer saw him. As he was normal, the medical officer considered that there was no evidence for the need for a therapy and the patient was decompressed, the whole cycle taking 25 minutes. Over the next two hours he had slight diszinces and headache for which he obtained some aspirin before leaving the

-137-

establishment with a friend. On a ferry he had an epileptiform attack and his friend parsuaded the arev to take the patient back and 20 minutes after boarding the ferry FB was back in a compression chember on his way to 165 feet. He was seen by a second medical officer who found him drowsy end compleining of a frontal headache but co-operative. No other abnormal signs were found. After 5 minutes, decompression to 100 feet was cannied out and a more intensive interrogation displosed emnesia from the time of his fit till he was in the chamber. Therapeutic Table II was started from this depth but when 60 feet was reached 50 minutes after the entry into the chamber, FB had another major enilentiform convulsion in which he bit his tongue slightly but during which he remained continent. Pressure was immediately increased to 165 feet but for helf-an-hour the patient's personality was inaccessible; he was restless, with ell his limbs flootid, his reflexes exaggerated, and his toes upgoing. He then responded to questions and Thorspeutic Table III decompression was connenced (Shorter). Re-examination at 120 feet showed recovery of all abnormal signs except that he had a further amesic period for the second fit and about half-am-hour afterwards. Intramyscular parallelyde was used to sedate the patlent and Table III was modified for the final 3 stops which took 3 hours. 5 hours and 8 hours successfully instead of the schedules 12 hours. 2 hours and 2 hours. In hospital, clinical examination and chest X-ray showed no abnormal signs but the B.E.G. was arossly abnormed especially in the posterior right hemisphere. Follow up E.E.G. one month later showed no abnormalities.

-130+

This case undoubtedly had cevebral air embolism and demonstrated how inadequate initial care complicated the subsequent therapy.

Traince AA had not carried out an entirely successful escent from 100 feet. on this his first course, when his red face and very firm grip on the ladder led to his being immediately recompressed to 165 feet. On exemination there he appeared fit and well so decommonsion commenced on a standard air diving table. At 100 feet interstitial emphysema was found behind the might clavicle and signs of a right pnoumothorar noted. Using his condition as a guide, decompression continued, taking 12 hours (Showter). On reaching surface, the superficial emphyseme was present over both sides of the face and neck, the right shoulder, and the right pectoral area. X-rays disclosed a right memothorar confined to the bese by adhesions, and added retroportioneal emphysema to the known area involved. Nowover 14 days convelescence led to the diseppearance of all abnormal signs and symptoms. This case of pulmonary berotrauma, as shown by interstitial emphysema and pheumothores (and perhaps air embolism if his initial grip was spectio in nature) demonstrated that a short routine could be used provided mediartinal emphysems and proumothomax were either contined strictly or else released to ereas so that no candigo or respiratory embarrasement occurred.

<u>Trained MD</u> had completed the course successfully once before. On this occasion he had a mild trachetis which he did not report and which was not detected on excaination. He cannied on the course successfully and had left the tank top area to get dressed after his 100 foot escent

-139-

when he noticed increasing numbers and lack of control initially over his logs and then his axes. He felt confused and disay and appeared disordentated although he responded to questions and was co-operative. He was recompressed to 165 feet 17 minutes after leaving the water and examination at pressure disclosed no abnormality; he was decompressed to 100 feet and re-examined and bilateral aural benotrauma was found. Questioning revealed partial amesia for the short interval between the development of dissiness and recompression. Decompression was then carried out on standard air diving tables for a dive to 165 feet of 27 minutes (Shorter). Further investigations were negative throughout but the initial picture was that of cerebral air embolism.

<u>Traince MF</u> opparently surfaced feeling well after the 100 ross ascent on his second course. He did not report for several minutes but he had noticed numbress, loss of control, and weekness in his right thigh just after he climbed out of the water. These symptoms radiated down the log and up into the trunk and he reported when he needed help to stand up. While he was being helped to the compression chamber, the numbress spread to his right shoulder and ann. Seven minutes after leaving the water he was compressed to 165 feet and other than annal banotranna bilaterally, no abnormality was found on examination. Decompression was commenced on standard air diving tables but on reaching 70 feet, the symptoms recurred now involving the right side of the face as well. The patient was recompressed to 165 feet with relief of his signs and symptoms. Therapeutic Table III decompression was started and completed

-140-

successfully (Recommended). Further investigations were negative. Clinically this was a case of cerebral air emboldsm.

Trainee WD felt a little confused on reaching the surface after a 100 foot ascent on his first course. He started to climb out. looking normal, but he suddenly crumpled to the floor whereupon he was immediately put in the compression chamber and within two minutes of surfacing, he had reached the equivalent of 165 feet. At this depth the patient remained quiet and seemed asleep but no efforts were successful in rousing him for about 10 minutes when he suddenly recovered. There was bilateral otitic barotrauma and a left opistexis but otherwise examination revealed no abnormality. Decompression on Therapeutic Table III was started (Recommended). On reaching 100 feet, after 8 minutes, the patient complained of difficulty in vision and bilateral central contomata were found. He was recompressed to 165 feet where the visual defect decreased and there was found pupil inequality, with the left larger than the light and accommodation convergence reflexes were sluggish; and richta there was transient nystagous to right and left. In addition he had developed frontal headache with nausea and vomiting of altered blood. Decompression on Therapeutic Table IV (Recommended) was commenced and there was steady improvement with time as judged by the print he was able to read with each eye centrally. Some slight pupil inquality percipted in hospital and a very transient nystagmus could be elicited. but other examinations and investigations disclosed no abnormality. This case was clinically one of corebral air embolism. Perhaps on

initial higher pressure would have been beneficial but the chamber was not designed for such pressures.

Trainco PP carried out his first escape patisfectorily until the 60 foot accent when he made a groaning expiration on arrival at the surface. He started to climb out but collansed back into the water apparently unconscious. Even on compression to 165 feet, he remained unconscious and showed laboured respiration with mild evenosis of his face and neck. Over 10 minutes, the patient passed through a restless condition to one of co-operation and rationality in which he complained of retrosternal pain. Dramination disclosed signs of weakness in his left ann, poor co-ordination of all his limbs and possible diminution of the area of cardiec duliness as well as bilateral stitle barotrauma and mild opistaxis common in rapid therepoutle compressions. Becompression on Thorapeutic Table III was started after his chest pain had almost cleared over a period of helf-an-hour (Shorter). At 80 feet over a period of 12 minutes the patient rapidly and progressively became disorientated and unco-operative and developed jargon aphasis and misunderstanding of simple commands. Recompression to 165 feet led to very slight improvement in his response to forceful commands but his moods rapidly alternated between a childish supports and aggressive restlepsness. (He was a big athletic individual) His tendon reflexes became very brisk. After 2 hours, decompression on Therapeutic Table IV was started and intranuscular paraldehyde was used to control the restleseness. 12 hours later, at 40 feet, he began to respond more to simple commands

-142-

and questions and to appreciate his surroundings and at 20 feet his mild suphoris and final traces of sphasis disappeared. His brisk reflexes were the only abnormalities on clinical examination before transfer to heapital; E.E.G. showed an abnormal record with slow theta and delta activity in the right hemisphere; chest X-ray disclosed a small left spical pneumothorax, a localized air cyst at the left base, and a well marked pneumo-pericardium. 3 days later, a chest X-ray showed decreased pneumothorax and pneumopericardium only and in another 5 days, these findings had gone. A repeat of the E.E.G. after an interval of 1 week showed a normal record. This patient had pulmonary barotrauma, as shown by the triad of pneumothorax, pneumopericardium and corebral air embolism, which was too serious to consider the use of a short therapy.

<u>Trainee NJ</u> was on his first escape course and his first accent from 30 feet was satisfactory after considerable initial difficulty in exhaling to the patisfaction of the instructor (he re-entered the air lock three times before he was released and each time he was keen to try again). His second accent, also from 30 feet started well but his exhalation became intermittant in the shallower depths. On climbing out of the tank he collapsed with loss of consciousness which did not apparently respond during compression till he reached 165 feet. Examination disclosed no abnormalities so decompression on standard air diving tables started (Shorter). This had to be halted at 56 feet because of a completint of inability to see. He was immediately

-11,3-

recompressed to 165 feet where he also mentioned the new developments of retrosternal pain and acho in his neek. On examination. there was no ovidence of moningism and all movements of his neck were full and painless. No abnormality was found on examination of his chest or fundi. There was no response to light of his dilated pupils and he had a blink verlex but blindness seemed to be complete. His co-operation decreased as he became drowsy and eventually his respirations became stortorous and rapid and he seemed to be in coma. Trial decompression to 100 feet was carried out and his come was replaced by semi-conscious activity which was violent if he was disturbed. An attempt to recompress him again to give him a prolonged pressure exposure failed because of obvious pain in his facial sinuses, at a depth of 122 feet. Therapeutlo decompression on Table IV was started with intranuscular paraldehyde as a sedative, which however was not completely effective till 16 fl.oz. of urine were dreined from his bladder by eatheter. He slept for 7 hours and examination of him then showed lack of avereness of his surroundings and total lack of response. Physical examination was negative but he became restlene until he was incontinent of urine and facces whereupon he settled back to sleep. For the next 30 hours, physical examination showed no change and he had alternating short outbursts of violence and long periods of sleep. At 10 feet some subcutaneous emphysems was detected in the right side of his neck. The picture of cerebral irritation and interstitial omphysems was confirmed in hospital where X-rays disclosed pneumopericardium and eir tracking into

the upper mediastium and the neck and E.E.G. showed a generally disorganised vecord with excess slow activity in the left hemisphere. He made allow steady progress and physical and mental recovery on clinical examination was complete in three weeks though another three weeks was needed for the E.E.G. record to become normal. This case of cerebral air embolism and mediastinal emphysical failed to respond to a short therapy and was not entirely successfully treated by a very long routine; this poor response might have been due to inadequate initial treatment allowing reactionary damage to occur.

Induce: 10 had reached the 100 feet escent stage of his first escape course. His own story was one of hitting his head as he left the look - not confirmed by the two attendant instructors - and that his next memory is reaching the surface, vague recollection of people talking to him, and then clearing his cars during compression to 165 feet. The attendants reported poor exhibition during the ascent and failure to respond to instructions on arrival at the surface so he was compressed. His attendant during the compression reported that he seemed fully orientated and rational. Physical examination was negative, even to signs of head injury, and after 30 minutes at maximum depth the patient was decompressed on standard air diving tables uneventfully (Shorter). In hospital, E.E.G. examination showed epicodic theta activity in all areas and a slow clearing delta response produced by overbreathing. A repeat examination in 4 days showed diminution of these findings and three weeks later the E.E.G. trace was within normal limits. Under the

w14.5-

circumstances, it is wiser to regard this case as one of corebral air embolism which responded to a short routino.

## Discussion

Many of those mineteen cases showed signs and symptome suggestive of neurological damage. In view of the short period of time possibly available before death, the patients were recompressed as soon as possible after suspicion had been raised, without waiting for the medical officer to make a diagnosis. Over the years the medical officer in the building changed. While the medical officer for the first few of these cases was called in to advise in the remainder, each new doctor would exercise his perogetive to treat his patient initially to the best of his belief pending arrival of assistance, sometimes the case was not seen by a more experienced doctor till the routine was well underway. This led to errors of diagnosis e.g. traince FB or errors of assossment e.g. traince NJ.

It is possible that only 8 of these men suffered from pulmonary barotrauma; none of the other men had abnormal findings on X-rays or electroencephelographic examination but with the proceeding activity and similar clinical ploture, it was the most probable diagnosis. When no physical signs were present the diagnosis could only be made on the history. The case of trainee FB and to some extent that of trainee CH show that any alternative diagnosis may lead to considerable complications. During the 5 year period covered by these cases, only two other cases were compressed; one was decompressed on the standard air diving table

-146-

with a diagnosic of disorientation under water and the other complained of pain in his foot - on being further questioned at 165 feet he added that he had accidentally kicked the tank well wile climbing out efter his escent. While there may be doubte about the diagnosis, any benefit must be given to the patient and inconvenience and discomfort to the staff are not important.

13 cases were considered suitable for trial decompression on a short routine taking a few hours. Of these 4 eventually had to be further treated, two by routines taking 20 hours, a third by a routine taking 40 hours, and the fourth by a modification of a 6 hour routine. 4 men were treated, from the beginning by Therapeutic Table III and of these, 2 had to have further treatment by Therapeutic Table IV. One man was treated on Therapeutic Table II which should not normally have been considered for this but it was effective. The remaining case sprang from an error in diagnosis and had a complicated compression history.

From this it can be seen that the routine treatment of neurological cases by a long therapeutic decompression is not always the best treatment. If proper selection of cases is made then a trial at a short routine may be justified provided the physician is prepared to return to higher pressures and a much longer routine when the condition of the patient warrants it. Only those cases whose signs and symptoms have completely cleared by the time maximum pressure is reached should be considered for a short therapy. Experience suggests that cases which still show some troubles, for however short a period, on the conclusion of

-147-

compression will later require to be treated with further compression or transferred to a long routine. In addition, there is evidence here for always using pressures of 6 atmospheres absolute right at the beginning and which should be maintained for some time, probably 30 minutes, before decompression commences. The possible use of higher pressures may be considered, if equipment limitations are overcome.

-148-

The discovery of phoumotherax or interstitiol emphysema does not necessarily lead to further compression as the former can be treated by orthodox means 12 needed - and has been on one occasion in the past and the latter usually tracks into the soft tissues of the neck so releasing the tension and the dangers of both are avoided. The dangers of pneumoperioardium may not be avoided so capily and in such a situation the compression and decompression would have to be controlled by the degree of cardiac embarrasement produced, with a very slow release of pressure over days being theoretically one solution. These conditions are basically mechanical in their action in producing signs and symptoms. Air embolism has however the secondary offect of producing local hyporia and the later appearance of signs and symptoms may be due either to the recrudescence of the embolus or to collular hypoxic effects or to reaction by the tipeues to the traumatic insult. In all cases, an increase in pressure should lead to improvement by reducing the size of embolup or increasing the amount of oxygen discolved in the tissues. The risk of the effects of hyporia are less if adequate pressure is applied repidly enough.

Any therapy for pulmonary barotnama should involve inmediate recompression to high pressures. The chembers were designed for an absolute working limit of 7 ats. abs. as Duffner and his colleagues (51) had shown that 6 ets. abs. was quite adequate and that a further relative alight increase in pressure was not worthwhile (although later work here shows that in fact 8 or 9 ats.ebs. may be valueble). Thus 165 feet as quickly as possible without too much regard for the tympenic membranes and sinuses of the patient is indicated. Thig pressure is 2 stmospheres more than the maximum to which the traince is usually exposed on his course. To encourage gas in any bubbles or pockets to go into solution, not only is a pressure gradient between bubble and tissue gas tensions required but also a certain time element and this period has been fixed arbitrarily at 30 minutes. The diffusion of gas in and out of the body tissues has varied with the pressure exposures of the individual during the proceeding training and account has to be taken of this in devising suitable decompression tables to evoid the production of decompression sickness. To make a dogmatic instruction, it has been accepted that the sum of time spent under pressure that day, the time from the last escent to compression, and the period spent at 165 feet - usually 30 minutes - will be regarded as the total time spent at that depth and hence the appropriate diving table can be found and used with some confidence in the right cases.

Examination of mineteen cases of pulmonary barotrauma lends support to the treatment of cases whose signs and symptoms clear during

-149-

compression to 6 ats.abs. by the use of a standard diving table for an artificial aggregate time. Other cases should be treated by the standard therapeutic tables strictly applied. Under both circumstances a few cases will not respond completely and subsequent therapy has to be directed on an ad hoc basis by experienced doctors.

## -151-

### GHAPTER XI

#### DEEP DIVING EXPERIMENTS

## Baokaround

Deep diving is regarded in the Navy as diving to depths in excess of 200 feet. This figure is chosen because it is a convenient whole number near the depth beyond which only very experienced individuals can dive efficiently using air as their breathing medium, although dives to 400 feet or so, using air, have been recorded. The main reason is the loss of efficiency and manual desterity due to depression of the higher intellectual centres that occurs with nitrogen either by direct action or indirectly as a result of carbon dioxide retention as discussed earlier. These problems can be avoided by using helium to provide the bulk of the respirable gas as theoretically, similar effects should not be expected till depths of the order of 1.000 feet or more are attained. Air at 300 feet or 10 atmospheres absolute, contains toxic quantities of oxygen as well as nitrogen. One solution is the provision of an artificial atmosphere where the proportion of oxygen ic reduced the diluont being helium.

### Notes on Techniques

a) Chambor divos

In chamber experiments, the subjects were usually compressed in pairs and they either donned the breathing equipment - usually a mouthplece mounted demand value - supplied from various banks of artificial atmospheres stored outside the chamber; or they continued to breathe the chamber atmosphere which was supplied directly from the banks (Fig.4). Once compression was completed the divers exercised alternately, on a rowing machine or other device, till just before the end of the dive when they dressed in warm clothing (Fig. 15). Decompression commenced and the breathing mixture was changed to avoid severe hypexia on reaching surface. Varying mixtures of oxygen and holium were used so that the proportion of helium was kept as low as possible consistent with safety in respect of the oxygen content; this was done to keep the pressure gradient as steep as possible between the tension of holium in the tissues and the partial pressure of holium in the respired gas to encourage elimination of the gas and hence reduce the risk of decompression sickness. During the decompression the men relaxed as much as possible to avoid stimuli to bubble formation.

b) Sea dives

In sea dives, the men used the submersible decompression chamber (S.D.C.) as a lift and donned their breathing equipment at 100 feet during the descent; if the water level in the chamber was being kept down by supplying the correct oxy-helium mixture for the dive they continued to breathe the chamber atmosphere (this supply was intended originally in case of equipment failure, but was later done for comfort and economy). After a spell of swimming or working (Fig. 16) the divers were recalled into the chamber and the ascent commenced. The selection of the various breathing mixtures in the course of a sea dive was

-152-

sometimes affected by the ease of using different mixtures for the diving set supplied from the cylinders mounted on the S.D.C.; the submersible chamber atmosphere supplied from storage cylinders in the ship; and the transfer chamber supplied from other storage banks on board the ship. The decompression continued in a similar manner to that described for chamber dives once the divers had transferred satisfactorily. Approach to the Trials

General principles, were that the number of dives to be done on each safe schedule was to be 10 and that minor cases of decompression sickness would signify caution but a major case would normally entail change either in total decompression time, oxygen percentage in the mixture, reduction in dive time, or complete recalculation depending on the course of action previously agreed. As the trials progressed they became more experimental in nature then potential operational routines. This resulted in changes in such respects as numbers of dives; number of men to do them; and in the action to be taken if serious cases developed. Such changes can best be followed by describing them in the various stages as they occurred.

### THE DIVES

a) First Period (103)

(1) Preparation

In this initial phase of optimism it was anticipated that the team would be diving within 2 months to 1,000 feet with shorter decompression times than ever previously envisaged. Several chamber dives

+153-

on the new concepts were carried out on air to 300 feet to test both schedules and rates of accent. This decision arose from theoretical and practical work abroad which suggested that oxygen and holium sixtures were safer then oxygen and nitrogen. Similar calculations were used to produce a schedule for a dive to 250 feet on air in the sea which would serve as a final work-up dive as well as acquainting the divers with the effects of narcosis in case air had to be breathed in an emergency during deeper dives. <u>Table XI. 1</u> Record of Dives. First Experimental Period

.

	Reclaim	Second 300	250	Vernon 300	300	250	Reclaim 400	First 300	UU <sup>C</sup>		Stage Depth Pt.	
d Total	Chamber Chamber	Chamber	Sea	Chamber	Sea	Sea	Chember	Chamber	Unamper	2	<u>Environment</u>	Dive
		0 <sub>2</sub> /He	Air	0 <sub>2</sub> /He	02∕/∄⊜	Ыr	$0_2/He$	0 <sub>2</sub> /He	A17		Gas	
	ം നാറ	° 7	5	G	3	J	÷	لا	N3 ==	5	Timper a	2) ) ) ) )
139	# # J	6	25	10	6	12	4	N	ತೆ	5	Dives	Numbe
27	∞ರೆರ	2 2	14	10	σ	12	4	N	18 18	5	Divers	ж Ог
<del>1</del> 72		<u>ه</u> ۸	<b>7</b>	2	I	1	<b>I</b> :	8	1 ~	3	láinor	Decompressi
20		4 UJ	N	N	۶,	*	N	ŧ	` <b>∳</b> ⊷	•	liajor	on Sickness

(ii) Results

-155-

The main burden of diving fell on 12 men who carried out 110 dives and who suffered 21 minor and 12 major cases between them. Of this group 3 men had one major attack each but no minor attacks while enother 3 yea had no major attacks but 8 minor incidents between them. Of the support group of 15 men 11 wore not permanent members of the. experimental team or the ship's team and were available only for a very limited period; 3 who were permanent members had had a serious attack of decompression sickness early in the trial and had not recovered fitness to dive before diving stopped in this period; and the last individual was not a fully trained diver and, as he had other responsibilities in the organisation, he was not permitted to dive often. One man who had the major attack on the first air chamber dive. (his first triels dive) in the first Reclaim stage, had enother attack on his first dive in the second stere, also on air but this time in the sea. The other diver who had a major attack after an air dive in this second stage was also carrying out his very first dive of the trials. The cause of these attacks was debatable but the schedules involved an unusually fast rate of ascent to the first atop which was slowed in subsequent periods. Table XI. 1 shows the serious effects of these dives. In the later dives of the series 100% oxygen was used instead of air during the shallower stages of decompression.

-156-

# b) <u>Second Period</u> (79. 104)

## (i) Preparation

It can be seen that the first experimental period resulted in a lack of success as far as the ability to provide safe diving tables for 300 feet, much less 1,000 feet, was concerned. Much information was obtained and further work was started to prepare schedules for a second period of chamber and sea dives. It was intended that all dives would finish with a spell of oxygen decompression and that no breathing mixture containing nitrogen would be used after the start of oxy-helium breathing till the diver reached surface again. The intention was to carry out dives at increasing depths to 450 feet of 10 and 20 minutes duration at each selected depth, using 10% oxygen and 90% helium as the main dive gas. Table XI. 2 Record of Dives; Second Experimental Period

		Beclaim .		Vernon	Stage			
Second Per	4,00 4,00 4,50 300 300	450 400	250 300	4,00 300 4,50 300	300		Depth Ft.	
	700060 7000	10 20	10 10	338 <u>33</u>	20	10	Duration Dins.	ų
lod Total	See	Chamber	29 <b>2</b>	Chambe <b>r</b>		Chamber	Environ-	ļve
	<u>ଟ</u>	Īie	4ir He	He Air He		He	Nein Gas	
	25 25 25 25	20 21	12 19	お あ コ あ み	13 14	12 13 0	Number	20%odulo
179	N& ด <b>อี่</b> N N	10 5	4t 4	๑๑๑๖อื	66	20¢	Dives	Ilumbo
21	NF 9 9 9 9 9	10 5	11 11	໑໑໑ຩ <b>ວັ</b>	IJ G	-104	Divers	Η̈́ P.
72	1 1 1 10 1 mb		51		42	<del>(</del> ه بر) <del>د</del> ه	Minor	Decompress
12	-1 # N) # 1 -++	W F	งง	# F I B B	<b>1</b>	8 8	llajor	ion Sickness

.

(11) Progress

-158-

The results recorded in Table XI. 2 show why efforts were eventually concentrated on the dives of shorter duration. None of the schedules had been used proviously in the experimental work but schedule 16 which was based on published tables (132) and which was well established in the past, was used in this poriod to introduce 5 mon to depths greater then they were accustomed to visit. It can be seen that eventually a schodule was produced for 300 feet for 10 minutes at the maximum depth which could be considered satisfactory in its chamber stages but in the sea it was not quite so satisfactory. The timing of schedule 12 yes netained but the oxygen content was raised from 10% to 13% and the few dives carried out on the schedule renumbered as schedule 25 scened at worst to be no more dangerous. A similar pattern of successful chamber dives more risky when tried in the sea and then becoming safer when the orvern content was adjusted, can be seen in the dives to 400 feet for 10 minutes. No satisfactory explanation could be found for the more serious outcome of schedule 21 which was supposed to be adjusted to be less risky then schedule 17. The most probable explenation lay in a misunderstanding of the rate of gas absorption during longer dives but es time was short it was decided to concentrate on the shorter dives to show that the technique was possible and that the risk in diving to 400 feet was acceptable if a real necessity to dive to such depth arose. However even adjusting the breathing mixture before the first sea dives did not ensure complete success on the dives to 450 feet for 10 minutes.

-159-

(111) Results

9 experienced non carried out the 60 dives in the Vermon stage. It was felt that the problems had been nearly mastered when only 1 major case and 8 minor ones occurred. The chamber dives in the Reclaim stage were not so satisfactory. 16 men carried out 49 chamber dives for 10 minor and 4 major cases between them. (No cases occurred in the 6 dives carried out by 3 men who were much less experienced then the others). 19 men took part in the 70 sea dives with 9 minor and 7 major cases between them. One very experienced man had a major incident on his only sea dive and he remained unfit to take further part underwater; 6 men with relatively little experience did 9 dives between them with only 1 minor case occurring in this novice group; and 3 of the experienced men did 35 dives with 4 minor and 7 major attacks between them but 2 of them withdrew from the team during this stage of the trials. c). Third Period(60, 105)

- (A) Laboratory Stage
- (1) Proparations

The third experimental period started with an attempt to check or establish certain basic points first on the assumed curres of gas elimination of dives which did not require decompression; second on the pressure drop for safe decompression by ascent to the first and subsequent steges; and third on the duration of these intermediate steges. The approach was based on the experiences in the previous periods, and on animal experiments. The use of 10% oxygen/90% helium for the dive and 100% oxygen for the shallow stages of decompression was augmented by the use of various oxygen and helium mixtures richer in oxygen such as 20/80 and 40/60. All dives were intended to be 16 minutes duration from leaving surface to leaving maximum pressure, with the descent at a rate of 100 feet per minute; this time was selected as giving a 10 minute working interval at 600 feet. Finally if a schedule was considered unacceptably risky, minor changes only were to be made to define the area of risk if possible.

(11) Progress and Results

The progress and results are shown in Table XI. 3 which covers the laboratory stage only. After the initial work-up dives on air, various points on the assumed depth-time curve for helium absorption by the body were tested initially with minimum decompression time (i.e. time for ascent only) to strespheric pressure and then with a decompression stop at 10 feet. 125 feet was considered to be about 20 feet shallower than the risky depth as indicated in work elsewhere (50) and this depth was selected as the first one to be tried. The schedules were numbered in order of first use and the effect of cases of decompression sickness can be noted. After the depth of the no-stop dives was reduced to 90 feet progress was more orderly. One dive to 235 feet had to be shortened because of illness of a diver at depth. The dives at 340 feet were unexpectedly risky initially and 2 pairs diving to that depth also had their dives curvailed for unexplained illness of one non in each pair while at this depth. Subsequent progress was more hazardous but new

-161-

nen al constructions and a particular an and a particular states of the			A CONTRACTOR OF A CONTRACTOR O	and the second state of th				
	Divo		Schedule	Numb	or of	Decompression Sickness		
Depth Ft.	Duration Mins.	Gas	Number	Dives	Divers	Minor	Major	
250	16	Air	26	4	4		**	
<b>30</b> 0	16	Air	16	9	б	2	5	
125	16	Не	27	11	5	1		
165	16	lio	28	4.	4		<del></del>	
210	<b>†</b> 6	lie	29	3	3	3		
160	16	Нө	30	<i>l</i> į.	4	-	5	
160	16	He	31	5	3	Silvest	ĩ	
185	16	He	32	lų.	l <sub>t</sub> ,		1	
100	16	Ho	33	9	7	1	1	
160	16 *	Ile	34	12	5	-	-ș <del>a</del>	
220	16	He	35	10	lų.	1	-	
90	16	Ho	36	17	11			
135	16	He	37	12	5	2		
280	16	He	38	8	L <sub>l</sub> .	1	-	
235	16	He	39	20	10	1	610	
235	14	He	39	2	2	3.00		
340	16	Ho	40	3	3	تبو	f I	
340	16	lio	41	2	2	year.	1	
34.0	16	He	42	12	6.	2		
340	5	Ho	42	2	2			
340	16	Air	43	2	2		ant.	
400	16	Ho	44	10	6	1		
400	16	Ио	45	10	8	3		
300	16	He	46	121-	10		1	
500	16	IIo	47	L.	4	1	2	
500	16	Ho	48	4	L <sub>l</sub> .	1	1	
<b>50</b> 0	16	Ho	49	10	7	Ļ	-	
400	16	IIo	50	6	6	1000 1010 - 1010 - 1010 - 1010 - 1010 - 1010 - 1010 - 1010 - 1010 - 1010 - 1010 - 1010 - 1010 - 1010 - 1010 - 1010	-1 meaning beams a sub-statement	
	All chambe	ər div	9	213	25	21;	17	
	(Main äiv:	ing te	am)	(186)	(12)	(22)	(10)	

# -162-

## Table XI. 3

Result of Dives: Laboratory Stage, Third Experimental Period

----

schedules were calculated and it seemed reasonable to proceed to the sea stages. The main burden of the diving was carried out by 12 men. Among the 13 men who carried out a small number of dives were 8 men who had two acquaintance helium dives each in the chamber before going on the sea jump.

- (B) Sea Stage
- (i) Preparation

To make maximum use of the time available to the ship for this sea period, it was not intended to recalculate the schedules in case of unacceptable incidence of decompression sideness but to reduce the durations of the stay at maximum depth progressively by 1 minute intervals while retaining the same decompression routine. Handling difficulties meant that the first modifications had in fact to be made to the decompression routines as the rates of ascent to the first stop demanded in the schedule could not be met by the winches. The times in the tables used in the laboratory were delayed by a half-minute from arrival at the first stop upwards, to meet the problem.

(11) Results

The operating team aboard H.M.S. Reclaim included 6 men in the phips' diving team; 8 men from the experimental diving team; and 4 diving officers. In addition 3 men were released, from their particular responsibilities, to experience the actual routine, and, together with 1 of the ship's divers and 1 diving officer carried out 8 dives between them, the bulk of the work being done by the remaining 16 men. The results are shown in Table XI. 4. (Note: 1 pair were deprived of their oxy-helium gas supply while at 300 feet and carried out the decompression on emergency air tables).

	Div	e	n na san kana sa	Sabaaria	Wizmio	er of	Decompression Sidness		
Dopth Ft.	Duxation Mins.	Ges	Enviro- nment	Mambon	Dives	Diver:3	Minor	Mejor	
250	16	Air	Sea	26	18	17		4444	
300 300	16 . 12	Ho Ho/ Alr	Sea Sea	46 43	16 2	<b>1</b> 6 2	2.	6400 	
300	16	lio	Chamber	46	2	2	ander 164 - party andere se andere van de state se andere en andere	a vezdavý se sa stára fa díter a stára stára teleforma a se	
400 400 400	16 16 15	fic Ho He	Sor Sor Sor	50 51 51	2 4 6	2 4 6	1	- 2 1	
300 400 300	16 14 16	fio Ne No	Soa Soa Chember	52 51 52	4 2 2	4 2 2	1 2 -	4956 1466. 	
400 300 500 500	14 10 16 15	Re He Ne Re	Soa Sea Sea Sea	53 52 54 54	8 2 2 12	8 2 2 10	1		
(	Mird peri	oda	Total, SI	119	88	. 21	12	Q,	

# Table XI. A

Result of Dives: "Recleim" Stage. Third Experimental Period

## d) Fourth Period (81)

This period is incomplete in this work as only dives supervised till the demission of office by the writer are considered. It is relevant to this project as the cases of decompression sickness that developed could not be treated by previously published therapeutic tables because of the depth of occurrence of signs and symptoms, and the complications of the use of high partial pressures of dry oxygen. In essence, the experience gained as described in previous sections on air diving and submarine escape together with the experience gained in the earlier deep diving periods was invaluable in the minute by minute changing circumstances of the cases that are now reported.

## (i) Preparation

From animal work that had been continued in parallel by others at the laboratory it became apparent that carlier theories of helium absorption and elimination were unreliable. In particular it appeared that the body became saturated with helium more quickly with the result that the short duration dives that had been attempted had been on the steepest part of the exponential curves for gas uptake and hence there were too many variables to afford a reasonable prospect of finding a reliable decompression routine. The whole plan of attack was changed and it was decided to expose divers first for a series of two hours and then for a series of four hour no-(decompression)-stop dives which with the established 90 feet for 16 minute no-stop dive would show the probable shape of the curve. As usual all dives would be exercise dives, on a

-165-

rowing machine. A carbon-dioxide absorption system was to be used which needed a foot-operated bellows for circulation of gas, as all the dives would be done breathing various chamber atmosphere mixtures of oxygen in helium (100% oxygen was provided by the usual domand valve to prevent bypoxia on initial compression as well as during certain parts of the decompression).

(ii) Progress

The animal experiments had suggested that 45 feet would be the danger depth for men on 2 hour exposures but Table XI. 5 shows that 70 feet was accomplished reasonably easily by the small diving team. As a consequence of this experience it was decided to start the 4 hour dives at 55 feet - i.e. 10 feet deeper than the depth previously thought the limit for 2 hour dives; the table shows the results, the effect of leave, and of lack of practice.

As a result a curve was found and considered established. It was decided to carry out a further series of dives aimed more to check theoretical points than to produce a praoticable schedule for sea dives. Hence the requirement to carry out a fixed number of dives was changed to a requirement to dive each of the individuals in the small team once on each schedule. In addition it was not so important to consider the total decompression time as it was better to remain at various decompression stops for a time long enough for any signs and symptoms of decompression sickness to appear; this pause would then point to the safety of the preceeding part of the schedule. The dives selected wore 300 feet for

-166-

Dive			Number of		Decompression Sickness		
Depth Ft•	Duration Mins.	Schedule No.	Dives	Divers	Minor Cases	Major Cases	Notes
90 35 40 45 48 51 54 57 60 63 66 69 72 75 55 60 300 500 300 500 300 500 300 500 300 500 300 55 45 300	16   120   240   120   120   120   120   240   120   120   120   120   120   120   120   120   120   120   120   120   120   120   120   120   120   12	36 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81	1266664 1 66 66666 4640000000 1 88	1266667 66 666666 46422222224 22		1	Case: Diver FK. Case: Diver FE. Case: Diver FT Post-Leave Diver LB locum for Diver BF. Diver CR locum for Diver PT. Cases: Diver BF,HI. Diver BF first dive on return. Case: Diver CR. Cases: Divers FE, LS, BF. Cases: Divers FE, LS, BF. Cases: Divers HI, BF. Cases: Divers HI, BF. Cases: Divers HI, BF. Cases: Minor - Divers BF, WF. Major: Diver PT, first dive on return to team. Case: Diver BF. Diver CR locum for Diver BF. Cases: Divers FF, HI. Divers BF, WF. Both Chesty. Divers CR, HI. (CR Chesty). Divers CR, HI. (CR Chesty). Diver WF. (Diver FE well). Diver BF. (Diver LB well). Divers FF, LS. Divers CR, HI. Both Chesty. Divers WF, LB. Divers WF, LB. Diver CR, major. Diver CR well).
-companyable and a second	 Total		130	12	19	15	ATAGE AN MATTI-

### Table XI. 5

## Result of Dives: Initial Part of Fourth Experimental Period

(All carried out in compression chambers at Royal Naval Physiological Laboratory using several mixtures of oxygen and helium as well as 100% oxygen. Programme continued under other Medical Officers).

. . .

. . .

. . .

( ) \* 20

1,81

-167-

4 hours, 500 feet for 2 hours, and because it was the maximum depth of the facilities, 800 feet for 2 hours. Various alterations to the tables were made as the programme advanced, mainly to the exygen content of the breathing mixtures at the various stops but after the failure of the 800 feet schedule, the same schedule was also used for the next 700 feet dive.

(iii) Convient

As well as showing that a man like diver FE could vary by as much as 20 feet between dives of similar duration that produced transitory symptoms of decompression sickness, Table XI. 5 shows the effect of a lay off in a man like diver FF. This lay off effect is difficult to assess later because each dive then took so long that any man had up to 14 days between his dives - the last 8 dives in the table were spread over 2 months - and there was much more tranble then. In these last dives, the men surfaced on occasion complaining that their cheets folt tight end raw retrosternally on inspiration and clinically the best diagnosis eppeared to be "pneumonitis" because of the slight changes in vocal resonance and occasional coarse rales. One man had radiographic changes described as "pneumonitis". Three episodes occurred in the heaviest smoker in the group.

## Decompression Sickness

It was not possible to compare the records of individual divers to find the differences between men who only had major attacks and men who only had minor attacks as the decompression mutine would be altered

-169-

after such episodes and thus the men would not carry out similar dives. Even the various periods cannot be reliably contrasted with each other because of modifications made in the basic assumptions and differing degrees of overlap of membership of the experimental team from one period to the next. 66 divers carried out 734 dives which were distributed over 81 different decompression schedules. 170 cases of decompression sidness occurred of which 62 wore treated by some recompression routine. 1 men was left with a permenent disability end one other had a very slight needbuck impairment to his hearing which he did not notice. A factor that may have masked other cases was the occasional recompression of mon in a pair of divers if one developed sions and symptoms which indicated a need for such treatment. This masking may have occurred in those few cases where there were no fecilities to separate the men and the need for recompression erose during the decompression before arrival at atmospheric pressure. Several times however the second diver also developed less severe symptoms which might have responded to less intensive therapy then that needed by his partner but it was accepted that he would be over-treated and he would act as attendant to his less fortunate collegate.

3 men dived on over 50 occasions each, with 19, 12, and 5 incidents in decreasing order of attacks but recompression was required on 5, 1, and 3 occasions respectively. The range of frequency of dives was from 1 to 59 depending on whether the individual was a regular member of the team or a visitor - for example 2 men of the ship's diving team during

-169-

the sea steges later joined the experimental team for subsequent laboratory and sea stages. 19 men dived and did not have any attacks but none of these dived more than 4 times each. 10 men had 13 attacks of decompression sideness between them, none treated by recompression while 12 men with 16 attacks were always treated by recompression and had no other attacks. The remaining 25 men had 141 attacks of decompression sideness of which 46 required recompression. The only deduction to be made was that the oftener a men dived, the more likely he was to dive on a schedule which had inadequate decompression.

It is worth zecalling that the difference between minor and major cesses depended on the treatment given and in turn this depended on the assessment by the same doctor of the viotim's meestions in the light of experience and of the severity and nature of the symptoms. No man who wished to be recompressed was over denied such treatment - several men had to be bullied into going into the compression chamber. The reporting of minor complaints by the man or the inquisition for them by the doctor was always a vexed point and depended to a great extent on the doctor patient relationship. This naturally fluctuated with daily close contact but on the whole an atmosphere of trust was engendered. This meant that the doctor would make the treatment suitable while considering his patient as well as the injury. Inevitably in such a situation some cases arose which in retrospect might have been more effectively treated in some other way.

-170-
There was the customary variety of signs and symptoms in keeping with the hypothetical variety of sites of bubble formation. Signs could occasionally be found either as direct consequence of the specific trauma or as result of reaction such as local tissue ordema. Assessment was often complicated by the multiplicity of sites involved concurrently for instance pain occurring in the right elbow and left knee - or consecutively as in pain occurring first in the right shoulder then clearing, then developing in the right hellux then clearing then settling in the right knee for a more prolonged spell. The difficulties were compounded when therapeutic recompression was carried out for a particular symptom or sign but in the later stages some new symptoms or sign would appear to pose the question of recurrence or new attack.

The type of onsist of symptoms and signs may be dramatically sudden or insidious to such an extent that only later is it possible to say when uncase marked the start of the incident. Such a slow onset may be aggrevated by wishful thinking on the part of the victim. As the dives were experimental in nature, the onset of decompression sickness could occur at any pressure. 50 of the cases first developed some indication of future troubles while still under pressures which varied from the equivalent of 490 feet to 10 feet. In many of these occurring at shallower depths no report was made till arrival at atmospheric pressure. Of these cases, 28 did not require recompression, 13 were started on therapeutic recompression before reaching atmospheric pressure, and 17 were recompressed later. It was important to the divers to reach

-171-

atmospheric pressure as this would permit transfer to more comfortable facilities as well as providing a "smoking-break".

The time of onset - and of reporting varied considerably. It wes thought that most urgent cases of collapse would occur within 10 minutes of the reduction of pressure and most other cases within 4 hours. This approach was a guide on how near to recompression feellities the divers should remain after they had reached surface and they had no complaints. The type of onset made timing difficult but cases with neurological damage could usually be assessed more accurately. The longest period between reaching surface and the onset of the first symptoms was 82 hours, and 5 had intervals longer than 3 hours. Some cases had early troubles which cleared and later developed a constant symptom not necessarily in the same site. The time of onset during decompression is vital as it indicates the stage where the mutine has been inedequate - a point exemplified by the case that occurred after 1 hour at the first stop after leaving maximum pressure. First stops previously had always been short. sometimes as short as 2 minutes and may well have been inadequate in the past to show where the defect in the decompression schedule was to be It will be seen that there was no difference in its character found. whother helium or nitrogen was the inert gas used.

Pain in some form was the commonest complaint. The "niggle" was a consciousness of pain which was more of nuisance value than causing disability or discomfort; it was usually transient rather than persistent. It could develop however to "ache" which was more constant and distressing being described as dull, deep seated, throbbing, drawing, or nagging; there was more persistence with this symptom and the severity varied from responding to time and mild analgesics to a syncopal attack. "Pain" was more intense, more localised, and charper than an ache and usually indicated that recompression would be needed. 109 of the 170 cases had pain in some form varying in severity and in site (Table XI. 6).

S2to	No. of Cases
Uppor limbs and Shoulder girdle	61
Lover limbs and Hindquarters	86
Trunk (excluding chest)	8
Generalised headache	5
Pein in chest	5

### Table XI. 6

#### Sitos of Pain

When exercise was weight-pulling upper limb sites were more frequent than later on when a rowing machine was used. Pain in the chest was a problem as At was usually associated with marked difficulties in respiration. It was always treated by recompression. Among the divers it seemed to be the worse degree of "tightness of the chest" rather than the muscular and skeletal pains of the rest of the body. No assessment was possible on whether the sensation had a cardiac, respiratory, or parietal origin.

An unusual blotchy reddish-purple <u>rash</u> was reported on 13 occasions.

The rash did not have a raised edge and was more suggestive of vescular statis in the skin vessels with cyanosis. It did not cause any inconvenience and it was usually noticed on changing clothes or while having a shower. The distribution tended to be confined to the upper trunk and arms. It did not indicate severity of the condition as rarely was it found in cases needing further treatment. As it responded to a hot shower or simple passage of time, no particular importance was attached to the appearance of this lesion.

<u>Liching</u>, mainly of a prickly nature, was reported on 22 occasions. This symptom was nost common in the beard area, the forearms and the trunk. It also occurred on the thighs. At no time did it lead to such scratching as to cause abrasions and the duration was short, being often out even shorter by a hot shower. It appeared to be associated with chamber dives where the body was more exposed to large volumes of air and the large temperature changes than when the diver was in his formfitting suit in the sea.

The <u>respiratory system</u> was apparently involved in 23 cases. 17 of them being recorded as simple "tightness of the chest". The significance could not be assessed because of individual variations. For example, one diver had this symptom recorded on 4 occasions but he claimed that he had some degree of difficulty of expiration after more than 40 of his 50 dives. In none of the cases did the symptom persist more than a few minutes and it cleared without treatment. The possibility remained that the sensation was really an indication of the extra effort required

-174-

of the respiratory muscles in the ventilatory dynamics under pressure. The remaining 6 cases had a retrostornal sensation of fullness associated with a short dry ticklish cough and basel râles; one man had X-ray changes suggestive of a patch of pneumonitis but the X-ray films of the others were reported as normal. These cases all occurred in men carrying out long deep dives who developed decompression sickness during the decompression necessitating therapeutic recompression without a surface interval. The duration of exposure to pressure varied from 22 to 52 hours when the partial pressures of oxygen were 1 atmosphere or greater. It is a debateable point whether the signs and symptoms were decompression sickness or not: and if not, whether they were due to oxygen or dry gas. The signs and symptoms lasted 3 days to 6 weeks. It should be borne in mind that after an exposure which could lead to decompression sickness, most cases of tightness in the chest and a slight cough either clear quickly or rapidly lead to a deterioration in the patient's condition with the consequent urgent need for recompression.

10 cases complained of some <u>disorder of nower</u> which was sometimes associated with pain as in cases of paralysis and encesthesia with a band of pain immediately proximal to the level of encesthesia. On other occasions weakness seemed to follow pain and it was then found as a reluctance to move a joint rather than intrinsic loss of power. On four occasions, there was general weakness associated with exhaustion of the individual. One man had a paraplegia about the level of the eleventh

-175-

thoracic segment with encesthesia of the same distribution which left him a permanent cripple in spite of recompression therapy. Both lower limbs were found to be weak on eight other occasions, while in a further three cases the right lower limb only was involved twice and the left once. The left upper limb was involved in the remaining two cases, once in the whole limb and once confined to the forearm. In only one case, involving the lower limbs, were there any tendon reflexes changes and then the knee and ankle reflexes were more brisk and bilaterally equal, after the dive than before.

<u>Sensations</u> were disturbed in 14 cases. 2 of these cases were associated with paralysis - one the paraplegic mentioned above and the other when the paralysis involved the left upper limb from the shoulder distally. In this latter case the disturbance took the form of complete loss of light touch, pressure and pinpricks; proprioception was also disturbed. 4 cases had localised areas of diminished response to light touch and pinpricks on the anterior aspects of both thighs (once), the dorsal aspect of the right foot (once). A seventh case had loss of discrimination of the terminal pade of all the digits of each hand = he also had possible exygen demage to his lungs (similar sensory loss in association with exygen convulsions has been described (15)). 3 cases complained of paraesthesia, one in the right shoulder and two as a generalised phenomenon. 3 cases complained of vertigo, it being confirmed later in one case who was considered to have transient vestibular codema 10 men had "<u>synconal</u>" attacks. By this is meant a rapid onset of weakness, pallor, cold sweat, nausea (with vomiting on three occasions) and transient loss of consciousness (twice) very suggestive of syncope except that the radial pulse was steady, rather slower than expected, and of good volume. Immediate recompression restored the normal physical state dramatically and on 6 occasions no other signs or symptoms appeared. The remaining four cases had either complained of some limb pain shortly before the attack or on recovery and were treated accordingly. All were considered to be cases of decompression sickness.

<u>Personality</u> was affected on 4 occasions, twice as an extreme degree of apprehension which developed during the stay at maximum pressure and persisted throughout the decompression and for some time thereafter; no adequate explanation is avilable for this clinical picture. The remaining 2 cases were classical cases of depression arising during the decompression and who returned to their normal state of lively, alert integest during recompression as if they were statues being unveiled.

In retrospect 6 of the men involved should have been recompressed as the quickest end most comfortable treatment. None of them appeared to have any permanent damage but the duration of symptoms was much longer than a therapeutic recompression might have been. Several of the milder cases of aches and pains and some of the cases of general apprehension were almost certainly not due to inedequate decompression but they had to be considered as such as the worst possibility. Notes on Major Cases: First Period

Diver DA had dived in a compression chamber to 300 feet for 10 minutes using air before; on this occasion a shorter decompression routino was used which involved on ascent rate of 120 feet per minute to the first decompression stop. On reaching 40 feet, he first felt slight difficulty on inspiration and this increased at the surface where he also developed aches in his chest. He had used "controlled deep breathing" during the ascent on his own initiative. He went about his duties for 20 minutes before he developed. first difficulty in controlling and then loss of power in his legs. He was recompressed without further exemination and he had made apparently, a complete recovery by the time the pressure had reached 100 feet. He was further compressed to 165 feet and a therapeutic decompression routine was started on Therapeutic Table III but as physical examination was still negative and his condition was so good, it was decided to reduce the time at 30 feet to 2 hours thus converting the therapeutic routine to Therapoutic Table II (Shorter). No complaints were made end examination was still negative. Several days later he mentioned a slight woolly sensation in his right foot which had persisted since the therapy and which he had concealed. No further treatment was thought worthwhile and gradually the symptoms cleared. His attack was thought to be due to pulmonary barotrauma due to his controlled respiration, aggravated

-178-

by considerable exercise immediately after surfacing.

Divers UL and AN together carried out a chamber dive to 400 feet for 10 minutes using 5% oxygen in 95% helium. Both men had been trained to equalise the pressure in their Bustachian tubes by swallowing and neither men had previously dived deeper than 300 feet or used a helium mixture. The dive proceeded according to plan with the mixture being breathing from 100 feet on compression to 140 feet on decompression, otherwise the chamber etmosphere of eir was breathed. On reaching 20 feet, Diver US complained of abdominel disconfort, backache redicting wound the abdomen. and he looked distressed: Diver AN complained of a mild ache over the left shoulder area. Examination suggested that gestric distension was the main trouble in each man but in any case recompression was required. In both men symptoms were relieved - and in Diver UL retching followed by vomiting, probably helped as much as the pressure increase - and Therepeutic Table II was used with apparent complete success (Recommended). It was considered at the time that leck of familiarity with depth and ges as well as apprehension and their carclearing technique had lod to excessive gas swallowing rather than that inadequate decompression was the fault. That explanation was too facile in retrospect and it is now considered that they were genuine cases of decompression sideness.

<u>Divers CK and MR</u> had the honour of carrying out the first sea helium dive under the new experimental programme after a work-up to ensure that the drill was correct. Both men breathed the 5% oxygen/

-179-

95% helium mixture from 100 feet on the descent to 300 feet for the period of 10 minutes at that depth and then to 90 feet on ascent and thereafter they breathed air. The submersible chamber was being transferred on to the receiving chember at 20 feet - the deepest stop of duration long enough to permit the manoeuvre - when the divors requested recompression. The transfer was completed first and it was then seen that diver MR was writhing on the chamber floor and that he had great difficulty in breathing. Becompression to 100 feet was immediately carried out. Examination of both men at this depth was negative and they each complained of a very slight general sensation of stiffness. It now became known that he serious mistake in depth keeping had occurred and that the men had been 14 feet shallower at each of the designed 40 feet and 20 feet stops; and that both men had developed slight difficulty in breathing during the ascent between these stops which had become more marked on arrival at the shallower depth. In Diver MR this dysphoes had been accompanied by a tightness in the chest which developed into a severe pain which radiated generally while in Diver CK a similar start was followed by radiation to the left shoulder and to the occiput. As soon as pressure began to be increased, the symptoms had ameliorated. In view of the speed with which pressure was increased; the rapid response to recompression; and the negative examination at 100 feet. it was decided to depart from standard practice and to try using Therapeutic Table I (Shorter) under constant observation instead of the recommended Table III. If any signs or symptoms recurred

-180-

then the alternative of Table IV would be immediately used. On reaching the surface, the only complaint was that the sensation of stiffness had persisted unchanged. These two cases were undoubtedly the type of decompression sickness referred to as "chokes" and essured to be due to multiple pulmonary artery gas embold initiated by the gross error in decompression.

Diver WI and his partner later carried out the same dive planned for Divers OK and MR after another pair who had uneventfully carried out the same dive. As soon as Diver WI transforred to the T.U.P. chamber, he was seen to be distressed. Preseure was immediately increased from 20 to 60 feet where he seemed to have recovered. Examination was negative; the history was similar to Diver MR. He and his partner were further compressed to 100 feet and it was decided to use Therapeutic Table I again. However after 25 minutes at this depth and 5 minutes before decompression was due to commence. Diver WM complained of coldness of his feet and numbress of the front of his thighs, he was immediately further compressed to 165 feet with complete relief of his coldness and a decrease in the area of numbross. The intended therapy now was to consist of a 2 hour exposure at this depth and then decompression on Therapeutic Table IV. His condition gradually deteriorated at this depth and he developed a paraplopia with paralysis and encesthesia to the nipplo area with a constricting band of pain and a narrow band of pareesthesia oranially to this level. Decompression was eventually commanced with no further deterioration. General nursing

-181-

procedures were used along with antibiotics and sedetives. Attempts to use closed circuit exygen breathing equipment to increase the oxygen content of the patient's inspired gas caused severe distress and laboured respiration and wore abandoned. It was folt that the pause in recompression at 60 feet to allow examination was the main fault as some of the pulmonary arterial emboli of "chokes" were presumed to have been compressed enough to pass through the pulmonary shunts thus reaching the central nervous system where growth of the bubbles restarted and each further increase in pressure only temporarily alleviated the condition. If the pressure had been increased initially to 165 feet or subsequently to 250 feet then the bubbles might have been compressed below the critical size for re-dissolving. In addition. the use of a less dense 20% oxygen/80% helium as a breathing medium during the therepeutic decomprospion might have been more tolerable for the patient as well as playing some part in encouraging any nitrogen which had diffused into the original helium bubbles to diffuse out thus making a less persistent traumatic agent. (2 years later, WN. atill a paraplogic. has some movement in one legs he has managed to remain standing without support for a short periods and to walk about 30 feet unsupported but then loses his balance because of clonic spasse. His progress is continuous but very slow - in keeping with observation by foreign colleagues with experience of similar diving cases).

<u>Diver HI</u> carried out a chamber dive to 300 feet for 10 minutes using 5% oxygen in 95% helium on a decompression routine which had a

-182-

slower accent rate than before and in which the transfer under pressure was simulated during an 8 minute stop at 100 feet. He surfaced with a stiff right ann which rapidly became very painful. Regrettably personality problems overshadowed the case and he was not recompressed. Strictly, this incident does not satisfy the criteria for classification as a major case but in every other respect of onset and severity it was a serious incident and it would be grossly micleading to omit it. His symptoms slowly waned over 3 days.

<u>Diver AD</u> also carried out this same chamber dive some days after Diver IV. He had slight sensation of stiffness of both his shoulders which steadily developed as an aching pain worse on the right and he was recompressed when he reported 25 hours after reaching surface. His symptoms were relieved at 120 feet and decompression on Therapeutic Table II was carried out (Recommonded). Shortly after surfacing, he had a very mild ache in his vight knee and examination disclosed a small quantity of fluid in the knoe-joint; conservative treatment was prescribed. (24 hours later, a diving officer seeing Diver AD limping. and on his own initiative, recompressed him again to 165 feet with no relief and used Therapoutic Table II to return to the surface - and then Diver AD informed the diving officer that he thought there had been some relief to his mild acho). These incidents showed that the cases were now not so dramatic. The latter case showed that long delayed recompression for minor symptoms is not necessarily the most effective treatment.

-183-

Divers DA and AD both carried out the work-up see dives on air to 250 feet for 10 minutes on the bottom. Neither men had carried out this dive on the provious trip. Diver DA because he was unfit to dive following a case of decompression sickness (q.v.) and Diver RB because of mib injuries. [The work-up decompression schedules involved a fast rate of ascent but as 12 individuals had used them without incident on the previous trip they were retained for the work-up on this occasion. On this trip 14 mon used them but only 7 uneventfully and 5 others with minor complaints. ] On surfacing Diver DA had a sharp pain in his chest, mild dyspnoon, generalised paraesthesia and a marbling skin rash of his trunk. Examination suggested a local area of hyperresonance in his left chest posteriorly compared with the right but no other abnormalities. As his condition was improving, rest and observation vere ordered. He had made a complete recovery in 12 hours. Diver NB started with an ache in his left shoulder which cleared completely within 10 minutes of curfecing and he complained of a patch of diminished sensation over the right forefoot. Exemination disclosed a transient hyper-resonance similar to that in Diver DA and no other abnormality and as symptoms cleared rapidly no further treatment was used. No radiographic facilities were available for either case. As it seemed that both men had some julmonany abnormality which might be causing pulmonary barotzauma, it was decided that they should be considered serious cases who might well have required recompression - at any rate it was decided that the ricks were too great for them to continue

-184-

experimental work and they were given other duties.

Diver ME was also a man who was not recompressed although his incident in retrospect after a chember dive to 300 feet for 10 minutes on 5% oxygen in 95% helium should have been considered "major". At the 20 foot stop. Niver WE developed an ache in the low lumbar area. which waxed over 3 minutes and then waned to disappear over the next 15 minutes. Exemination disclosed no abnormality. 5 minutes after reaching surface, his symptoms returned in a milder form and examination showed that there was loss of power of both legs on resisted extension. As his general condition was good, and more severe symptoms carlier had cleared, it was decided to wait and watch his progress. After 2 hours, no weekness could be detected but he still felt slightly week end this condition persisted for 24 hours. During this time he remained in a comfortable bed instead of an uncomfortable recompression chamber where the return of power would probably have been quicker but the therapy would have taken as long. On reconsideration this case was considered as a major one as recompression was the correct therapy and there was no certainty of recovery without it indeed the delay could cosily have resulted in permenent neurological demage.

<u>Diver WF and Diver EO</u> carried out the same dive in the chamber os Diver FE later the same day. 5 other men had also previously carried out this dive, one of whom had had very mild niggles in his shoulders. At 20 feet Diver EO developed a dull ache in his coccygeal area with slight diminution of sensation in his right foot. After clearing while

-185-

still at that depth, his symptoms recurred with increased severity on reaching surface; he also developed a typical trunk mash. His pain worsened and spread to the perineum and he developed a pain in his left shoulder. Recompression was advised and while he was being taken into the chamber he developed extreme weakness of his legs. Meanwhile Diver WT developed severe pain in his left calf and upper left thigh about 10 minutes after reaching atmospheric pressure. 18 minutes after reaching the surface, recompression of both men started. Diver EO's signs and symptoms, although apparently more serious, cleared at 100 feet while Diver WT did not respond till shortly before reaching 165 feet. Therapeutic Table II was used with complete recovery in the case of Diver EO (Shorter) and a stiff sensation in the legs of Diver WT (Recommended). (Divers FE and EO had cenced for over 2 hours the previous evening and it is tempting to escribe the unusual sites of their pains to this).

<u>Diver WP</u> caused out enother chamber dive to 300 feet for 10 minutes using the same gas as before. This dive followed another dive, which had resulted in therepeutic recompression, after an interval of two clear days. 12 other men carried out uneventfully this particular routine. Diver WP compleined while at the 20 foot stop of pain in his chest and mumbress and stiffness of his hands, wrists, buttocks, thighs, and feet. His condition rapidly deteriorated and numbress spread to the lumbar area. He was recompressed to 165 feet where examination showed that the patient was very depressed, and, had some scattered

-186-

zhonchi. Decompression was started on Therapeutic Table III (Recommended) but his condition at 100 feet suddenly caused anxiety so he was again recompressed to 165 feet. Re-examination showed that the patient's morale was better and that he had had a syncopal attack apparently vaso-vagal in origin. After 2 hours, decompression on Therapeutic Table IV was started (Recommended) and, although he had some generalised aches at 50 feet which responded to tablets of codein compound, he eventually neeched atmospheric pressure well but entremely fatigued. It was considered that Diver WT had not really been fit enough to dive. The case showed the routine of changing from one therapeutic table to another. The cause of the depression was debateable as long afterwards it was disclosed that he had been up till 3 a.m. drainking beer the night before - a fact deliberately concealed from the doctor who examined him as he knew he would not be allowed to take any further part if detected, in the trials.

<u>Diver DR</u> dived to 300 feet for 10 minutes and the dive vent according to plan. 20 minutes after surfacing, he had mild pains in his shoulders, a marked body rash and he felt faint. On examination he was pale and had a cold sweat but his pulse was good; his rash was the typical marbling kind. It was decided to treat him by resting in his bunk, with his head slightly lower than his feet. Recovery of his shocklike state took place over two hours and his shoulder discomfort was replaced by a mild low back ache. Next day his legs felt a little weak but examination disclosed no abnormality. This patient should have been

-107-

necompressed but it was considered that recovery was probable in as short a period when treated by observation as by recompression. The case was of value in that it showed the possible progress of the untreated condition when improvement begins rapidly. If this case had occurred in the later stages of this trial he would have been rapidly recompressed. The immediate access to treatment chambers and an unwilling patient aborted the decision to observe the patient, but later experience of unpredictability of such cases led to recompression.

<u>Diver EQ</u> carried out the same dive as Diver DR. During the stop at 20 feet, Diver B0 complained of a mild ache in his right knee which eased, wowened, and then eased again but examination was negative. Decompression continued and within 5 minutes of reaching surface the ache in his knee spread throughout the whole night lower kimb and up to the same-iliae neglon. He was then recompressed and as his symptoms had cleared before 60 feet was reached, Therapeutic Table I decompression was carried out (Recommended). After surfacing, he complained that the affected limb felt rubbery but after a nights' sleep, this sensation had cleared.

Diver PM and Diver CC twice repeated the dive of the two preceeding oases with an interval of ) days. On the second ocdasion, Diver FM developed a mild right shoulder pain at the 20 foot stop which eased and decompression continued. At the surface, it worsened and radiated across to the left shoulder and early signs of shock appeared. Meanwhile Diver CO, on surfacing, complained of stiffness and mild aching in his

-188-

hamstrings in both thighs. Both men were recompressed. Diver EA, apparently the more seriously affected was free of symptoms by a depth of 60 feet but Diver CC, after an initial improvement deteriorated egain and eventually cleared completely only at 165 feet. Therapeutic Table III was then carried out and noither man had any residual symptoms on completion of the decompression (Recommended Diver FA. Longer Diver CC).

Diver MR and Diver WC also repeated the dive after a three day interval and again decompression sickness developed on the second occesion. The first signs appeared at 20 feet where Diver MR had a mild algele inchis right knee. Diver WO appeared restless and apprehensive and, although he protected he was well, he repeatedly asked for the chamber to be ventilated. On surfacing the latter complained of pain in front of both thighs; so he was asked to leave the chamber for a short period while wet diving suits were removed and e mattmess was put inside prior to recompression. In this very short period he suddenly fainted. He was immediately put into the chember and recompressed. Diver MR was also recompressed in case his complaint recurred later elthough he had no symptoms at the sotual time. Diver WC did not improve till a depth of 165 feet was reached and then Thenapeutic Table III was started (Recommended). At 40 feet, he developed dizziness, nausea, netching and vomiting with impalpable radial or brachial pulses. The occupants were again recompressed to 165 feet where Diver MR remained well but Diver WC did not improve as much as before, remaining

-189-

vory weak. lethargic and retching and weakly co-operative. Therapeutic Table IV was started (Recommended) but in addition a semi-closed circuit breathing apparatus was provided for him which used 30% oxygen/ 70% holium. He used this when he felt the need and his total spells amounted to 92 minutes cach spell ending when the advantages of the less dense breathing medium were counteracted by the discomfort of the equipment. It was remarkable how his general condition improved while using the equipment initially. Any attempt to sit up however, restarted the nausea, retching and dizziness. When an intravenous infusion was about to be started to restore hydration, he began to retain fluids orally and generally to improve. On arriving at 50 feet. examination showed an exhausted patient with slight weakness of his right thigh. After surfacing and a night's sleep the weakness had cleared and his fetigue had improved. Diver MR was decompressed once 30 feet was reached to relieve him of attendant's duties; he developed mild aches in both knees. which cleared with rest and a night's sloop.

Diver MR dived in the chamber again to 300 feet for 10 minutes. 10 minutes after surfacing, he complained of numbress in the secrococcygeal area and a feeling of heat radiating down the left log. This consetion changed character to severe pain which sapped the power of his logs, worse on the left. On recompression, his symptoms were relieved at 100 feet and Therapeutic Table II was used but on surfacing he felt stiff generally and later reported that this cleared with rest. (Shorter).

-190-

# (ii) Second Period

<u>Diver SD</u> complained on surfacing after his very first experimental chamber dive, one to 300 feet for 10 minutes, of slight stiffness in his right shoulder but this closered rapidly. 15 minutes after surfacing, he suddenly developed a sensation of loss of veight in both legs and examination disclosed a diminution of left knee jerk and loss of extensor power of his left thigh. During the examinations he complained of a sudden sharp pain in his left hip, and he was recompressed. His symptoms did not clear till 100 feet so he was further compressed to 165 feet and decompression on Therapoutic Table II started (Shorter). Although there was a short period when he complained of stiffness in his legs while at 60 feet decompression was concluded successfully.

-191-

<u>Diver MR</u> carried out a chamber dive to 450 feet for 10 minutes which followed the planned routine. On surfacing he reported an ache in both thighs which had just developed although he had had a mild ache in both forearms which had appeared at the 80 foot stop and each stop thereafter but which had cleared at each stop. While he was being questioned he felt faint and giddy and examination disclosed a hypotension and bradycardia which seemed to be recovering during the examination, so it was decided to keep him at rest under observation. Within an hour, he had recovered completely and he was allowed to join his measures. 2% hours after surfacing, he developed a slight ache in his right upper arm and in both knees which, over the next 3/4 hour before he reported, worsened and radiated to both thighs and hips. On examination, he was found not to be exerting full power of his limbs because of pain. Recompression commenced. His symptoms were not relieved when the pressure reached 165 feet so further compression was used and the symptoms cleared dramatically at 230 feet. Further compression to 250 feet was done before ad hec decompression was commenced to 140 feet where standard therapeutic tables were joined. In view of his condition, Therapeutic Table II was used successfully to carry out the shellower stops (Special). Diver MR was not compressed initially as his vagel symptoms were improving and his muscular symptoms had cleared at each stop. With previous experience in mind, it was decided to try a greater depth to speed recovery.

<u>Diver SD and Diver AS</u> together carried out a work-up see dive on air to 250 feet on the routine used in previous sea trips. Diver SD reported with aches in his left arm 25 minutes after reaching surface and in the next 5 minutes he reported blurwing of vision in his left eye, pain in his right loin, buttock, and grein so he was immediately recompressed with his symptoms, other than visual, clearing at 30 feet and the visual ones at 60 feet. Diver ES surfaced with some itching and, when Diver SD reported, it was found that Diver ES had had an urticarial rash over his left arm and forearm which had flared and cleared over a period of 10 minutes. It was decided to put Diver ES in with Diver SD in case of further trouble although at the moment of recompression he had no signs or symptoms. Therapeutic Table I was used and after surfacing, after intensive questioning Diver SD mentioned

-192-

a very slight stiffness of his right upper arm, 2 hours later this stiffness had increased and the mid humeral area was found to be swollen, painless, of normal colcur and non-pitting. This tissue swelling cleared over the next 2 days (Shorter Diver SD. Longer Diver RS).

Diver SD and Diver RS were again partners one week later for a chamber dive to 400 feet for 20 minutes. On surfacing. Diver SD mentioned an ache in his left wrist while at depth which had eased on decompression: this had occurred in the site of an old known malunited scephoid fracture. 40 minutes later, he complained of loss of power of his left arm as well as pain in both knees and uppor legs, spreading to his left hip. His symptoms gradually cleared throughout the compression till at 165 feet. he had completely recovered. Diver RS developed fleating niggles 10 minutes after surfacing which eventually settled as a dull steady cohe in his logs. It was decided to try the effect of breathing 100% exygen at atmospheric pressure and 20 minutes later he was sign and symptom free. However it was decided that once again Diver RS would be compressed with Diver SD and the niggles recurred as he was helping his colleague into the chamber. Decompression on Therepoutic Table II was uneventful for both men (Shorter Diver SD. Longer Diver RS). 9 hours after leaving the compression chamber, both men were permitted to go on a specially arranged bus trip as they insisted they were fully recovered. It was not realised that the bus trip would be at an altitude of more than 6,000 feet for some periods; at that altitude Diver SD developed blurring of vision and Diver RS developed

-193-

mild niggles in both knees. Both men recovered completely on the rapid return of the bus to sea level.

<u>Divor SC</u> carried out the same dive as Divers RS and SD on the same day. 3 minutes after reaching surface, he developed a nigele in his right shoulder where a tender spot was found on the posterior aspect of the joint at the border of the deltoid muscle. This cleared in about 40 minutes. 2 hours after surfacing a spagmodio stabbing pain developed in his left upper arm which recurred at longer intervals and less severe. It was decided to try the effect of 100% orygen at stmospheric pressure and after 40 minutes the residual sensations were clearing rapidly and therapy stopped. 4 hours after surfacing, the left arm ache returned and 3 hours later after local heat and massage had been ineffective there was a sudden increase in the severity of his pain. He was recompressed but the relief at 165 feet was so slight that it was decided to compress him to 250 feet. The pain cleared leaving a sensation of bruising, and this remained while a similar routine was used to that tried with Diver MR (q.v.) which finished with the later stages of Therapeutic Table II (Special). Diver BR who had not dived that week was his attendant and 4 hours after surfacing, the latter was allowed on the bus trip, mentioned in the previous case, it being assumed that as he had only been an attendent there was no need to keep him on board. Because he had not been specifically ordered to remain on board Divor SE also went on the trip. It later transpired that two hours after surfacing Diver BR had developed a mild ache in his left

-194-

knee with a sensation of weakness in his left thigh and that Diver SD had developed a very mild ache in his left knee as well as the persistent bruised feeling in his left shoulder. During the trip each man developed agonising pain in all limbs which was treated by rapid descent to sea lovel and intravenous morphine. In spite of printed slips in Spanish. language difficulties led to a 5 hour delay in transport to the nearest compression chamber (which was the ship) as the bus descended the other side of the mountain. Neither on arrival nor over the next 12 hours was any sign or comptom found to suggest the need for compression. This case showed that needless delay in therapy leads to decreased likelihood of complete and rapid success; that occasionally the attendants can develop decompression sickness; and that 250 feet may only alloviate rather then cure. Finally, it emphasizes that activities over the next 24 hours must be closely supervised both in their intrinsic nature and the distance from treatment facilities.

<u>Diver SR</u> dived in the sea to 300 feet for 10 minutes. He had previously carried out this dive uneventfully. This time he surfaced with a body rash and itching and an acho in his upper left thigh which cased over 10 minutes when an ache in his left upper arm stump was reported. This latter pain waxed and waned in severity over a period of 20 minutes, recurring at longer and longer intervals and no specific treatment was given. 17 hours after his dive - next moving - he reported because of a sharp pain in his left upper arm severe enough to bring tears to his eyes. This pain had lasted half-an-hour and he

-125-

admitted that the provious evening he had a constant dull cohe in the same area which he had not reported. On examination the area of the posterior border of the left deltoid was tender. It was decided to recompress him as a test of cure to 165 foot and initially all his signs and symptoms cleared but after helf-an-hour the same troubles recurred undiminished in intensity and it was obvious that his condition was intensitient in character. He was decompressed on a routine diving table and treated thereafter with strong analgesies and hypnotics. The attacks returned 4 times over the next two days - it being of interest that the sensation of phentom fingers returned in their correct spatial relationship although the intervening forearm was not sensed. This case was undoubtedly decompression sickness but the long delay in recompression made that line of therapy valueless in this case.

<u>Divor WS</u> was considered to be very experienced and relatively resistent to decompression sidmess so he was asked, as the 30th diver in the sea on the eighth schedule for 300 feet for 10 minutes to carry out an empirical test. It was intended in later dives to fill the submersible chamber with the breathing mixture so that the divers could rest from their equipment during this stage. Some work (11.94) suggested that helium had a rapid percutaneous diffusion and in such a delicate gas absorption situation as decompression was thought to be, this might possibly lead to complications. Accordingly, to avoid skin pinches his diving suit was kept inflated by an ory-helium mixture instead of the customary air. About 1 hour after surfacing in good

-196--

spirits and having had his midday meal with the measure of run customary in the Navy, he folt tired and, as men were excused duties after such a dive, he turned in in his bunk. 3 hours later, it was decided to waken him to make certain that he remained well. There was some difficulty in wakening him and he immediately complained of severe aching pains in both knees and both shoulders. He was assisted to the compression chamber. On recompression, his knee symptoms cleared first and his right shoulder felt normal by 65 feet. His left shoulder did not clear till 100 feet so he was further compressed to 165 feet and then decompressed on Therapoutic Table II (Recommended).

<u>Hver SR</u> dived to 400 feet for 10 minutes in the sea with Diver B0 as his partner. There was a slight error in the decompression when the submensible chamber was raised 10 feet past the 100 foot stop, but the error was corrected within one minute. Diver SB regularly complained of tightness in his chest after oxy-belium dives but this time both he and his partner requested to stay in the transfer chamber on arrival at surface because of vague generalised aches, slight difficulty in breathing and approhension. The symptoms cased so both men were assisted over the 15 feet distance to a larger and more confortable chamber to lie down. Diver SB then reported fleeting aches first in his right knee, then his left knee, then both shoulders and then steady dull aches in his right knee and then also in his left knee and he developed some neuses. Diver E0 developed a burning sensation in his perincum which oleaned repidly, then niggles in both knees and a sensation of weakness

-197-

in his left log all of which cleared during examination leaving him symptom free. It was decided to recompress both men but Diver BO spiritedly requested that he be not so treated and he was kept under observation in the chember lock with no deterioration in his condition. Diver SB was clear of all symptoms by 65 feet. so it was decided to try the effect of Therepeutic Table I from 100 feet (Shorter). 24 hours after surfacing, he reported with a mild dull ache in his left knee diagnosed as part of the residual bruised sensation which was to be expected and which would not respond to further recompression. In spite of hypnotics he had a restless night and had pain when weight bearing on that leg next morning so that his gait was affected. No abnomality could be found to locate the lesion and local heat was used es the condition aleaned over the next three days. Without doubt this case was a genuine recurrence and would probably have responded to a further compression, in face of the obvious inedequate initial compression.

<u>Diver WS and Diver RE</u> formed one of the three pairs who canned out the final dives of this stage, to 450 feet for 10 minutes at depth. Both men surfaced tixed but in excellent spirits as a result of their achievement and were permitted to move about the ship freely immediately after their dive. 15 minutes later Diver WS developed niggles in his left knee which rapidly became severe. Diver RE also developed niggles in his left knee in the same time interval but not so seriously and initially demuzzed at being recompressed with his partner. The symptoms cleaxed from Diver WS at 30 feet and from Diver NE at 45 feet and both men were decompressed from 100 feet on Therapeutic Table I with complete success - although Diver WS had a very mild niggle in his right knee 4 hours after surfacing which responded to mild analgesics and a night's rest. These cases showed the imprudence of allowing too much exercise too soon - as well as the unpredictability of the pressure required to relieve symptoms subjectively assessed in different individuals (Recommended for both men).

## (111) Third Period

(A) Laboratory Stage

<u>Diver N5 and Diver 52</u> carried out a chamber dive to 160 feet for 16 minutes. 20 minutes after reaching surface, Diver N5 developed a dull ache in his right shoulder, in both his forearns and in his right knee successively and all clearing fairly quickly. He then had a sharp pain behind his left knee which cleared with a suddon involuntary kick and one hour after reaching surface his only complaint was thredness. 2 hours later he reported a peculiar sensation in the front of his left thigh when weight bearing and an apprehensive look had reappeared on his face. 1 hour later still his left knee jerk was found to be diminished. Diver 52, at this late stage developed verses of namese and it was decided to recompress both men to 30 feet and to give them 100% oxygen for one hour at that depth and then to continue its use during 35 minutes decompression to surface. Diver 52 had recovered after the initial 30 minutes but there was no dramatic

-199-

improvement in Diver NS. At surface, the latter had a sensation of tightness in his left thigh but his reflexes were brisk. Next day, Diver RS mentioned that he had had mild niggles in his elbow which he did not report as he wished to get to his bed; he now felt "worked over" but examination showed a pyrexial illness with pheryngitic and he was treated for this in the Sick Day. These two men were puzzling in their progress and the recompression was carried out as much for the sake of doing something as for specific therapy (Shorter).

Diver MR dived to 160 feet for 16 minutes in the chamber. He had dived twice before to this depth uneventfully once on the same routine as his current dive, and the other occasion on the routine carried out by divers NS and SE. On surfacing, he reported a slight niggle in his right upper jay and his right thigh which had appeared at the 10 foot stop and which persisted till arrival at the surface where it cleared during 5 minutes. 5 minutes later still he reported that his right eye was beginning to close - external examination did not confirm this and that he had abdominal discomfort with nausea and dizziness. While the letter symptoms could be attributed to distension from swallowed ges - as had happened on previous occasions in minor degrees - the quickest way to provide relief was to recompress him and his symptoms cleared at 15 feet. He was kept at this pressure for 8 minutes and then pressure was slowly reduced over the next 28 minutes. He seemed fully recovered and examination disclosed no abnormality. He reported some days later that he had had difficulty climbing the stairs to his flat

-200--

due to a mild niggle in his knees. This latter niggle then moved to give a low back ache and inability to stamp his feet - a state that lasted about 24 hours. The pressure to relieve the abdominal symptoms probably diminished the symptoms and signs of decompression sickness and as the patient concealed such symptoms as might be present, the treatment was not as effective as it appeared (Shorter).

Diver RS carried out a chamber dive to 185 feet for 16 minutes. On surfacing he developed an itch over the left costal mangin and enother over his loft elbow. He then reported a dull ache in his right hip gradually becoming more severe and radiating to the low back area and down into both thighs. On recompression his symptoms cleared at 8 feet so a shallow therapy commenced from 20 feet where he stayed for 6 minutes and had 36 minutes decompression. On re-surfacing his left upper arm and both thighs felt weary but no abnormality was found on examination (Shorter).

<u>Diver F5</u> cerried out his first helium and his first experimental dive in a chamber to 100 feet for 16 minutes when the only decompression was the ascent at 50 feet per minute, breathing 100% oxygen. 5 minutes after surfacing, he developed a sensation of mumbress in his right thigh which increased while he walked to the therapeutic chember for examination but only a small steadily decreasing area of diminished sensation to pinprick was found over the right calf. He begen to feel that his right leg was becoming heavy and on re-examination weakness on resisted hip flexion was found, so recompression started. His symptoms cleared at

-201-

100 feet, but compression was continued to 165 feet and decompression carried out according to Therepeutic Table II (Shorter).

<u>Divor SE</u> carried out one of the first chamber dives to 340 feet for 16 minutes. On coming to 10 feet and again on coming to the surface, he developed slight niggles in his left shoulder and left knee. Although these disappeared while at 10 feet, and again at the surface, he slowly developed a rubbery feeling in his left leg with a sensation of stiffness on the lateral aspect of his left thigh. On examination, he was seen to have a rolling gait and while his only reflex disturbance was a diminished left knee jerk, he was found to have difficulty initiating straight leg raising in his left leg. Muscle power was normal. He was recompressed 1 hour after surfacing but his symptoms did not clear till he had spent 20 minutes at 165 feet. Decompression on Therapeutic Table II was successful (Shorter).

<u>Diver JN</u> dived to 340 feet for 16 minutes in the chamber on a schedule expected to be less testing than that used by Diver SE. During the stop at 70 feet he developed a prickly sensation over his right shoulder which became an ache on arrival at 50 feet where he started to breathe 100% oxygen and his ache cleared. At the 20 foot stop he developed itchiness over his left knee and at the next stop this became generalised over both thighs. 15 minutes after reaching surface he had a sensation of stiffness in both thighs and, 5 minutes later still, he complained of a burning constion in the sole of his right foot. On examination both feet were thought to be unusually cold; while his am

#### -202-

and might leg reflexes were normal, his left leg reflexes were equivocal; his pulse was 65 per minute with frequent extra-systoles. He was recompressed to 30 feet and he was given 100% oxygen to breathe. Initially, he developed a patchy pallor of both feet but after 10 minutes this cleared and was replaced by paraesthesia and his pulse became regular at 60 beats per minute. After a further period of half-an-hour his only complaint was a tingle in the plantar aspects of both helluces and examination disclosed bilaterally equal reflexes. After one hour at depth, he was decompressed over the next hour and surfaced with no complaints or abnormalities (Shorter).

<u>Diver D4</u> carried out the first chamber dive to 500 feet for 16 minutes. Due to a catarrhal condition he was under treatment with an oral anti-histomic "Eskomade" to enable him to dive. Generally his dive was unsatisfactory. While he was at the 30 foot stop he reported a slight pain in his right thigh which waned during his stay at that pressure but waxed and waned at 20 feet, at 10 feet and again at the surface. No abnormality was found on examination although he complained of residual stiffness in the thigh. Then the pain returned and gradually vorsened so recompression was started. His symptoms cleared at 30 feet; he was further compressed to 100 feet; and decompression carried out successfully on Therapeutic decompression Table I (Recommended).

<u>Diver FS</u> carried out a similar chamber dive to 500 feet for 16 minutes as that done by Diver DN. He surfaced in good spirits but 8 minutes later while having a cup of coffee and a cigarette, he suddenly

-203-

felt faint and giddy and he became very pale; his radial pulse was moderate in rate but almost imperceptible in volume. He was assisted to the therapeutic chamber where, as soon as he key down, his colour improved and he felt better - his pulse was now 74 per minute and easily perceptible at the wrist. He developed a severe ache in his left log down the whole lateral aspect so recompression was started 11 minutes after reaching surface. All his symptoms cleared at 21 feet, and as it was suspected that the syncopal attack was not due to decompression sickness, it was decided to treat the limb condition by recompression to 30 feet for 1 hour and then decompression over the next hour. At 3 feet, he reported an ache in his right knee but during the 2½ minutes the decompression stopped while the doctor was called, the ache cleared and 10 minutes later at the surface he had only a slight sensation of stiffness of his knee (Shorter).

<u>Divor MR</u> carried out a chamber dive to 500 feet for 16 minutes on a schedule presumed to be safer than that used by Diver DN and Diver FS. He developed a slight ache in his right knee at each stop from 40 feet to the surface which cleared during the stay at each stop. At the surface his symptoms progressed to flitting oches around his elbows and shoulders within half-an-hour and then cleared. 3 hours later, he suddenly felt faint and complained of a very severe pain behind his right knee and he became neuscated during his transfer to the therapeutic chamber. He admitted that he had had a persistent mild ache in the knee since surfacing but that he did not report it, in the hope he would be

-204-

allowed home but salving his conscience by reporting it as short-lived. His symptoms cleared at 100 feet on compression, and he was decompressed on Therapeutic decompression Table II (Recommended).

Divor CI carried out his first helium dive in a chamber dive to 300 feet for 16 minutes on the schedule that was considered past the development stage and was expected to be the operational routine. At the 30 foot stop, he suddenly developed pain in his right hip spreading to his knoe and rapidly becoming more severe. When the medical officer was locked through to examine him, he was found to be obviously in pain. sweating profusely, and there was slight tendemess in his right loin. No other abnormality was found. He began to improve so he was kept at the same pressure for 6 minutes longer than planned, then when it was decided to continue the original routine, he was decompressed to 20 feet. A further reduction in pressure led to aggravation of his symptoms at 18 feet so he was returned to 20 feet to continue oxygen breathing for 20 minutes. It was decided to decompress him slowly with the intention of transforring him to the large therapeutic chamber if this was indicated end again about 18 feet his symptoms deteriorated so he was rapidly transferred. On recompression, all his signs and symptoms cleared at 33 feet and, after compression to 100 feet, Therepeutic Table I was successfully used (Recommended).

<u>Diver WS</u> carried out the probable operational dive schedule to 400 feet for 16 minutes in a chamber. 3<sup>1</sup>/<sub>2</sub> hours after reaching surface, he developed - but did not report - slight twinges in his left knee which

-205-

changed into a persistent mild dull eche. 2 hours later he became eware of rapidly developing numbress and weakness of his loft arm and hand and some pain in his loft shoulder - and then he reported. He was recompressed 10 minutes later with no relief on arrival at 165 feet except for his knee ache which he had not noticed had disappeared till he was questioned. On examination he was found to have a glove encesthesic up to his elbow; he had a partial motor paralysis below his left shoulder in that if his ann was raised, intense effort did allow him to maintain the position for a very short time. After 65 minutes. there was almost negligible improvement so it was decided to compress him further to 250 feet. Between 200 and 230 feet. the pain cleared completely first then full power and movement returned and on arrival at 250 feet only a residual small patch of diminished sensation on the lateral aspect of his left forearm remained - and that cleared muckly. After a 30 minute stay at this pressure, he was decompressed to join the therapeutic tables, the decision on whother to use Therapeutic Table II or III being delayed till 30 feet when it would be taken as indicated by the diver's condition. His condition was excellent so Therapeutic Table II was selected (Special). However 14 minutes before he was due to be decompressed from 20 feet he developed an extremely severe pain in his left knee and he was immediately recompressed. When his symptoms cleared at 70 feet decompression was then transferred to Therapeutic Table IV and eventually he reached surface in a rather exhausted state (Recommended for meanrence). 3 hours later he developed

+206+
stiffness in both legs which progressed to pain over the next 2 hours and he was recompressed again. His symptoms were relieved at 155 feet while on his way to 165 feet and decompression was commenced after 30 minutes at 165 feet on Therapeutic Table IV (Recommended). On reaching 155 feet, Diver WS felt apprehensive about his legs so decompression was halted for 12 hours and he was given 20% oxygen/80% helium to breathe during this pause. As usually happens under such prolonged therapies under pressure, the diver's psychological attitude became one of reluctant co-operation and irrasoibility. At 50 feet his left leg reflexes were equivocal and his arm reflexes were bilaterally absent. He was treated by hypnotics and mild analgesios and he eventually surfaced 3 hours over 4 days after he started his original dive. As he was weak and unsteady, he was put to bed for a rest and re-examined the next day. He complained of aching in both thighs and fairly severe pain in his knees on movement. He was found to have weakness on resisted movements of both lower limbs; diminished proprince ption of both his big toes but more so on his left side where there was also diminished vibration sense; and a shuffling gait. Further recompression was rejected as suitable treatment. After one week with hydrotherapy followed by one week of active physiotherapy, he was fully recovered. This case showed the problems that can arise if a delay occurs in reporting and how a recurrence may occur in the less prominent sites even if more dramatic signs and symptoms are completely cleared.

-207-

(B) Sea Stage

Diver FZ and Diver HD carried out a sea dive to 400 feet for 16 minutes. Diver HD developed a niggling ache in his right shoulder on ascending to 40 feet which cleared at that depth; this pattern was repeated at 30 feet, 20 feet and 10 feet. It recurred on reaching surface, but this time it did not clear and was joined 12 minutes later by on ache in his right hip. Diver FZ reported an ache in his left knee which started as a niggle at the 10 feet stop and which became more severe during the first 10 minutes at the surface. It then began to improve. It was decided to recompress both men. Diver FZ was clear of all symptoms at 15 feet where the shoulder pain cleared in Diver HD but the latters' pain in his hip did not clear till 23 feet. Both men were further compressed to 100 feet and Therapeutic Table I was used successfully (Recommended).

<u>Diver CI</u> dived in the sea to 400 feet for 15 minutes on the same decompression schedule as Divers HD and FZ. On reaching surface, he complained of such severe pain in both knees that he could not move them, let alone stand up. A niggle had appeared in both knees on breaking down to 30 feet, 20 feet and 10 feet and had cleared at the first 2 of these stops. However at 10 feet it had persisted and became very severe on breaking down to surface. He was transferred to a more comfortable chamber and he was recompressed within 2 minutes of reaching surface. His symptoms cleared dramatically at 23 feet and a sensation of discomfort in his left knee disappeared at 30 feet. At his request, It was decided to use 100% oxygen as his breathing medium for 1 hour et this depth and for one more hour that it would take to decompress him to surface. This routine was successful (Shorter).

Diver CR was one of the first pair to dive in the see to 500 feet for 16 minutes. On surfacing, he complained that he had been having spasms of giddiness since leaving the 60 foot stop when he also had noticed a sensation of disorientation in space of his body. No physical abnormality could be found but his personality was subdued, lacking his normal wit and humour and with startling loss of interest in food or a cigarette. When he complained of a headeche, it was concluded that his depressed personality was due to decompression sickness and he was recompressed. At 60 fect, he recovered as if he was being unveiled. He was further compressed to 100 feet-and decompression on Therapeutic Teble I started (Shorter). As a precention it was decided to increase his tissue ovvenation by giving him 100% ovven to breathe for the lest half hour at 30 feet and 20 feet and for the last hour at 10 feet. On exemination 10 minutes after surfacing, there was diminished sensation to pipprick on the back and front of his right forefinger which improved overnight. This case emphasized the protean menifestations of decompression sidmess.

### Unnanal Case

<u>Diver M9</u> dived in the sea to 500 feet for 15 minutes. He surfaced in very good spinits which alloyed the wornies of the observers while he sat in a chair with a signative and a hot cup of coffee. During this

-209-

10 minute period of observation while at nest, he had a second eigenstte. lle and his parimor had just been told they could remove their subber suits prior to an examination and their midday meal, when Diver PS had an oplicptiform convulsion. He was immediately mushed to the compression chember and the supply valves opened. No compressed gas was delivered because permission had been given by the doctor for the supply to be diverted for a minor task. Permission had been given in view of the first-class appearance of the men. In the short time before the supply was restored. Diver FS recovered and recompression was not started pending examination. There were no symptoms and no signs other then a short duration ameria from seconds before his fit. He remained in the chamber under close attention for 4 hours and he remained well. He had had a light breakfast on hour before his dive which had lested 54 hours. Hypoglyceenie and chain smoking seemed to be precipitating factors. At a later date, a fasting electroencephalograph showed some slow activity largely within normal limits which disappeared after glucose administration. This attack and his two earlier episodes  $(q \cdot v \cdot)$ seemed to be similar and not basically due to decompression sickness although in the other two attacks he had signs and symptoms at least of minor cases. He carried out 18 experimental helium dives eltogether.

(iv) Fourth Period

<u>Diver PT</u> cauried out a chamber no-stop experimental dive to 72 feet for 2 hours. His previous dive had been of a similar duration to a depth of 51 feet, 5 weeks carlier when he had had minor

-210-

pain symptoms in his left knee. This particular dive was carried out deliberately with his full knowledge to see the effects of a layoff period without subsequent work-up as it was a dive which his colleagues had carried out without incident. 10 minutes after reaching surface, he felt some pain over his xiphistermum which worsened. -7 minutes later, when he was recompressed he was having some difficulty in breathing and he was sweating profusely. At 34 feet all his symptoms cleared, but he was found to have a slight typical trunk resh. He was kept at this depth for one hour breathing air, and he was slowly decompressed over the next hour (Shorter). One hour after surfacing, he developed magging aches deep in both knees which he bore for 3/4 of an hour before reporting. He was egain compressed with relief at 50 feet and subsequent decompression was on Therapeutic Table I (Shorter). Next day he had a sensation of stiffness in his right knee but no further treatment was required.

<u>Diver BF</u> dived in the chamber on an experimental no-stop dive to 75 feet for 2 hours. At each of the 3 preceeding dive depths he had had minor incidents. **15** minutes after surfacing, while having a shower, he felt some pain in his left shoulder which made drying himself difficult. Due to pain, there was some weakness of resisted movements of the shoulder on examination 5 minutes later. He was recompressed with relief of ell his signs and symptoms at 14 feet. He was then taken to 30 feet for one hour and decompressed over the next hour, air being breathed throughout (Shorter). Next day his shoulder felt as if

-211-

he had slighly strained a muscle.

<u>Diver HT</u> carried out an experimental no-stop dive in the chamber to 60 feet for 4 hours. At depth, he mentioned some sharp transient pains in his right knee but examination when he surfaced was negative. 15 minutes later he developed severe sharp pains in his right knee of a different character to the earlier ones. The new pains became constant and radiated up the right thigh and 30 minutes after surfacing he was recompressed. His pain was relieved at 37 feet but he was taken to 100 feet and then decompressed on Therapeutic Table I (Recommended). Next morning, an initial stiffness in his right knee on leaving his bed, cleared with walking.

<u>Diver PF</u> cannied out a similar dive to Diver HI. As on a previous occasion, he developed tightness in his chest about 10 minutes after reaching surface. This progressed to complaints of difficulty with respiration, particularly on inspiration but observation did not confirm this. He was recompressed with relief of his symptoms at 15 feet. He was taken to 30 feet for 1 hour and then decompressed over one hour with air as the breathing medium, with no after effects (Shorter).

<u>Diver BF and Diver WF</u> dived in the chamber to 300 feet for 4 hours, both men surfacing on the planned decompression schedule after a total time at high pressures of 9 hours 17 minutes. Diver BF had a slight pain in his left shoulder at this time which flitted to both ankles and within 5 minutes had settled in his left knee where it gradually became more severe. Recompression started 15 minutes after he had reached surface

-212-

end at 50 feet he was relieved of his symptoms. At this moment, Diver WF weiching the recompression, developed pain in his right knee and rapidly took on the appearance of shock, so he was locked through to the main compariment where he too recovered. Both men were then taken to 100 feet and decompression on Therapeutic Lable I started (Recommended Diver BF. Shorter WF). On arriving at the 30 foot stop. Diver BF compleined of a recurrence of the niggle in his left knoe and slight tightness across his chest and shoulders which had been constant from the first stop. This condition persisted and the doctor on duty decided to increase the pressure again, first to 40 feet, then to 50 feet and in view of the lack of response, eventually to 165 feet to carry out decompression on Therepeutic Table IV (Recommended) as his knee was now slightly better. Abnormal sounds were present in the chests of both men with scattered monohi and altered vocal responses. Eventually they reached surface where Diver DF still had pain but now in his right knee which affected his gait. Radiographic examination disclosed lower right mid zone patch of pnoumonitis in Diver WF who was clinically much better then Diver BF. Over the next few days both men tired easily on exertion. It was one month before Diver WP was radiologically clear, although he felt and was clinically completely recovered in about one week. The routine of this dive is given in Appendix 4.

<u>Divor BP</u> again dived in the chamber to 300 feet for 4 hours after an interval of 1 month. The dives were similar for the first 8 hours but thereafter the second decompression schedule was designed for a

-213-

further 8 hours instead of 12 hours thus there were more decompression stops. Diver DP first reported a slight pain in his right knee on arrivel at each stop from 65 feet but clearing while at the stop. At 25 feet the pain only eased a little during the 3 hours at this depth. On ascending to 10 feet the pain became more severe and did not alleviate much over the next hour. He then claimed that he had improved rapidly so an attempt was made to reduce the pressure but at 6 feet, his pain recurred severaly. He was recompressed with complete relief of his symptoms at 18 feet and he was taken to 30 feet with the intention of staving there for one hour end then slowly decompressing over the next hour. On arrival at 16 fost. Diver DF reported a pain in his right endle and in his right knoe and decompression was halted for a pause of 10 minutes. He also compleined of nausea and tightness of his chost on inspiration. Due to a power out affecting the lighting system, the pause lested one hour during which all his complaints cased. Decompression was then restarted with an extremely slow bleed till 8 feet was reached when there was a recurrence of his right knee pain. After a pause of 30 minutes an even slower bloed was started without telling him. This was halted at 4 feet because the petient became restless. The chamber doors at this depth are not airtight and the excess pressure was slowly vented. On examination his gait was affected by sohing in his right knee end enkle; his chest had an increase in fine moist sounds on auscultation; during the observation period at rest in a chair, he scened to be in considerable pain and it was decided to recompress him again. However

**-**214+

on walking to a bigger chamber, his pain eased and his gait improved so it was decided to let him have a hot shower while continuing the observation. His improvement continued to complete recovery. The routine of this dive is given in Appendix 4.

<u>Diver 15 and Diver FR</u> carried out a chember dive to 900 feet for 2 hours. the greatest pressure to which any man had ever been exposed for such a period of time. At depth, the men reported that food tested terrible; that they had some difficulty in moving; that sudden or modd movements were painful; and that they had impairment of control of their legs on welking. 6 minutes were taken in decompressing to the first stop at 490 feet, - well within the expected safe pressure drop. After 1 hour, Diver LS compleined of newsee and of a spinning sensation of his head and shoutly afterwards he vomited. It was decided to continue the decompression routine with Diver LS remaining flat and taking sips of water. Whenever he moved, he developed an attack of giddiness which led to some retching. At the 340 foot stop, there wes some suspicion of his hearing and shortly afterwards Diver FE reported that he was also feeling a little giddy if he moved around. It was now decided to recompress the men and they were taken back to 400 feet 5 hours and 20 minutes after the start of the dive. Diver FE was cleared of symptoms and Diver LS seemed to improve and vent into a peeceful sleep. After 2 hours, decompression to 340 feet was carried out to repeat that stop. Diver IS still had some dizziness and a tendency to retch if he moved too much. Decompression was carried out to 260 feet after 2 hours

-215-

and the exygen was enriched to 10% in the helium chamber etmosphere which was breathed for 2 hours. On decompression to 190 feet, - a drop in pressure planned in the original schedule, - both men developed nain: Divez WE had twinges in both elbows and a more persistent pain in both knees; Diver 15 developed pain in his right knee to add to his other symptoms. Both men were recompressed to 220 feet and their mains disappeared at 204 feet. After 30 minutes, both men were decompressed egain to 190 feet this time without aggrevation or reappearance of symptoms. After 90 minutes, decompression to 165 feet was carried out. Both men had a short recurrence of pain in both knees on arrival at this depth, at 140 feet, at 120 feet, at 110 feet end at 100 feet. Over 5 hours after leaving 165 feet, an attempt to reach 90 feet had to be stopped at 93 feet and both men were returned to 100 feet because of severe pain in both knees especially of Diver ME. After enother hour. 90 feet was achieved and after 2 hours 60 feet and then after a further period this time 6 hours. 70 foet was reached. His pain recurred in an axtremely severe form in Diver HE, so both men were recompressed to 165 feet with relief of pain at 100 feet for Diver FE and relief of Addiness for Diver IS. Both men graduelly became aware of a mild porsistent dull ache in their knees which cased during the decompossion to 140 feet after 2 hours at 165 feet. Decompression continued with 1 hour at 140 feet, at 120 feet, at 100 feet, at 80 feet, at 70 feet, and at 50 feet where Diver DE became restless because of disconfort in his right knee. On moving to 40 feet, this disconfort became a niggling

-216-

pain which lasted 3 minutes. At this depth, the oxygen content of the chamber simosphere was increased to 40%. After 1 hour, both men were decompressed to 30 feet and after some initial restlessness, they settled down for a stay of 6 hours at that depth. On subsequent bleed to 20 feet and again to 10 feet, where the stops were 1 hour long there was a recurrence of disconfort in the right knoe of Diver ME. On the last bleed to surface. this reappeared at 5 feet for Diver FE and Diver 19 reported a soreness in his night knee at 2 feet. On leaving the chamber Diver DI was much more exhausted then Diver LS but he was free of signs or symptoms (Fig. 17). Diver LS was found to have a lesion of the right cookles and vestibule. The cooklear lesion cleared in the course of a few days with the vestibular lesion improving more slowly. The therapeutic decompression was an empirical mixture of reducing pressure to keep the symptoms constant but minimal; of return to pressures higher then required for relief of symptoms; and of prolonged stays at pressures to equalise the saturation of the tissues at the new embient levels. The depth at which signs and symptoms appeared is one to or beyond which very few men have dived and it is remarkable that the whole episode took less than 50 hours. The routine of this dive is given in Appendix 4.

<u>Diver CR and Diver HT</u> carried out a chamber dive to 700 feet for 2 hours. The decompression schedule to be used was the one prepared for the 800 foot dive so it was expected that the initial pressure drops should be safely carried out. All went well till the men left the 190 foot

-217-

stop, 10 hours from the start of the dive and 4 hours after the oxygen had been envioled to 20% from 5%. During this decompression change, Diver HI slowly developed severe pain in both knees and Diver CR was dramatically sudden at 150 feet in reporting pain in both his knees. Doth men were recompressed, Diver HI recovering by 185 feet but Diver CR not till 260 feet was reached. Noth men were decompressed in stops of 2 hours duration and reached the surface with only en occasional reminder of pain in their knees which appeared on reaching each stop and this cleared at that stop. Both men had retrosternal pain on deep inspiration with find rales scattered throughout the lung In addition Diver OR had diminished sensation of the pads of arces . the tourinal phalanges of all digits of both hands - an abnormality which prevented the differentiation of coins. (This has been reported as occurring in prolonged oxygen breathing). These men had hed twinges from the beginning of decompression which they had not reported and their painful episodo was dealt with by a great increase in prossure and then a steady monitor of their symptoms to regulate each succeeding pressure drop. The routine of this dive is given in Amendix A.

<u>Diver PK and Diver MA</u> dived to 55 feet in a chamber for 2 hours at depth using 10% oxygen/90% helium for the dive and 100% oxygen for the direct escent to the surface at 50 feet per minute. 20 minutes after surfacing Diver MM reported with a mild ache of the flower aspect of his right forean which cleared within the next hour. Diver PK had an ache in the lateral espect of his left upper an distal to his

-218-

shoulder and reaching his albow. Over the next 2% hours it eased to a mild dull ache in the region of his left elbow. It then gradually became more severe again and in a further 3 hours developed a "bite". Recompression was effective at 7 feet, he was taken to 30 feet and a bleed decompression was carried out using air as the breathing mixture with complete success (Shorter).

Diver LC and Diver CH dived to 300 feet in a chamber for 4 hours at depth. Diver 10 reported on arrivel at the 45 feet stop that he had a severe pain in the medial side of his left knee which had been present at the provious stop of 70 feet. Both men were recompressed to 70 feet with relief of symptoms at 57 feet. The decompression continued with stops of 1 hour at 70 feet and 60 feet but the pain appeared again at 55 feet where there was a panse of 1 hour. On resuming the decompression the pain recurred at 45 foet and the chamber pressure was related to 50 feet. After 2 hours, decompression was restarted with a recurrence of symptoms at 33 foet, 22 feet, 8 feet and 2 feet there being a very slow bleed to the next point of 30 feet, 20 feet, and 8 feet where the pressure was maintained for 1 hour before proceeding. On the surface Diver LC was walking well and both men had retrostornal tightness on deep inspiration but no clinical or radiological abnormalities on exemination. This case was treated by an increase in pressure beyond that required for relief and by stops of 1 hour duration monitored by the patient's response, and changes in pressure were carried out by a slow bleed. The routine of the dive is given in Appendix 4.

-219-

#### Comments

There were many lessons learnt end relearnt in this triel. The vaniability of response enoug individuals is shown by the fact that the first see dives resulted in 1 permenent oripple, 3 men successfully treated and two men apparently unaffected. The value of keeping in reasonably regular practice and of an adequate work-up after a lay-off is appearent. The chronic effects of pure dry oxygen, so often overshadowed by the scute convulsive effects, hed to be recognized and borne in mind when considering prolonging a therapeutic contine. The points of decompression theory assumed from work done elsewhere or many years before, coupled with animal experiment results were found to be misleading. In fact many misconceptions had to be shed and the initial cuphoria became eventually a planned attack. Even then there was the surprise that extrapolation of curves and extension of tentetive theories was not streightforward in spite of an ultra conservative approach. There is some suggestion from the case histories that once decompression sickness occurred, there was no real banafit to be gained by the patient breathing oxygen enniched mintures, even pure oxygen at 30 feet, instead of a gas mixture intended to maintain the partial pressure of oxygen equivalent to normal etmospheric levels. While there is doubt whether mue oxygen or dry ges caused pulmoneny discomfort, it is better to keep the oxygen level down. The concept of using the depth of relief as the pressure required for therapy was not entirely satisfactory it being of value to add an increment to this figure. It is difficult to assess

the effect of long pauses at various depths and it is possible that shorter duration at shallower intervals might be more effective. The use of discomfort as a guide to therapy was never satisfactory but in such an empirical situation there was little choice. The fact that some cases recovered without recompression may disguise one asset of such action in that patients are more rapidly relieved of all signs and symptoms; this in turn means that one has a fit energetic young nan as a patient who remembers the subsequent boredom and informe confinement of his therapy; this then leads to a reductance to accept recompression therapy at a later date. The spectre of asoptic bone necrosis is always present in deep diving, but never more then enound cases of decompression sickness because of queries on the adequacy of any therapeutic regimen. Finally it is obvious that this type of deep diving will always require to be a major undertaking and not to be tackled by a small ill-equipped term.

-221-

## CHAPTER XII

## DISCUSSION

Inevitably several aspects have already been mentioned in the preceeding chapters. Broadly, 1t is submitted that the established compression treatment of decompression sidness of whetever origin. diving or tunnelling, commercial, military, or for sport, must be It is tempting to include evictor's decompression sickness reviewed. but there is very little experience recented as yet by men who have seen cases in both low pressure and high pressure situations. In essence the evolution of the theories leading to measures for the prevention of decommension sickness also led to measures for the treatment of decompression sidmess. The response of various species has not enabled a reliable experimental animal to be found except perhaps, in some limited especis for the gost. There is a great tendency anonast workers in this field to promote for use with man, treatment routines based on their enimal cases whether the enimals be dogs, or goats. This proferred work appears to be unusual first in the number of cases in divers, cases which were the responsibility of one individual, sometimes in collaboration with less experienced colleagues; and second in the variety of types of exposure to high pressures which resulted in such cases.

# Critorion for Discovels

It is probably apparent that there has only been one criterion required to establish a prime facto diagnosis of decompression sickness and that has been the complaint of unusual symptoms or signs from a man who had a recent history of exposure to pressures greater than atmospheric pressure. The diagnosis has been seperated into two groups depending on the duration of exposure and to some degree on the development of the case. By this is meent that cases axising after a fairly rapid return to atmospheric pressure after a short exposure to maximum pressure and which then presented symptoms of neurological disturbance. motor or sensory, were regarded from the start as "air embolism" consequenton lung demage mather than as classical decompression sickness. The classical label was considered when the cases developed after prolonged exposures to pressures with a decompression routine and were more often associated with symptoms and signs of the locomotor system. The coincidental development of both types is always possible as is the possible occurrence of one type when the other is more likely. One wonders if ceses showing lung cysts as reported in men from tunnel sites elthouch arising after long pressure exposures are not welly cases of pulmonary baratuuma rether than decompression sielness, as indeed is suggested by Campbell Golding and his colleagues (28).

#### Differential Diamosis

The question of differential disgnosis from conditions not caused by pressure changes is sometimes difficult. Hysteria used to be put forward commonly in submarine escape training cases and was difficult to refute when no radiological, clinical, or electro-encephalographic evidence of the lesion was present. In diving, muscle and joint injuries are always possible as well as malingering and, again in the absence of positive

=223+

signs, some alternative diagnosis to decompression sickness may be made. An example was given in the chapter on deep diving of a case that almost certainly had a hypoglycaemic cause but the same man on carlier occesions, (also recorded) probably had a hypoglycaenic attack es well as decompression sickness. Griffiths (67) has described a tunnel worker with classical clinical signs and symptoms of perforated peptic ulcer a short time after he had completed a shift but who on recompression recovered dramatically and remained well. As there is no evidence or expectation that patients not suffering from decompression sidness will dramatically improve on recompression while those so afflicted do improve the disgnosis can always be checked by a test of cure by recompression. This is particularly the case where the prospect of permanent disability due to decompression sickness may occur as there is no record, so far, of a person dying of enother condition when recompressed for possible decompression sideness. A case has been described (1) where a patient died while in a compression chember from multiple causes when a mistaken diagnosis of air embolion was made without a history of decrease in ambient pressure, the immediate cause of death being oxygen poisoning complicating septic abortion. Problems of Treetment

Having advanced to the point where the diagnosis is presumed because of the history and where recompression as the correct treatment is indicated directly, or after a test of cure, for the prevention of serious disability of a temporary or a permenent nature, an explanation

-224-

must be given for the fact that the majority of cases are not recompressed. To this end the circumstances of treatment must be considered. A compression chamber is a confining cell of such a size that very rarely can an occupant stand up inside it and, due to the cylindrical shape of narrow diameter, even a confortable sitting position is not always attainable (Fig. 3). Great strides have been made in comfort for example in providing mattresses so that lying down is the most acceptable activity. The patient is not "ill" and once his signs and symptoms are relieved he is once more a healthy active individual. The temperature inside the chamber varies with the season and environment there being a marked increase on compression followed by a sharp drop each time the pressure is subsequently released. The cold is aggrevated by cloud formation as the temperature drops below the dev point of the chamber stancephere (Fig. 15). In a minor way, replenishing the atmosphere also has a cooling effect on the internal conditions of the chamber.

Associated with temperature is the problem of noise. The release of gas from a higher pressure to a lower pressure is always noisy: this varies with the volume involved and the rate of release. A skilled operator can change the atmosphere with only a slight sibilant sound and, if the decompression is a bleed, at a very slow rate, he can sometimes carry out a fairly eilent progression to lower pressures. In addition, the metal construction of the chambers transmits noise to the inside, especially such scratching or tapping sounds which may be carelessly made by attendants on the chamber itself. Lighting is important so that

-225-

the patients may be observed but by switching off all but a soft light, it is possible to encourage relaxation and remove the uncomfortable sensation of being "on the spot", but still observe such important indications as restlessness or restricted movements. In any long period in a compression chamber, enmity can be generated fairly easily towards the attendants who regularly observe through the portholes - as they must - but who thus deprive the occupants of privacy even for natural functions; this emotion is aggravated by visitors, however well-intentioned.

The patient and his attendant may pass the time reading, playing board games or dozing and they can carry out normal activities such as feeding and attending to personal hygiene. The depressent effect of high pressure attributed to nitrogen - but also noticed to a lesser extent when helium is the inert gas - leads to a lack of concentration so that reading requires too much concentration and is usually abandoned. Games played are simple ones with cards or ludo but never chess or other games requiring high intellectual performance - perhaps because the men cannot play such games. Food tends to be unappetising however wellprepared and the taking of fluids has to be actively encouraged. Because of the inflammability risks of high exygen tensions, smoking in a chamber is banned and in smokers this can become a troublesome matter.

Some people may consider that no action should be taken till the man volunteers a complaint, thus avoiding a dilemma in diagnostic skill and treatment. This was not the practice, first because of the risks

-226--

of delay when examination disclosed signs of which the patient was unaware, and second, because of the experimental nature of so much of the work when the number of cases of varying severities formed the main part of the assessment of success or failure. In addition, questioning could lead to continuous observation of the man so that any deterioration was rapidly spotted. In at least two men who reported, the benefit of the doubt was not given because of personality clashes causing suspicion of "financial" decompression sickness but the case records show the error in retrospect.

#### Alternative Treatment

Alternative therapy to recompression was usually labelled "symptomatic". A deep hot bath was often the first measure in suspected or minor cases and was always welcomed as the men wanted to clean themselves after their work. It was hoped that the traumatic bubble might expand in such a situation causing aggravation of symptoms thus confirming the diagnosis; and that such heat could ease the muscle aches and strains sometimes occurring in helmeted diving. Next was more a gentle pastime of sitting with friends, eating, talking, smoking, and similar activities rather than bed reat; this routine often led to permanent as well as temporary improvement. Analgesies were not used often to avoid masking of any developing signs and symptoms but if they were used then it was usually after several hours had passed with a nuisance type of pain remaining constant. If the patient vished to sleep he was allowed to do so under observation.

-227-

## Pressure Required for Recompression

Once the decision to recompress was taken. the minimum pressure required was that where relief of symptoms and signs occurred. Sometimes, the improvement of the presenting symptoms was so marked that the patient was temporarily unaware of minor discomforts that might remain but these often led to subsequent trouble. This principle was woll established in the past in divers and in tunnel workers but it had fellen into disrepute because of suggestions that if prolonged exposure to 6 Ats.Abs. did not result in improvement the diagnosis was wrong. It is true that as a rule pressures greater than 6 Ats. Abs. are unlikely to be required except for cases that appear after dives to greater pressures but this is by no means invariable. Thus in deep diving, therapeutic chambers should at least match in maximum working pressure the pressure of the dive. It is important that the considerable improvement in the patient be due to increasing the pressure and not to pausing at a pressure. It was remarkable how often the persistence of signs or symptoms. for however short a period, at the pro-selected depth of recompression seemed to lead to trouble during the subsequent decompression. Equally in cases of submarine escape training air embolism, high initial pressures were always required even if the subsequent decompression was curtailed. In prectice an increment of pressure was usually added to the pressure of relief - not as a fixed amount but by increasing the pressure to 2 Ats.Abs., 4 Ats.Abs., 6 Ats.Abs., or 82 Ats.Abs. whichever was the next greater figure. During the decompression of deep diving cases, it seemed to be

-228

of benefit to increase the pressure again whenever the symptoms recurred or became persistent or became unbearable.

# Duration of Stey at Pressure

As well as the pressure of relief, and the increment of pressure, three points had to be considered, the stay at maximum pressure, the gas to be breathed; and any adjuvant therapy before decompression commenced. All previous work in divers and compressed air workers had suggested that a period of helf-an-hour was the minimum required at maximum pressure and this recommendation is endorsed by the experience in these various trials. This duration seemed to permit a stabilising of the patients' condition - or it was long enough for signs and symptoms to reappear. For the routines involving an increase in pressure to 2 Ats.Abs. it became standard practice to wait there for 1 hour and this seemed satisfactory.

### Breathing Cas

Air was the gas usually selected as the breathing mixture because it was the atmosphere in the chamber for most treatments and because no breathing equipment was needed to use it, in spite of the effects of nitrogen and density. In some of the experimental chamber deep dives, the chamber atmosphere contained holium and in those cases that developed during the decompression, the patients continued to breathe exygen and helium mixtures. On two occasions, when cases occurred after deep dives but which were treated with air breathing, exygen and helium was administered intermittently by mask with improvement in the morale of

-229.-

the patient. 100% exygen was used in half of the cases treated with the shallow routine as well as in some cases with prolonged therapy. In addition two men used it at atmospheric pressure in an attempt to see whether improved exygenation led to alleviation of their symptoms but in one of them temperary improvement led to a more resistant case later so that no further cases were so treated. In the cases following deep dives and needing prolonged recompression there were physical signs of lung irritation in men confirmed once only by X-ray. This effect was possibly due to prolonged inspiration of high partial pressures of exygen although successful attempts to keep these low indicated that the dryness of the gas supply might have been as important as the exygen content in producing chest symptoms. In the shallower and shorter routines, there was no difference in the success of treatment as judged by the incidence of residual mild disconforts after decompression,

whether air or 100% oxygen was used.

The impression was obtained that whereas a high oxygen content and thus a low inert gas content - in inspired gas enhanced the rate of elimination of the inert gas during a routine decompression, once the diver reported signs or symptoms this physical process was considerably slowed. This might have been due to tissue reaction to the noxious agent and the consequent local ordems postulated then led to impaired circulation and thus impaired gas exchange. It was thought that the partial pressure of oxygen in the mixtures at high pressures was adequate to maintain oxygenation without incurring the risks of using 100% oxygen.

-230-

[It was assumed - perhaps unjustifiably - that the tissue oxygen tensions would reflect the inspired levels in the saturation situation of diving] Experience tended to confirm opinions that the use of 100% exygen during the decompression of developed cases did not accelerate recovery and, because of equipment limitations, often was an added stress to the patient. With these views, it is not surprising that so few of the cases reported here were actively treated with this gap. Comparison of cases treated at some stage with pure exygen with the other cases is not of value because of small numbers.

The early deep dives suggested that air was undesirable as breathing mixture for decompression following oxygen and helium dives without an intermediate period of oxygen breathing. This opinion arose out of ideas that helium formed multiple micro bubbles whenever decompression started. bubbles which closed autokly, when the pressure steadied again unless nitrogen was breathed. The diffusion and solubility characteristics of the two inert gases are such that it is feasible that nitrogen diffuses into the helium bubble to make it more stable whereas the oxygen content decreases as it is used up in metabolism. In later dives, the treatment on air elways followed a period of oxygen decompression so that there vas no repeat of the routine used in the one case that resulted in permanent neurological demage. Thus for routine air diving, it is suggested that air is the best therapoutic breathing modium, with the oxygen and helium mixtures being kept available for intermittent administration at the wish of the patient to lighton his depression.

-231-

After deep dives, the main therapeutic breathing mixtures should be the various oxygen and helium mixtures unless 100% oxygen has already been used for a reasonable period during the decompression in which case air may be used. 100% oxygen would seem to have little place as a breathing mixture of choice in therapy as its efficiency is doubtful and its dangers under high pressures are real. In addition there is the discomfort of administration of any mixture which depends on the type of equipment used.

### Therapeutic Adjuvants

The use of other therapeutic agents at pressure, other than breathing gases and pressure has to be considered when signs and symptoms are not entirely relieved by such recompression as may be carried out. This situation may arise in one of two ways, either the treatment chamber is limited in its maximum working pressure or the treatment may be delayed deliberately or, more often, because of transport difficulties, both these latter causes producing a more resistant case. It has happened that cases have been treated in chembers where the maximum working pressure of 2 or 3 Ats.Abs. was inadequate to relieve symptoms rapidly (67. 154) but the routine used then was to keep the patient at the maximum pressure possible till eventually signs and symptoms cleared before successful decompression. The situation did not arise in the diving triels as the working limits were always adequate even if not fully used although such a situation could arise in submarine escape training. As mentioned earlier hypothermia has been used inside chambers

-232-

in cases grossly damaged by transport delays, and in the later deep diving periods, facilities to carry out this adjuvent therapy were prepared but were not required. The use of saline or dextran infusion was considered in one case but an improvement in his condition made this operation unnecessary. The use of plasma volume expanders in a most unusual case of decompression shock (31) was of great interest and shows the sort of situation where benefit can be expected. It is understood that intravenous urea to reduce corebral codema was tried in a case, which had had a long delay to reaching treatment facilities. with considerable recovery but opinion on the efficiency of this mechanism is doubted by some neurophysicians and credit is given to recompression. time. and nursing (154). It is instinctive in some doctors to gild the lily by using drugs or some other evidence of their expertise. even when recompression has been successful; the intention being presumably to encourage gas - and theoretically bubble - elimination. If there is a risk of masking of recurrence of signs and symptoms, it is better to avoid unnecessary measures in the absence of specific indications. It is worth remembering at this point, that the petient should always be asked to carry out active full movements including weight-bearing on his less at geveral points in the decompression lest the confining effect mask signs and symptoms by imposing restrictions on mobility (145). Where there are indications, then suitable drugs may be used e.g. paraldehyde in the case with epileptiform fits and codium anytal during a prolonged stay at a constant pressure on Therapoutic Tables III and IV

(especially recurring cases) thus allowing the patient to rest more effectively. Such drugs should be selected as far as possible for rapidity of action, shortness of duration, and minimal respiratory depression. Similarly, mild analgeside may be used.

In the cases described, one patient was not completely relieved by recompression in that he progressed to a paraplegia. His case was the major stimulus to investigation of therapeutic routines and later he would have been treated by much higher pressures. It would encer that he had multiple air embold in his spinel cord, relieved by the Initial pressure increase but hypoxic effects then appeared when the amboli grow again and this sequence was repeated a second time. No drugs were used and the condition did not deteriorate during decompression - although hypoxic areas might well have been rendered completely enoxic. Oxygen and helium mixtures might have been used during his therapeutic decompression to make life less miserable and perhaps increase tissue oxygen and reduce tissue carbon dioxide levels. In two cases in which it was decided to continue mutine decompression after symptoms appeared. there was a disappearance of the symptoms and a negative examination et the next lower pressure; again both were in the deep diving programe. There was no explanation for this unusual progression. When a case developed signs referable to the balancing mechanisms while at very meet depths during decompression it was decided to continue the decompression as long as the signs remained constant till more familiar pressure levels were reached using the experience of the first case as a guide

-234-

and hoping for an improvement as in the other two. Eventually the appearance of other signs at much shallower depths led to an increase in pressure to an intermediate level with improvement in the impaired sense of balance, suggesting that the hypothetical bubble had in fact been-kept fairly constant in size as planned. No other therapeutic agents were used and the patients condition remained good.

# Therepeutic Decompression Tables

As a sule the excepter the pressure required for treatment, the longer the decompression routine will take. As oxygen does not seen to have much effect in speeding up the process compared with eir or oxygen and helium mixtures, there does not appear to be enything to be gained by using potentially toxic concentrations; this means however that higher quantities of mort gas are evailable for absorption or to retard elimination. Thus one must be avaue of the gas tensions in the body es a whole as well as at the site of the lesion. These tensions may differ because of circulatory impairment locally. This may be important in deep prolonged diving where it is possible to reach lower pressure levels during a therepeutic routine quicker then intended for the original dive and usually with a longer period at much greater intermediate pressures as well. (This situation arises when the decompression is carried out using the signs and symptoms as a guide and not using the calculated tissue levels). Previous diving therapeutic tables used the stage method of decompression as against the bleed of compressed air vorkers end, for the more serious cases, the stages could be prolonged so that the

#235-

body could reach equilibrium with the inert gas partiel pressures et each stage. It is suggested that once decompression sidness occurred the assumptions on equilibration no longer hold. The occurrence of decompression sickness in previously uninjured attendents suggest that these assumptions might have been felsely besed in any case. The first departure from standard routines was to reduce the 12 hour duration of the 30 foot stop of Therapeutic Table III to the 2 hours of Therapeutic Table II although the indications for the use of the table remained. In effect, Therapeutic Table III fell into disuse. The next development occurred in the treatment of submarine escape training accidents where the vital point of adequate initial pressure exposure on recompression became obvious but, in the special pressure history, where it was found that the use of the standard diving table provided an adequate therapeutic decompression routine in selected cases. The next departure was to treat all cases, inrespective of the symptoms, on the dopth of relief and to use the appropriate tables so that cases at one time treated on Therapeutic Table III could be treated on Therapeutic Table I. After 2 drematically successful cases treated thus, there came the case of Diver MN which was not really a failure of this approach as in fact he did not start the Therapeutic Teble I decompression but was further compressed from his initial depth of relief plus increment to what would have been the recommended depth, although perhaps not as quickly as he might have been. The development then continued by instituting the routine of depth of relief and a little bit more for cases relieved at

+236-

shallow depths, followed by a bleed decompression as used in tunnel workers. There was some adjusting of times at pressure and wates of bleed before simple instructions were devised. In the same period, there were occasions when a need arose to use higher pressures than previously recommended up to  $6\frac{1}{2}$  Ats.Abs. and which was initially devised on the spot with the first case. The table, developed in emergency, very quickly reached the shellower stages of Therapoutic Table II was fairly successful on the 4 occasions it was used. For later trials tables were prepared at leisure in readiness for cases which required such pressure levels and which either responded quickly or which had a longer resolution. Neither prepared table was used. These results suggested that either Therapoutic Table II was excessively long for most occasions or that a decompression routine did not have to have longer and longer stages the lower the pressure levels reached.

The stage was now set for the really deep dives and some principles had been established as guide lines for therapy. It was decided not to prepare a complete range of therapeutic tables but the scientists were asked to produce a decompression routine to cover slips in drill, equipment failure or other error at the maximum depth which could also act as a framework for the therapy. In the result, this table was used for emergencies twice but never, not even as a framework, for therapy. The evolution of a recommended routine is still not complete but the cases in the fourth deep diving period show the following points and modifications to illustrate the principles established. Pain was used

-237--

as a guide in that a mild bruised sensation was noted but any increase in severity was an indication for recompression to depth of relief. Recompression usually occurred if any other signs or symptoms developed. The rate of decompression between stages was a slow bloed and each stage tended to be of similar duration rather than longer than the preceeding The pressure interval between steges was variable, depending on one the appearence of or change in symptoms and signs. Any subsequent increase in pressure varied from considerable to minimal and the decompression started again from whatever new level was reached. It is still a matter for trial and error whether recurrence of trouble should be treated by recompression only to depth of relief or to depth plus increment. Experience would suggest that the addition of an increment followed by a pause might allow a more confortable and a more rapid subsequent decompression than one being guided by signs and symptoms; and that the continuous "bleed" descent is an advantage compared with fixed stage intervals. The reason for the attempt to speed up the process is the four of respiratory abnormalities already discussed/

## Signs and Symptons

There has been little discussion or comment on the actiology of the signs and symptoms in this thesis. In the absence of special investigative procedures ranging from X-rays and ultra sonies to blood coagulation studies, the eyes and hands can only have a limited value. The symptoms have been mainly pain but the flitting nature both in site and timing sometimes accompanied by local signs of tenderness and swelling.

-2.78-

suggested a central focus at one time and a peripheral one at another. The localised swelling is mysterious in that the bilateral malar swelling should suggest a serious intra-oranial venous lesion belied by the condition of the patient although compatible with restricted venous return when it occurs in a limb. As a practitioner, one is forced to accept multiple gas bubbles intra-vascular and extra-vascular, central and peripheral, and get on with treating the patient so that the evidence disappears as quickly as possible.

# Double-Blind Trial

At one stage in the deep diving trials, suggestions were made that symptoms might be due to increased apprehension on the part of the divers as the depth of the chamber dives increased. Equally, it was thought that the assessment of success and the choice of treatment was influenced by knowledge of the pressure of the traumatic dive. The programme that was being prepared at that time involved dives on oxygen and helium to depths shallower than 300 feet - a possible psychological barrier as few divers have been deeper than this level. There were also intended to be several fluctuations in depth in several depth ranges so that some departure was possible from the normal progression of dives gradually getting deeper. A medical officer took part in the actual preparation of the divers so that a qualified opinion was available in the planning stages. The divers did not know the depth of their dive and the assessing medical officer, also in ignorance, remained cutside the chamber house till the final few minutes unless he was required

+239-

carlier. The system failed because the divers could estimate the time of compression at a steady rate and compare it with their own trained subjective feelings and thus be aware of their depth within narrow limits. The doctor deliberately had to exclude the estimation of depth by the total duration of the dive and, in the event, his assessments were made with a fair idea of the depth obtained by several stray clues. It was perhaps relevant that all those taking part, had misgivings from the start about the practicability of such a trial.

### Acolimatisation

This phenomenon has been noticed by medical men and others who have worked with compressed ein undertakings and has been often described in the literature. An ettack of decompression sickness may occur in a man in one of his first three shifts, or after a break such as a holiday or a strike. The man can thereafter tolerate pressure exposures thought to be more stronuous or causing attacks in never employees. In this work, the dives during the several air-breathing trials and the first three stages of the expendent trial, were carried out with a rapid frequency of diving for each individual so that he might dive daily or every second day. The last deep diving stage required that the man dived less often than once a week. No clear effect was noticed in the air dives but this possibility was one of the reasons for the recommendation that divers should remain in regular practice or should steadily work-up to a desired level of operational readiness after a lay-off. There was a strong feeling that about one man in ten diving for the first

-240-

time with helium had a likelihood of a dramatic major attack of decompression sidmess. With the numbers involved and the interruptions for leave and travel, no clear evidence for or against this impression was obtained. The analysis was complicated by the abandonment of any decompression schedule which resulted in a major case. Over the two years, the first helium dive became an indectrination dive and as the occasional case appeared, succeeding candidates carried out shallower dives of the same duration for the same decompression schedule. When a trial air routine was promulgated for use by the operational teams. certain minor changes were incorporated to take into account less accurate depth-accoping or possible errors in drill and to simplify the routine as much as possible. As these teams are in constant practice. acolimatication may be as good or better than in the trials team. Little controlled work has been done on divers in this aspect and some research is indicated to determine the optimum interval between dives and thus the appropriate size of a diving team for a particular task.

-241-

#### CHAPTER XIII

#### CONCLUSIONS AND RECOMPLETIDATIONS

Diving will always carry a risk to those who choose this activity for their sport or employment. This risk can be reduced considerably. The need for economical use of manpower on the productive and preventive phases of a dive often compromises the best advice. As retribution is not as frequent as a doctor might wish for those who ignore his recommendation, there is a tendency, to consider rules as unnecessarily restrictive till it is too late. Even the sportsman often adopts this attitude to his subsequent regret.

There is no evidence which might lead to better selection of divers on physical or psychological grounds. Volunteering is a reasonable way to become qualified in the basic skills. For experimental work, this action is vital and must be made in as full knowledge as possible of the risks and without direct or indirect pressures. The status of being a member of a select group is obviously attractive, but direct financial inducements are not necessarily helpful. One suggestion, that these should take the form of "income protection" insurance in the event of disability causing loss of carnings, has some advantages in these Hays of hire purchase and easier credit without the penalties of volunteers more interested in money than the work - and avoids the problem of assessing the magnitude of an incentive payment.

The reasons for the variability between individuals and in the case
individual from day to day are not understood at present. This means that the risk of decompression sickness, pulmonary barotrauma, or oxygen toxicity is always present in any dive. This risk may be ignored for short and shellow exposures to raised pressures but as the exposure is prolonged or is to higher pressures the risk will increase for each condition at different rates depending on the basic physiological mechanism. Greater attention may have to be paid to such factors as changes in circulatory and respiratory dynamics with immersion of the individuals. It is only recently that plans have been developed to investigate with modern techniques and equipment the physiology of man under pressure to discover any alterations from his functions at atmospheric pressure. There is a case for suspending further research on decompression and on prolonged exposures to pressures till such investigations have been made into that part of the field already explored.

 $\mathcal{Q}_L$ 

While decompression sickness occurring in standard air diving may often be satisfactorily treated by recompression under the supervision of an experienced diver, this is not always the case. When the condition arises in deep diving especially if helium is being used or the possibility of pulmonary barotreum is present then the prosence of a medically qualified individual is important and, he should be experienced in the field. Most doctors have a very vague idea on causation and almost no idea on treatment; even those with a little knowledge may misdiagnose a case; while those with several years experience have each their own records of mistakes. It is important for any diver to know the whereabouts of the nearest treatment facilities, the nearest source of skilled advice, and the quickest way to reach and obtain these services. When in doubt, any signs and symptoms arising after an exposure to higher pressures must be assumed to be due to that activity. Very few conditions are aggravated by recompression therapy - and most of these should have been diagnosed before the initial dive. In this situation the best treatment is recompression, wheever is supervising, and to keep the patient at adequate pressures till further advice can be obtained. Such action may save life or disablement and be more comfortable for the patient; if it is wrong, the penalty is a little inconvenience and wasted time. When skilled advice is present, loss degratic therepeutic routines may be instituted always with the possibility of recompression as more positive treatment if indicated initially or subsequently.

For ordinary diving, the therapeutic routines in the various diving handbooks are usually adequate. Often they may be unnecessarily prolonged, but they will suffice for the rare case occurring in ordinary supervised operations and treated by experienced lay divers. It is suggested that these routines could be augmented by another for these cases which respond to small increases in pressures. It is further suggested that in all these routines only air should be used as the standard breathing mixture for the patient. If an experienced medical officer is evailable then various departures may be possible, even in routine air diving. These departures include the use of greater pressures than these recommended in the manuale; the use of experienced helium

-21µ14-

mixtures to relieve the neroosis of nitrogen; and the use of 100% oxygen where the doctor considers the theoretical advantages outweigh possible toxic manifestations. It follows that these departures are used where the affected person has not made or sustained a complete recovery with the initial procedures.

In deep diving and submarine escape casualties. recompression therapy can be complicated and even skilled medical aid may be handicapped by pressure limitations of treatment equipment or by delay in treatment. In pulmonary barotrauma an initial adequately high pressure is the most important need; thereafter fairly rapid decompression may be carried out - as rapid as a dive decompression schedule instead of a theropeutic decompression schedule. In deep diving decompression sickness the subsequent decompression is unlikely to be straight-forward so that a detailed schedule is of little value. The need for a high initial pressure to relieve signs and symptoms is still vital but thereafter a very slow release of pressure between stages of such duration and depth as is considered wise ( and always one hour or more) will be dependent on the appearance or aggravation of further trouble. In addition, the partial pressure of oxygen has to be carefully watched. This means that other problems such as dehydration, temperature control, and carbon dioxide accumulation must also be considered as the therapontic routine might be prolonged.

Many individuals consider the treatment automatic for specified signs and symptoms and should follow "tables" - on whatever basis these

-245-

tables were devised. A plea is made that the cases are treated as human beinge; that the published tables be not regarded as the invariable right enswer; and that fundamental research is needed into the illness and its treatment as well as into its astiology. .

# REPRESIDENT

1. Anonymours	Contribution to discussion. Proceedings 2nd International Conference on Hyper- baric Ozygen, Glasgow, September 1964, Ed. I. M. Ledinghem; publisher E.& S. Livingstone Ltd. (in preparation).
2. Armstrong, N. G.	"Principles and practice of aviation medicine". Williams and Williams, Baltimore 1939, 1st edition.
3. Barthelomy, L.	"Blood coagulation and chemistry during experimental dives and the treatment of diving cooldents with heparin". Rof. no. 96, pp 46 - 56.
4. Belinko, A. R.	"Analysis of accidents occurring in training with submarine "lung" ". Nav. med. Bull. 1932, <u>30</u> : 177 = 185.
5. Behnke, A. R.	"Decompression sickness following exposure to high pressures". Ref. no. 59, pp 53 - 89.
6. Behnke, A. R.	"Decompression sickness incident to deep sea diving and high altitude ascent". Medicine, Baltimore 1945 24 : 359 - 402.
7. Bohnke, A. R.	"Effect of high pressures, prevention and treatment of compressed air illness". Mod. Clins. N. 20m. 1942, <u>26</u> : 1213 - 1237.
8. Bohnke, A. R.	"Decompression Sickness". Milit. Med.1955. 117 : 257 - 271.

- 9. Belmke, A. R. and "The use of oxygen in the treatment of compressed air illness". Nav. med. Bull. L. A. Show 1937. 35 : 61 - 73.
- 10. Bohnko, A. R. and "Gaseous nitrogon and holium elimination from the body during rost and exercise". T. L. Willmon Am. J. Physicl. 1941, 131 : 619 - 626.
- 11. Bohnko, A. R. and "Cutaneous diffusion of holium in relation to peripheral blood flow and the absorption T. L. Willmon of atmospheric nitrogen through the skin". Am. J. Physiol. 1941, 131 : 627 - 632.
- 12. Bohnko, A. R. and "Physiologic studies of helium". Nev. med. 0. D. Varborough Bull. 1938, 36 : 548 - 550.
  - "Neuropharmacologic and neurophysiologic ohangos in inert gas narcosis". Ror. no. 96. pp 209 - 225.
  - "The actiology of compressed air intoxication and inert gas narcosis". Pergamon Press, London. In pross.

"Investigation into the actiology of inert gas narcosis". Thesis for Ph.D. University of Southampton, approved 1964.

"Investigation into the actiology of inert gas narcosis" (Thesis summary). Mod.Res. Council. R. Nevel Personnel Res.Committee. Report U.P.S. 233. 1964.

"Electroencephalographic and other changes 17. Bennett, P. D. and induced by high partial pressures of A. Glass nitrogen". Electroenceph. clin.Neutophysiol. 1961, <u>13</u> : 91 - 98.

- 13. Bennett. P. B.
- 14. Bennett, P. B.
- 15. Dennett, P. B.
- 16. Bennett. P. B.

18.	Dort,	P.	"La Pression Barometrique", 1878.
			Translation by Mitchoook, M. A. and
			F. A. Hitchcook, College Book Company,
			Columbus, Ohto, 1943.

19. Bjurstedt, H. and G. Severin C. Severin

20. Bond, G. F. Personal communication, 1959.

21. Boothby, W. and"Oxygen in Aviation". J. Aviat. Med. 1938.R. Lovelace2:172 - 198.

Boycott, A. E., "The prevention of compressed air illness".
 G. C. C. Damant, and J. Hyg. Camb. 1908, <u>9</u> 342 - 443.
 J. S. Heldane

23. Boyle, R. "New pneumatical experiments about respiration". Philosoph. Trans. 1670, <u>5</u>: 2011 - 2058.

24.Brunner, F. P.,<br/>P. G. Frick and<br/>A. A. Buhlmann"Post-decompression shock due to extra-<br/>vasation of plasma". Lancet 1964, 1:<br/>1071 = 1073.

25. Buhlmann, A. A. "Deep diving" in"the Undersea Challenge" Proceedings of the Second World Congress of Underwater Activities. B. Ecton editor. British Sub-Aqua Club, London 1963, pp 52 -57.

26. Buhlmann, A. A. "Respiratory resistance with hyperbaric gas mixtures". Ref. no. 96, 98 - 107.

- 27. Burnett, W. A. "Four cases of suspected air embolism from the Submarine Escape Training Tank" (H.M.S. DOLPHIN) Med. Res.Coun. R.Neval Personnel Res.Committee. Report U.P.S. 159. 1956.
- 28. Compbell Golding, F., "Decompression Sickness during construction P. Griffiths, H. V. of the Dartford Tunnel". Br.J.Ind.Med. Hempleman, W.D.M.Paton 1960, <u>17</u> : 167 - 186. and D. W. Walder
- 29. Carpenter, F. G. "Inert Gas Nercosis". Ref. no. 64. pp 124 -130.
- 30. Cetchpole, H. R. and "Physiological factors affecting the production of gas bubbles in rabbits decompressed to altitude". J. coll. comp. Physiol. 1946, 27 : 15 26.
- 31. Cockett, A. F. K. and R. M. Nakamura
   "A new concept in the treatment of decompression sickness (Dysbarism)" Lancet 1964, 1 : 1102.
- 52. Crocker, W. E. "Survey of medical experience in free ascent training". Med. Res.Coun. R. Naval Personnel Res. Committee. Report U.P.S. 147, 1954.
- 33. Crocker, W. E. "Principle and technique of free ascent in submarine escape". Jl. R. Nav.Med.Serv. 1955 XLL: 133 - 140.
- 34. Grocker, W. E. "Investigation into Decompression Tables Report VII. Sea Trials of proposed new diving tables." Med. Res.Coun.R.Naval Personnel Res.Committee. Report R.N.P. 57/885, 1957.

35.	Crocker, W. E.	"Investigation into Recompression Tables Report IX : Revised Tables". Med. Res.Coun. R. Naval Personnol Res.Committee. Report R.N.P. 58/902, 1957.
36.	Crocker, W. D.	"Proposals for applying the new standard tables to surface decompression and combined dives". Med. Res.Coun. R. Naval Personnol Res. Committee. Report U.P.S. 175, 1958.
37.	Grocker, W. H. and H. V. Hempleman	"The decompression problems of diving to 600 feet". Med. Res.Coun. R. Maval Personnel Res. Committee. Report 57/887, 1957.
<b>3</b> 8.	Crollin, R. Q., W. B. Crocker, and H. Hollis	"Investigation into Decompression Tables Report IV : Trials of the proposed new tables". Med. Res. Coun. R. Naval Personnel Res. Committee. Report R.N.P. 54/791, 1954.
39.	Doment, G. C. C.	"Notes on the "Laurontio" salvage operations and the prevention of compressed air illness". J. Hyg. Camb. 1926, <u>XXV</u> : 26 - 49.
40 <b>.</b>	Remant, G. C. C. and B. R. Lockwood-Thomas	"A case of compressed air illness cured by recompression". Brit. Med. J. 1909, 2 : 881 - 882.
44 <b>.</b> ,	Devidson, J. K.	"Radiology in Decompression Sickness". Thesis submitted to University of Edinburgh for the Degree of Dector in Hedicine, 1964.
42.	Devlo, R. II.	"Deep Diving and Submarine Operations". St. Catherine's Press, London, 6th ed. 1955, p 130.
43.	Devic, R. H.	tbid y 156.

44.	Donald, K. W.	"Variation in the oxygen tolerance of human subjects in the wet and in the dry". Med. Res. Coun. R. Naval Personnel Res. Committee. Report R.N.P. 44/95, 1944.
45.	Donald, K. W.	"Oxygen poisoning in man". Br. Med. J. 1947 1 : 667 - 672, 712 - 717.
46.	Donald, K. W.	"Oxygen bends". J. Appl. Physicl. 1955. J : 639 - 644.
47.	Donald, K. W. W. M. Davidson and W. O. Shelford	"Submarine Eccape breathing cir". J. Hyg. Camb. 1948, <u>46</u> (2) 176 - 183.
48.	Budloy, S. P.	"Caisson Disease" in the British Encyclopaedic of Medical Fractice 1936, 2 : 730 - 736. Butterworth & Co. London.
49.	Duffner, G. J. and H. H. Snider	"Effects of exposing men to compressed air and helium-oxygen mixtures for 12 hours at pressures of 2 - 2.6 atmospheres". U.S. Navy Experimental Diving Unit Research Report 4 - 59, 1958.
50.	Duffner, G. J., J. P. Snyder, and L. L. Smith	"Adaptation of helium-oxygen to mixed gas SCUBA". U.S. Nevy Experimental Diving Unit Research Report 3 - 59, 1959.
51.	Duffner, G. J., O. E. Van dor Aue, and A. R. Bohnke	"The treatment of decompression sickness : an analysis of 113 cases". J. Ind. Hyg. Toxicol. 1947, 22 : 359 - 362.
52.	Dwyer, J. V.	"Calculation of air decompression tables". U.S. Navy Experimental Diving Unit Research Report 4 - 56, 1956.

53.	Baton, W. J.	"Survey of one hundred and ten cases of compressed air illness in goats which required therapeutic recompression". Med. Res. Coun. R. Naval Personnel Res.Committee. Report U.P.S. 183, 1958.
54•	Raton, W. J. and H. V. Hempleman	"The incidence of bends in goats after direct surfacing from reised pressures of air". Med. Res.Coun. R. Naval Personnel Res. Committee. Report R.N.P. 62/1025, 1962.
55•	Bad, E.	"The use of new equipment and helium gas in a world record dive". J. Ind. Hyg. Toxicol. 1938, <u>20</u> : 511 - 520.
56.	Erde, A.	"Experience with moderate hypothermia in the treatment of neurological signs and symptoms of decompression sickness". Ref. no. 96, pp 66 - 81.
57.	Erdman, S.	"The acute offects of caisson disease or acropathy". Am. J. Med.Sci. 1913, <u>145</u> : 526 - 542.
58.	Fulton, J. F.	"Aviation medicine in its preventive aspects". Heath Clark Lectures 1947, Oxford University Press, London, 1948.
59,	Fulton, J. F. editor	"Decompression Siekness". Sub-committee on Decompression Siekness of Committee on Aviation Medicine, Division of Medical Sciences, National Research Council, Washington. W. D. Saunders Co. Philadelphia & London, 1951.
60.	Gensh, I. and H. R. Catohpole	"Appearance and distribution of gas bubbles in rabbits decompressed to altitude". J. Cell. Comp. Physicl. 1946, <u>28</u> : 253 - 269.

-254-

61.	Gersh, I. and H. R. Catchpole	"Physical factors and pathological consequences" in Ref. no. 59, pp 165 - 181.
62.	Gersh, I., G. N. Havkinson, and E. H. Jenney	"Comparison of vescular and extravascular bubbles following decompression from high pressure atmospheres of oxygen, helium- oxygen, argon-oxygen and air. J. Cell. Comp. Physiol. 1945 26: 63 - 74.
63.	Gorsh, I. ot al	Reports 1 - 10 Research project X - 284, Navel Medical Research Institute, National Navel Medical Center, Bethesda, Maryland 1945.
64.	Goff, L. G. ed.	"Proceedings of the Underwater Physiology Symposium". National Academy of Sciences - Mational Research Council, Washington D.C. 1955.
65.	Gordon, J. O. and C. H. Heacock	"Roentgenologic demonstration of localised gas in calsson disease". J. Am. Med. Ass. 1940, <u>114.</u> : 570 - 571.
66.	Gray, J. S.	"Constitutional factors affecting suscept- ibility to decompression sickness" in Ref. no. 59, pp 182 - 191.
67.	Griffiths, P.	personal communication 1964.
68.	Narvey, E.M.	"Physical factors in bubble formation" in Ref. no. 59, pp 90 - 114.
69.	Harvey, E. M.	"Bubble formation" in Ref. no. 64, pp 53 - 60.
70.	Hargey, N. M. et al.	"Bubble formation in animals". J. Cell. Comp. Physiol. 1944, <u>24</u> : 1 - 24, 117 - 146, 257 - 290.

- 71. Heller, R., W. Meger, and von Schotter, H.
  72. Helsinki, Decleration of
  73. Helsinki, Decleration
  74. Holder Vienne 1900 quoted by Boycott, Dament, and Haldene (Ref. No. 22) interalia.
  75. Helsinki, Decleration
  76. Code of Ethics of the World Medical Association, English Translation. Br. Med. J. 1964 <u>II</u>: 177 - 180.
- 73. Hompleman, H. V. "Investigation into Decompression Tables : Report VIII : Further basic facts on Decompression Sickness". Med. Res. Coun. R. Naval Personnel Res. Committee. Report R.N.P. 50/896, 1957.
- 74. Hempleman, H. V. "The unequal rate of uptake and elimination of tissue nitrogen gas in diving procedures". Med. Res. Coun. R. Naval Personnel Res. Committee. Report R.N.P. 62/1019, 1962.
- 75. Hempleman, H. V. "Investigation into Decompression Tables Report X : An extension of the experimental findings of Report V of this series". Med. Res. Coun. R. Maval Personnel Res. Committee. Report R.N.P. 62/1020, 1962.
- 76. Hempleman, H. V. "Tissue inert gas exchange and decompression sickness". Ref. no. 96 pp 6 - 13.
- 77. Hempleman, H. V. personal communication (1965).
- 78. Hemplemen, H. V., W.E. Crooker, and H. J. Taylor
  "Investigation into Decompression Tables Report III : Part A: A new theoretical basis for the calculation of decompression tables. Part B: A new method of calculating decompression stages and the formulation of new diving tables". Med. Res. Coun. R. Naval Personnel Res. Committee. Report R.N.P. 52/ 708, 1952.

- 79. Nempleman, H. V. and "Deep Diving Trials". Med. Res. Coun. C. Trotter R. Naval Personnel Res. Committee. Report U.P.S. 225, 1963.
- 80. Hempleman, H. V. and
   C. Trotter
   Deep diving experiments at R.N.P.L. (May
   October 1963) and Sea testing (November
   December 1963). Mcd. Res. Coun. R. Naval
   Personnel Res. Committee. Report U.P.S. 242, 1964.
- 81. Hempleman, H. V. and "Deep diving experiments 1964". In C. Trotter preparation.
- 82. Henderson, Y. "Effects of altitude on aviators". Aviat. and Aeronaut. Engineering 1917, 2: 145 - 7 quoted in Ref. No. 58.
- 83. Hildebrand, J. H., "Heldum in deep diving end calsson work".
   R. R. Sayers and Neture, London, 1928, <u>121</u>: 557 8 and 591.
   W. P. Yant
- 84. Hill, L. "Caisson Sickness". Edward Arnold, London, 1912.
- 85. Hoff, H. C. "A bibliographical sourcebook of compressed air, diving and submarine medicine". Office of Naval Research and Bureau of Medicine and Surgery. Washington, Vol. I. 1948.
- 86. Hoff, E. C. and Abid. Bol. II, 1954. L. J. Greenbaum

87. Hoff, R. C. et al ibid. Vol. III in press.

88. Hoppe-Seyler, F. "Ueber den Einfluss welchen der Wechsel des Luftdruckes auf des Blut ausübt". Arch. Anet. Physiol. Lpz 1857, 24:63 - 73 quoted by Ref. No. 84 Inter alia.

89 o	Jones, R. R. J. W. Grosson, F. E. Griffith, R. R. Sayers, H. H. Schrenk, and E. Levy	"Administration of oxygen to compressed air workers using decompression: prevention of occurrence of severe compressed air illness". J. Ind. Hyg. Toxicol. 1940, 22 : 427 - 444.
90.	Keays, F. L.	"Compressed air illness with a report of 3692 cases". Puble. Cornell Univ. Med. Coll. Dept. Med. 1909, <u>2</u> : 1 - 55.
91.	Kollar, H.	"Towards the limits of the Continental Shelf", in "The Undersea Challenge". Proceedings of the Second World Congress of Underwater activities. B. Raton ed. British Sub-Aqua Club, London, 1963, pp 162 - 169.
92.	Kiessling, R. J. and G. J. Duffner	"The development of a test to determine the adequacy of decompression following a dive". U.S. Nevy Experimental Diving Unit Research Report 2 - 60, 1960.
93.	Eleosling, R. J. and C. H. Maag	"Performance impairment as a function of nitrogen narcosis". J. Appl. Psychol. 1952, 16.: 91.
94.	Klocks, R. A., G. H. Gurtner and L. D. Ferhi	"Gas transfer across the skin in man". J. Appl. Physicl. 1963, <u>18</u> : 311 - 316.
95.	Lanbert, R. J. W.	"Submarine Escape". Jl. R. Nev. Med. Serv. 1958, <u>XLIV</u> : 156 - 161.
96.	Lembortsen, C. J. and L. J. Greenbaum editors	"Droceedings Second Symposium on Underwater Physiology". National Academy of Sciences - National Research Council Publication 1181, Washington, 1963.

\_\_\_\_

97*	Lawrence, J. H. C. R. Tobias, W. R. Lyons, H. F. Helmhols and A. R. Sweeney	"A study of aero-medical problems in a Liberator bomber at high altitude". J. Aviat. Med. 1945, <u>16</u> : 286 - 310.
98.	Liebow, A. A., J. E. Stark, J. Vogel, and K. E. Schaoffer	"Intrapulmonery air trapping in submarine escape training casualties". U.S. Armed Forces Med. J. 1959 X : 265 - 289.
99.	Littleton, T.	"Effect of submarine descent". Ass. Med. J. 1855 <u>3</u> : 127 - 128.
100.	Lundin, G. and C. D. G. Lundgron	personal communication 1964.
101.	Maokay, D. E.	"Sea trials of new surface decompression routine". Med. Res. Coun. R. Naval Personnel Res. Committee. Report U.P.S.181, 1958.
102.	Maokay, D. E.	"Memorandum on combined dives". Med. Res. Coun. R. Naval Personnel Res. Committee. Report U.P.S. 193, 1960.
103.	Nackey, D. R.	"Report on medical aspects of deep diving trials 1962". Appendix I to Admiralty Experimental Diving Unit Report No. XXIX 1962.
104.	Mackay, D. R.	"Reep Diving Trials - medical aspects". Med. Res. Coun. R. Neval Personnel Res. Committee. Report U.P.S. 226, 1963.
105.	Mackay, D. E.	"Deep Diving Trials - April - December 1963 : Medical Aspects". Med. Res. Coun. R. Naval Personnel Res. Committee. Report U.P.S. 230, 1964.

- 106. Mackay, D. E. "Comments on Therapeutic Recompression". Rof. No. 96, pp 57 - 65.
- 107. Mackay, D. E. "Investigation into Decompression Tables : Report XI : Trials of Decompression Table II and comparison with two other tables". In press 1964.
- 108. Maokay, D. E. and unpublished observations. E. E. P. Barnard

109. Malhotra, M. S. and H. C. Wright "Air emboliem during decompression and its prevention". Med. Res. Coun. R. Naval Personnel Res. Committee. Report R.N.P. 60/996, 1959.

110. Malhotra, M. S. and H. C. Wright

"The effects of a raised intrapulmonary pressure on the lungs of fresh unchilled cadavers". Med. Res. Coun. R. Naval Personnel Res. Committee. Report U.P.S. 189, 1960.

111. Medical Research Council "Statement on Responsibility in Investigations on Human Subjects" in "Annual Report for 1962 - 63". Comnd. 2382, 1964.

- 112. Miles, S. "Underwater Medicine". Steples Press London, 1962, p. 161.
- 113. Miles 8. 1bid p 172.
- 114. Miles, S. ibid pp 244 247.
- 115. Miles, S. personal communication 1959.

- 116. Miles, S. "A simple presentation of the neval decompression tables". Med. Res. Coun. R. Naval Personnel Res. Committee. Report U.P.S. 186, 1959.
- 117. Miles, S. and D. E. Mackay Personnel Res.Committee.Report R.N.P. 60/ 995, 1959.
- 118. Miles, S. and"Pulmonary Barotraume". Mod. Med. Gt. Br.H. C. Wright1963, 8 : 358 66.
- 119. Ministry of Lebour
   Statutory Instrument 1958 No. 61. Factories

   and National
   "The Work in Compressed Air Special

   Service
   Regulations".
- 120. Moir, E. W. "Tunnelling by compressed air". Jl. R. Soc. Arts. 1896 - 7. <u>44</u> : 567 - 585.

121. Molumphy, G. G. "Computation of helium-oxygen decompression tables". U.S. Nevy Experimental Diving Unit Research Report 7 - 50, 1950.

122. Nime, L. F. "A physical theory of decompression sickness". Ref. No. 59 pp 192 - 222.

123. Peirano, S. H. "Submarine escape training experience.
N. J. Alvis, and
G. J. Duffner
Bunnary of 25 year period July 1929 December 1954.". U.S. Navy, Bureau of
Medicine of Medicine and Surgery, Project
NH 002 015 0803, 1955.

124. Pol, B. and J. J. Watelle Memoire sur les effects de l'uir applique au creusement des puits à bouille". Anmls. Hyg. Publ. Méd. 16g. 1354. Ser. 2.1 : 241 - 279.

- 125. Folak, B. and "Traumatic alr embolism in submarine escape H. Adams training". Nav. Mod.Bull. 1932. 30 : 177-185.
- 126. Rashbass, C. "Investigation into Decompression Tables : Report V. A consideration of basic theories of decompression sickness". Mcd. Res. Coun. R. Naval Personnel Res. Committee. Report R.N.P. 54/789, 1954.

5

- 127. Rashbass, C. "Investigation into Decompression Tables ; Neport VI : New tables". Med. Res. Coun. R. Naval Personnel Nes. Committee. Report R.N.P. 55/847, 1955.
- 128. Rashbass, C. "The unimportance of carbon dioxide as a cause of nitrogen narcosis". Med. Res. Coun. R. Naval Personnel Res. Committee. Report R.N.P. 55/854, 1955.
- 129. Reshbase, C. "The actiology of itching". Mod. Res. Coun. N. Nevel Personnel Res. Committee. Report R.N.P. 57/894. 1957.

130. Reshbass, C. and"The effect of oxygen concentration on the<br/>occurrence of decompression sickness".W. J. Eaton"Med. Res. Coun. R. Naval Personnel Res.<br/>Committee. Report R.N.P. 57/897, 1957.

- 131. Rivera, S. C. "Decompression Sickness among divers : an enalysis of 935 cases" : Milit. Med. 1964, <u>129</u> : 313 - 334.
- 132. Royal Naval Diving Manual; B.R. 1550, 1956 as amonded by amondment No. 3 A.F.O. P 109/58, 1958, Admiralty, London.
- 133. Royal Navy : The Diving Manual; B.R. 155. 1964. Admiralty, London.

-262-

134.	Sayers, R. R. and W. P. Yant	"The value of helium-oxygen atmosphere in diving and caisson operations". Ourr Res. Anesth. Analg. 1926, <u>5</u> : 427 - 138.
135.	Sayers, R. R., W. P. Yant, and J. N. Hildebrand	"Possibilities in the use of helium oxygen mixtures as a mitigation of caleson disease". Report of Investigations, Dept. of Int. R. I. No. 2670, February 1925, quoted in Bef. No. 137.
136.	Seusing, J. and H. Drubo	"The importance of hypercapnic in depth intoxication". Klin. Wschr. 1960, <u>38</u> : 1088.
137.	Shilling, C. W.	"Compressed-air illness", Way Med. Bull. 1938, 36 : 9 - 17, 235 - 259.
158.	Slark, A. G.	"Treatment of 137 cases of Decompression Sickness". Jl. R. Nev. Med. Serv. 1964, L : 219 - 225.
139-	Smith, A. H.	"The effects of high atmospheric pressure including the caisson disease". Eagle Print, Brooklyn 1873, quoted by Hoff in Ref. No. 85.
140.	Smith, J. L.	"The pathological effects due to increase of exygen tension in the air breathed". J. Physicl. 1899, <u>24</u> : 19 - 35.
14 <b>4</b> *	Snydor, J. P.	"Dive Reaction Scale Study". U.S. Navy Experimental Diving Unit Research Report 5 - 58, 1958.
142.	Snydor, J. P. and G. J. Duffnor	"A methodological test of resistance of divers to decompression". U.S. Navy Experimental Diving Unit Research Report 4 - 59, 1958.

143. Swindle, P. F. "Occlusion of blood vessels by agglutinated red colls mainly as seen in tadpoles and very young kangaroos". Am. J. Physiol. 1937, <u>120</u>: 59.

- 144. Triger -, mentioned in ennotation Annls. Publ. M6d. 16g. 1845, 33 : 643.
- 145. United States Nevy Diving Menual. NevShips 250 538, Washington, 1958.
- 44.6. Van der Aue O. E., R. J. Koller and E. S. Brinton
   "The effect of exercise during decompression from increased barometric pressures on the incidence of decompression sickness in man". U.S. Navy Experimental Diving Unit Research Report 8 - 49. 1949.
- 11.7.Van der Aue, O. E."Physiologic factors underlying the preventionW. A. White,and treatment of decompression sickness".R. Hayter,U.S. Navy Experimental Diving Unit ResearchE. S. Brinton,Project X 44,3 Report No. 1, 1945.
- 148. Walder, D. N. "A possible explanation for some cases of severe decompression sickness in compressed air workers" in "The Regulation of Human Respiration" edited by Cunningham D.T.C. and D. B. Lloyd. Blackwell Scientific Fublications Oxford, 1963. pp 570 - 2.

A. R. Behnke

- 149. Webster, A. P. "Some theoretical aspects of the use of multiple gas mixtures for doep-sea diving" in Ref. No. 64 pp 79 - 83.
- 150. Wise, D. A. "Constitutional Factors in decompression sickness". U.S. Nevy Experimental Diving Unit Research Rport 2 - 63, 1963.

-264--

151. Woodward, C. M. "A history of the St. Louis Bridge". G. I. Jones & Co. St. Louis 1881, quoted by Hoff in Ref. No. 85.

152. Workman, R. D. "Calculation of air saturation decompression table". U.S. Navy Experimental Diving Unit Research Report 11 - 57, 1957.

153. Workman, R. D. personal communication 1963.

154. Workman, R. D. personal communication 1964.

- 155. Workman, R. D.,
   G. F. Bond and
   W. F. Mazzone
   "Prolonged exposure of animals to pressurised normal and synthetic atmospheres". U.S.
   Newy Submarine Base Medical Research Laboratory Report 374, 1962.
- 156. Workman, R. D. and personal communication 1964. M. Goodman

157. Yarboxough, O. D. and "The treatment of compressed air illness A. R. Behnke utilising oxygen". J. Ind. Hyg. Toxicol. 1939, 21 : 213 - 218.

158. Xerborough, O. D. and "Report on a case of decompression sidness W. Welhem following an oxygen dive". Nav. Med.Bull. 1945, 2 : 607 - 609.

159. Zettezstrom, A. "Deep-see diving with synthetic gas mixtures". Milit. Surg. 1948, 103 : 104 - 106.

160. Zuntz, N.
"Zur pathogenese und therapue der durch rasche Luftdruck underungen erzengten Krankheiten". Forschr. d. Med. 1897, <u>15</u>: 632 - 639, quoted by Shilling in Ref. No. 137.

-265-

#### ACKNONLEDCELENTS

This work depended first and foremost on the 164 officers and men who organised and carried out the dives and who suffered the attacks of decompression sickness. Without their co-operation and encouragement the various trials and therapeutic experiments would not have been possible. In particular, on the executive side, special thanks are due to Commander J. R. Carr, O.B.E., R.N. and Commander E. C. Haunen, R.N. successive Superintendents of Diving, Admiralty Experimental Diving Unit, for their tolerance and support during my tenure of the post of Unit Medical Adviser; and to Licutenant Commander G. M. H. Drummond, R.N., one-time Commanding Officer of H.M.S. Reclaim and later Deputy Superintendent of Diving for his encouragement and confidence during actual trials, particularly the deep diving phases.

The groundwork was laid by the Scientific staff of the Royel Nevel Physiological Laboratory, Gosport with whom many hours of discussion and argument at all times of the day and night helped to plant and develop ideas so that any original thought expressed was the product of several minds rather than any one individual. Special thanks are due to Mr. H. V. Hempleman, M.A., Frincipal Scientific Officer, Royal Neval Scientific Service for many years of happy collaboration and to Surgeon Licutemant Commander E. E. P. Barnard, R.N., my colleague and successor, for his help, most appreciated during the very long dives and therapies.

Surgeon Lieutenant G. Wray, R.N. aboard H.M.S. Reclaim end

Surgeon Lieutenant M. H. Parsons, R.N. of H.M.S. Dolphin were outstanding amongst the several medical officers with whom I had the pleasure of working on actual cases. The assessment, treatment and result of the cases and the outcome of the trials came under the scrutiny of the Underwater Physiology Sub-committee of the Royal Naval Personnel Research Committee - a joint body of the Admiralty and the Medical Research Committee.

Professor K. W. Donald, D.S.C., M.A., M.D., D.Sc., F.R.C.P., Cheirman and Professor W. D. M. Paton, D.M., F.R.S. and Surgeon Captain S. Miles, M.D., Ch.B., M.Sc., D.T.M.& H., R.N. were especially holpful and understanding while the Secretary,

Mr. F. E. Smith smoothed out problems in a most admirable fashion.

On more personal aspects, my thanks are gratefully rendered to Surgeon Lieutenant Commander A. G. Slark, Royal New Zealand Navy for his purposeful prodding, and to Surgeon Lieutenant Commander R. J. J. Gray, Royal Australian Nevy for his encouragement, to record the experiences and trace the pattern of development of ideas that I have endeavoured to show. Surgeon Captain J. Glass, O.B.E., R.N., Assistant to the Medical Director General for Research, maintained the pressure and I am grateful to him for his efforts. Mrs. M. Holloway has coped in a most remarkable fashion with the many soribbled pages and I appreciate the effort she has put into the production of the typesoript.

Finally, I wish to thank Sungeon Vice Admiral Sir Derek Steele-Perkins, C.B., C.V.O., Q.H.S., F.R.C.S. (Eng.), F.R.A.C.S., D.L.O., Medical Director General (Navy) for his permission to use naval modical records for this thesis.

### -267-

### LIST OF FIGURES

Figure	1.	Stendard 2-compartment compression chamber.
	2.	3-compartment chember for experimental work.
	3.	Interior of main compartment of standard compression
		chamber.
	4.	Breathing equipment and noving machine inside a chamber.
	5.	Submersible compression chamber over side of H.M.S.Reclaim.
	6.	a)Diagram of transfor-under-pressure system.
		b)Submersible compression chamber during transfer operation.
	7+	Diver dressed in Standard Equipment.
	8.	Mon dressed in Surface Demend Diving Equipment.
	9.	H.M.S. Reclaim at diving station.
	10.	Diving Flat H.M.S. Reclaim.
	41.	Resh in decompression sickness a. Back view.
		b. Front view.
	12.	Facial ocdema in decompression sickness.
		a)after treatment.
		b)2 weeks later.
	13.	Diegram of Submarine Escape Training Tower and of air lock.
	14.	Submarine escaper in submarine escape immersion equipment.
	15.	Below the devpoint during decompression.
	16.	Sketch of possible use of dives from submerisible
		compression chamber.
	17.	After the deepest "bend" ever.





# Fig. 2.

3-Compartment Compression Chamber for Experimental Work at the Royal Naval Physiological Laboratory





Chamber shown in Fig. 2.



A.E.D.U.

# Pig. 5.

Submersible Compression Chamber suspended opposite access to diving flat H.M.S. Reclaim



1000 FT TRANSFER UNDER PRESSURE SYSTEM

ć

-273-



Fig. 6b.

Submersible Compression Chamber being lowered on to Compression Chamber aboard H.M.S. Reclaim





Diver dressed in Standard Equipment





Fig. 9.

H.M.S. Reclaim at Diving Station with Submersible Compression Chamber hoisted outboard and Diving Door open






Fig. 13. Submarine Escape Training Tank at H.M.S. Dolphin

and an an ample of a loss

A. . . . . . . . .

-281-



Fig. 14.

Submarine escaper showing life jacket fully inflated with escape valve to permit expansion of gas during ascent; and exposure suit inflated as it would be after arrival at surface.



R.N.P.L.

## Fig. 15.

Below the dew point

Diver FE wrapped in blanket during decompression to keep warm: Note mist in the chamber



Fig. 16.

Sketch of possible use of divers operating from Submersible Compression Chamber



R.N.P.L.

## Fig. 17.

After the deepest "bend" ever Diver LS follows Diver FE out of the Compression Chamber at the end of compression therapy

## -286-

### APPENDIX I

Statement by Surgeon Commander D. E. Hackey with reference to paragraph 2 (a) of Appendix A to U.P.S. 213 dated 12th October 1962. (for Med. Res. Coun. R. Naval Personnel Research Committe Underwater Physiology Sub-committee)

## 1. Signs and Symptoms of Decompression Sickness

The problem is best described in the following quotation: "Decompression sickness manifests itself in a variety of symptoms each ranging from mild to severe. The evaluation of these or any symptom is subjective by nature. The measure is influenced by the subject's description and the examiner's observation and evaluation of the ailment ... Severe symptoms and signs of decompression sickness must be considered relatively objective and valid indices of decompression adequacy. However, both objectivity and validity decrease with the decrease in symptom severity so that the transient pain, the dult muscle ache, etc. is almost always open to question". (Kiessling & Wood) (1).

Behnke (2) has described the nescent gas bubble as replacing the spirochaete as the "great imitator" and this property of the bubble causes difficulty in accurate diagnosis when symptoms are fairly mild in character and no signs are present or can be detected. If signs and symptoms are present in combination then the total picture may be more serious than any single item might indicate and the following list of symptoms and signs should be considered with this point in mind.

## Tiching

The significance of this symptom is controversial. However it is definitely minor in character, rarely lasting more than 15 minutes after a dive (which is usually one in a dry chamber) and no treatment is needed though a hot shower reduces the duration.

## Rash

This blotchy lesion distributed mainly on the upper trunk is a fairly serious sign when allied to other troubles (other than itching). Again no specific treatment is indicated for rash alone - although recompression leads to its disappearance - as it fades fairly quickly, and even faster if a hot bath is taken.

## Fabiane

The degree of severity of this symptom is important and only an undue tiredness unexplained by diving conditions and state of training should be considered as a form of decompression slokness. On its own, treatment is rest under observation and rarely does recompression have any place in treatment of fatigue.

## Poin

This is the most common symptom and requires experience both of decompression sickness and the individual diver to assess its importance, from the transient negging pain of 10 - 15 minutes duration, through the dull constant type completely cleared by aspirin tablets, to the worsening ache which may or may not cause secondary signs such as collapse. The differential diagnosis from strained muscles or other injuries is not easy in inexperienced men and local heat may help in sorting out cases. In the more severe cases therapeutic recompression is essential and as in other cases any deterioration in the patient's condition, or increase in severity of the pain is a clear indication for recompression.

Pain is commonly located in the limbs in the region of joints and may radiate or even migrate and may be either single or multiple in location. Pain elsewhere such as low back ache, or pain in the perineum, abdominal pain or colie due to distension is unusual and so is regarded with more suspicion in the assessment of a case although recompression is not mandatopy. In the case of abdominal pain, recompression is perhaps the quickest way of relieving symptoms due to intestinal gas but this is not a reason for enforcing a long therapeutic recompression. Great care has to be taken to diagnose the severe pain in the segmental distribution at the level of the lesion sometimes met in cases of paralysis; immediate recompression is the vital step in treatment.

## Panaesthesia

Like pain, this symptom may be present in minor or major form.

The sensation of hot-and-cold must be closely watched. I have usually treated such cases by recompression without waiting for any worsening but where there has been the possibility of pressure damage to a nerve - such as excessive tight cuff on the radial nerve - then I have doubted the diagnosis of decompression sickness.

## Central Nervous System

All lesions are serious when they involve cerebral structures, the spinal cord, cranial nerves, and the roots or peripheral nerves. While recompression as soon as possible is the normal treatment, it is not invariable. For example visual disturbance which improves during examination may be treated by a policy of waiting as long as improvement is occurring. Behaviour abnormalities are more difficult to detect but they also may respond to treatment, even 40 hours later. Vestibular damage is a less rare manifestation of decompression sickness and must always be recompressed.

## Cender-respiratory Systems

The problem here is that known as "chokes" where substernal distress and difficulty in breathing may lead to asphysia or the "shock" syndrome. Some cases have occurred of tightness of the chest on respiration which have not needed recompression. Some cases closely resemble the straightforward vaso-vagal attack and respond to simple measures. As a rule this group of symptoms and signs however should be recompressed rapidly.

There may be other symptoms or signs in decompression sickness but the guide would always be the condition of the patient after his dive. If he had little trouble, rest and constant skilled observation would be adequate; if his symptoms deteniorate or cause him distress or anxiety then recompression is the best treatment.

## 2. <u>Criteria for Treatment</u>

a) If there is doubt as to the diagnosis:

The routine should be to keep the patient under close observation at rest or else gentle exercise and measures such as local heat or baths could be used to determine the effect on the symptoms and signs. Any worsening of old symptoms or the appearance of new symptoms or signs would indicate that the patient should be assumed to have decompression sickness. The history of the dive and diver should be rechecked for possible injury, wrong gas etc.

- b) If the diagnosis is substantiated or assumed:
  - (i) Gases with minor symptoms only e.g. itching, rash,
     niggles and other symptoms not causing the patient any
     distress, these cases would be treated by close
     observation and local measures such as heat and massage
     and perhaps mild analgesics. If the symptoms worsen or

change then the case would be negarded as serious. (ii) Cases with serious symptoms such as show involvement of the central nervous system or the candic-respiratory system, these cases would usually be treated by necompression on the therapeutic tables published in the diving manual. If this were not adequate then consideration would be given to increasing the pressure even to the maximum the recompression chamber will take; to changing the gas mixture being breathed from air to onyhelium, pure oxygen, or even oxygen enriched air; or to a combination of greater pressure and different breathing mixture.

REFERENCES 1. Kieseling, R. J. and W. B. Wood, "The Development of a test to determine the adequacy of decompression following a dive Phase II". U.S. Navy Experimental Diving Unit, Research Report 3 - 61, 1961.

Behnke, A. R. "Decompression Sickness". Milit. Med.
 1955, <u>117</u> : 257 - 271.

# APPENDIN II

## THERAPEUTIC TABLES

## SECTION T

# Extract from: U.S. Nevy Diving Manual NevShips 250 - 538 January 1958 "General Principles of Diving"

## Notes of Recompression

Explanation: All references to TABLES indicate parts of Table 1 - 21 "Treatment of Decompression Sickness and Air Embolism".

- 1. GENERAL CONSIDERATIONS
  - a. FOLLOW TREATMENT TABLES (table 1 21) accurately.
  - b. Pennit no shortening or other alteration of tables except on advice of trained diving medical officer or in extreme emergency.
- 2. RATE OF DESCENT IN CHAMBER
  - a. Normal rate is 25 feet per minute.
  - b. Serious symptoms: repid descent is desirable.
  - c. If pain increases on descent: stop, resume at a rate tolerated by patient.
- 3. TREATMENT DEFTI
  - a. Co to full depth indicated by table required.

b. Do not go beyond 165 feet except on decision of medical officer.

- A. EXAMINATION OF PATIENT (see article 1.6.2. (14)).
  - a. If no serious symptoms are evident and pain is not severe, examine thoroughly before treatment.

- b. If any serious symptom is noted, do not delay descent for examination or for determining depth of relief.
- c. In "pain only" cases where relief is reported before reaching 66 feet, make sure it is complete before deciding on TABLE 1.
- d. On reaching maximum depth of treatment, examine as completely as possible to detect.
  - 1) Incomplete relief
  - 2) Any symptoms overlooked
  - NOTE: At the very least, have patient stand and walk length of chamber.
- e. Recheck before leaving bottom.
- Ask patient how he feels before and after coming to each stop and periodically during long stops.
- g. Do not let patient sleep through changes of depth or for more than an hour at a time at any stop. (Symptoms can develop or recur during sleep).
- h. Recheck patient before leaving last stop.
- 5. PAPIENT GEPTING WORSE
  - a. Never continue bringing a patient up if his condition is vorsening.
  - b. Treat as a recurrence during treatment (see 6).
  - c. Consider use of helium-oxygen as breathing medium for patient, (see 8).

- 6. RECURRENCE OF SYMPTOMS
  - a. <u>During</u> treatments
    - Take patient to depth of relief (but nover to less then 30 feet; and not deeper than 165 feet except on decision of medical officer) (If recurrence involves serious symptom not previously present, take patient to 165 feet).
    - 2) Complete the treatment according to TABLE 4.
  - b. <u>Bollowing</u> treatment:
    - 1) Recompress to depth giving relief.
    - 2) If depth of relief is less then 30 feet,
      - a) Take to 30 feet.
      - b) Decompress from 30-foot stop according to TABLE 3.
    - 3) If relief occurs deepor than 30 feet,
      - a) Keep patient at depth of relief for 30 minutes.
      - b) Complete remaining stops of TABLE 3.

NOTE. - If original treatment was on FABLE 3, USE FABLE 4.

4) Examine carefully to be sure no serious symptom is present. If the original treatment was on TABLE 1 or TABLE 2, appearance of a serious symptom requires full treatment on TABLE 3 or TABLE 4.

## 7. USE OF OXYGEN

- a) Use oxygen wherever pennitted by tables unless:
  - 1) Patient has not had oxygen tolerance test, or
  - 2) Is known to tolerate oxygen poorly.

- b. Be sure mask fits snugly.
- c. Take all precentions against fire.
- d. Tend carefully, being alert for symptoms of oxygen poisoning such as
  - 1) Twitching
  - 2) Dizziness
  - 3) Neusee
  - 4) Bluering of vision
- e. Know what to do in event of convulsion. Have mouth-bit evailable.
- f. If symptoms appear, remove mask at once.
- g. If oxygen breathing must be interrupted -
  - 1) on TABLE 1, proceed on TABLE 1-A.
  - 2) on TABLE 2, proceed on TABLE 2-A.
  - 3) on TABLE 3, continue on TABLE 3 using air.
- h. At medical officer's discretion, onygen breathing may be resumed at 40-foot stop. If this is done, complete treatment as follows:
  - Resuming from TABLE 1-A: breathe oxygen: at 40 feet for 30 minutes at 30 feet for 1 hour
  - 2) Resuming from PABLE 2-A: breathe oxygen: at 40 feet for 30 minutes
    - at 30 feet for 2 hours

- 3) In both cases, then surface in 5 minutes still breathing oxygon.
- 4) Resuming from TABLE 3: breathe oxygon:
  at 40 foot for 30 minutes
  at 30 foot for first hour
  (then finish treatment with air)
- 8. USE OF HELTUM-OXYGEN
  - a. Helium-oxygen mixtures (xatio about 80:20) can be used
     <u>instead of air</u> (not in place of oxygen) in all types of treatment and at any depth.
  - b. Use of holium-oxygen is especially desirable in any patient who
    - Has serious symptoms that fail to clear within a short time at 165 feet.
    - 2) Has recurrence or otherwise becomes worse at any stage of treatment.
    - 3) Has any difficulty in breathing.



# Table 1-21. Treatment of Decompression Sickness and Air Embolism.

		Y					
Stops		Bends-Pain only				Serious Symptoms	
Rate of de- scent-25 ft. per min. Rate of as- cent-1 minute between stops.		Pain relieve less than Use table 1 not availa	<ul> <li>elieved at depths han 66 ft.</li> <li>ble 1-A if O<sub>2</sub> is vailable.</li> <li>If pain does not improve within 30 min. at 165 ft. the case is probably not bends. Decompress on table 2 or 2-A.</li> </ul>		<ul> <li>Serious symptoms include any o of the following:</li> <li>1. Unconsciousness.</li> <li>2. Convulsions.</li> <li>3. Weakness or inability to use arms or legs.</li> <li>4. Air embolism.</li> <li>5. Any visual disturbances.</li> <li>6. Dizziness.</li> <li>7. Loss of speech or hearing.</li> <li>8. Severe shortness of breath c chokes.</li> <li>9. Bends occurring while still under pressure.</li> </ul>		
						Symptoms re- lieved within 30 minutes at 165 ft. Use table 3	Symptoms not re lieved within 3 minutes at 165 ft. Use table 4
Pounds	Feet	Table 1	Table 1-A	Table 2	Table 2-A	Table 3	Table 4
73.4	165	•••••		30 (air)	30 (air)	30 (air)	30 to 120 (air
62.3	140			12 (air)	12 (air)	12 (air)	30 (air)
53.4	120			12 (air)	12 (air)	12 (air)	30 (air)
44.5	100	30 (air)	30 (air)	12 (air)	12 (air)	12 ( <b>a</b> ir)	30 (air)
35.6	80	12 (air)	12 (air)	12 (air)	12 (air)	12 (air)	30 (air)
26.7	60	30 (O <sub>2</sub> )	30 (air)	30 (O <sub>2</sub> )	30 (air)	30 (O <sub>2</sub> ) or ( <b>air</b> )	6 hrs. (air)
22.3	50	30 (O <sub>2</sub> )	30 (air)	30 (O <sub>2</sub> )	30 (air)	30 (O <sub>2</sub> ) or (air)	6 hrs. (air)
17.8	40	30 (O <sub>2</sub> )	30 (air)	30 (0 <sub>2</sub> )	30 (air)	30 (O <sub>2</sub> ) or (air)	6 hrs. (air)
13.4	30		60 (air)	60 (O <sub>2</sub> )	2 hrs. (air)	12 hrs. (air)	First 11 hrs. (a Then 1 hr. (O or (air)
8.9	20	5 (O <sub>2</sub> )	60 (air)		2 hrs. (air)	2 hrs. (air)	First 1 hr. (air Then 1 hr. (O <sub>2</sub> or (air)
4.5	10		2 hrs. (air)	5 (O <sub>2</sub> )	4 hrs. (air)	2 hrs. (air)	First 1 hr. (air Then 1 hr. (O or (air)
Surface		1	1 min. (air)	]	1 min. (air)	1 min. (air)	1 min. (O <sub>2</sub> )
		1	L	I	J	1	

Time at all stops in minutes unless otherwise indicated.

`~\_ ~

| i

#### SECTION II

## Retract from: R.N. Diving Manual, 1956, B.R. 1550

"0314. Procedure for treatment of Decompression Illness.

It has been found in practice that if a diver gets decompression illness and is recompressed to enable any nitrogen bubbles to redissolve, his subsequent decompression must be at a rate slower than the normal or there is every chance that the bubbles will re-form. This longer decompression is known as THERAPEOPIC DECOMPRESSION.

(1) PROCEDURE FOR REHERAPEUTTC DECOMPRESSION

- (a) The patient should be instantly placed in a Recompression Chambor and the pressure should be run up at a rate of 25 feet per minute.
- (b) The pressure at which pain is relieved should be noted and from this figure the correct Therapeutic Decompression Table to use can be discovered. The pressure should then be allowed to mount to the maximum figure indicated by the correct table, and decompression should be carried out according to the prescribed stages.
- (c) The diver should be treated in this way even if the symptoms are only suggestive of Decompression Illness. Recompression can do no hasm and it may prevent serious results".

(2) The Medical Officer should examine the patient as soon as possible and satisfy himself that the correct table is being used. He should not hesitate to change from Table 1 or 2 to Table 3 or 4 if he suspects that bubble formation has caused a lesion of the Central Nervous System. 0315. RECORRENCE OF SYMPTOMS

If symptoms return DURING treatment, recompress to dopth of relief but never to a depth of less than 30 ft., and complete decompression from this depth according to Table 4.

If symptoms return FOLLOWING treatment, recompress the divor to a depth giving relief.

- (a) If relief occurs at a depth of loss than 30 ft., take the diver to 30 ft. and decompress from the 30 ft. stop according to Table 3.
- (b) If relief occurs deeper than 30 ft., remain at the depth of relief for 30 minutes and then complete remaining stops of Table 3 using air throughout.

It is highly improbable that the attendant as a result of being under pressure with the patient, will himself develop decompression illness. Should however, the symptoms be such that the Medical Officer decides to place the attendant under pressure within 12 hours of arriving at surface his subsequent decompression must be on the Therapeutic Table. Cramp is commonly experienced after a long period in the confined space of a recompression chamber.

0316. USE OF OXYCIN

Oxygen should be used when indicated by the tables, if available. The use of oxygen should not be allowed to shorten the total therapeutic decompression time. Oxygen should not be administered unless a Medical Officer or qualified Diving Officer is in attendence.

If diziness, names, muscular twitching or blurring of vision occur while breathing oxygen, remove monthpiece and complete the rest of the treatment on air.

"BEN Pain reliev at depths LESS than 66 ft.		"BEND"-F	AIN ONLY	SERIOUS SYMPTOMS		
		Pain relieved at depths LESS than 66 ft.	Pain relieved at depths GREATER than 66 ft. If pain does not im- prove within 30 mins. at 165 ft. the case is	<ol> <li>Spinal "bends"-weakness or numbness of limbs.</li> <li>"Staggers"-staggering, vomiting, dizziness, vertigo nystagmus.</li> <li>"Chokes"-pain in chest with shortness of breath, cyanosis, collapse, uncon- sciousness.</li> </ol>		
			probably not bends.	Symptoms RELIEVED with- in 30 mins. at 165 ft.	Symptoms NOT RELIEVED within 30 mins. at 165 ft.	
STO.	PS	TABLE 1	TABLE 2	TABLE 3	TABLE 4	
Lb.	Ft.		Time in minutes wise indicated	unless other-		
73•4	165		30 (Air)	30 (Air)	30-120 (Air)	
62.3	140		12 (Air)	12 (Air)	30 (Air)	
53•4	120		12 (Air)	12 (Air)	30 (Air)	
44•5	100	30 (Air)	12 (Air)	12 (Air)	30 (Air)	
35.6	80	12 (Air)	12 (Air)	12 (Air)	30 (Air)	
26.7	60	30 (Air)	30 (Air)	30 (Air)	6 hrs. (Air)	
22.3	50	30 (Air)	30 (Air)	30 (Air)	6 hrs. (Air)	
17.8	40	30 (Air)	30 (Air)	30 (Air)	6 hrs. (Air)	
13.4	30	60 (Air)	2 hrs. (Air)	12 hrs. (Air)	First 11 hrs. (Air). Then 1 hr. (O <sub>2</sub> or Air)	
8.9	20	60 (Air)	2 hrs. (Air)	2 hrs. (Air)	First 1 hr. (Air). Then 1 hr. (O <sub>2</sub> or Air)	
4•5	10	2 hrs.(0 <sub>2</sub> or Air)	2 hrs.(0 <sub>2</sub> or Air)	2 hrs. (0 <sub>2</sub> or Air)	First 1 hr. (Air). Then 1 hr. (O <sub>2</sub> or Air)	
SURFACE		1 minute (O <sub>2</sub> or Air)	1 minute (O <sub>2</sub> or Air)	1 minute (O <sub>2</sub> or Air)	1 minute (O <sub>2</sub> or Air)	

#### SECTION III

Extract from: R.N. Diving Manual, 1964, B.R. 1550

"0313. THERAPEUPIC DECOMPRESSION - TABLE 5

1. Any diver who has decompression sickness or even symptoms not directly attributable to some other cause is to be treated strictly in accordence with the instructions in this article except on the advice of a medical officer or in extreme emergency. Recompression is most unlikely to do harm and it may prevent serious consequences.

2. If a Modical Officer is available and no serious symptoms are present, the patient is to be thoroughly examined before entering the recompression chamber. If, however, no Medical Officer is available or if serious symptoms are present, the patient is to be immediately placed in the chamber, with an attendant whenever possible, and the chamber pressurised as follows:

(a) IF THE SYMPTOMS ARE PAIN ONLY. The chamber is to be pressurised at a normal steady rate of 25 ft. per min. until the patient's symptoms are relieved, subject to a maximum recompression of 165 ft. which must not be exceeded except on the advice of a Medical Officer. The pressure at which rolief is complete is to be noted and the chamber pressure increased at the same rate to the depth of the first stop given in Part A or Part B of Table 5 according to whether relief is experienced at a depth of less than 66 ft. or at a depth of 66 ft. or more. If pain is not relieved within 30 min. at 165 ft. (Part B),

the case is probably not 'bends'.

- (b) IF THE SYMPTONS ARE SERIOUS:
  - (1) 'Spinal bends'. Weakness or numbress of the limbs.
  - (ii) 'Staggers' Staggering, vomiting, dizziness, vertigo,
     nystagmis.
  - (111) \*Chokes\*. Pain in chest with shortness of breath, cyanosis, collapso, unconsciousness.

For such symptoms the chamber is to be pressurised to the maximum depth of 165 ft. at the highest rate that can be tolerated by the patient and the patient then decompressed receiving the stops given in Part C or Part D of Weble 5 according to whether or not the symptoms are relieved within 30 min. at this depth. If pain increases on descent, the descent must be halted and then continued at a rate acceptable to the patient. The maximum depth of 165 ft. is not to be exceeded except on the advice of a Medical Officer.

3. Rate of Ascent and Timing Stops. The time for the ascent between all successive stops is to be 5 min. The time for the first stop is to commence on leaving maximum pressure and the time for each subsequent stop is to commence when the diver leaves the preceeding stop.

4. Treatment in the Chamber. The Medical Officer is to examine the patient as soon as possible and satisfy himself that the correct table is in use. He must not hesitate to change from one table to another if he suspects that bubble formation has caused a lesion in the Central Nervous System. While in the chamber, the following checks are to be applied by the Medical Officer or by the attendent if no Medical Officer is present:

- (a) On reaching the maximum depth of treatment the patient is to be examined as completely as possible to detect incomplete relief and any symptoms previously overlooked. The patient must, at least, be stood up and made to walk the length of the chamber.
- (b) A reacheck as in (a) is to be made before commencing the ascent.
- (c) The patient is to be asked how he feels before and after arrival at each stop and periodically during long stops.
- (d) The patient must not be allowed to sleep through changes in depth or for more than an hour at any time during a stop (symptoms can recur during sleep).
- (c) The patient is to be re-checked as in (a) before leaving the last stop. If the patient's condition vorsens during the ascent, the ascent must be stopped and the patient treated as for a recurrence of symptoms (para. 5). The use of a heliumoxygen mixture as a breathing medium (para. 7) should be considered.
- 5. Recurrence of Symptoms. If symptoms returns
  - (a) DURING TREATMENT. The patient is to be recompressed to the depth at which relief of the symptoms occurs but never to a depth less than 30 ft. or to a depth greater than 165 ft.

except on the advice of a Medical Officer. Decompression is then to be carried out on Part D of Table 5.

- (b) FOLLOWING TREATMENT. The patient is to be recompressed to a depth giving wellef and:
  - (1) If relief occurs at a depth LESS than 30 ft. Take the patient to 30 ft. and decompress from the 30 ft. stop in accordance with Part C of Table 5.
  - (ii) If relief occurs at a depth GREATER than 30 ft. Remain at the depth of relief for 30 min. and then complete the remaining stops of Part C, Table 5, or, if the original treatment was on Part C, then Part D is to be used.

6. Use of Oxygen. Oxygen should be used as indicated in the Tables whenever it is available but must not be administered unless a Medical Officer or Qualified Diving Officer is in attendance.

If dizziness, nauses, muscular twitching or blurring of vision occurs while breathing oxygen, the mouthpiece is to be removed and the rest of the treatment given breathing air.

7. Use of Helium-Oxygen. When available, helium-oxygen mixtures (ratio about 80:20) can be used instead of air (not in place of oxygen) in all types of treatment and at any depth but only in the presence of a medical officer. The use of helium-oxygen is especially desirable in any patient who:

(a) has sorious symptoms which fail to clear within a short time

at 165 ft.

- (b) has recurrence, or otherwise becomes worse at any stage of treatment.
- (c) has difficulty in breathing.

8. Attendant - Decompression Sickness. It is highly improbable that the attendant will develop decompression sickness as a result of being under pressure with the patient, but should this occur and the symptoms be such that the Medical Officer decides to place the attendant under pressure within 12 hours of arriving at surface, his subsequent decompression must be therapeutic. It should be noted that eramp is commonly experienced after a long period in the confined space of a recompression chember. Note. Part B. If pain is not relieved within 30 min. at 165 ft., the case is probably not 'Bends'.

A		E			
Part D Symptoms NOP re- Lioved within 30 min. at 165 ft.	Part C Symptoms RELIEVED within 30 min. at 165 ft.	Part B Pain relieved at depths GHEATER than 66 ft. (See Note)	Part 1 Pain relieved at depths LISS than 66 ft.	Pepth of Stops (ft.) Chamber Pressure 10./sq.in.	
30 to 2 hrs.	elr. So	et in S		165 73•4	SPOPPAGE
35 eir	17 min. eir	eir		140 62•3	S III H
adr.	17 11. 21.	84. 17		120 53-4	ours and
35 air	17 nin. eir	ein.	30 eir	# 100 5.00	D MINUTE
23 24 24	eir.	217 217	17 211. 212	9.55 39.68	S AT DI
o hrs.	35 min. air	ein.	alr.	60 26•7	I ZMBRAN
6 hrs.	35 ein. eir	35 ein.	e i s	22•3 22•3	DEFINS BRI
6 hrs. air	35 11. 12.		air. Sir	40 17.8	ATHING AL
First 11 hrs. air Then 1 hr. air or 02	12 hrs. air	2 hrs. air	60 sir	30 13.4	COR OXYGEN
First 1 hr. air Then 1 hr. 21r or 02	2 hrs. air	2 hrs. air	adn. Str	20 8.9	(0 <sub>2</sub> )
First 1 hr. air Then 1 hr. air or 02	2 hrs. air or 02	2 hrs. air or <sup>0</sup> 2	2 hrs. air or O <sub>2</sub>	10 4•5	
1 min. air or 02	1 min. air or <sup>0</sup> 2	1 min. air or <sup>0</sup> 2	1 min. air or 02	Surface	
					T

# TABLE V

THERAPENTIC DECORPRESSION

-307-

## SECTION IV

## Thorapeutic Tables devised during trials

A. Shallow Therapy Table

## Table X

## Assumptions

- 1. Symptoms and signs were relieved rapidly and before 30 feet was reached.
- 2. The symptoms end signs were minimal in severity.
- 3. 100% oxygon is available and can be cesily given.
- 4. The patient does not object to breathing from breathing apparatus.
  <u>Routine</u>
  - 1. Compress to 30 feet.
  - 2. Patient breathes 100% oxygen or air for 1 hour.
  - 3. Decompress the chamber at 1 foot per minute to 20 feet.
  - Decompress the chamber at 1 foot per 2 minutes from 20 feet to 10 feet.
  - 5. Decompress the chamber at 1 foot per 3 minutes from 10 feet to atmospheric pressure.

## B. Long Deep Therapy Table

## Table Y

## Therapeutic Recompression for exceptional cases of decompression Mickness

## Assumptions

1. The diver has been to 165 feet for 30 minutes and there has been no appreciable improvement in his condition, and it is considered that

further recompression to a deeper depth would be beneficial.

- 2. No depth loss than 265 feet is worthwhile as the next pressure to be used. At 165 feet the bubble is reduced to 1/6th its initial size and at 265 feet to 1/9th its initial size.
- The diver will not remain longer than 30 minutes at the new depth.
   The U.S.N. system of therapy is reasonably satisfactory and the same ideas are capable of extension to 265 feet.

## NOTES

- (1) As ½ hour has been spent at 165 feet and ½ hour has been spent at 265 feet, together with 1 hour (approx.) at depths in excess of 165 feet during the decompression, it is clear that join-ing up with Table 50 may well be inadequate.
- (ii) Table 5D is itself not adequate and has been known to give the attendant an attack of decompression sickness.
- (iii) In a recent case, residual troubles were still present after the treatment. These may have been due to tissue damage and therefore unavoidable by any decompression routine, or they may display an inadequacy in the decompression. As there is doubt, and in view of (4) and (ii) above it is clear that some additional safety must be added to Table 5D.
  - (iv) Experience with therapy at 250 feet has, of necessity, been limited, but recurrences are too frequent. Attempts to join up with the accepted therapy tables don't appear to succeed.
    (v) In view of the narcotizing effect of air at 265 feet and the

possibility of errors of judgement occurring both in the diver and attendant, it is especially beneficial to breathe an oxy-helium (20/80) mixture whilst at depths in excess of 165 feet.

- (vi) In view of the density of air at 265 feet and the possibility of breathing difficulties it is also beneficial to breathe oxy-helium gas, whilst at depths in excess of 165 feet.
- (vii) Unless the diver is in a state of complete exhaustion the breathing of oxygen at the lower depths cannot be other than beneficial, provided always that there is no risk of oxygen poisoning.
- (viii) Prolonged breathing of dense air at depths of 60 feet and over can cause distress. It is desirable to breathe oxy-helium as much as possible.

## THE SCHEDULE

- 1. Raise pressure from 165 feet to 265 feet at 25 feet per minute.
- 2. Commence breathing oxy-helium mixture (20/80) immediately upon reaching 265 feet and continue oxy-helium breathing until arrival at 140 feet. Change to air.

12

20

20

30

40

30 minutes

3.	Rate of ascent,	1 minuto	between stops.
	DEFDH		TIME
	265 fee <b>t</b>	30 m <b>in</b> u	
	020		49

2,50 200

170

140

120

100 1 Hour

- 80 2 Hours
- 6 Hours 60
- 50 \* 8 Hours
- 40 \* 10 Hours
- 12 Hours 30 \*
- 20 \* 2 Hours
- 10 \* 2 Hours

Total Time 45 Hours 46 Minutes.

OXYGEN BREATHING at the end of each stop as follows:-

- 50 feet last 10 minutes
- 40 feet last 20 minutes

30 feet lest 40 minutes

20 feet last 60 minutes

10 feet last 60 minutes

 $\mathcal{X}$ 

C. <u>Short Deep Therapy Table</u> <u>Teble Z</u> <u>Shorter alternative version for decompression from 265 fect.</u>

Assumptions

- 1. The diver has been taken to 165 feet for a short period (less than 10 minutes).
- 2. Symptoms (or signs) were not relieved at 165 feet.
- 3. No more than 30 minutes is to be spent at 265 feet when it must be assumed that all troubles have ceased.
- 4. The diver's initial 'bends' were not accompanied by chokes or C.N.S. involvement - this is the U.S.N. classification i.e. pain or serious symptoms.

Schedule based on U.S.N. table 2-A which it must be realised is <u>NOT</u> the same as R.N. Table 5B.

DEFINI	TIME
265	30
230	12
200	12
170	12
140	12
120	30
100	30
80	30
60	30

DEPTH	<b><u>921</u></b> Me
50	60
40	90
30	120
20	180
10	240

Total Decompression time 15 hours 28 minutes.

## NOTES

- (4) Table 5B has 9 hours 14 minutes decompression time and Table 50 has 19 hours 44 minutes decompression time. As may be seen this table is intermediate but more towards Table 50. This tendency towards extra time could be checked if the R.N. Table 5B principles were adopted instead of the U.S.N. Table 2A, which it must be noted is 10 hours 30 minutes in duration.
- (11) It is advisable to breathe oxy-helium (20/80) at 265 feet, and all stops deeper then 140 feet.

#### APPENDIX III

SHORF THERAPETFIC RECOMPRESSION FOR S.E.T.T. ACCIDENTS Statement by Surgeon Lieutenant Commander D. E. Mockey, 7th Merch 1962 and 4th Decomber 1962 to Command Escape Officer, Flag Officer Submarines and his medical advisors.

Some confusion has arisen both about the reasons that I had for introducing a shortened form of therapeutic recompression, for the few accidents that occur during submarine escape training; and about the routine used.

2. I arrived at the training tank with some experience of decompression sickness and its treatment. I was aware that there was a recurrence rate; that treatment was unpleasant in so far as sleeping, feeding, and hygiene were concerned; and that there were residual troubles such as exhaustion and sensetions of bruising.

3. Cases at the tank were quite different in their symptomatology to the majority of cases of decompression sickness as one would expect theoretically; in fact they resembled either the very rare severe cases obviously involving the central nervous system or they had signs in the chest or under the skin (or of course various combinations of these signs) which are not seen in decompression sickness.

4. The causation of all these cases is assumed to be the presence of a bubble in the appropriate areas to produce the relevant signs and symptoms. However, one must try and find the source of the bubbles as treatment is aimed at making the bubbles regress and disappear. It is accepted that bubbles may arise in situ in a tissue which becomes supersaturated with a gas to such an extent that bubble formation may occur (there is a degree of supersaturation of the order of double saturation which is allowable for all tissues - some may have more and other variations may occur but this doubling factor is a cornerstone in that bubble formation will only occur if supersaturation exceeds this amount). It is also accepted that gas may be rapidly introduced to a tissue through the surrounding membrane (usually alveolar) and once through, it will take up the form of any gas so introduced to a fluid i.e. a bubble.

5. For decompression sickness two things are important - the time of exposure to the gas and the ambient pressure: these determine the amount of gas absorbed and hence the degree of asturation of the tissues. Previous work has shown that the risk of decompression sickness after 20 minutes at 100 feet (i.e. 4 atmospheres absolute) is infinitesimal, in fact 30 minutes is accepted by many as quite safe: i.e. the degree of saturation of the tissues is such that a person can return to atmospheric pressure without much worry about bubble formation in situ. True, there is the question of speed of return to atmospheric pressure but consideration of circulation times and volumes of blood through the lungs would show that the difference in quantities released in 20 seconds or 100 seconds is not of such great significance as to affect tissue saturation materially.

-315-
6. In the case of the introduction of gas as bubbles through a lining membrane, the duration of exposure is immaterial as it is thought that the cause is a local area of gas causing distension of the containing membrane and either rupturing it or otherwise passing through it and so forming bubbles in the tissues (regarding blood as a tissue). The vital points have are the rate of expansion and route of escape of the gas.

7. In training for submarine escape, the duration of exposure for the man with the longest exposure is of the order of 12 = 15 minutes and the maximum emblent pressure is equivalent to 100 feet of sea water. Thus the likelihood of decompression sickness occurring is extremely remote. As the rate of escent may be up to 7.5 feet per second (or as slow as 2.2 feet per second) the expansion of gas in the lungs may be great enough to cause alveolar rupture and any of the main forms of pulmonary barotrauma (air embolism) pneumotherax, or intervititial emphysema) may occur. It is always possible that a case may occur where decompression sickness and pulmonary barotrauma develop concurrently.

8. As has been said before, treatment aims at regression of the bubble. Some forms of pulmonary barotrauma need more than this simple aim as pneumothorax and interstitial emphysema do not reabsorb rapidly and other means of treatment may be needed. Air embolism however responds on the theory that Boyle's Law applies and if enough pressure is used the volume of the bubble will decrease till either the bubble continues on in the circulation and is trapped in the lungs or else the bubble tends to

-316-

dissolve in the blood stream - in either case the presumed blockage is relieved. This routine of shrinking the bubble is the idea behind treatment in decompression sickness also, but here one is starting with tissues supersaturated with the same gases and as such the resolution of the bubble is bound to be slower for comparable pressures; subsequent release of pressure must make certain that such supersaturation is not increased.

9. At this point, let me restate the position. There are two serious conditions due to a decrease in ambient pressures, which have different mechanisms for producing the same result. I considered that it was justified for the general benefit of the patient to apply two types of treatment depending on my interpretation of the symptoms as a clue to the mechanism involved. There are several other benefits but these are in the form of bonuses and in my opinion should never influence the treatment used.

10. The standard method of treatment is to regard the patient as suffering from decompression sickness and treat him on the appropriate therapeutic table i.e. 19 hours 57 minutes for one table, or a minimum of 37 hours 33 minutes on the other. My experience with the method has been detailed earlier, in particular with regard to the condition of the patient at the ond of such memocuyres.

11. Recompression to 6 atmospheres absolute would reduce the volume of the bubble to about 1/6th of its size at the surface. As this pressure is 2 atmospheres in excess of any exposure in the S.M.T.T., it is assumed

-317-

that the gas is encouraged to go into solution in the tissues. The speed of such an action would depend a great deal on the amount of gas already present in the tissues. In the case of decompression sickness one would anticipate a slow rate but as this possibility is most unlikely, (under the training conditions) in the case of air embolism a fast rate can be expected. In fact, I presumed that a rate similar to one in a dive to those depths would be reasonable, and hence adequate decompression for such a dive would suffice. As time at maximum pressure is important, 30 minutes at least should be spent at 465 feet so that there is a reasonable period for absorption of the bubble.

12. Therefore I advised a shorter therapeutic regime for cases which I considered to be due to pulmonary barotrauma. In the cases with interstitial emphysema, the deciding point was the appearance of subontaneous bubbles in the neck as this usually relieved modiastinal signs and symptoms; if they did not appear then long decompression was indicated to prevent cardiac embarrassment. Those cases with pneumothorax varied whether there was a possibility of a tension pneumothorax or not; if present then paracentesis thoracis enabled fairly rapid decompression; if it was not a tension type then decompression depended on the effects of the pneumothorax acting as a guide. In the third bread type, air embolism, again the short routine was used unless it became obvious that a local lesion persisted as shown by the continued presence of any sighs or symptoms by the time of arrival at 165 feet in which case the full routine was used. (A clinical observation is that

-318-

the original injury very rarely persists i.e. the lesion is solf scaling on most occasions).

13. The procedure entailed automatic recompression to 165 feet where the patient was examined. (There did not seem to be any purpose in stopping at a shallower depth as treatment would always involve a descent to 165 feet). Depending on the results of examination, advice was given on the use of a full therapeutic table on the requirements for a shorter technique. If the latter was indicated, the R.N. Diving Manual was consulted in the schedule for 170 feet for a duration equal to the sum of a) the time the man had been under pressure that day; b) the time from his last exposure to reaching 165 feet on treatment; and o) an addition of 30 minutes. This would entail a decompression of at least 2 hours and probably nearer 3 hours.

14. During the shorter decompression, the patient was kept under close observation by a doctor who re-examined him at every stop. Any recurrence of symptoms was a definite indication for a full therapeutic routine. If there was no recurrence, the patient would be kept under observation for two hours and then sent to R.N. Hospital, Haslar, for X-rays of chest and B.N.G. and admission for observation. It should be emphasized that this routine is only applicable to cases whose symptoms are completely cleared on recompression to 165 feet, as any persistance would indicate that the bubble had not disappeared.

15. Following some experience of such methods, it is fair to say that the victims left the chamber relatively fresh, without any residual eches, and frequently only a little suboutaneous emphysema of the neck to prove that pulmonary barotrauma had occurred. The bonuses consisted of (1) the good effect on the morale of other trainces; (2) the reduction in requirements of the staff; and (3) the minimal interruption of training; but again, the patient's needs have complete priority over any advantage to others. It is essential that constant medical observation is kept, that recurrence of symptoms - usually of neurological origin - be treated on a long table without delay; and that a period of observation at the tank be maintained post-treatment.

## APPENDIX IV

-321-

## SOME INTERESTING THERAPEDERO-ROUFINES

A. Special Routino 1.

Divers BF and WF.

Chamber dive

Deroise - rowing mechine.

Position		Time	Notes	
left	surface	00,00,00		
Arrived	300 feet	00.07.35	Broathing 10% oxygon/90% helium	
loft	300 feet	04+00+00	Ascend 2½ min. Change to 20% oxygen/ 80% helium on arrival at first stop.	
loft	170 foot	05-03-00	Ascend 1% min. Change to 40% oxygen/ 60% helium on arrival at next stop.	
Loft	100 foot	08.05.00	Ascend 2 min. Change to 100% oxygen on arrival at next stop.	
left	40 feet	09.16.00		
Arrived	eurícoe	09-17-07	BF complained of niggle L.Shoulder.	
At	surface	09.22.00	Niggle moved to both ankles and then sottled in L. knee.	
left	guríaco	09•33•00	Therapeutic recompression started.	
Arrived	50 Loot	09.34.00	MF sign and symptom free. WF at surface suddenly had severe pain R.Knee and looked in shock. Recompressed and looked through to Main compartment to join BF at 50 feet. Recovered.	

on		Time
50	fest	09.35.00
100	feet	09.36.00

10.05.00

10.23.00

10.58.00

11.33.00

12.08.00

12.13.00

13.26.00

Position

100 foet

80 feet

60 feet

50 feet

40 feet

30 feet

30 feet

Loft

Loft

Loft

Left

Left

Loft

Left

Arrivod

Arrived

Notes

Decided to use Therapeutic Table I.

DF complained of slight niggle in L.knee on arrival. He also reported slight tightness across chest present since first therapeutic stop. Examination - no abnormality either man. Niggle persisted - recompression decided to case pain.

••••			
Arrived	40 feet	13.28.00	
loft	40 feet	13.41.00	
Arrived	50 feet	13.43.09	
Loft	50 feet	13.56.00	No change in pain or chest compleint.
Arrivod	165 feet	14.01.00	Knee recovered. Chest no change. Decided to use Therapeutic Table IV. Diver WF - continue as attendant.
lost	165 feet	14.31.00	
Loct	140 feet	15.06.00	
loft	<b>1</b> 20 foot	15.41.00	
Left	<b>1</b> 00 feet	16.16.00	

-322-

P	<u>osition</u>	<b>Time</b>	Notes
Loft	80 feet	16.51.00	a constant for head ready ( and
Loft	60 feet	22.51.00	Examined 23.20.00. BF - altered voice sounds in chost on auscultation. WF - abnormalities. Both men irritative cough.
loft	50 feet	28.51.00	
<b>Left</b>	40 foot	34.51.00	Examined 46.00.00. DF - chest clear, but still complained of tightness on very deep inspiration. WF - southered rhonchi but no complaint. Cough persists in both men.
Left	30 feet	46.51.00	
Loft	20 feet	48.51.00	
Left	10 feet	50.51.00	
Arrived	surface	50.56.00	
Doors of	ened	50,57.00	

-323

After a short period of observation, both men had full plate X-rays. BF - no abnormality. WF - "small area pneumonitis present lower aspect R.M.Z. peripheral occlusion left costo - phrenic angle not present ... (6 weeks earlier)". Continued under direct observation and seemed recovered. Permitted to return to their own messes 6 hours after surfacing. After 4 days, BF stated that later in the day of his surfacing, he had had a pain in his <u>right</u> knee which had affected his gait but which eased with several hot baths - he had in fact warned his friends to be ready to take him to the chamber but he had refused to report to any responsible individual; over the next three days he had had only a rapid tiring on exertion. (He had been examined daily). WF had no complaints and felt fully recovered but his X-ray plate did not show resolution of the shadow till 5 weeks later - without any treatment.

в.	Speciel R	outine 2			
	Dviers BF	and LB.		Chamber dive	
				Exercise - roving machine	
	Position		<u>IN me</u>	Notes	
Left	BUR	laco	00.00.00		
Arrivo	ad 300	feot	00.05.36	Breathing 10% oxygon 90% holium.	
loft	300	feet	02++00+00	Ascend 2% min. then change to 20% oxygen 80% holium.	
loft	170	feet	06.00.00	Ascend 15 min. Change to 40% oxygon 60% helium at 07.30.00.	
loft	100	feet	00,00,00	Accond 1 min.	
Loft	65	îeot	10.00.00	Ascend 1 min. Change to 100% oxygen at 11.45.00.	
Loft	40	foet	12.00.00	Ascend 1 min. Change to 40% oxygen 60% helium at 12.06.00.Then to 100% oxygen at 14.30.00. At 12.08.00 <u>Divor DF</u> reported slight transient niggle in right knee.	
Loft	25	foct	15.00.00	Ascend 1 min. Diver BF reported ache in right knee. (Diver 18 continued dive routine) Diver BF examined, no abnormality found.	
Loft	10	foot	16.12.00	Slow decompression started.	

-32),--

• •

Post	Position		Notes
Arrivod	6 foet	16.12+.00	Pain unbearable, recompressed.
Arrived	18 feet	16.19.15	Recovered.
Loft	30 feet	17.20.15	
Arrived	15 foot	17.42.00	"Bleed" halted as mild recurrence of pain in right knee, development of pain in right ankle, nausea and tight- ness of chest on respiration.
Loft	15 Loot	18.50.00	All gradually eased. Slow bleed.
Arrivod	8 feet	20.17.00	Recurrence of ache in right knee. Diced halted till eased.
Left	8 feet	20.50.00	Mlood restarted.
Arrived	<b>4 S</b> eet	21.15.00	Restless. Pressure could not be maintained and "bleed" to surface. Exact time not known.
Opened doo	rs	22.06.00	Examined, gait affected; basal râles;
At .	SULLEO	22 <b>, 30,</b> 00	seemed in pain. Reported transient pains on arrival at every stop from 170 feet. Decided to recompress but on walking to chamber all signs and symptoms (except rales) cleared completely.

6.	Special	Routine 3		
Divers MD and LS.				Chambor dive
				Exercise - rowing machine
	Positio	<u>ù</u>	<u>Time</u>	Notes
Loft	BI	urfeoo	00.00.00	
Arriv	ed 8	teet ·	00.23.15	Breathing 5% oxygen 95% helium.
Loft	8	00 feet	02-00-00	
Arriv	ed. 49	90 feet	02.06.12	02.58. Diver 18 nanses and vertigo then vomited several times in next hour after moving.
loft	4	90 foot	04+00+00	Condition unchanged 04.34 deafness
Arriv	ed 3	0 fect	04.03.00	suspected 04.40. Diver FE complained of giddiness on mising and later vomited.
Loft	ÿ	teet 04	05.20.00	Recompressed.
Arriv	od 4(	00 fect	05.21.20	ME recovered. IS improved steadily and went to sleep. Before accent both seemed recovered.
Left	240	00 Loot	07.22.00	
Arriv	od. 3	+0 200t	07.24.00	
Loft	3	40 feet	09.22.00	
Arriv	od 21	60 foet	09.23.30	Ohanged to 10% oxygen 90% helitum at 09.33.30.
Left	2	60 feet	11.33.00	

•

Post	ltion	<u>Timo</u>	Noten	
<b>Arriv</b> ed	190 feet	11.34.24	Change to 20% oxygen 80% helium IE 11.35 reported twinges both elbows, both knees; LS pain in right knee. Both shivering. Pain in LS persisted. Recompressed.	
Lott	190 Leet	11.50.00		
Arrived	220 feet	11.51.07	IS rolieved at 204 feet.	
Loft	220 feet	12.21.00		
Arrivod	190 foot	12.26.00		
Loft	190 feet	13.56.00		
Arrived	165 foot	14.01.00	W tringes both knees.	
Loît	165 feet	14.31.00		
Arrived	140 feet	1436.00	Both divers pain in knees. 15.32 both divers pain free. IS still has dizziness on rapid movement of head.	
Loft	140 feet	16.36.00		
Arrived	120 feet	16.42.00	Both men twinges in both knees, which cleared rapidly.	
Loft	120 feet	17.41.00		
Arrivod	110 feet	17.47.00	Both men twinges in both knees which cleared rapidly.	
Left	110 feet	18.46.00	• •	
Arrived	100 feet	18.51.30	Similar optoodo. 1905 LS vomited	
Loft	100 feet	19.51.00		
Arrivod	93 foot	19.54.00	Twinges recurred and became severe.	

-328-

E	osition		Timo	Notes
Arrivod	. 100	foot	19.56.00	
Loft	100	feet	20.56.00	
Arrived	. 90	feet	21.01.30	Twinges again present and cleared in about 1 hour.
Loft	90	feet	23.01.00	
Arrived	. 80	feet	23.06.00	Similar opicodo. Examined 26.00.00.
Loft	80	feet	29.06.00	
Amived	70	feet	29.16.00	Recurrence of severe pain. Recompressed
Arrived	165	foot	29.21.55	Relieved at 100 feet.
Loft	165	feet	31.22.00	
Arrived	. 140	feet	31.27.00	
Loft	140	foot	32.27.00	
Arrivod	120	feet	32.32.00	Both divers well except for slight dizziness in IS.
Loft	120	feet	33.32.00	
Arrived	100	feet	33-37-00	
loft	100	foot	34-37-00	
Arrived	. 80	feet	34.42.00	ME suspicious of right knee.
Loft	80	feet	35.42.00	
Arrived	. 70	icet	35.47.00	
Left	70	feet	36.47.00	
Arrived	60	feet	36.52.00	
loft	60	feet	37.52.00	

Position		Time	Notes
Arrived	50 feet	37-57-00	ME still restless after each reduction of pressure.
Loft	50 feot	38.57.00	
Arrivod	40 feet	39.02.00	FE reported niggle right knee which cleared in 3 minutes. Changed to 40% oxygen 60% helium.
loft	40 feet	40.02.00	
Arrivod	30 feet	40.08.00	Both men trouble with knees - FE in right, LS in left but cleared rapidly.
left	30 feet	46.08.00	
Arrived	20 feet	46.30.30	TE conscious of his right knee.
loft	20 feet	47.28.00	
Arrived	10 feet	47.49.00	
Loft	10 feet	48.48.00	
Arrived	surface	49.11.00	
Doors oper	boi	49.21.00	Both had niggles. IS unsteady gait.

.

Both had niggles. IS unsteady gait. Examined by ENT. specialist - Right cochlear and vestibular damage.

-329-

D.	Special Routine 4				
	Divers C	R and HI.		Chamber dive	
				Exercise - roving machine	
	Position		Time	Notes	
Left	su	rface	00.00.00		
Arrive	ed 70	) feet	00.17.14	Breathing 5% oxygen/95% helium	
loft	70	) feet	02.00.00.		
Arrive	od 48	) feet	02.04.24.		
Left	48	) feet	04.00.00.		
Arrive	od 350	) feet	04++02-36		
Left	35	) feet	06.00.00		
Arrivo	a 260	) feet	06.01.48	Changed to 20% oxygen/80% helium	
loft	200	) foot	08.00.00		
Arrivo	od 19	) feet	08.01.24		
Left	190	) feet	10.00.00		

10.01.48

HI developed pain in both knees steadily and OR suddenly hit by pain. Recompressed HI relieved by 185 feet, CR by 240 foot.

Arrived	260 feet	10.06.50
Left	260 feet	12.06.00
Arrived	225 feet	12.11.30
Left	225 feet	14.11.00
Arrived	185 feet	14.18.00
loft	185 feet	16.11.00

150 foot

Arrived

CR transient pain during ascent

•	
~?;	51 met

· · ·

Posi	tilon.		Timo	Notos
Arrived	155 9	fest	16.16.00	
Loft	155 s	feet	18.11.00	
Arrived.	125 1	feet	18.17.00	19.15 CR twinges in both legs
Left	125 :	feot	20.11.00	
Arrived	100 :	řeot	20.16.00	21.54 ohanged to 60% oxygen/40% helium.
Left	100 1	feet	22.11.00	
Arrived	75 :	feet	22,16,00	
Lort	75 1	feet	24.11.00	
Arrivod	55 t	feet	24.16.00	GR tingly both hands became numbress.
loft	55 <b>:</b>	feet	26.11.00	HI slight tightness in chest before ascent.
Arrived	35	feet	26-21-00	HI niggle during accent in left ankle.
Loft	35 f	feet	28.11.00	
Armived	20 i	feet	28.21.00	NI slight ache loft knee radiating
				down to ankle. Chest tightness cased a little. CR slight ache right knec on arrival at each stop. Hands still reduced sensation.
Loft	20 1	feet	30.11.00	
Arrived	10 :	foot	30.21.00	HI similar but a little more severe. CR similar in severity as previous.
lort	10 3	feet	32.11.00	
Arrived	aurf	2 <b>0</b> 0	32.24.145	

Doors open	eđ.	32.30.00	HI very slight niggles in both knees which cleared in $\frac{1}{2}$ hour. Chest hurts on coughing. CR fairly severe pain both knees steadily easing, loss of fine touch and thermal sense both hands. Chest hurts on coughing.
		34.30.00	HI dull ache both knees. CR limping on right log. Numbress cleared to palmar aspects distal phalanges all fingers.
		48.00.00	HI minimal tightness in chest also in CR whose numbress is q.
D. Special	Routine 5		
Divers	ic and ev.		Chambor dive
			Exercise - rowing machine
Post	tion	<u>Tino</u>	Notos
Left	eurleco	00.00.00	
Arrived	300 200t	00+07+50	Breathing 10% oxygen/90% helium
left	300 feet	04.00.00	
Arrived	170 foet	04.02.36	Changed to 20% oxygon/80% helium.
Loët	170 Seet	06.00.00	
Arrived	<b>100 feet</b>	06+01+00	07.30 changed to 40% oxygen/60% holium.

Arrivod 70 feet 08.01.00

100 feet

Loft

08.00.00

Position

× 1 - -

Posit	<u>ilon</u>	Time	Notos
left	70 feet	10.00.00	
Arrived	45 feet	<b>10.01.</b> 00	LC reported not too severe pain left knee but lookod in severe pain.
left	45 feet	10.17.45	Examined. Recompressed.
Arrived	70 feet	10.21.00	Pain relieved 57 feet.
left	70 feet	11.22.00	
Arrived	60 feet	11.32-30	LC pain recurred but tolerable.
Left	60 feet	12.32.00	Slow bleed commenced with pauses.
Arrived	54 Leet	13.00.00	
beft	54 Leet	13.48.00	
Arrived	45 feet	14.05.30	LC slight increase in pain.
Left	45 feet	14.12.00	Increase in pressure to case pain.
Arrived	50 feet	14.14.00	
Left	50 feet	16.15.00	In fact 48 by loak.
Arrived	40 feet	16.27.00	
Left	40 feet	16.27.00	
Arrived	34 feet	17.43.00	LC reported pain had moved round knce and more of an ache.
Azrived	30 feet	17.53.00	Bleed by leak.
left	30 feet	18.53.00	
Arrived	20 feet	19.18.00	LC slightly wouse but tolerable and cleared by 19.36.
left	20 feet	20.18.00	
Arrived	10 feet	20.46.00	Slight leak. LC moving easily and weight bearing.

Posi	tion	Time	Notes
Loft	8 feet	21.22.00	
Arrived	2 feet	22,02,00	LC slight increase in pain.
Azrived	surface	21.54.00	By leak.
Doors open		22.02.00	Both men slight chest pain on deep
			inspiration, otherwise well.