



University
of Glasgow

<https://theses.gla.ac.uk/>

Theses Digitisation:

<https://www.gla.ac.uk/myglasgow/research/enlighten/theses/digitisation/>

This is a digitised version of the original print thesis.

Copyright and moral rights for this work are retained by the author

A copy can be downloaded for personal non-commercial research or study,
without prior permission or charge

This work cannot be reproduced or quoted extensively from without first
obtaining permission in writing from the author

The content must not be changed in any way or sold commercially in any
format or medium without the formal permission of the author

When referring to this work, full bibliographic details including the author,
title, awarding institution and date of the thesis must be given

Enlighten: Theses

<https://theses.gla.ac.uk/>
research-enlighten@glasgow.ac.uk

MAINLY ABOUT EMPHYSEMA OF THE LUNGS

Volume II

(This Volume Should Be Consulted In Conjunction With Volume I)

ProQuest Number: 10984291

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



ProQuest 10984291

Published by ProQuest LLC (2018). Copyright of the Dissertation is held by the Author.

All rights reserved.

This work is protected against unauthorized copying under Title 17, United States Code
Microform Edition © ProQuest LLC.

ProQuest LLC.
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106 – 1346

CONTENTS (Vol.II)


Figures 1 - 68.

Tables I - XXVI

Appendix I

Appendix II - Tables of Results

Bibliography.



R. T. H. LAENNEC,

Professeur Royal de Médecine au Collège de France. Prof. de
Clinique à la Faculté de Médecine de Paris. Chev. de l'Ordre
Royal de la Légion d'honneur, &c.

Figure 1. Sketch of Laennec by C.J. Williams showing him as he appeared when making his visits to La Charité in 1825-26 with Laennec's signature. (Taken from 'Memoirs of Life and Work' by Williams, London 1884).




Figure 2. Classification of emphysema by Dr. L.Mc.A. Reid illustrating variety of types presented clinically according to the presence or absence of airways obstruction. The types in boxes are those with which the thesis is mainly concerned. (Reproduced with the authors' permission).

1. Excess of air in the Lungs.

- * Low flat diaphragm
- * Large retrosternal translucent area

2. Cardiovascular changes.

- * Narrow vertical heart
- Prominence of pulmonary trunk
- Large hilar and small lung vessels

3. Bullae.

- Avascular transradiant area
- May be demarcated by a white line
- May have no definite border

Figure 3. Criteria for radiographic diagnosis of emphysema
by Dr. George Simon.

* These values can be measured (Reproduced with
the author's permission).




Figure 4. The position of the measurements described in the text on the postero-anterior and lateral chest radiographs are shown.

In the postero-anterior view:-

T.H. - the transhilar diameter measured at the point where the upper lobe vein crosses the lower lobe artery.

T.C. - transverse diameter of the heart.

A-B - the movement of the hemidiaphragms between inspiration and expiration.

The dotted line shows the position where the size of the mid-lung vessels is observed.

In the lateral view:-

X - depth of retrosternal translucent area.

Y - downward extension of retrosternal translucent area.




Figure 5. This is a perspective drawing of the electrode water bath. A, B, and C are the pH, oxygen and carbon dioxide electrodes respectively. Heaters H₁ and H₂ are shown, the first being used for rapid boosting of heat initially and the other is used for maintenance purposes. The temperature of the water in the bath is controlled by the contact thermometer regulator K and further checked by readings from the thermometer J. F and G are the entry and exit tubes for the water flow through the bath. Note that water circulation of the pH electrode is in a separate jacket. The tubes E contain distilled water equilibrated with different percentages of gases. They have direct access to the CO₂ and O₂ electrodes through the sleeve valve, D. L is a tonometer which has been used for other projects but is not generally employed.




Figure 6. This is a diagram of the multicathode oxygen electrode. The blood sample is contained in the cuvette A and is in contact with the polyvinyladine membrane B. The platinum multicathode C is on the other side of the membrane surrounded by electrolyte solution in the electrolyte chamber E. D is the silver anode, F the electrode housing and G the electrode retaining screw.




Figure 7. This is a diagram of the carbon dioxide electrode. The blood sample is contained in the cuvette A where it is in contact with the outer Teflon membrane B which is in apposition with the woven fibreglass membrane C. The membranes are kept in place with 'O' rings. D is a glass electrode and E is the calomel reference anode. F is the housing of the whole electrode.




Figure 8. A diagram of the pH replaceable capillary electrode supplied by Electronics Instruments Ltd. The sample is sucked through replaceable tube A into J. A, B and C comprise the glass electrode. B is a chamber containing HCl with a filling point at F₁, and C is a silver/silver chloride electrode. D is a porous plug separating the sample from E which is filled with Molar KCl filled from the reservoir at F₂. G is a reference electrode and H is the heated circulated water jacket.




Figure 9. Diagram of the tonometer used for storing ^{133}Xe and the apparatus used during transfer of the radioactive gas from the ampoule A to the tonometer bulb E. To transfer the ^{133}Xe , the bag B containing 100% CO_2 is connected to the tonometer and the dead space of the bulb and T piece, D is flushed with CO_2 leaving the bulb half full of CO_2 . Mercury then fills the T piece and the seal of the ampoule is broken by tip of the metal cannula C. The radioactive gas is displaced by the mercury into the tonometer diluted by CO_2 . Stopcocks T_1 , T_2 , T_3 , T_4 and T_5 are used during this procedure. F is a manometer and G is a mercury reservoir, both of which are required to maintain the stored ^{133}Xe under negative pressure.




Figure 10. This is a drawing to show the fixed geometric system used for counting the activity in the syringe A of ^{133}Xe in saline before perfusion. The counting head C is one of those from the gantry. B₁, B₂ and B₃ constitute an alarm bell system with microswitch and transformer and bell respectively. These warn the operator if the procedure with the patient is began with a counting head missing from the gantry.




Figure 11. This is a diagram of the closed circuit spirometer system containing the ^{133}Xe in air. The patient breathes through the mouth-piece and 3-way tap A. The gas is circulated by the mixing pump C in the direction of the arrows at 36 litres/min. through a soda lime canister B which absorbs CO_2 during rebreathing. The radioactivity in the circuit is monitored by the counting head E. Inspiration and expiration are shown on the spirometer F and recorded through potentiometer on the spirometer pulley G. Oxygen is introduced at A during the rebreathing procedure sufficient to keep the volume of the circuit constant.




Figure 12. A perspective drawing of the gantry supporting the counting heads used in the detection of ^{133}Xe . The patient sits upright in the chair and the anterior and posterior battery of counters are lowered and advanced into position for the procedure. The position of the counting heads is shown.




Figure 13. A block diagram of the circuit used in counting and recording the surface activity of ^{133}Xe .

Figure 14 (A) Photograph to show position of patient between batteries of counters, with shielded spirometer in position.

(B) Photograph of the bulky tower containing electronic equipment, ratemeters and recorders. The yellow trolley is the spirometer and closed circuit contained in a lead shield. The gantry and counting heads are shown.




Figure 14 A.



Figure 14 B




Figure 15. A tracing of a chest radiograph in inspiration in which the position of the counters are shown. They are placed equidistant from the mid-line in the upper, middle and lower zone of each lung. The semicircle between the clavicles represents the suprasternal notch which was used as a reference point.




Figure 16. A tracing of the regional ventilation and perfusion ^{133}Xe procedure in a normal subject reduced approximately ten times. The right and left upper, middle and lower zones counting rates are shown at appropriate attenuations also the spirometer volume and count rates from 'hot' spirometer circuit. At A ^{133}Xe was perfused into the superior vena cava in saline. At B ^{133}Xe in air was inspired. C to D shows the rebreathing procedure ending with a breathholding period. On this particular occasion duplicates were not done and the spirometer counting rate had to be increased before the rebreathing procedure.

Figure 17. Photograph of the right upper, middle and lower zone and spirometer volume tracings during perfusion of ^{133}Xe in solution and a breathholding manoeuvre. The plateaus, after perfusion, are marked with horizontal lines as are the background counts before perfusion. Two points should be noted:

- 1.) The off scale deflection in the right upper zone before breathholding as the radioactivity passes that counter in the superior vena cava.
- 2.) The rise in the background count immediately prior to injection as the syringe of isotope is brought near the patient.







Figure 17.




Figure 18. Illustration of a good 'wash in' mixing curve in the right middle zone recorded simultaneously with a poor 'wash in' curve in the left lower zone in the same patient. "A" represents the 90% mixing time in the right middle zone and "B" the 90% mixing time in the left lower zone.

T A B L E I.

A statistical analysis of pairs of observations
of ventilation and perfusion indices.

| | <u>Ventilation Indices</u> | <u>Perfusion Indices</u> |
|---|----------------------------|--------------------------|
| Number of pairs | 408 | 390 |
| Mean value of first and second estimate. | 1.004 0.998 | 0.989 0.987 |
| Standard deviation of means | 0.122 0.122 | 0.133 0.184 |
| Correlation coefficient | 0.964 | 0.986 |




Figure 19. The height and weight of 46 men in Group I is shown. The dots represent patients with no bronchitis and the triangles patients with bronchitis. The straight line shows the average weight according to height for men from 40-50 years (Society of Actuaries 1959). All except 2 of these patients are underweight and 35 are 20 lbs. or more below their expected weight.

T A B L E II

Summary of respiratory complaints in patients in Group I.

| | With Bronchitis | Without Bronchitis | Total |
|---|--------------------|-----------------------|--------------------------------------|
| <u>Initial Complaint</u> | <u>32 Patients</u> | <u>18 Patients</u> | <u>50 Patients (46male;4females)</u> |
| Cough | 23 " | nil | 23 " |
| Breathlessness | 7 " | 17 Patients | 24 " |
| Cough and Breathlessness | 2 " | nil | 2 " |
| No Complaints | nil | 1 Patient | 1 " |
| <u>Grade of Breathlessness</u> | | | |
| I | 0 Patients | 1 Patient | 1 Patient |
| II | 0 " | 3 " | 3 " |
| III | 4 " | 3 " | 7 " |
| IV | 2 " | 2 " | 4 " |
| V | 26 " | 9 " | 35 " |
| <u>Recent Chest Infections.</u> | | | |
| Grade 0 | 5 Patients | 10 Patients | 15 Patients |
| 1 | 9 " | 3 " | 12 " |
| 2 | 18 " | 5 " | 23 " |
| <u>Length of history of initial complaint</u> | | | |
| Mean Value | 7.7 years | 11.1 years | 8.8 years |
| <u>Age of Onset of Breathlessness.</u> | | | |
| Mean Value | 47.5 years | 42.8 years | 46.1 years |
| <u>Length of history of breathlessness</u> | | | |
| Mean Value | 7.0 years | 10.9 years | 8.4 years |
| Mean Age | 55.0 years | 54.9 years | 54.9 years |

T A B L E III

Comparison of smoking habits of patients in Group I with data from other studies. The results are presented as percentages of the total population studied.

| | <u>Group I</u> | <u>Higgins (1957)+</u> | <u>Olsen and Gilson (1960)*</u> |
|-----------------|----------------|------------------------|---------------------------------|
| Never Smoked | 4% | 3.8% | 7.4% |
| Stopped Smoking | 66% | 11.5% | 13.0% |
| 1-14 cigs/day | 8% | 27.0% | 44.4% |
| 15-24 cigs/day | 16% | 57.5% | 25.8% |
| 25+ cigs / day | 6% | | 9.3% |

+ Chronic bronchitic subjects.

* Population survey data.




Figure 20. The grades of intensity of the breath sounds heard anteriorly and posteriorly are compared between two observers in 34 patients in Group I. The diagonal lines indicate complete agreement or only one grade of difference in their observations.

The observers are J.R. Nairn and M. Turner-Warwick.




Figure 21. A comparison between the intensity of the breath sounds heard anteriorly in 47 patients in Group I and the dynamic distribution index in different zones.
The observer is J.R. Nairn.




Figure 22. A comparison between the intensity of the breath sounds heard anteriorly in 47 patients in Group I and the volume of air entering different zones.

The observer is J.R. Nairn.




Figure 23. A comparison of the difference in the grade of intensity of the breath sounds between the top and the bottom of the lung as recorded by two observers, J.R. Nairn and M. Turner-Warwick. Thirty-four patients in Group I were examined. The line of complete agreement between the observers passes at a 45° angle through the crossing dotted axes shown from below left to upper right. The 2 points shown in the upper left quadrant are instances of complete disagreement between observers.




Figure 24. The difference between the grades of intensity of the breath sounds from top to bottom of the lungs is compared with the difference measured in the D.D.I. between these zones. The observer is J.R. Nairn. The results in 46 patients from Group I (dots) and 7 patients from Group II (squares) are shown.




Figure 25. A comparison of the difference in the grade of intensity of the breath sounds from top to bottom of each lung and the difference in the volume of air calculated to enter the upper and lower zones.

The dots are patients in Group I and the squares depict patients in Group II.



Figure 26. Postero-anterior radiograph of J.W., a patient with generalised emphysema who complained of severe breathlessness (Grade V) but denied having a chronic productive cough. The diaphragm is flat and low (7th rib). The transverse diameter of the heart is 10.0 cms. and the transthoracic diameter 11.5 cms. The pulmonary and hilar arteries are not dilated. There are small mid-lung vessels and vascular loss in all zones except the left upper zone. This patient had an F.E.V.₁ of 500 ml. and a vital capacity of 1.80 litres. Intrapulmonary mixing was poor in all zones of the lung.



Figure 27. Postero anterior radiograph of a patient (A.P.) with extensive localised emphysema, who complained of breathlessness (Grade V) but had no bronchitis. The right dome of the diaphragm is curved at the level of the 7th anterior rib. The transverse diameter of the heart is 14·0 cms. and the trans hilar diameter 11·0 cms. The pulmonary artery is dilated but the hilar arteries are of normal size. The mid-lung vessels are small in the upper and middle zones and there is evidence of vascular loss in the upper zones. This patient (A.P.) had very poor overall lung function ($F.E.V_1 = 600 \text{ ml.}$) but the lower zones showed normal intrapulmonary mixing.



Figure 28. Postero anterior radiograph of a patient (R.G.) with extensive localised emphysema who admitted to some breathlessness (Grade III) which was not severe and a productive cough. The diaphragm is curved between the level of the 6th and 7th anterior ribs. The transverse diameter of the heart was 11.5 cms. and the transthoracic diameter 11.0 cms. The pulmonary artery is dilated but the hilar arteries are of normal size. The mid-lung vessels are small in both upper zones and the left middle and lower zones. Vascular loss is also evident in these zones. There is a demarcated bulla in the left lower zone (4 x 12 cms.) in size. This patient had an F.E.V.₁ of 2 litres and a vital capacity of 3.80 litres. Only the right lower zone had normal intrapulmonary mixing.

T A B L E IV

The mean values, standard deviations and significance levels of lung function tests in patients in Group I classified according to the radiological extent of emphysema.

| | Generalised | Extensive Localised | Localised |
|--|--------------------|---------------------|---------------------|
| Number of Patients | 22 | 13 | 14 |
| F.E.V. ₁ (ml.) | 611.3 (± 243) | 884.6 (± 1367) | 1078.5 (± 640) |
| V.C. (litres) | 2.20 (± 0.674) | 3.04+ (± 0.988) | 3.33* (± 0.0846) |
| % V.C. | 53.8 (± 14.2) | 69.3+ (± 17.8) | 78.9* (± 14.1) |
| R.V./T.L.C. | 69.7 (± 10.53) | 59.2+ (± 10.48) | 53.1‡ (± 10.7) |
| F.R.C. (litres) | 6.11 (± 1.83) | 5.58 (± 1.18) | 5.26 (± 1.19) |
| % F.R.C. | 166.8 (± 40.99) | 145.8 (± 27.05) | 127.4+ (± 34.21) |
| T.L.C. (litres) | 7.52 (± 1.99) | 7.40 (± 1.16) | 7.06 (± 1.54) |
| % Pred. T.L.C. | 119.3 (± 24.21) | 115.0 (± 16.6) | 114.1 (± 20.2) |
| D _{LCO} mLCO/min/mm.Hg. | 6.5 (± 1.96) | 8.6 (± 3.05) | 8.8 (± 2.12) |
| Rest Pa _{O₂} mm.Hg. | 73.2 (± 13.93) | 89.1+ (± 6.58) | 84.5* (± 8.09) |
| Rest Pa _{CO₂} mm.Hg. | 45.7 (± 8.27) | 40.9 (± 6.13) | 41.7 (± 5.73) |

+ = Significantly different ($p = 0.01$) from the generalised group.

* = Highly significantly different ($p = 0.001$) from the generalised group.




Figure 29. The values of forced expired volume in one second in patients in Group I according to the extent of emphysema radiologically. The horizontal bars represent the means in each group. Black dots are patients without bronchitis and empty circles are patients with bronchitis.




Figure 30. The values of vital capacity in patients in Group I according to the extent of emphysema radiologically. The horizontal bars represent the mean in each group. Black dots are patients without bronchitis and empty circles are patients with bronchitis.




Figure 31. The values of the ratio of the residual volume to the total lung capacity expressed as a percentage in patients in Group I according to the extent of emphysema radiologically. The horizontal bars represent the mean in each group. Black dots are patients without bronchitis and empty circles are patients with bronchitis.




Figure 32. The diffusing capacity for carbon monoxide in ml.CO/minute/mm.Hg. in patients in Group I according to the extent of emphysema radiologically. The horizontal bars represent the mean in each group. Black dots are patients without bronchitis and empty circles are patients with bronchitis.




Figure 33. The resting arterial oxygen tension in patients in Group I according to the extent of emphysema radiologically. The horizontal bars represent the mean in each group. Black dots are patients without bronchitis and empty circles are patients with bronchitis.

T A B L E V.

Mean values, standard deviations and significance levels of the functional residual capacity and total lung capacity expressed as a percentage of the predicted value and % RV/TLC in patients in Group I according to the shape of the diaphragm.

| | Shape of Diaphragm | |
|--------------------|--------------------|------------------|
| | Curved | Flat. |
| Number of Patients | 15 | 28 |
| % RV/TLC * | 55.1 ± 10.68 | 66.6 ± 11.44 |
| % Predicted F.R.C. | 145.1 ± 35.21 | 154.2 ± 44.05 |
| % Predicted T.L.C. | 117.5 ± 19.49 | 119.6 ± 23.83 |

* p = 0.05.




Figure 34. The relationship between the shape of the diaphragm seen radiologically and lung volumes. The residual volume to total lung capacity ratio is expressed as a percentage. The functional residual capacity and the total lung capacity are shown as a percentage of the predicted value. The patients are from Group I. The horizontal bars represent the mean in each group.

T A B L E VI

Mean Values, standard deviations and significance levels of the functional residual capacity and the total lung capacity expressed as a percentage of the predicted value and % RV/TLC in patients in Group I according to the level of the diaphragm.

| | Level of Diaphragm | |
|---------------------|--------------------|------------------|
| | Normal | Low |
| Number of Patients | 23 | 26 |
| RV /TLC % + | 55.6 ± 11.18 | 68.0 ± 12.53 |
| % Predicted F.R.C.+ | 128.3 ± 28.41 | 169.1 ± 38.6 |
| % Predicted T.L.C.* | 105.0 ± 17.72 | 125.0 ± 21.12 |

+ = p = 0.001

* = p = 0.01




Figure 35. The relationship between the level of the diaphragm (according to the anterior ribs showing above the right dome of the diaphragm) and lung volumes in patients in Group I. The volumes are expressed as in figure 34. The horizontal bars represent the mean in each group. The diaphragm is lower than normal if the 7th or more anterior ribs are above the right dome of the diaphragm on inspiration.

T A B L E VII

The mean values, standard deviation and significance levels between the lung volumes and the shape and level of the diaphragm are shown. Diaphragms which had one dome flat and the other curved are not included.

| Diaphragm Shape and Level. | | | | |
|----------------------------|------------------------|---------------------------|--------------------------|-----------------------------|
| | Flat and low (1) | Flat and Normal (2) | Curved and low (3) | Curved and Normal (4) |
| Number of Patients | 16 | 12 | 8 | 7 |
| % Predicted F.R.C. | 171.0 ± 44.3 | 132.1 ± 35.5 | 168.6 ± 35.9 | 119.7 ± 13.9 |
| % Predicted T.L.C. | 126.1 ± 23.3 | 102.2 ± 20.0 | 127.7 ± 20.7 | 105.8 ± 14.7 |
| % RV/TLC | 71.4 ± 10.6 | 60.1 ± 11.1 | 60.1 ± 9.1 | 49.3 ± 10.8 |

Between 1 and 4 % Predicted F.R.C. p = 0.001

% Predicted T.L.C. p = 0.050

% RV/TLC p = 0.001

Between 1 and 2 % Predicted F.R.C. p = 0.050

% Predicted T.L.C. p = 0.050

% RV/TLC p = 0.050

Between 1 and 3 No significant difference

Between 2 and 3 % Predicted F.R.C. p = 0.050

% Predicted T.L.C. p = 0.050

% RV/TLC No significant difference

Between 2 and 4 No significant difference

Between 3 and 4 % Predicted F.R.C. p = 0.001

% Predicted T.L.C. p = 0.050

% RV/TLC p = 0.050

T A B L E VIII.

Mean values, standard deviations and significance levels of the percentage of predicted F.R.C., T.L.C. and the $\frac{RV}{TLC} \%$ in patients in Group I according to the depth of the retrosternal translucent zone.

| | Depth of the Retrosternal Translucent Zone | |
|-------------------------|---|-------------------|
| | Normal | Large |
| Number of Patients | 15 | 31 |
| % $\frac{RV}{TLC} \%$ * | 60.0 ± 10.58 | 64.4 ± 11.96 |
| % Predicted F.R.C. * | 136.8 ± 40.74 | 159.3 ± 36.74 |
| % Predicted T.L.C. * | 113.5 ± 25.74 | 119.2 ± 19.08 |

* None of these values are significantly different.




Figure 36. The relationship between the depth of the retrosternal translucent zone measured on a lateral film and lung volumes in patients in Group I. The volumes are expressed as in figure 34. The horizontal bars represent the mean in each group. (The retrosternal space is larger than normal if it is more than 3 cms.).

T A B L E IX

The relationship between abnormally low zonal dynamic distribution indices in zones and the appearance of the mid-line vessels and evidence of vascular loss radiologically. Patients from Group I with and without bronchitis and patients from Group II are considered separately.

| Diagnosis (no. of patients) | Zone | Percentage of Zones with a low D.D.I. | | | |
|--------------------------------------|--------|---|------|--|--------|
| | | Small Mid-Lung Vessels and Vascular Loss (No. of zones) | | Normal Mid-Lung Vessels With No Vascular Loss (No. of Zones) | |
| Emphysema (18) | Upper | {(23)} | 78·0 | 20·0 | {(10)} |
| | Middle | {(21)} | 71·5 | 30·0 | {(15)} |
| | Lower | {(17)} | 100 | 46·7 | {(15)} |
| Emphysema and Bronchitis. (32) | Upper | {(29)} | 72·5 | 29·0 | {(24)} |
| | Middle | {(27)} | 85·0 | 80·0 | {(25)} |
| | Lower | {(32)} | 94·0 | 79·0 | {(24)} |
| Bronchitis (8) | Upper | - | - | 25·0 | {(16)} |
| | Middle | - | - | 75·0 | {(16)} |
| | Lower | - | - | 81·0 | {(16)} |




Figure 37. The relationship between poor intrapulmonary mixing and the radiological appearance of the mid-lung vessels and evidence of vascular loss in upper, middle and lower zones of the lung. Mixing is measured by the dynamic distribution index and is abnormal if this index is below the normal value for the zone in question. The black areas represent zones where there were normal mid-lung vessels and no vascular loss and the empty rectangular areas represent zones where there were small mid-lung vessels and evidence of vascular loss. The blocks labelled EMPHYSEMA are 18 patients in Group I who had no bronchitis. Those labelled EMPHYSEMA and BRONCHITIS are 31 patients in Group I with bronchitis and the BRONCHITIS patients are from Group II. There are no black zones in the latter group as there were no small mid-lung vessels or vessel loss in these patients.

T A B L E X.

The relationship between poor perfusion and the radiological appearance of the mid lung vessels and vascular loss. Poor perfusion is judged to be present if the perfusion index is below normal for that zone. The patients with emphysema are from Group I, divided according to the presence or absence of bronchitis. The patients with bronchitis in this table are from Group II.

| Diagnosis (number of patients.) | Zone | Percentage of zones with poor perfusion | |
|--|--------|--|--|
| | | Small Lung Vessels and Vascular Loss (no. of zones*) | Normal Lung Vessels and No Vascular Loss (no. of zones*) |
| Emphysema (18) | Upper | {23} 4.3 | {10} 0 |
| | Middle | {21} 0 | {10} 0 |
| | Lower | {17} 76.0 | {15} 0 |
| Emphysema and Bronchitis (31) | Upper | {29} 0 | {24} 4.2 |
| | Middle | {27} 3.7 | {25} 0 |
| | Lower | {32} 78.1 | {24} 29.2 |
| Bronchitis (8) | Upper | {-} - | {16} 0 |
| | Middle | {-} - | {16} 0 |
| | Lower | {-} - | {16} 68.8 |

* The reason for the discrepancy in the number of zones shown is the occurrence of small lung vessels and no vascular loss or normal lung vessels and vascular loss in the zones omitted from this table.

T A B L E XI

The mean and ranges of values of the forced expired volume in one second (F.E.V.₁) static lung volumes and diffusing capacity at rest and during exercise in patients in Group I.

| Test | Mean Value | Range | Number of Patients |
|--|------------|------------|--------------------|
| F.E.V. ₁ (ml.) | 824 | 250 - 2300 | 50 |
| F.E.V. ₁ /F.V.C.% | 34.10 | 19 - 64 | 50 |
| V.C. (litres) | 2.76 | 1.2 - 4.7 | 50 |
| F.R.C. (litres) | 5.60 | 3.11-9.80 | 50 |
| R.V. (litres) | 4.84 | 2.19-9.00 | 50 |
| T.L.C. (litres) | 7.32 | 4.19-10.5 | 50 |
| R.V./T.L.C. % | 61.50 | 34 - 88 | 50 |
| D _L CO at rest ml.CO/min/mm.Hg. | 7.89 | 3.0 - 16.1 | 50 |
| D _L CO during exercise ***** | 12.17 | 4.5 - 24.2 | 40 |
| % Extraction CO at rest | 30.14 | 17 - 48 | 50 |

T A B L E XII

The mean values of arterial blood gas tensions at rest and after exercise* in patients in Group I. Derived values are also shown.

| | Mean Value | Range | Number of Patients |
|---|------------|-----------|--------------------|
| PaO_2 (mm.Hg.) at rest | 83.5 | 53- 99.5 | 45 |
| PaO_2 (mm.Hg.) after exercise | 79.5 | 56-104.5 | 33 |
| PaCO_2 (mm.Hg.) at rest | 43.2 | 32.5-62.0 | 45 |
| PA-aO_2 difference (mm.Hg) at rest | 17.33 | 2.5-48.7 | 25 |
| V_A (l/min) at rest | 4.8 | 2.4- 7.8 | 25 |
| V_D/V_T at rest | 0.552 | 0.36-0.70 | 25 |

* The PaCO_2 after exercise has not been considered for reasons described in the text.

T A B L E XIII.

The mean, standard deviation and significance of the difference between arterial blood gas tensions at rest and after exercise in patients in Group I considered according to the presence or absence of bronchitis.

| | With Bronchitis | | | Without Bronchitis | | | Significance Level. |
|---|-----------------|--------|--------------------|--------------------|--------|--------------------|---------------------|
| | Mean | ± S.D. | Number Of Patients | Mean | ± S.D. | Number Of Patients | |
| Pa _{O₂} (mmHg) at rest. | 82.4 | 12.29 | 30 | 84.6 | 12.45 | 15 | Not Signif. |
| Pa _{O₂} (mmHg) after ex. | 77.7 | 12.69 | 21 | 82.7 | 16.25 | 12 | Not Signif. |
| Pa _{CO₂} (mmHg) at rest | 45.2 | 8.85 | 30 | 39.4 | 4.65 | 15 | p 0.01 |
| V _A l/min at rest | 4.36 | 1.40 | 15 | 5.54 | 1.37 | 10 | p 0.05 |
| V _D /V _T at rest | 0.584 | 0.25 | 15 | 0.506 | 0.10 | 10 | Not Signif. |
| P _{A-a} O ₂ mm.Hg | 18.03 | 12.60 | 15 | 16.87 | 24.20 | 10 | Not Signif. |




Figure 38. The arterial carbon dioxide tension ($P_{a}CO_2$) in mm.Hg. and the forced expired volume in one second in patients in Group I with emphysema. The black dots are patients with no chronic bronchitis and the triangles patients with chronic bronchitis. The straight line depicts the upper limit of the normal range of $P_{a}CO_2$ in this laboratory.




Figure 39. A comparison of the ventilation (V) and perfusion (Q) indices in patients in Group I. The empty triangles are measurements from the upper zone, the black triangles the middle zone, and the black dots from the lower zone.




Figure 40. A comparison between the ventilation (V) index and the dynamic distribution index (D.D.I.) in all zones in patients in Group I.

T A B L E XIV.

The distribution of impairment in regional function
in patients in Group I using the dynamic distribution index.

| Regional Function Impairment | No. of Patients | Coding * |
|---|-----------------|----------|
| All Zones | 8 | 222 } |
| All Zones Except 1 Upper Zone | 5 | 122 } |
| All Zones Except Both Upper Zones | 13 | 022 |
| Lower Zones and 1 Middle Zone | 5 | 012 } |
| Lower Zones Only | 9 | 002 } |
| Upper and Middle Zones and 1 Lower Zone | 8 | 221 } |
| Upper and Middle Zones only | 2 | 220 } |

* The coding is for ease of reference. The first, second and third number in the series refers to the upper, middle and lower zones respectively. The number itself refers to the number of zones in the lungs with a lower D.D.I. than normal.

For the purposes of comparison with total lung function data these patterns of impairment regionally have been considered in 4 approximately equal groups according to the brackets of the coding shown.

T A B L E XV.

Mean Values, ranges (in brackets), and standard deviations of total lung function tests in 4 groups of patients from Group I defined according to impairment in regional lung function.

| Total Lung Function Tests | GROUPS ACCORDING TO REGIONAL IMPAIRMENT IN FUNCTION | | | |
|--|---|-----------------------------|-----------------------------|--------------------------------|
| | 222 122 } } | 022 | 221 220 } } | 012 002 } } |
| F.E.V ₁ (ml) * | 661 ± 212.3 (250-2,000) | 715 ± 205.7 (400-1050) | 1230 ± 252.0 (350-2300) | 817 ± 411.1 (300-1100) |
| V.C. (l) | 2.58 ± 1.014 (1.2-4.5) | 2.61 ± 0.762 (1.4-3.8) | 3.24 ± 1.180 (1.6-4.7) | 2.65 ± 0.805 (1.4-4.4) |
| F.R.C.(l) | 6.01 ± 1.44 (3.11-7.66) | 5.26 ± 1.50 (3.32-8.40) | 5.90 ± 1.35 (4.05-7.80) | 5.34 ± 1.86 (3.08-9.80) |
| RV/TLC % | 64.6 ± 12.49 (38-79) | 60.4 ± 11.66 (45 - 80) | 59.0 ± 14.39 (41-82) | 61.4 ± 14.25 (38 - 77) |
| D _{LCO} Rest mlCO/min/Hg | 6.64 ± 2.97 (3.0-11.4) | 7.43 ± 1.85 (4.6-11.8) | 8.76 ± 2.45 (5.7-12.6) | 8.56 ± 3.13 (4.7-13.5) |
| D _{LCO} Ex. ml.CO/min/Hg | 10.9 (8.4-15.2) | 11.7 (4.5-18.2) | 13.4 (5.8-24.2) | 12.4 (5.9-19.8) |
| * p > 0.001 between 221 } and 022 and 012 } 220 } 002 } | | | | |
| PaO ₂ (rest) mm.Hg.* | 77.6 ± 16.09 (53.0-96.5) | 79.1 ± 10.67 (72.5-99.5) | 88.9 ± 9.20 (72.0-98.5) | 79.4 ± 10.54 (64.0 - 99.0) |
| PaO ₂ Ex. + mm. Hg. | 75.3 ± 9.70 (58.0-85.0) | 74.5 ± 10.39 (57.5-100) | 92.0 ± 7.86 (56.0-104) | 79.3 ± 16.2 (58.5- 104) |
| PaCO ₂ rest mm.Hg. | 43.7 ± 9.83 (32.5-62.0) | 45.5 ± 9.06 (33.0-62.0) | 43.4 ± 6.35 (36.0-56.0) | 41.8 ± 9.29 (33.0-56.0) |
| P _{A-a} O ₂ diff. | 18.4 (7.4 - 24.9) | 18.3 (2.5 -25.4) | 13.2 (3.0-29.3) | 17.2 (6.8-27.8) |
| V _D /V _T 0 | 0.52 ± 0.089 (0.37-0.62) | 0.62 ± 0.100 (0.41-0.70) | 0.49 ± 0.084 (0.36-0.62) | 0.58 ± 0.029 (0.56-0.64) |
| * p > 0.05 between 221 } and other 3 groups. 220 } | | | | |
| + p > 0.01 between 221 } and 222 } and 022. 220 } 122 } | | | | |
| 0 p > 0.5 between 221 } and 022 221 } | | | | |




Figure 41. Postero anterior radiograph of R.T., a patient from Group II with chronic obstructive bronchitis. This is "chronic bronchitis gone wrong". The right dome of the diaphragm is curved and at a level midway between the 6th and 7th anterior rib. The transverse diameter of the heart was 12.5 cms, and the transtilar diameter 10.5 cms. The pulmonary artery is dilated (the transverse diameters of the arteries basalis on 24 and 23 mm. left and right, respectively). The mid-lung vessels are normal except for dilated vessels in the apical zones and there is no vessel loss.

T A B L E XVI.

A summary of the symptoms and history of 9 patients selected from Group I and 7 patients selected from Group II for the 'emphysema' and 'bronchitis' groups.

| | EMPHYSEMA | BRONCHITIS |
|--|------------|------------|
| Average Age | 54.7 yrs. | 57.7 yrs. |
| Average age at onset of breathlessness | 41.8 yrs. | 51.3 yrs. |
| Number of years of breathlessness | 12.8 yrs. | 6.4 yrs. |
| First Symptom Cough | - | 6 Patients |
| Breathlessness | 9 Patients | 1 Patient |
| Average grade of breathlessness | 4.7 | 4.1 |
| Incidence of chest infections in past 3 years, i.e. more than a week incapacitated, and with sputum. | | |
| None | 5 Patients | 1 Patient |
| 1 | 2 Patients | 0 Patients |
| 2+ | 2 Patients | 6 Patients |
| History of episode of heart failure | 0 Patients | 4 Patients |




Figure 42. Height and weight of 7 patients with bronchitis from Group II and 9 patients from Group I with emphysema selected as described in the text. The stars show the bronchitic patients and the circles the emphysematous. Males are the solid symbols and females are depicted by outlined symbols. The dashed line represents the average weight for height in males and continuous line applies to females (Society of Actuaries, 1959). These lines apply to the same age group as the patients studied.

T A B L E XVII

The patients selected from Group I and II for the 'emphysema' and 'bronchitis' groups with their age, sex, smoking habits, haemoglobin concentration and E.C.G. findings.

| Object | Age | Sex | Smoking Habits | Hb (gms.) | E.C.G. Findings* |
|-------------------------|-----|-----|----------------|---------------------|-------------------|
| <u>EMPHYSEMA GROUP</u> | | | | | |
| J. | 59 | M | Moderate | 15.3 | Normal Pattern |
| H. | 55 | F | Moderate | 14.9 | Normal Pattern |
| J.M. | 63 | M | Ex. Moderate | 13.1 | Normal Pattern |
| J. | 44 | F | Ex. Light | 13.4 | - |
| H. | 55 | F | Ex. Light | 15.4 | RV + |
| J. | 40 | M | Ex. Moderate | 15.0 | Normal Pattern |
| V. | 55 | M | Ex. Moderate | 14.4 | RV + |
| J. | 58 | M | Light | 14.0 | Possibly RV+ |
| Mah. | 62 | M | Ex. Moderate | 17.5 | RV + |
| Mean (\pm SD) | | | | 14.78 (\pm 1.27) | |
| <u>BRONCHITIS GROUP</u> | | | | | |
| J. | 60 | M | Ex. Moderate | 16.8 | RV + |
| J. | 64 | M | Light | 16.1 | RV + |
| L. | 62 | M | Moderate | 13.8 | Ischaemic changes |
| J. | 48 | M | Ex. Heavy | 17.2 | RV +++ |
| Sc. | 60 | F | Non Smoker | 17.8 | RV + |
| Pat. | 51 | F | Moderate | 14.3 | RV + |
| J. | 59 | M | Light | 16.1 | Normal Pattern |
| Mean (\pm SD) | | | | 16.01 (\pm 1.36) | |

The E.C.G. evidence of right ventricular hypertrophy (RV+) in cor pulmonale is according to the criteria of Towers (1966).

T A B L E XVIII

Mean values and standard deviations, and significance, of total lung function tests in 9 patients from Group I and 7 patients from Group II selected for the 'emphysema' and 'bronchitis' groups.

| | <u>Emphysema</u> | <u>Bronchitis</u> | <u>Significance</u> |
|--|------------------|-------------------|---------------------|
| F.E.V. ₁ (ml.) | 628 | 636 | |
| F.E.V. ₁ / F.V.C. | 31.2 | 35.7 | |
| V.C. (litres) | 2.54 ± .94 | 2.22 ± 0.42 | - |
| % Predicted V.C. | 62.2 ± 16.12 | 62.9 ± 6.31 | - |
| F.R.C. (litres) | 6.05 ± 1.60 | 4.76 ± 0.47 | - |
| % Predicted F.R.C. | 172.3 ± 34.79 | 149.4 ± 22.63 | - |
| T.L.C. (litres) | 7.64 ± 1.62 | 6.46 ± 0.68 | - |
| % Pred. T.L.C. | 126.8 ± 15.33 | 116.1 ± 11.66 | - |
| R.V./T.L.C. | 66.6 ± 11.83 | 66.1 ± 5.45 | - |
| <u>At Rest</u> | | | |
| D _L C ₀ (ml.CO/min/mm.Hg) | 6.9 ± 3.09 | 7.6 ± 1.97 | - |
| Minute Ventilation (l/min) | 10.12 | 8.64 | - |
| % Extraction CO | 28.7 ± 9.3 | 32.0 ± 3.74 | - |
| <u>During Exercise</u> | | | |
| D _L C ₀ (ml.CO/min/mm.Hg) | 10.83 ± 5.41 | 10.68 ± 5.06 | - |
| Minute Ventilation (l/min) | 18.42 | 15.22 | |
| <u>At Rest</u> | | | |
| PaO ₂ (mm.Hg) | 75.2 ± 12.25 | 66.5 ± 10.05 | - |
| PaCO ₂ (mm.Hg)* | 42.4 ± 3.24* | 53.1 ± 4.48* | * |
| pH | 7.39 | 7.36 | |
| % Satn. Hb | 92.81 ± 4.15 | 89.74 ± 4.24 | - |
| A-a P _O ₂ gradient (mm.Hg) | 15.9 ± 4.09 | 13.8 ± 11.27 | - |
| V _A l/min. | 5.1 ± 1.10 | 3.8 ± 0.67 | - |
| V _D /V _T | 0.57 ± 0.256 | 0.62 ± 0.075 | - |
| Minute Ventilation L/min. | 12.3 | 9.42 | - |

* p = 0.001




Figure 43 The vital capacity expressed as a percentage of the predicted value in 9 patients with emphysema and 7 patients with bronchitis. The mean value is shown by a horizontal bar in each group.




Figure 44. The functional residual capacity expressed as a percentage of the predicted value in 9 patients with emphysema and 7 patients with bronchitis. The mean value is shown by a horizontal bar in each group.




Figure 45. The total lung capacity expressed as a percentage of the predicted value in 9 patients with emphysema and 7 patients with bronchitis. The mean value is shown by a horizontal bar in each group.




Figure 46. The ratio of the residual volume to the total lung capacity in 9 patients with emphysema and 7 patients with bronchitis. The mean value is shown by a horizontal bar in each group.




Figure 47. The steady state diffusing capacity in 9 patients with emphysema and 7 patients with bronchitis measured at rest and during stepping exercise. The mean value is shown by a horizontal bar in each group.




Figure 48. The arterial carbon dioxide tension at rest in 8 patients with emphysema and 7 patients with bronchitis. The mean value is shown by a horizontal bar in each group.




Figure 49. The arterial oxygen tension at rest and after exercise in 8 patients with emphysema and 7 patients with bronchitis. Some of the patients were unable to do any exercise. In them only the resting values are shown. The horizontal bar shows the mean value in each group.




Figure 50. The dynamic distribution indices in the upper, middle and lower zones of both lungs in 9 patients with emphysema. The rectangular blocks indicate the range of normal values. The numbers 1-9 refer to individual patients so that the values in the left and right lungs can be compared in each case. Radiologically there were small mid-lung vessels and vascular loss in the upper and middle zones of patient no. 1, 2, 5 and 6. These changes were only in the lower zones of patient no. 3 and 4. Nos. 7, 8 and 9 had no normal peripheral vessels visible on a chest radiograph.




Figure 51. The perfusion indices in 9 patients with emphysema in the upper, middle and lower zones of each lung. The rectangular blocks indicate the range of normal values in these zones.




Figure 52. The dynamic distribution indices in the upper, middle and lower zones of both lungs in 7 patients with bronchitis. The rectangular blocks indicate the range of normal values in these zones.




Figure 53. The perfusion indices in 7 patients with bronchitis in the upper, middle and lower zones of each lung. The rectangular blocks indicate the range of normal values in these zones.




Figure 54. The arterial carbon dioxide tension ($P_a\text{CO}_2$) in mm.Hg. and the forced expired volume in one second in 8 patients with chronic obstructive bronchitis (Group II) and patients from Group I with no chronic bronchitis. The empty circles are the former group and the solid dots the latter.




Figure 55. The arterial carbon dioxide tension (PaCO_2) and the forced expiratory volume in one second in patients from Group I who have bronchitis in addition to emphysema. The straight line depicts the upper limit of the normal range of PaCO_2 in this laboratory.

T A B L E XIX.

Clinical pre-operative assessment and subjective change after surgery in patients in Group Ia.

| Patient | Age | Sex | + C.B.* | ECG findings* | Grade Of Breathlessness Before Op | Grade Of Breathlessness After Op. | Subjective Improvement |
|---------|-----|-----|---------|---------------|--------------------------------------|--------------------------------------|------------------------|
| R.K. | 56 | M | + | Normal | V | II | 3 |
| A.S. | 60 | M | + | Normal | V | II | 3 |
| E.Sh. | 64 | M | - | Normal | IV | II | 2 |
| W.Sh. | 59 | M | - | RV+ | V | III | 2 |
| W.P. | 44 | F | - | Normal | V | III | 2 |
| J.W. | 55 | M | - | RV+ | V | III | 2 |
| Gl.H. | 55 | F | - | Normal | V | II | 3 |
| A.V.M. | 63 | M | - | Normal | II | II | 0 |
| T.P. | 44 | M | + | Normal | IV | III | 1 |
| H.B. | 50 | M | + | Normal | V | IV | 1 |
| E.Pl. | 55 | F | - | RV+ | V | V | 0 |
| Geo. H. | 53 | M | - | RV+ | II | II | 0 |
| R.H. | 49 | M | + | Normal | V | V | 0 |
| J.E. | 39 | M | + | RV+ | III | III | 0 |

* C.B. = Chronic Bronchitis, + = present; - = absent.

+RV+ = Right ventricular hypertrophy according to the criteria of Towers (1966).

T A B L E XX.

Radiological findings and localisation with ^{133}Xe before operation
with subjective improvement after surgery in Group I patients.

| Patient | Rad.+ Diagnosis | Compression | Site and Size Demarcated Bullae | Radiologically Hypovascular Zones* | Zones with Low D.D.I.* | Subjective Improvement |
|---------|-----------------|-------------|---------------------------------|------------------------------------|--------------------------|------------------------|
| R.K. | Gen. | - | - | All Except LU | Mids and Lowers. | 3 |
| A.S. | Loc. | + | RL 15x15 cms. | Lowers | All Except LU | 3 |
| E.Sh. | Gen. | - | RL 15x12 cms. LL 7x7 cms. | Uppers R.M. & Lowers | R.M. and Lowers | 2 |
| W.Sh. | Gen. | - | - | All Except LU | R.M. and Lowers | 2 |
| W.P. | Loc. | - | RL 8x8 cms. | R.M. & Lowers | All Except RU | 2 |
| J.W. | Gen. | - | - | All Except LU | All Except LU | 2 |
| Gl.H. | Gen. | - | - | Uppers & Mids Uppers & Mids | All Zones R.Lung Only | 3 0 |
| A.V.M. | Loc. | - | R.L. 9x9 cms | Lowers | R.U. and Lowers | 1 |
| T.P. | Loc. | - | LL 15x15 cms | Left Lower | L.M. and Lowers | 1 |
| H.B. | Normal | - | - | None | Mids and Lowers | 1 |
| E.Pl. | Loc. | - | - | Uppers & Mids | All Zones | 0 |
| Geo.H. | Loc. | + | LL 18x18 cms | Uppers & L. Lower | L. Upper & Lowers. | 0 |
| R.H. | Gen. | + | - | L.M. and Lowers | Middles & Lowers | 0 |
| J.E. | Loc. | - | - | Lowers | L.U. L.M.& Lowers | 0 |

+ In this column Gen. - generalised. Loc. - Localised emphysema.

* L.U. - Left Upper Zone.
L.M. - Left Middle Zone.

R.U. - Right Upper Zone.
R.M. - Right Middle Zone.

T A B L E XXI.

Vital Capacity and Forced Expired Volume in 1 Second before operation and the change in these values after operation in Patients in Group Ia.

| Patient | V.C.*(l.) before operation | Change in V.C. after operation(ml) | FEV ₁ % Before Op. | Change in FEV ₁ after operation(ml.) |
|---------|----------------------------|------------------------------------|-------------------------------|---|
| R.K. | 3.60 (89) | +0.30 | $\frac{600}{2400} 25\%$ | nil |
| A.S. | 2.20 (59) | +0.30 | $\frac{500}{1670} 30\%$ | + 100 |
| E.Sh. | 2.40 (55) | +1.30 | $\frac{900}{2090} 43\%$ | + 400 |
| W.Sh. | 3.25 (74) | +1.20 | $\frac{1000}{3000} 33\%$ | + 100 |
| W.P. | 2.40 (60) | +0.20 | $\frac{600}{1500} 40\%$ | nil |
| J.W. | 1.80 (41) | +0.40 | $\frac{500}{1800} 28\%$ | nil |
| G1.H. | 1.40 (50) | +0.70 | $\frac{300}{1000} 30\%$ | + 200 |
| | 2.10 (14) | +0.05 | $\frac{500}{2280} 22\%$ | - 100 |
| T.P. | 3.20 (60) | +0.10 | $\frac{900}{2700} 33\%$ | - 150 |
| A.V.M. | 4.40 (100) | -0.80 | $\frac{900}{2400} 38\%$ | + 300 |
| H.B. | 2.90 (72) | +0.10 | $\frac{600}{2400} 25\%$ | + 200 |
| E.Pl. | 1.20 (51) | -0.05 | $\frac{250}{1100} 23\%$ | + 50 |
| Geo. H. | 3.40 (92) | -0.30 | $\frac{800}{3300} 24\%$ | -200 |
| R.H. | 2.15 (53) | +0.15 | $\frac{1050}{2500} 42\%$ | -650 |
| J.E. | 3.80 (72) | -0.05 | $\frac{1050}{3600} 29\%$ | +150 |

* % of the predicted value is shown in parenthesis.




Figure 56. Lateral chest tomograph taken 7 cms. from the table on the left side of R.K. before operation.



Figure 57. Lateral chest tomograph taken 7 cms. from the table on the left side of R.K. after operation. Note difference in vessels behind heart.




Figure 58. A comparison between the change in the F.E.V.₁ and subjective improvement after thoracotomy in 15 patients with emphysema (Group Ia). The subjective improvement is judged by the change in the grade of breathlessness.




Figure 59. A comparison between the change in the F.R.C. and subjective improvement after thoracotomy in 15 patients with emphysema (Group Ia).




Figure 60. The relationship between the change in resting arterial oxygen tension (PaO_2) after thoracotomy and the subjective improvement in 9 patients with emphysema from Group Ia.




Figure 61. A comparison between the change in the D.D.I. in the operated zone and the subjective improvement after thoracotomy in 15 patients with emphysema (Group Ia).




Figure 62. A comparison between the change in the D.D.I. in the zone adjacent to that mainly operated upon and the subjective improvement after thoracotomy in 15 patients with emphysema (Group Ia).




Figure 63. A comparison between the sum of the changes in the D.D.I. in the operated and the adjacent zone ("Total D.D.I.") and the subjective improvement after thoracotomy in 15 patients with emphysema (Group Ia).




Figure 64. Postero anterior chest radiograph of patient (M.E.) with unilateral transradiancy. The right lung is transradian. Note also the smaller pulmonary artery on that side. breath sounds in the transradiant lung and the smaller bronchies are breath sounds heard in the contralateral lung.




Figure 65. This shows the relationship between the grade of intensity of the breath sounds and the D.D.I. in 7 patients with unilateral transradiancy (Group III). The circles are breath sounds heard in the transrariant lung and the smaller triangles are breath sounds heard in the contralateral lung.

T A B L E XXII.

Grading of Symptoms in 7 Patients with Macleod's
Syndrome, (Group III).

| Patient (Age & Sex) | Cough | Sputum | History (Recent) | Breathlessness | History of Childhood Illness. |
|------------------------|-------|--------|---------------------|----------------|--|
| M.E. (18;F.) | 2 | 0 | 1 | 1 | "Born with Bronchitis" |
| A.K. (46;M.) | 0 | 0 | 2 | 3 | Serious Chest Illness, aet. 9/12 |
| A.S. (53;M.) | 2 | 2 | 2 | 3 | Cough Since age of 2 years. |
| E.L. (47;M.) | 2 | 2 | 2 | 4 | Double pneumonia & bronchitis as baby |
| D.C. (41;M.) | 2 | 2 | 2 | 3 | Double pneumonia aet. 8 years. |
| M.D. (28;M.) | 1 | 1 | 2 | 2 | Pneumonia, aet. 2 years |
| H.S. (54;F.) | 1 | 1 | 0 | 5 | Pneumonia and pleurisy when "very young". |

T A B L E XXIII

Lung volumes, flow rates and airway conductance measurements
in 7 patients with Macleod's Syndrome (Group III). The
predicted values are shown beneath the observed values in brackets.

| Patient | F.R.C. litres | RV/TLC | V.C. litres | $\frac{FEV_1}{FVC}$ ml. | Airway Conductance (C) and Lung Vol.(V) at which it was measured | |
|---------|------------------|--------------|----------------|-------------------------|---|----------|
| | | | | | C (litres/sec/cmH ₂ O) | V.litres |
| M.E. | 2.18 (2.67) | 41% (26%) | 2.60 (3.60) | $\frac{1800}{2600}$ 69% | 0.260 (0.263) | 2.01 |
| A.K. | 4.03 (3.48) | 55% (33%) | 3.10 (4.05) | $\frac{1000}{2600}$ 38% | 0.121 (1.001) | 7.65 |
| A.S. | 4.90 (3.74) | 48% (33%) | 3.50 (4.36) | $\frac{900}{2900}$ 31% | 0.267 (0.770) | 5.92 |
| E.L. | 4.28 (2.98) | 60% (34%) | 2.40 (3.40) | $\frac{650}{2050}$ 32% | 0.235 (0.780) | 6.32 |
| D.C. | 4.81 (4.25) | 54% (31%) | 3.50 (5.30) | $\frac{1000}{2500}$ 40% | 0.192 (1.050) | 8.10 |
| M.D. | 4.32 (3.99) | 54% (28%) | 2.80 (5.28) | $\frac{900}{2450}$ 37% | 0.105 (0.690) | 5.30 |
| H.S. | 2.12 (2.90) | 44% (34%) | 2.05 (3.22) | $\frac{1200}{1800}$ 67% | 0.247 (0.314) | 2.42 |

T A B L E XXIV.

The diffusing capacity for carbon monoxide (D_{LCO}) and the minute ventilation (Vent.) at rest and during exercise with the percentage extraction of CO (% Extr.) at rest in 7 patients with Macleod's Syndrome (Group III). The predicted values are shown beneath the observed values in brackets.

| Patient | D_{LCO} at rest | Vent. l/min. | % Extr. | D_{LCO} Exercise | Vent. l/min. |
|---------|-------------------|-----------------|------------|-----------------------|-----------------|
| M.E. | 17.7 (21.9) | 11.0 | 49 (55) | 17.9 (25.0) | 15 |
| A.K. | 9.4 (15.5) | 13.0 | 30 (40) | 15.2 (28.0) | 21.6 |
| A.S. | 8.9 (16.9) | 8.7 | 37 (40) | 12.0 (29.0) | 19.0 |
| E.L. | 10.4 (19.3) | 7.0 | 44 (46) | 13.2 (30.0) | 15.4 |
| D.C. | 9.1 (20.0) | 10.1 | 39 (46) | 27.8 (35.0) | 27.6 |
| M.D. | 20.9 (23.2) | 10.4 | 48 (53) | 22.4 (45.0) | 20.9 |
| H.S. | 9.8 (14.6) | 11.3 | 33 (44) | 13.4 (28.0) | 23.8 |




Figure 66. The partial pressure of oxygen in the arterial blood at rest and after exercise in 7 patients with unilateral transradiancy (Group III). The dashed line at the level of 95 mm.Hg. represents the normal mean value.




Figure 67. The dynamic distribution indices in the upper, middle and lower zones of the lungs in 7 patients with unilateral transradiancy (Group III). The rectangular blocks show the range of normal values. Points connected by a solid line are observations from the transradiant lung whereas points connected by a dotted line are observations from the contralateral or better lung.




Figure 68. The perfusion indices in the upper, middle and lower zones of the lungs in 7 patients with unilateral transradiancy (Group III). The rectangular blocks show the range of normal values in each zone. Points connected by a solid line are observations from the transradiant lung, whereas points connected by a dotted line are observations from the contralateral or better lung.




Figure 69. Calculated and predicted values for the vital capacity of the transradian and other remaining lung in 7 patients with unilateral transradiancy (Group III). The predicted value for each lung is calculated from the total predicted vital capacity and Svanberg's (1957) data. The observed vital capacity for both lungs is apportioned according to the sum of the counting rates in each lung after maximal inspiration.

T A B L E XXV.

The calculated distribution of the observed vital capacity of 7 patients exhibiting Macleod's syndrome (Group III). The normal partition of the predicted vital capacities are shown for comparison.

| Patient | Side Affected | Vital Capacity 1. | | | |
|---------|------------------|-------------------|-----------|-------------|-----------|
| | | Transradian Lung | | Better Lung | |
| | | Calculated | Predicted | Calculated | Predicted |
| M.E. | Right | 0.94 | 1.94 | 1.66 | 1.66 |
| A.K. | Right | 0.84 | 2.19 | 2.23 | 1.86 |
| A.S. | Right | 1.23 | 2.35 | 2.27 | 2.01 |
| E.L. | Right | 0.62 | 1.89 | 1.78 | 1.61 |
| D.C. | Left | 2.21 | 2.44 | 1.29 | 2.86 |
| M.D. | Left | 1.10 | 2.26 | 1.70 | 3.02 |
| H.S. | Left | 1.23 | 1.53 | 0.82 | 1.79 |

T A B L E XXVI.

The observed lung volumes and conductances measured at the same time in 7 patients exhibiting Macleod's syndrome (Group III).

The volume of the contralateral lung is calculated from the total observed lung volume according to the sum of the counts in each lung at equilibrium. The predicted conductances are calculated from the lung volumes according to Briscoe and Dubois, (1958).

| Patient | Side Affected | Observed Lung Volume (l.) | Observed total conductance (1/sec./cm. water). | Predicted total conductance (1/sec/cm. water). | Calculated volume of contralateral lung (l.). | Predicted conductance of contralateral lung (1/sec/cm water). |
|---------|---------------|---------------------------|--|--|---|---|
| M.E. | Right | 2.01 | 0.260 | 0.263 | 1.00 | 0.130 |
| A.K. | Right | 7.65 | 0.121 | 1.001 | 4.16 | 0.540 |
| A.S. | Right | 5.92 | 0.267 | 0.770 | 3.15 | 0.410 |
| E.L. | Right | 6.32 | 0.235 | 0.780 | 3.46 | 0.450 |
| D.C. | Left | 8.10 | 0.192 | 1.050 | 4.73 | 0.615 |
| M.D. | Left | 5.30 | 0.105 | 0.690 | 2.70 | 0.350 |
| H.S. | Left | 2.42 | 0.247 | 0.314 | 1.26 | 0.150 |

Questionnaire on Respiratory Symptoms

COUGH

Yes

No

1. Do you usually cough first thing in the morning in winter?
2. Do you do this in summer?
3. Do you usually cough during the day or at night in winter?
4. Do you do this in summer?
5. Do you cough like this on most days for as much as 3 months each year? (If "yes" to questions 1-4).

PHLEGM

6. Do you usually bring up any phlegm from your chest first thing in the morning in winter?
7. Do you do this in summer?
8. Do you bring up any phlegm from your chest during the day or at night in winter?
9. Do you do this in summer?
10. Do you bring up phlegm like this on most days for as much as 3 months each year? (If "yes" to any of questions 6-9).
11. How long have you had this phlegm?
Is it less than 2 years or more than 2 years?

BREATHLESSNESS

12. Are you ever troubled by shortness of breath, when hurrying on the level or walking up a slight hill?
13. Do you get short of breath walking with other people at an ordinary pace on the level?

Yes No

14. Do you have to stop for breath when walking at your own pace on the level?
15. Are you short of breath on washing or dressing?

WHEEZING

16. Does your chest ever sound wheezy or whistling?
17. Do you get this with colds?
18. Do you get this apart from colds?
19. Do you get this most days or nights?

CHEST ILLNESSES

20. During the past 3 years have you had any chest illness which has kept you off work, indoors, at home or in bed?
21. If "yes", ask details of each illness:-

| Year | Duration of Incapacity | | Increased Phlegm | | Doctor's diagnosis |
|------|------------------------|--------------------|------------------|----|--------------------|
| | One week or more | Less than one week | Yes | No | |
| | | | | | |

OPERATIONS:

Yes No

22. Do you smoke?
23. Have you ever smoked?
a) No. of cigarettes per day
b) Time of stopping smoking

PREVIOUS HISTORY

- a. How old were you when you first began to bring up phlegm most days in winter? Age
- b. Did this come on gradually? Yes / No
- c. Did this come on after an illness? Yes / No
- d. Did this come on in some other way? Yes / No
If "yes" to c. or d. describe
.....

Onset of breathlessness:

- e. How old were you when you first noticed shortness of breath? Age
- f. Did this come on gradually? Yes / No
- g. Did this come on after an illness? Yes / No
- h. Did this come on in some other way? Yes / No
If "yes" to g. or h. describe
.....
- i. Did you have any cough or phlegm at all before you first became breathless? Yes / No
- j. If "no", ask: Not even a smoker's cough? Yes / No
- k. If "yes" to j: Did you bring up any phlegm with this cough? Yes / No

System of Grading.

| | | |
|--------------|---------|---------------------------------|
| <u>Cough</u> | Grade 0 | "No" to questions 1-4 (incl.) |
| | Grade 1 | "Yes" to questions 1 or 2 and 5 |
| | Grade 2 | "Yes" to questions 1-5 (incl.) |

| | | |
|---------------|---------|----------------------------------|
| <u>Phlegm</u> | Grade 0 | "No" to questions 6-10 (incl.) |
| | Grade 1 | "Yes" to questions 6 or 7 and 10 |
| | Grade 2 | "Yes" to questions 6-10 (incl.) |

For Cough and Phlegm grades 1 and 2 also required the answer "yes" to question 11.

Breathlessness

| | |
|---------|----------------------------------|
| Grade 1 | "No" to question 12-15 (incl.) |
| Grade 2 | "Yes" to question 12 |
| Grade 3 | "Yes" to questions 12 and 13 |
| Grade 4 | "Yes" to questions 12-14 (incl.) |
| Grade 5 | "Yes" to questions 12-15 (incl.) |

Chest Illnesses

| | |
|---------|--|
| Grade 0 | "No" to question 20 |
| Grade 1 | "Yes" to question 20 and history of one illness with increased phlegm with 1 weeks incapacity. |
| Grade 2 | "Yes" to question 20 and history of two or more illnesses with increased phlegm with 1 weeks incapacity. |

Smoking

| | |
|--------------------|------------------------------|
| Non-smoker | - never smoked at any time |
| Ex-smoker | - stopped smoking 3 months |
| Light smoker | - (0-14) cigarettes per day |
| Moderate smoker | - (15-24) cigarettes per day |
| Heavy smoker | - 24 cigarettes per day |

A P P E N D I X II.

Tables Of

Physical Characteristics and Symptomatology

Radiological Observations

F.E.V.₁, Static Lung Volumes and D_LCO

Arterial Blood Gases

Regional Ventilation and Perfusion Data

Physical Characteristics and Symptomatology in Patients in Group I.

| Patient | Sex | Age (yrs) | Ht. (ins) | Wt. (lbs.) | Grade B | Grade C/S | Chest Illness | Smoking Ops | Chest Ops | Age Onset B | First Symptoms |
|---------|------|-----------|-----------|------------|---------|-----------|---------------|-------------|-----------|-------------|----------------|
| F.B. | Male | 57 | 70 | 133 | 3 | 2/2 | 0 | Light | 0 | 50 | B |
| R.K. | Male | 56 | 68 | 120 | 5 | 1/1 | 0 | Mod. | 1 | 50 | C.S. |
| E.St. | Male | 62 | 68 | 126 | 5 | 1/1 | 0 | Ex.Light | 0 | 55 | B |
| G.B. | Male | 57 | 72 | 168 | 5 | 2/2 | 1 | Non | 1 | 44 | C.S. |
| H.B. | Male | 50 | 68 | 140 | 5 | 1/1 | 2 | Ex.Mod | 1 | 43 | B |
| J.E. | Male | 39 | 73 | 131 | 3 | 2/2 | 2 | Ex.Heavy | 1 | 34 | B |
| R.G. | Male | 46 | 64.5 | 120 | 3 | 2/1 | 1 | Heavy | 0 | 46 | C.S. |
| S.G. | Male | 58 | 67 | 118 | 5 | 1/1 | 2 | Mod. | 0 | 42 | C.S. |
| R.H. | Male | 49 | 67 | 143 | 5 | 2/2 | 2 | Ex.Light | 1 | 41 | C.S. |
| T.J. | Male | 60 | 70.5 | 108 | 5 | 1/1 | 2 | Ex.Mod | 2 | 45 | C.S. |

Ht. - Height (inches)
Wt. - Weight (lbs.)
B - Breathlessness

C - Cough
S.- Sputum
Ops.- Operations

Light - < 15 cigarettes per day
Mod - 15-24 cigarettes per day
Heavy - > 24 cigarettes per day

Physical Characteristics and Symptomatology in Patients in Group I.

| Patient | Sex | Age (yrs) | Ht. (ins) | Wt. (lbs) | Grade B | Grade C/S | Chest Illness | Smoking Ops. | Age Onset B | First Symptoms. |
|---------|------|-----------|-----------|-----------|---------|-----------|---------------|--------------|-------------|-----------------|
| T.F. | Male | 52 | 66 | 133 | 5 | 2/2 | 2 | Light | 0 | 49 |
| J.MCM | Male | 44 | 73 | 133 | 5 | 2/2 | 2 | Mod. | 1 | 40 |
| W.MCD | Male | 65 | 67.5 | 155 | 5 | 1/1 | 0 | Ex.Heavy | 0 | 60 |
| C.M. | Male | 53 | 73 | 138 | 5 | 2/1 | 2 | Ex.Light | 0 | 43 |
| T.P. | Male | 44 | 72 | 140 | 4 | 2/2 | 2 | Ex.Mod | 1 | 37 |
| L.Car | Male | 59 | 67 | 144 | 5 | 1/1 | 0 | Ex.Mod | 0 | 24 |
| A.S. | Male | 60 | 67.5 | 137 | 5 | 1/1 | 2 | Ex.Mod | 1 | 59 |
| G.P. | Male | 55 | 71 | 138 | 5 | 1/2 | 1 | Light | 0 | 48 |
| W.Sm | Male | 70 | 70.5 | 120 | 5 | 1/1 | 0 | Ex.Mod | 0 | 61 |
| J.K. | Male | 65 | 67 | 124 | 4 | 2/2 | 2 | Ex.Heavy | 0 | 60 |

* Key to abbreviations are at beginning of appendix.

Physical Characteristics and Symptomatology in Patients in Group I.

| Patient | Sex | Age (yrs) | Ht. (ins) | Wt. (lbs) | Grade B | Grade C/S | Chest Illness | Smoking | Chest Ops. | Age Onset B | First Symptoms |
|---------|------|-----------|-----------|-----------|---------|-----------|---------------|----------|------------|-------------|----------------|
| A.H. | Male | 61 | 70 | 154 | 4 | 1/1 | 0 | Ex Heavy | 0 | 60 | B |
| J.Her | Male | 62 | 71 | 136 | 5 | 2/0 | 2 | Ex Light | 0 | 58 | C.S. |
| N.S. | Male | 51 | 70.5 | 112 | 5 | 2/2 | 1 | Light | 0 | 37 | B |
| G.T. | Male | 48 | 65 | 94 | 5 | 2/2 | 2 | Ex Mod | 1 | 33 | C.S.B. |
| A.V. | Male | 57 | 69 | 142 | 5 | 2/1 | 1 | Ex Heavy | 0 | 50 | C.S.B. |
| S.B. | Male | 62 | 67 | 126 | 3 | 2/2 | 2 | Ex Mod | 0 | 60 | C.S.B. |
| C.B. | Male | 42 | 68.5 | 105 | 5 | 2/2 | 2 | Ex Heavy | 0 | 38 | B |
| T.C. | Male | 61 | 74 | 159 | 5 | 2/2 | 0 | Heavy | 0 | 56 | C.S. |
| R.F. | Male | 37 | 70.5 | 177 | 5 | 2/2 | 2 | Ex Heavy | 0 | 32 | C.S. |
| J.Hud | Male | 60 | 68 | 121 | 4 | 2/2 | 1 | Mod | 0 | 59 | C.S. |

Physical Characteristics and Symptomatology in Patients in Group I.

| Patient | Sex | Age (yrs) | Ht. (ins) | Wt. (lbs) | Grade B | Grade C/S | Chest Illness | Smoking | Chest Ops | Age Onset B | First Symptoms |
|---------|--------|-----------|-----------|-----------|---------|-----------|---------------|----------|-----------|-------------|----------------|
| L.Cun | Male | 66 | 67 | 109 | 5 | 2/2 | 1 | Light | 0 | 58 | C |
| D.C. | Male | 53 | 67 | 109 | 5 | 2/1 | 2 | Mod | 0 | 47 | C |
| A.M. | Male | 55 | 68 | 146 | 3 | 0/0 | 0 | Ex Heavy | 0 | 50 | C |
| J.C. | Male | 59 | 72.5 | 163 | 3 | 0/0 | 0 | Mod | 0 | 52 | B |
| G.F. | Male | 56 | 73.5 | 203 | 2 | 0/0 | 0 | Ex Heavy | 0 | 46 | B |
| G.H. | Female | 55 | 61 | 93 | 5 | 0/0 | 0 | Mod | 2 | 58 | B |
| A.V.M. | Male | 63 | 72 | 173 | 4 | 0/0 | 0 | Ex Mod | 1 | 50 | B |
| Geo.H. | Male | 53 | 64.5 | 147 | 3 | 0/0 | 2 | Ex Light | 1 | 47 | B |
| E.P. | Female | 55 | 60 | 92 | 5 | 0/0 | 1 | Ex Light | 1 | 40 | B |
| A.P. | Male | 40 | 67 | 153 | 5 | 0/0 | 0 | Ex Mod | 1 | 31 | B |

Physical Characteristics and Symptomatology in Patients in Group I.

| Patient | Sex | Age (yrs) | Ht. (ins) | Wt. (lbs) | Grade B | Grade C/S | Chest Illness | Smoking | Chest Ops | Age Onset B | First Symptoms |
|---------|--------|-----------|-----------|-----------|---------|-----------|---------------|----------|-----------|-------------|----------------|
| W.Sh | Male | 59 | 70 | 167 | 5 | 0/0 | 2 | Ex Light | 2 | 48 | B |
| J.W. | Male | 55 | 71 | 128 | 5 | 0/0 | 2 | Ex Mod | 1 | 45 | B |
| W.P. | Female | 44 | 69 | 146 | 5 | 0/0 | 0 | Ex Light | 1 | 30 | B |
| R.P. | Male | 60 | 74 | 134 | 1 | 0/0 | 1 | Heavy | 0 | - | - |
| P.J. | Male | 58 | 73 | 120 | 5 | 0/0 | 0 | Light | 0 | 48 | B |
| H.K. | Male | 54 | 71 | 176 | 5 | 0/0 | 0 | Ex Mod | 1 | 45 | B |
| E.Sh | Male | 64 | 71 | 140 | 4 | 0/0 | 0 | Ex Light | 1 | 59 | B |
| J.S. | Male | 54 | 71 | 132 | 2 | 0/0 | 0 | Ex Heavy | 0 | 52 | B |
| C.Mah | Male | 62 | 72 | 110 | 5 | 0/0 | 0 | Ex Mod | 0 | 42 | B |
| N.H. | Female | 42 | 65 | 96 | 4 | 0/0 | 0 | Ex Heavy | 0 | 29 | B |

Physical Characteristics and Symptomatology in Patients in Group II.

| Patient | Sex | Age (yrs) | Ht. (ins) | Wt. (lbs) | Grade B | Grade C/S | Chest Illnesses | Smoking Ops. | Chest Ops. | Age Onset B | First Symptoms |
|---------|--------|-----------|-----------|-----------|---------|-----------|-----------------|--------------|------------|-------------|----------------|
| E.N. | Male | 60 | 67 | 141 | 5 | 2/2 | 2 | Ex Mod | - | 56 | C.S. |
| L.M. | Male | 64 | 65 | 158 | 5 | 2/2 | 2 | Light | - | 62 | C.S. |
| E.G. | Male | 53 | 71.5 | 180 | 3 | 2/2 | 2 | Heavy | - | 48 | C.S. |
| F.K. | Male | 62 | 68 | 126 | 3 | 2/2 | 2 | Mod | - | 58 | C.S. |
| R.T. | Male | 48 | 68.5 | 111 | 5 | 2/2 | 2 | Ex Heavy | - | 37 | C.S. |
| E.Sc | Female | 60 | 61.5 | 114 | 5 | 2/2 | 2 | Non | - | 41 | C.S.B. |
| E.P. | Female | 51 | 63 | 157 | 5 | 2/2 | 2 | Mod | - | 49 | C.S. |
| F.R. | Male | 59 | 68.5 | 145 | 3 | 2/2 | 0 | Light | - | 56 | C.S. |

KEY TO ABBREVIATIONS IN RADIOLOGICAL TABLES.

Diaphragm Sh - Shape (F-flat, C-curved).
Ll - Level (no. of anterior ribs).
Move L/R- Movement of left and right
hemidiaphragms (cms.).

RsS - Retrosternal space.
Dp - depth (cms.)
D.E. - downward extension (cms.)

T.C./T.H. - Transcardiac and transhilar
diameters.

P.A. - Pulmonary artery } dilated (dil)
H.A. - Hilar artery } or normal (N)

Comp. - Compression
fiss. - fissure displacement
V.Cr. - vessel crowding
* N - Normal size vessels, no vessel loss.
- Small size vessels, vessel loss.
+ - Dilated vessels
Shown in Zones.

Dem. Bullae - Demarcated bullae lobe
and diameters (cms.)

Rad. Diagnosis- Gen. - Generalised emphysema
Ext.loc. - Extensive localised
emphysema.
Loc. - Localised emphysema.

a - before operation.
b - after operation.

Radiological Observations and Measurements in Patients in Group I.

| Patient | DIAPHRAGM Sh./LJ | Move L/R | ReS. Dp/D.E. | T.C/T.H. diameter | PA/HA | Comp. Riss/V.Cr | Mid Lung* Vessels | Vessel* loss | Dem. Bullae | Rad. Diagnosis |
|---------|---------------------|----------|-----------------|----------------------|-------|--------------------|----------------------|-----------------|----------------|--|
| F.B. | C.7. | 0/1.0 | 4.0/7.0 | 11.0/9.5 | dil/N | nil | — | — | — | Ext. loc. |
| R.K. a) | F.7. | 1.0/1.0 | 2.0/7.0 | 10.0/10.0 | dil/N | nil | — | — | — | Gen. |
| b) | F.7. | 2.0/3.5 | 2.0/7.0 | 11.0/10.5 | dil/N | nil | — | — | — | Gen. |
| E.St. | LF 64 RC | 0.5/1.0 | — | 12.0/10.0 | N/N | + / + | — | — | — | RLL, 10x10 LLL, 8x4 17x4 Ext. loc. |
| G.B. | F.7. | 0.5/0.5 | 4.0/5.0 | 11.5/NM | dil/N | + / + RML | N | N | — | Gen. |
| H.B. a) | F.7. | 1.0/2.5 | 2.5/6.0 | 12.5/9.5 | N/N | nil | N | N | N | Normal |
| b) | F.7. | 0/2.5 | 2.5/6.0 | 12.0/9.5 | N/N | nil | N | N | N | Normal |

Radiological Observations and Measurements in Patients in Group I.

| Patient | DIAPHRAGM Sh/LL | Move L/R | RsS. Dp/D.E. | T.C/T.H. diameter | PA/HA | Comp. fiss/V.Cr | Mid lung Vessels | Vessel loss | Dem. Bullae | Rad. Diagnosis |
|---------|--------------------|----------|-----------------|----------------------|---------|--------------------|---------------------|----------------|------------------------|-----------------------|
| J.E. a) | C.7. | - | 4·0/12·0 | 9·0/9·5 | dil/N | N N - | N N - | N N - | nil | Loc. |
| b) | F.6½ | 3·0/2·0 | 4·0/12·0 | 9·0/9·5 | dil/N | N N - | N N - | N N - | nil | Loc. |
| R.G. | LF 6½ RC | 2·5/1·9 | 4·0/6·0 | 11·5/11·0 | dil/N | N - | N - | N - | L.I.L. 4 x 12 | Ext. Loc. |
| S.G. | LF 7½ RC | 3·5/2·5 | 5·5/1·5 | 11·5/10·0 | N/N | N N - | N - | N - | L.U.L. 13 x 6 | Ext. Loc. |
| R.H. a) | F.7. | - | 6·0/2·0 | 13·5/11·0 | N/dil | + /+ RML | + N | N N - | Gen. Kyphosis | I.L.L. 4x4 |
| b) | F 6½ | 2·5/1·5 | 6·0/2·0 | 14·0/11·0 | N/dil | + /+ Ling | + + | N N - | nil | Gen. Kyphosis |
| T.J. | F.6. | 0/0·5 | 5·5/4·0 | 11·5/ | N/dil | + /+ RUL | - | N N - | RLL 10x10 RUL 10x10 | Gen. |
| T.F. | F 6½ | - | - | 15·5/13·5 | dil/dil | nil | + N - | N N - | RLL 11x12 | Loc. Pulm. Hyper * |
| J.M.M. | F 6½ | 0/0 | 6·3/3·0 | 10·5/ | N/N | + /+ Ling | N N - | N N - | LLL 26x6 RUL 7x7 | Gen. Old T.B. |
| W.M.D. | C 7½ | 1·0/1·0 | 2·5/5·0 | 14·0/13·0 | dil/dil | + /+ RUL | - N - | - N - | nil | Loc. Pulm. Hyper |

* Pulm. Hyper. - Pulmonary Hypertension.

| Patient | DIAPHRAGM Sh/Ll | Move L/R | RsS. Dp/D.E. | T.C/T.H. diameter | PA/ HA | Comp. fiss/V.Cr. | Mid Lung Vessels | Vessel loss | Dem. Bullae | Rad. Diagnosis |
|---------|--------------------|----------|-----------------|----------------------|-----------|---------------------|---------------------|----------------|----------------------|-----------------------|
| C.M. | F.7. | 0.5/0.5 | 5.0/2.0 | 11.5/11.0 | dil/dil | nil | - - | - N - | nil | Gen. |
| T.P. a) | FL 6 RC | 1.0/1.0 | 5.0/3.5 | 12.5/12.0 | dil/dil | nil | N N N | - N - | LLL 15x15 | Ext. Loc. |
| b) | C 6½ | 1.5/3.0 | 3.0/4.0 | 13.0/12.5 | dil/dil | nil | N N N | - N - | LLL 4x4 | Pulm. hypert. |
| L.Car. | C.7. | 2.0/1.5 | 4.5/5.0 | 11.0/12.0 | N/N | + /+ RLL | - - | - - | nil | Ext. Loc. |
| A.S. a) | F.7½ | 0/0 | 3.0/6.0 | 9.0/11.0 | N/dil | RML | N N N | N N - | RLL 10x10 | 20x20 Loc. |
| b) | F.7. | 2.5/2.0 | 3.0/6.0 | 11.5/11.5 | N/dil | nil | N N N | N N - | RLL 12x12 | Loc. |
| G.P. | C.7½ | 1.5/1.5 | 4.0/4.0 | 10.5/10.5 | N/N | nil | - N N | - N N | nil | Loc. |
| W.Su. | F.7. | 1.5/2.0 | 3.5/2.0 | 10.0/10.5 | dil/dil | nil | - - | - - | nil | Gen. |
| J.K. | F.7. | 0.5/0.5 | 4.5/4.0 | 14.5/13.5 | N/dil | + /+ RLL | - + | - N - | nil | Gen. Pulm. Hypert. |
| A.H. | C.6½ | 5.0/5.0 | 4.5/4.0 | 14.0/12.0 | N/N | nil | N N - | - N - | LUL 12x12 RUL 4x4 | Loc. |

Radiological Observations and Measurements in Patients in Group I.

| Patient | Sh/LL. | DIAPHRAGM Move I/R | ReS. Dp/D.E. | T.C/T.H. diameter | PA/HA | Comp. fiss/V.Cr | Mid Lung Vessels | Vessel loss | Dem. Bullae | Rad. Diagnosis |
|---------|--------|--------------------|--------------|-------------------|----------------------------|-----------------|------------------|-------------|-------------|--------------------|
| J.H. | F.6½ | 2·5/1·5 | 6·5/0 | 12·5/12·0 | dil/dil | nil | + | N | N | RLL 5 x 5 Gen. |
| N.S. | F.6½ | 2·0/1·0 | 5·5/0 | 9·5/- | N/N | nil | - | N | LLL - | Ext. Loc. Old T.B. |
| A.V. | F.7. | - | 4·0/0 | 11·0/9·0 | N/dil | nil | - | + | N N | nil Gen. |
| G.T. | F.8 | 1·0/1·0 | 6·0/0 | 10·5/12·0 | dil/dil | nil | - | N | LLL - | 15 x 15 Gen. |
| S.B. | | | | | Radiographs not available. | | | | | |
| C.B. | F.6½ | 2·0/1·5 | 2·0/5·0 | 9·5/9·5 | dil/N | nil | - | N | N | nil Gen. |
| T.C. | F.7. | - | 6·0/1·0 | 12·5/14·0 | dil/dil | nil | - | + | + | nil Gen. |
| R.F. | F.6½ | 3·5/3·0 | 5·5/1·0 | 13·5/11·5 | N/N | nil | N N | N N | N N | nil Loc. |
| J. Hud. | C.6. | 1·5/2·5 | 1·5/7·0 | 14·0/9·5 | N/N | nil | N | N | N N | RLL 4·5 x 5·0 Loc. |
| L. Cun. | F.7. | 0·5/0·5 | 4·0/1·0 | 9·0/10·5 | N/dil | nil | - | N | N | nil Gen. |

| Patient | Sh/L1 | DIAPHRAGM Move L/R | BS Dp/D.E. | T.C/T.H. Diameter | PA/HA fiss/V.Cr. | Comp. fiss/V.Cr. | Mid Lung Vessels | Vessel loss | Dem. Bulbae | Rad. Diagnosis |
|--------------|-------------|-----------------------|---------------|----------------------|---------------------|---------------------|---------------------|----------------|----------------|-------------------|
| D.C. | LF 7½ RC | 0·5/3·0 | 3·0/8·0 | 10·0/9·0 | N/N | nil | — | — | LUL 13x13 | Ext. Loc. |
| A.M. | C.7. | 3·0/3·0 | 2·5/- | 13·0/11·0 | N/N | nil | N/N | — | LLL 10x10 | Loc. |
| J.C. | C.6½ | 4·5/4·5 | 4·5/6·0 | 14·5/11·5 | dil/dil | +/+ | — | — | — | Ext. Loc. |
| G.F. | F.6½ | 4·0/4·5 | 2·5/3·5 | 15·5/12·0 | N/N | +/+ RAL | — | — | — | nil |
| G1.H a) | F.7. | 0·5/0·5 | 2·5/5·0 | 11·5/ 8·5 | dil/N | nil | — | — | RUL 10x10 | Loc. |
| b) | F.7. | 1·0/1·0 | 3·0/4·5 | 10·5/9·5 | dil/N | nil | — | — | — | Gen. |
| c) | F.7. | 1·0/0·5 | 3·0/4·5 | 11·0/11·0 | dil/N | nil | — | — | RLL 6x6 | Gen. |
| A.V.M. a) | C.6½ | 1·0/3·0 | — | 13·5/10·5 | N/N | nil | N/N | — | RLL 9x9 | Loc. |
| b) | C.6½ | 4·0/3·0 | — | 13·5/10·5 | N/N | nil | N/N | — | RLL 9x9 | Loc. |

Radio logical Observations and Measurements in Patients in Group I.

| Patient | Sh/LI | DIAPHRAGM Move L/R | ReS. Dp/D.E. | T.C/T.H. diameter | PA/HA | Comp. fiss/V.Cr vessels | Mid Lung vessels | Vessel loss | Dem. Bullae | Rad. Diagnosis |
|---------|----------------|-----------------------|-----------------|----------------------|---------|-------------------------------|---------------------|----------------|----------------|-------------------|
| Geo.H | a) LF 6½ RC | 3·5/2·0 | 4·5/5·5 | 13·0/ | N/N | nil/+LLL | — | — | LLL | 18x18 Loc. |
| b) | LF 6½ RC | 0·5/1·5 | 4·5/5·0 | 11·5/ | dil/dil | +/+LLL | — | — | — | |
| E.P. | a) F.7. | 0·5/0·5 | 3·5/5·0 | 9·0/10·0 | dil/dil | nil | — | — | — | Gen. |
| b) | F.6½ | 2·0/1·4 | 3·5/5·0 | 8·5/10·5 | dil/dil | nil | + | + | — | Gen. |
| A.R. | C.7. | 0·5/0·5 | 7·0/7·0 | 14·0/11·0 | dil/N | nil | — | — | — | Ext. Loc. |
| M.Sh | a) F.6½ | 1·0/0·5 | 5·0/0 | 11·0/9·5 | dil/N | nil | — | — | — | Gen. |
| b) | F.6½ | 0·5/0·5 | 5·0/1·0 | 13·0/9·5 | dil/N | nil | — | — | RUL | 5x5 Gen. |
| J.W. | a) F.6½ | 1·0/1·0 | 5·0/5·0 | 10·0/11·5 | N/N | nil | — | — | — | Gen. |
| b) | F.7. | 3·0/2·0 | 5·0/5·0 | 9·5/ | N/N | nil | — | — | RUL | 7x7/7x7 Gen. |

| Patient | Sh/Ll | DIAPHRAGM | RsS Move L/R | T.C/T.H. Dp/D.E. | PA/HA | Comp. fiss/V.Cr Vessels | Mid Lung Vessel loss | Dem. Bullae | Rad. Diagnosis |
|---------|-------|-----------|-----------------|---------------------|---------|-------------------------------|----------------------------|---------------------------------|------------------------|
| W.P. a) | C.6½ | 4.0/4.0 | 2.0/4.5 | 12.5/10.0 | N/N | nil | N N - | RLL 8 x 8 | Ext. Loc. |
| b) | C.6½ | 4.0/3.5 | 3.0/ | 12.5/10.0 | N/N | nil | N N - | nil | Ext. Loc. |
| R.P. | C.7. | 3.5/3.5 | 4.5/1.0 | 11.5/11.5 | N/dil | nil | N N - | nil | Loc. |
| P.J. | F.7. | 0.5/2.0 | 5.0/1.0 | 11.0/10.5 | dil/dil | nil | - - - | - | |
| H.K. | C.6. | 3.0/2.0 | 4.5/0 | 12.0/10.5 | N/N | +/+LLL | N N - | nil | Kyphosis |
| E.Sh a) | F.6½ | 2.0/0.5 | 2.5/1.0 | 12.0/ | N/N | nil | - - - | - | Multiple both lungs |
| b) | C.6 | 3.5/7.0 | 2.5/1.5 | 13.0/10.5 | N/N | nil | - - - | RUL 5x5 RLL 15x12 LLL 8x8 | Ext. Loc. |
| J.S. | C.6½ | 3.0/3.5 | 2.5/5.0 | 9.5/10.0 | N/dil | nil | - - - | RML 7x7 RLL 6x6 LLL 8x8 | Gen. |
| C.Mah | F.6½ | 1.5/2.0 | 2.0/3.0 | 9.5/11.0 | N/dil | nil/+ RLL | - - - | RLL 12x12 LLL 3x3 | Gen. |
| N.H. | F.7. | 0/3.5 | 3.0/0 | 9.0/9.5 | dil/dil | +/+ RUL | - - - | RLL 20x20 LLL 23x15 | Gen. |

Radiological Observations and Measurements in Patients in Group III.

| Patient | Sh/L1 | Move L/R | RsS. Dp/D.E. | Diameter T.C/T.H. | PA/HA | Comp. fiss/V.Cr Vessels | Mid lung Vessel Loss | Vessel Loss | Dem. Bullaæ | Rad. Diagnosis |
|---------|-------|----------|-----------------|----------------------|---------|-------------------------------|----------------------------|----------------|----------------|--------------------------------|
| E.N. | C/6½ | ½/0 | 2½/ 6 | 12·5/10·5 | N/N | nil | N N N N | N N N N | nil | Normal |
| L.M. | C/6½ | ¾/4 | 4/7 | 13·0/9·5 | N/N | nil | N N N N | N N N N | nil | Normal Kyphosis |
| E.G. | C/7 | ½/½ | 3/7 | 16·0/14·0 | N/dil | nil | N N N N | N N N N | nil | Normal |
| F.K. | F/7½ | 2/3 | 2½/5 | 15·5/12·0 | N/dil | nil | N N N N | N N N N | nil | Normal |
| R.T. | C/6½ | ½/½ | 3/10 | 12·5/0·5 | dil/N | nil | N N N N | N N N N | nil | Normal |
| E.S. | C/6 | 2/2 | 5/5 | 12·5/11·0 | dil/dil | nil | N N N N | N N N N | nil | Normal Pulmonary Hypert. |
| E.pat. | C/6½ | ¾/1 | 2½/10 | 14·0/- | dil/dil | nil | N N N N | N N N N | nil | Normal Pulmonary Hypert. |
| F.R. | C/6 | 3/3 | 2/8 | 12·5/11·5 | dil/N | nil | N N N N | N N N N | nil | Normal |

F.E.V.₁ Static Lung Volumes, Airways Resistance and
Diffusing Capacity in Patients in Group I.

| Patient | FEV ₁ | FEV ₁ / FVC% | V.C. | FRC | RV | RV/TLC% | A.R. | D _{LCO} | REST M.V. | %Extr. | EXERCISE D _{LCO} | M.V. |
|---------|------------------|----------------------------|------|------|------|---------|------|------------------|--------------|--------|------------------------------|------|
| F.B. | 2000 | 46 | 4.50 | 7.55 | 5.65 | 56 | - | 5.7 | 11.1 | 23 | 8.4 | 21.8 |
| R.K. a) | 600 | 25 | 3.60 | 6.03 | 4.42 | 55 | 4.85 | 7.8 | 10.0 | 30 | 11.5 | 18.0 |
| b) | 600 | 23 | 3.90 | 5.23 | 3.53 | 48 | 5.37 | 6.8 | 7.2 | 35 | 13.2 | 27.6 |
| E.St. | 600 | 25 | 2.65 | 5.00 | 4.15 | 61 | 5.05 | 6.9 | 10.5 | 26 | 8.7 | 19.8 |
| G.B. | 550 | 34 | 2.00 | 4.70 | 4.10 | 67 | - | 6.8 | 12.2 | 25 | - | - |
| H.B. a) | 600 | 25 | 2.90 | 3.66 | 2.36 | 45 | - | 7.4 | 8.9 | 34 | 11.7 | 20.8 |
| b) | 800 | 26 | 3.00 | 4.45 | 3.45 | 53 | 6.15 | 6.03 | 8.25 | 30 | 9.96 | 16.1 |

FEV₁ Forced Expired Vol. in 1 sec. (ml.)

FVC Forced Vital Capacity (ml.)

VC Vital Capacity

FRC Functional Residual Capacity (litres)

RV Residual Volume

A.R. Airways Resistance (cm₅H₂O/litre)

D_{LCO} Diffusing Capacity for Carbon Monoxide

(ml.CO/mm.Hg/min.)

M.V. Minute Ventilation (litres)

% Extr. Percentage Extraction of Carbon Monoxide

F.E.V.₁ Static Lung Volume, Airways Resistance and
Diffusing Capacity in Patients in Group I.

| Patient | FEV ₁ ml. | FEV ₁ % FVC | V.C. | FRC | RV | RV/TLC % | A.R. | D _{LCO} | REST M.V. | % Extr. | D _{LCO} | EXERCISE M.V. |
|---------|-------------------------|---------------------------|------|------|------|----------|------|------------------|--------------|---------|------------------|------------------|
| J.E. a) | 1050 | 29 | 3.80 | 5.31 | 3.81 | 50 | 3.24 | 7.1 | 7.8 | 37 | 15.3 | 32.2 |
| b) | 1200 | 35 | 3.75 | 4.92 | 3.72 | 50 | 2.94 | 4.9 | 6.6 | 30 | 11.0 | 24.0 |
| R.G. | 2000 | 59 | 3.80 | 4.05 | 2.65 | 41 | 2.15 | 12.0 | 13.3 | 26 | 12.0 | 25.0 |
| S.G. | 350 | 27 | 1.80 | 5.61 | 4.71 | 72 | 8.06 | 6.6 | 7.6 | 32 | 9.6 | 12.6 |
| R.H. a) | 1050 | 42 | 2.15 | 5.48 | 4.98 | 70 | - | 6.0 | 11.0 | 25 | 7.1 | 16.8 |
| b) | 400 | 29 | 2.30 | 6.15 | 5.30 | 70 | - | 5.8 | 8.1 | 29 | 8.6 | 15.3 |
| T.J. | 500 | 29 | 2.85 | 5.91 | 4.26 | 60 | - | 5.9 | 9.2 | 28 | - | - |
| T.F. | 700 | 41 | 2.00 | 3.32 | 2.19 | 52 | 7.80 | 5.5 | 10.9 | 23 | 8.1 | 15.3 |
| J.McM. | 350 | 37 | 1.60 | 7.40 | 7.10 | 82 | - | 7.7 | 7.1 | 40 | 9.3 | 9.0 |
| W.McD. | 600 | 25 | 2.50 | 7.66 | 6.56 | 73 | - | 8.2 | 12.0 | 38 | - | - |

F.E.V.₁, Static Lung Volumes, Airways Resistance and
Diffusing Capacity in Patients in Group I.

| Patient | mL. | FEV ₁ FVC % | V.C. | FRC | RV | RV/TLC% | A.R. | D _{LCO} | M.V. | % Extr. | D _{LCO} | EXERCISE M.V. |
|---------|------|---------------------------|------|------|------|---------|-------|------------------|------|---------|------------------|------------------|
| C.M. | 400 | 22 | 2.00 | 8.40 | 7.70 | 80 | - | 9.0 | 8.6 | 29 | 18.2 | 12.6 |
| T.P. a) | 900 | 33 | 3.20 | 5.12 | 4.50 | 59 | 4.08 | 9.1 | 11.6 | 30 | 16.4 | 18.4 |
| b) | 750 | 26 | 3.30 | 4.65 | 4.15 | 58 | 5.28 | 7.0 | 7.5 | 39 | 13.4 | 15.0 |
| L.Car | 700 | 28 | 2.60 | 6.87 | 5.93 | 70 | 10.95 | 5.7 | 8.0 | 26 | 12.1 | 16.8 |
| A.S. a) | 500 | 30 | 2.20 | 6.44 | 5.84 | 73 | 7.38 | 7.2 | 8.6 | 33 | - | - |
| b) | 600 | 33 | 2.50 | 6.25 | 5.05 | 65 | 4.83 | 8.4 | 9.7 | 33 | 10.4 | 16.8 |
| G.P. | 550 | 28 | 3.20 | 7.10 | 5.90 | 65 | 3.64 | 8.6 | 8.7 | 33 | 15.2 | 16.8 |
| W.Sa | 500 | 42 | 1.65 | 7.65 | 7.10 | 81 | - | 6.1 | 8.6 | 27 | 7.1 | 11.4 |
| J.K. | 800 | 37 | 2.20 | 5.78 | 4.85 | 69 | - | 6.1 | 11.3 | 26 | 7.6 | 17.1 |
| A.H. | 2300 | 58 | 4.30 | 4.68 | 3.25 | 43 | 4.6 | 8.5 | 12.6 | 26 | 19.9 | 22.5 |

F.E.V.₁, Static Lung Volumes, Airways Resistance and
Diffusing Capacity in Patients in Group I.

| Patient | F.E.V. ₁ ml. | FEV ₁ ,% FVC | V.C. | FRC | R.V | RV/TLC % | A.R. | D _{LCO} | REST M.V. | % Extr. | D _{LCO} | EXERCISE M.V. |
|---------|----------------------------|----------------------------|------|------|------|----------|------|------------------|--------------|---------|------------------|------------------|
| J.Her | 600 | 27 | 2.10 | 4.60 | 3.90 | 65 | 6.45 | 7.1 | 10.2 | 28 | 11.4 | 21.6 |
| N.S. | 900 | 36 | 2.45 | 4.94 | 4.04 | 62 | 4.60 | 6.6 | 7.5 | 33 | 9.7 | 15.4 |
| G.T. | 300 | 25 | 1.60 | 6.53 | 5.93 | 79 | 5.60 | 4.1 | 5.8 | 32 | - | - |
| A.V. | 550 | 24 | 2.80 | 7.80 | 6.85 | 71 | 4.53 | 5.7 | 7.5 | 32 | 5.8 | 10.5 |
| S.B. | 1600 | 64 | 2.50 | 3.72 | 2.82 | 53 | 4.68 | 11.4 | 12.3 | 32 | 19.2 | 20.0 |
| C.B. | 600 | 27 | 2.20 | 4.58 | 3.58 | 62 | 7.20 | 6.0 | 8.5 | 29 | 11.6 | 18.4 |
| T.C. | 1000 | 39 | 3.05 | 9.80 | 8.75 | 74 | - | 12.6 | 16.8 | 28 | - | - |
| R.F. | 1000 | 23 | 3.70 | 5.41 | 3.71 | 50 | 3.00 | 11.2 | 13.4 | 32 | 14.0 | 23.0 |
| J.Hud | 800 | 37 | 2.50 | 3.72 | 2.62 | 51 | 6.00 | 8.1 | 13.4 | 25 | 11.4 | 22.4 |
| L.Cun | 500 | 46 | 1.40 | 5.65 | 5.27 | 79 | - | 4.6 | 7.5 | 30 | - | - |

F.E.V.₁ Static Lung Volumes, Airways Resistance and
Diffusing Capacity in Patients in Group I.

| Patient | F.E.V. ₁ ml. | FEV ₁ % FVC | V.C. | FRC | R.V | RV/TLC% | A.R. | D _{LCO} | REST M.V. | %Extr. | D _{LCO} | EXERCISE M.V. |
|--------------|----------------------------|---------------------------|------|------|------|---------|-------|------------------|--------------|--------|------------------|------------------|
| D.C. | 350 | 19 | 1.80 | 5.22 | 4.42 | 71 | 10.40 | 10.4 | 8.8 | 33 | 11.0 | 14.4 |
| A.M. | 1100 | 48 | 3.15 | 4.50 | 3.50 | 53 | 3.05 | 13.5 | 14.4 | 36 | 17.7 | 25.6 |
| J.C. | 800 | 34 | 2.95 | 5.87 | 4.97 | 63 | 4.45 | 4.9 | 8.7 | 24 | 8.1 | 17.1 |
| G.F. | 1900 | 45 | 4.70 | 5.45 | 4.35 | 48 | 2.15 | 11.3 | 10.3 | 36 | 24.2 | 32.0 |
| G1.H. a) | 300 | 30 | 1.40 | 4.85 | 4.60 | 77 | — | 5.7 | 5.2 | 40 | — | — |
| b) | 400 | 22 | 2.10 | 4.39 | 3.49 | 63 | 11.80 | 5.3 | 7.1 | 31 | — | — |
| c) | 500 | 23 | 2.20 | 3.75 | 2.86 | 56 | 4.90 | 4.8 | 8.6 | 26 | 6.0 | 13.8 |
| A.V.M. a) | 900 | 38 | 4.40 | 4.30 | 2.70 | 38 | 3.80 | 7.4 | 12.1 | 26 | 11.0 | 22.0 |
| b) | 1200 | 38 | 3.60 | 4.05 | 2.95 | 45 | 3.70 | 8.5 | 12.0 | 29 | 12.9 | 26.4 |

F.E.V.₁, Static Lung Volumes, Airways Resistance and
Diffusing Capacity in Patients in Group I.

| Patient | F.E.V. ₁ ml. | FEV ₁ / FVC | V.C. | FRC | R.V. | RV/TLC | A.R. | D _{LCO} | REST M.V. | %EXTR. | D _{LCO} EXERCISE | M.V. |
|-----------|----------------------------|---------------------------|------|------|------|--------|------|------------------|--------------|--------|------------------------------|------|
| Geo.H. a) | 800 | 24 | 3.40 | 4.50 | 3.70 | 52 | - | 10.9 | 8.4 | 38 | 19.8 | 22.2 |
| b) | 600 | 24 | 3.10 | 5.97 | 4.87 | 61 | - | 10.5 | 10.6 | 34 | 10.8 | 12.8 |
| E.P. a) | 250 | 23 | 1.20 | 4.45 | 3.98 | 78 | - | 3.0 | 6.8 | 22 | - | - |
| b) | 300 | 23 | 1.15 | 4.00 | 3.60 | 78 | - | 4.9 | 6.2 | 29 | - | - |
| A.P. | 600 | 21 | 2.90 | 6.54 | 5.24 | 64 | - | 12.6 | 7.8 | 48 | 19.8 | 18.8 |
| W.Sh a) | 1000 | 33 | 3.25 | 4.59 | 3.54 | 52 | - | 6.0 | 11.0 | 25 | 8.3 | 20.0 |
| b) | 1100 | 28 | 4.45 | 4.85 | 2.85 | 42 | 3.17 | 5.2 | 10.6 | 23 | 9.2 | 22.4 |
| J.W. a) | 500 | 28 | 1.80 | 7.26 | 6.46 | 78 | 4.26 | 6.6 | 12.6 | 23 | - | - |
| b) | 500 | 30 | 2.20 | 8.80 | 8.00 | 78 | - | 4.0 | 7.9 | 24 | 5.7 | 12.0 |

F.E.V.₁, Static Lung Volumes, Airways Resistance and Diffusing Capacity in Patients in Group I.

| Patient | F.E.V. ₁ ml. | FEV ₁ % FVC | V.C. | FRC | R.V. | RV/TLC % | A.R. | D _L C _O | REST M.V. | % Extr. | D _L C _O | EXERCISE M.V. |
|----------|----------------------------|---------------------------|------|------|------|----------|------|-------------------------------|--------------|---------|-------------------------------|------------------|
| W.P. a) | 600 | 40 | 2.40 | 4.41 | 4.21 | 64 | - | 11.8 | 14.5 | 33 | 15.7 | 19.8 |
| b) | 600 | 33 | 2.60 | 4.23 | 3.53 | 58 | - | 10.2 | 9.1 | 39 | 12.9 | 24.3 |
| R.P. | 2300 | 56 | 3.85 | 5.46 | 3.91 | 50 | 1.50 | 8.5 | 9.5 | 30 | 13.3 | 15.6 |
| P.J. | 900 | 32 | 2.60 | 9.00 | 7.85 | 75 | - | 4.7 | 7.9 | 25 | 5.9 | 13.6 |
| H.K. | 800 | 25 | 4.20 | 4.39 | 2.59 | 38 | - | 11.4 | 14.2 | 33 | 14.4 | 20.8 |
| E.Sh. a) | 900 | 43 | 2.40 | 3.08 | 2.32 | 48 | 4.05 | 7.7 | 11.6 | 28 | 9.9 | 21.0 |
| b) | 1300 | 33 | 3.70 | 4.38 | 3.20 | 47 | 4.52 | 6.7 | 9.5 | 29 | 12.0 | 19.0 |
| J.S. | 900 | 26 | 4.30 | 5.02 | 3.90 | 48 | 5.20 | 8.6 | 6.4 | 38 | 12.9 | 18.0 |
| C.Mah | 800 | 25 | 3.20 | 7.75 | 6.05 | 62 | - | 5.0 | 15.5 | 17 | 4.5 | 19.2 |
| N.H. | 500 | 46 | 1.25 | 3.11 | 2.98 | 70 | - | 8.1 | 9.0 | 34 | 9.4 | 12.5 |

F.E.V.₁ Static Lung Volumes, Airways Resistance and Diffusing Capacity
in Patients from Group II

| Patient | F.E.V. ₁ (ml) | F.E.V. ₁ F.V.C. | V.C. [†] F.V.C. | F.R.C. [‡] | R.V. ⁺ | RV/TLC% | A.R. [†] | DLCO [†] | M.V. ⁺ | EXERCISE | | |
|---------|-----------------------------|-------------------------------|-----------------------------|---------------------|-------------------|---------|-------------------|----------------------|-------------------|-------------------|------|------|
| | | % | | | | | DLCO [†] | % Extr. [†] | DLCO [†] | M.V. ⁺ | | |
| E.N. | 600 | 40 | 2.00 | 5.45 | 4.95 | 71 | — | 7.9 | 8.0 | 35 | 8.6 | 13.7 |
| L.M. | 800 | 36 | 2.50 | 4.61 | 3.91 | 61 | 3.77 | 11.4 | 14.7 | 30 | 16.7 | 28.9 |
| E.G. | 1500 | 52 | 2.90 | 5.00 | 4.38 | 65 | 3.12 | 17.1 | 18.0 | 32 | 32.0 | 32.2 |
| F.K. | 600 | 32 | 2.45 | 5.08 | 4.18 | 63 | 5.10 | 4.8 | 5.6 | 29 | 6.6 | 9.4 |
| R.T. | 450 | 32 | 2.35 | 4.46 | 4.04 | 63 | — | 5.9 | 9.4 | 27 | 6.1 | 12.5 |
| E.Sc. | 500 | 33 | 1.60 | 3.90 | 3.30 | 67 | — | 7.3 | 8.4 | 31 | — | — |
| E. Pat. | 500 | 29 | 1.80 | 5.15 | 4.65 | 72 | 7.10 | 6.9 | 6.6 | 39 | 7.3 | 9.2 |
| F.R. | 1000 | 48 | 2.85 | 4.70 | 4.10 | 59 | 3.82 | 8.7 | 7.8 | 33 | 18.8 | 17.6 |

F.E.V.₁ Forced Expired Vol. in 1 sec. (ml.)

F.V.C. Forced Vital Capacity (ml.)

Vital Capacity

Functional Residual Capacity litres.

Residual Volume.

A.R. Airways Resistance (cmH₂O/litre)

DLCO Diffusing Capacity for Carbon

Monoxide (ml.CO/mm.Hg/min)

M.V. Minute Ventilation (litres)

% Extr. Percentage Extraction of Carbon

Monoxide.

F.E.V.₁, Static Lung Volumes, Airways Resistance and Diffusing Capacity in Patients in Group III.

| Patient | F.E.V. ₁ ml. | $\frac{FEV_1}{FVC} \%$ | V.C. | FRC | R.V. | RV/TLC% | A.R. | L.V.* | D _{LCO} | M.V. | REST M.V. | % Extr. | D _{LCO} | EXERCISE M.V. |
|---------|----------------------------|------------------------|------|------|------|---------|------|-------|------------------|------|--------------|---------|------------------|------------------|
| M.E. | 1800 | 69 | 2.60 | 2.18 | 1.78 | 41 | 3.85 | 2.01 | 17.7 | 11.0 | 49 | 17.9 | 15.0 | |
| A.K. | 1000 | 38 | 3.10 | 4.03 | 3.78 | 55 | 8.30 | 7.65 | 9.4 | 13.0 | 30 | 15.2 | 21.6 | |
| A.S. | 900 | 31 | 3.50 | 4.90 | 3.20 | 48 | 3.70 | 5.92 | 8.9 | 8.7 | 37 | 12.0 | 19.0 | |
| E.L. | 650 | 32 | 2.40 | 4.28 | 3.58 | 60 | 4.25 | 6.32 | 10.4 | 7.0 | 44 | 13.2 | 15.4 | |
| D.G. | 1000 | 40 | 3.50 | 4.81 | 4.06 | 54 | 5.20 | 8.10 | 9.1 | 10.1 | 39 | 27.8 | 27.6 | |
| M.D. | 900 | 37 | 2.80 | 4.32 | 3.32 | 54 | 9.54 | 5.30 | 20.9 | 10.4 | 48 | 22.4 | 20.9 | |
| H.S. | 1200 | 67 | 2.05 | 2.12 | 1.72 | 44 | 4.04 | 2.42 | 9.8 | 11.3 | 33 | 13.4 | 23.8 | |

* Denotes Lung Volume while panting in body plethysmograph (litres).

in Group I*.

| Patient | PaO ₂ rest mm. Hg. | PaO ₂ Ex. mm. Hg. | PaCO ₂ rest mm. Hg. | PaCO ₂ Ex. mm. Hg. | pH | % Satn. Hb | V _D /V _T | V _A l/min. | RESTING VALUES | Work Kgm/mins. |
|--------------------|----------------------------------|---------------------------------|-----------------------------------|----------------------------------|--------------|---------------|--------------------------------|--------------------------|----------------|-------------------|
| F.B. | 96.5 | 81.0 | 35 | 33 | 7.41 | 97.0 | 0.58 | 3.5 | 7.4 | 458 (3) |
| E.St. | 82.5 | 73.8 | 40 | 49 | 7.36 | 94.7 | 0.62 | 4.6 | 24.9 | 244 (2) |
| G.B. | 65.0 | - | 33 | - | 7.40 | 91.5 | - | - | - | - |
| H.B. b) | 90.5 | 84.0 | 37 | 32 | 7.40 | 96.5 | 0.47 | 5.2 | 12.3 | 180 (5) |
| J.E. a) J.E. b) | 74.0 88.0 | - 85.0 | 33 35 | - 37 | 7.38 7.37 | 93.5 96.0 | - 0.40 | - 6.8 | - 16.3 | - 266 (3) |
| R.G. | 85.0 | 91.0 | 36 | 40 | 7.41 | 96.0 | 0.62 | 3.7 | 16.8 | 328 (5) |
| S.G. | 88.0 | - | 46 | - | 7.38 | 96.0 | - | - | - | - |
| R.H. a) R.H. b) | 61.5 89.6 | - 64.0 | 41 40 | 47 | 7.59 7.36 | 89.5 96.0 | - 0.47 | - 5.1 | - 16.8 | - 245 (2) |
| T.J. | 48.0 | - | 64 | - | 7.33 | 81.0 | - | - | - | - |
| T.F. | 75.5 | 71.0 | 62 | 65 | 7.37 | 93.7 | 0.67 | 2.8 | 5.5 | 171 (3) |

* PaO₂ - arterial oxygen tension,
 PaCO₂ - arterial carbon dioxide tension,
 % Satn. Hb - Percentage saturation
 of oxyhaemoglobin.

V_D/V_T - ratio of dead space volume
 to tidal volume,
 PA-aO₂ - alveolar arterial oxygen
 tension difference,
 V_A - alveolar ventilation.

| Patient | PaO ₂ mm. Hg. | rest | PaO ₂ mm. Hg. | Ex. | PacO ₂ rest mm. Hg. | PacO ₂ Ex. mm. Hg. | pH | % Satn. Hb | VD/ VT | RESTING VALUES | | Work Kgm/mins |
|---------|-----------------------------|------|-----------------------------|------|--------------------------------------|-------------------------------------|------|---------------|-----------|--------------------------|------------------------------|------------------|
| | | | | | | | | | | V _A l/min. | PA-aO ₂ mm.Hg. | |
| A.H. | 98.5 | 104 | 41.5 | 41.5 | 7.37 | 97 | 0.48 | | | 7.8 | 3.0 | 350 (4) |
| J.Her | 72.5 | 57.5 | 55 | 62 | 7.34 | 92 | 0.62 | | | 3.4 | 24 | 213 (3) |
| N.S. | 94.5 | 95.5 | 45 | 45 | 7.37 | 96.5 | - | | | - | - | 155 (1½) |
| G.T. | 63 | - | 62 | - | 7.38 | 90.0 | - | | | - | - | - |
| A.V. | 98.5 | - | 40 | - | 7.43 | 97.5 | - | | | - | - | - |
| S.B. | 79.5 | 76.5 | 43 | 41 | 7.44 | 91.5 | 0.57 | | | 4.5 | 17.1 | 232 (3) |
| C.B. | 99.5 | 100 | 44.0 | 45.0 | 7.36 | - | - | | | - | - | 145 (3) |
| T.C. | 64.0 | 73.0 | 40.0 | 45.0 | 7.41 | 91.5 | - | | | - | - | 152 (2) |
| R.F. | 87.0 | 77.5 | 37 | 36.5 | 7.39 | 96 | 0.59 | | | 5.7 | 25.4 | 327 (5) |
| J.Hud. | 89.5 | 81.0 | 48.0 | 53.0 | 7.38 | 96 | 0.68 | | | 3.3 | 2.5 | 205 (2) |

Arterial blood gases, pH at rest and after exercise, and derived values in Patients in Group I.

| Patient | PaO ₂ rest mm.Hg. | PaO ₂ Ex. mm.Hg. | PaCO ₂ rest mm. Hg. | PaCO ₂ Ex. mm. Hg. | pH | % Satn. Hb | RESTING VALUES VD/VT 1/min | V _A /min | V _A -aO ₂ mm.Hg. | Work Kgm/mins. |
|---------------------|---------------------------------|--------------------------------|-----------------------------------|----------------------------------|------|---------------|----------------------------------|---------------------|---|-------------------|
| J.McM. | 72.0 | 56.0 | 56 | 52 | 7.34 | 92 | - | - | - | 184 (1½) |
| W.McD. | 86.0 | - | 41 | - | 7.42 | 96 | - | - | - | - |
| C.M. | 80.5 | 73.5 | 54 | 57 | 7.35 | 94.5 | 0.41 | 4.7 | 8.5 | 166 (2) |
| T.P. ^a) | 89.9 | 68.5 | 56 | 43 | 7.38 | 96.5 | - | - | - | 194 (3) |
| T.P. ^b) | 83.7 | 80.5 | 47 | 54 | 7.34 | 94.5 | 0.56 | 4.9 | 6.8 | 250 (3) |
| L.Car. | 91.0 | 76.5 | 37.0 | 44.0 | 7.46 | 97.2 | 0.47 | 4.9 | 17.5 | 194 (2) |
| A.S. ^a) | 82.5 | 77.0 | 45 | - | 7.39 | 95 | - | - | 11.0 | 190 (3) |
| A.S. ^b) | 95.0 | 82.5 | 42.0 | 47.0 | 7.38 | 96.5 | 0.54 | 3.6 | 13.7 | 190 (3) |
| G.P. | 94.5 | 83.5 | 42 | 42 | 7.41 | 95 | 0.61 | 3.35 | 16.2 | 204 (2) |
| W.Sm. | 85.0 | - | 52 | - | 7.37 | 95.5 | - | - | - | - |
| J.K. | 70.0 | 58.5 | 50 | 52.5 | 7.33 | 91.3 | 0.57 | 6.1 | 26.0 | 136 (1) |

| Patient | PaO ₂ rest | PaO ₂ Ex. | PaCO ₂ rest | PaCO ₂ Ex. | pH | % Satn. | RESTING VALUES | Work Kgm./Mins | | |
|-----------|-----------------------|----------------------|------------------------|-----------------------|------|----------------------|---------------------------------------|----------------|------|---------|
| | mm. Hg. | mm. Hg. | mm. Hg. | mm. Hg. | Hb. | V _A /min. | V _D /V _T 1/min. | | | |
| L.Cun. | 77.7 | 75.0 | 50 | 43 | 7.39 | 94.2 | 0.70 | 2.4 | 48.7 | 222 (2) |
| D.C. | 99 | 104 | 35 | 39 | 7.37 | 97 | 0.57 | 4.6 | 10 | 191 (3) |
| A.M. | 81 | 101 | 35 | - | 7.42 | 95 | 0.58 | 6.2 | 27.8 | 310 (3) |
| J.C. | 83.0 | 61.0 | 42.5 | 43 | 7.38 | 95.5 | 0.50 | 6.9 | 22.3 | 360 (2) |
| G.F. | 91 | 98 | 37.5 | 34.5 | 7.42 | 97 | 0.50 | 5.8 | 11.7 | 450 (3) |
| G1.H c) | 94 | - | 34 | - | 7.40 | 97 | 0.57 | 2.7 | 18 | - |
| A.V.M. a) | 79 | - | 39 | - | 7.48 | 96 | - | - | - | - |
| b) | 94.5 | 91.0 | 33.5 | 33.5 | 7.41 | 97 | 0.43 | 7.3 | 14.0 | 240 (3) |
| Geo.H. b) | 91.5 | 90.0 | 47 | 49 | 7.38 | 96.3 | 0.58 | 3.4 | 5.2 | 407 (2) |

Arterial Blood Gases and pH At Rest and After Exercise and Derived Values
In Patients in Group I.

| Patient | PaO ₂ rest | | PaO ₂ Ex. | | PaCO ₂ rest | | PaCO ₂ Ex. | | % Satn. | | RESTING | | VALUES | |
|-----------|-----------------------|--------|----------------------|--------|------------------------|------|--------------------------------|-----------------------|------------------------------|------------------|------------------|------------------|------------------|--|
| | mm.Hg. | mm.Hg. | mm.Hg. | mm.Hg. | pH | Hb | V _D /V _T | V _A l/min. | A-aP _O 2 grad.mm. | Work Done Kg.mm. | Work Done Kg.mm. | Work Done Kg.mm. | Work Done Kg.mm. | |
| E.P. a) | 60 | - | 45 | - | 7.42 | 90 | - | - | - | - | - | - | - | |
| b) | 68 | 50 | 45 | 46 | 7.39 | 92 | - | - | - | - | - | - | - | |
| A.P. | 93 | 97.5 | 45 | 43 | 7.35 | 96 | 0.49 | 4.2 | 11.2 | 240 (3) | | | | |
| W.S.B. b) | 92.8 | 72.5 | 33.5 | 29.5 | 7.39 | 96.5 | - | - | - | - | 308 (2) | | | |
| J.W. a) | 53 | 58 | 47 | 57 | 7.35 | 85 | - | - | - | - | 260 (1) | | | |
| b) | 73.5 | 61.5 | 45 | 43 | 7.39 | 93.5 | 0.63 | 3.1 | 21.8 | 248 (1½) | | | | |
| W.P. a) | 78.5 | - | 37 | - | 7.41 | 94.5 | - | - | - | - | - | - | - | |
| b) | 75.0 | 88.0 | 31 | 37 | 7.42 | 94.5 | 0.50 | 6.0 | 17.4 | 270 (3) | | | | |
| R.P. | 76 | 94.5 | 39 | 39 | 7.39 | 93.8 | 0.36 | 7.2 | 29.3 | 260 (3) | | | | |
| P.J. | 80.3 | 68.0 | 44 | 48 | 7.37 | 94.5 | 0.64 | 4.2 | 16.1 | 177 (2) | | | | |

Arterial Blood Gases and pH At Rest and After Exercise and Derived Values

In Patients in Group I.

| Patient | PaO ₂ rest mm.Hg. | PaO ₂ Ex. mm.Hg. | PaCO ₂ rest mm.Hg. | PaCO ₂ Ex. mm. Hg. | pH | % Satn. Hb. | RESTING VALUES | V _A / l/min. | A-aP _O 2 grad.mm. | Work Done Kg.mm. |
|----------|---------------------------------|--------------------------------|----------------------------------|----------------------------------|------|----------------|-------------------|----------------------------|---------------------------------|---------------------|
| H.K. | 81.5 | 82.0 | 35.0 | 35.0 | 7.39 | 95 | 0.49 | 5.7 | 23.2 | 336 (3) |
| E.Sh. a) | 78.0 | 85.5 | 38.5 | 38.0 | 7.38 | 94.5 | - | - | - | 285 (3) |
| b) | 91.0 | 92.6 | 35 | 39 | 7.39 | 96.5 | 0.39 | 6.7 | 20 | 346 (3) |
| J.S. | 95.5 | 100.5 | 42 | 40.5 | 7.35 | 96.5 | 0.46 | 3.1 | 7.3 | 290 (3) |
| C.Mah. | 74.5 | 61.5 | 40 | 40 | 7.37 | 93 | 0.67 | 5.1 | 14.0 | 150 (1) |
| N.H. | 88.5 | 85.0 | 32.5 | 33.0 | 7.37 | 95.7 | 0.37 | 7.0 | 17.4 | 165 (3) |

Arterial Blood Gases At Rest and After Exercise, and Derived Values in
Patients in Group II.

| Patient | PaO ₂ rest mm.Hg. | PaO ₂ Ex. mm.Hg. | PaCO ₂ rest mm.Hg. | PaCO ₂ Ex. mm.Hg. | RESTING VALUES | | | | A-aP _O 2 grad.mm. | Work Done Kg. mm. | Minutes Work. |
|---------|---------------------------------|--------------------------------|----------------------------------|---------------------------------|----------------|------|-------|--------------------------|---------------------------------|----------------------|------------------|
| | | | | | % Satn Hb. | pH | VD/VT | V _A l/min. | | | |
| E.N. | 52.3 | 58.5 | 52 | 52 | 7.39 | 85 | 0.50 | 4.4 | 16.5 | 285 | 3 |
| L.M. | 53.0 | - | 50 | - | 7.36 | 83 | 0.64 | 4.7 | 36.4 | - | - |
| E.G. | 73.5 | 80.5 | 47 | 52 | 7.40 | 93.5 | 0.72 | 2.8 | 13.0 | 315 | 3 |
| F.K. | 78.0 | 72.5 | 54 | 54.5 | 7.39 | 94.5 | - | - | - | 139 | 3 |
| R.T. | 62.0 | 51.0 | 59 | 73 | 7.32 | 87.5 | 0.71 | 4.0 | 4.0 | 143 | 12 |
| E.S. | 75.0 | 53.0 | 50 | 55 | 7.38 | 93.5 | 0.67 | 3.0 | 8.2 | 126 | 4 |
| E.Pat. | 69.5 | - | 60 | - | 7.33 | 91.2 | 0.54 | 2.9 | 3.0 | - | - |
| F.R. | 76.0 | 70.8 | 47 | 55 | 7.37 | 93.5 | 0.66 | 3.6 | 14.5 | 334 | 3 |

Regional Ventilation and Perfusion in Patients in Group I.

| Patient | Ventilation Index | | | | | | Perfusion Index | | | | | | Dynamic Distribution Index | | | | | |
|---------|-------------------|------|------|------|------|------|-----------------|------|------|------|------|------|----------------------------|------|------|-------|------|-------|
| | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. |
| F.B. | 0.81 | 0.88 | 1.85 | 0.69 | 0.56 | 1.10 | 1.27 | 0.71 | 1.69 | 0.64 | 0.58 | 1.17 | 23.8 | 20.8 | 23.2 | 33.3 | 29.7 | 47.6 |
| R.K. a) | 1.65 | 0.90 | 0.76 | 1.82 | 0.88 | 0.68 | 1.66 | 1.60 | 1.26 | 1.15 | 0.55 | 0.58 | 66.7 | 41.2 | 28.1 | 48.5 | 43.6 | 23.9 |
| b) | 1.20 | 0.83 | 0.52 | 1.36 | 0.97 | 1.32 | 0.77 | 0.79 | 0.86 | 0.99 | 1.09 | 1.44 | 96.0 | 35.0 | 20.2 | 135 | 76.0 | 59.0 |
| E.St. | 1.05 | 1.41 | 0.65 | 1.35 | 0.87 | 0.53 | 0.98 | 2.13 | 0.82 | 0.71 | 0.88 | 0.53 | 61.0 | 50.8 | 26.8 | 33.0 | 34.8 | 20.8 |
| G.B. | 1.24 | 0.37 | 0.34 | 1.56 | 1.34 | 0.74 | 1.20 | 0.47 | 0.79 | 1.19 | 1.18 | 1.01 | 83.3 | 25.2 | 25.2 | 111.0 | 96.3 | 50.5 |
| H.B. a) | 0.81 | 1.20 | 0.82 | 1.32 | 1.02 | 0.51 | 1.08 | 1.65 | 0.89 | 0.63 | 0.71 | 0.89 | 65.0 | 57.7 | 27.0 | 80.0 | 29.4 | 23.9 |
| b) | 1.55 | 1.10 | 0.46 | 1.59 | 0.67 | 0.22 | 1.48 | 1.26 | 0.75 | 0.89 | 0.68 | 0.45 | 180 | 29.6 | 20.5 | 54.5 | 37.0 | 19.6 |
| J.E. a) | 1.48 | 0.93 | 0.51 | 1.43 | 1.03 | 0.25 | 1.44 | 1.05 | 0.32 | 1.44 | 0.87 | 0.53 | 70.0 | 34.9 | 13.4 | 44.9 | 25.4 | 14.9 |
| b) | 1.02 | 1.09 | 0.64 | 1.44 | 1.35 | 0.40 | 1.08 | 1.22 | 0.58 | 1.39 | 1.21 | 0.58 | 88.5 | 60.0 | 19.3 | 154.0 | 65.0 | 17.7 |
| R.G. | 0.63 | 1.06 | 1.28 | 0.71 | 0.89 | 1.42 | 0.34 | 0.84 | 2.68 | 0.31 | 0.53 | 1.21 | 26.8 | 49.6 | 55.5 | 39.4 | 63.3 | 111.0 |

R.U. - Right Upper Zone.
R.M. - Right Middle Zone.
R.L. - Right Lower Zone.

L.U.- Left Upper Zone.
L.M.- Left Middle Zone.
L.L.- Left Lower Zone.

Regional Ventilation and Perfusion in Patients in Group I.

| Patient | Ventilation Index | | | | | | Perfusion Index | | | | | | Dynamic Distribution Index | | | | | |
|---------|-------------------|------|------|------|------|------|-----------------|------|------|------|------|------|----------------------------|------|------|------|------|------|
| | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. |
| S.G. | 0.66 | 1.00 | 1.54 | 0.87 | 0.98 | 0.62 | 1.00 | 0.92 | 1.26 | 0.64 | 1.05 | 1.04 | 49.0 | 44.0 | 29.4 | 52.0 | 72.5 | 35.0 |
| R.H. a) | 1.62 | 0.53 | 1.44 | 0.86 | 0.15 | 1.22 | 1.31 | 0.84 | 1.09 | 1.03 | 0.52 | 101 | 53.0 | 21.4 | 73.7 | 40.5 | 21.4 | |
| b) | 2.06 | 1.30 | 0.37 | 1.39 | 0.86 | 0.43 | 1.43 | 1.25 | 0.59 | 1.42 | 1.00 | 0.47 | 122 | 40.0 | 21.1 | 65.5 | 45.1 | 24.3 |
| T.J. | 1.41 | 0.80 | 0.50 | 0.70 | 0.99 | 1.49 | 1.20 | 0.51 | 0.25 | 0.56 | 1.15 | 1.89 | 23.4 | 15.7 | 14.0 | 15.2 | 19.3 | 36.4 |
| T.F. | 1.42 | 0.86 | 0.45 | 0.91 | 1.42 | 0.54 | 1.47 | 1.23 | 0.53 | 0.81 | 1.02 | 0.63 | 80.0 | 23.2 | 18.1 | 48.6 | 41.3 | 39.6 |
| J.M.M. | 0.50 | 0.98 | 1.52 | 0.62 | 1.15 | 1.03 | 0.52 | 1.31 | 2.09 | 6.22 | 0.56 | 0.48 | 43.4 | 61.3 | 112 | 43.5 | 43.7 | 38.0 |
| W.M.D. | 0.69 | 0.63 | 1.36 | 0.43 | 0.68 | 1.63 | 0.57 | 0.47 | 1.44 | 0.30 | 0.48 | 1.96 | 30.0 | 27.5 | 57.0 | 32.5 | 49.5 | 156 |
| C.M. | 1.09 | 0.63 | 0.75 | 1.37 | 1.48 | 0.71 | 0.81 | 1.07 | 1.48 | 0.70 | 1.17 | 0.58 | 46.8 | 26.4 | 27.4 | 65.3 | 68.0 | 26.0 |
| T.P. a) | 0.94 | 1.25 | 1.07 | 1.14 | 0.71 | 0.76 | 0.66 | 1.43 | 1.78 | 0.37 | 0.56 | 1.00 | 74.5 | 83.5 | 74.5 | 74.5 | 27.5 | 33.3 |
| b) | 1.02 | 1.23 | 1.15 | 0.81 | 0.78 | 0.96 | 0.60 | 1.34 | 1.61 | 0.63 | 0.70 | 0.89 | 48.5 | 78.0 | 49.2 | 37.6 | 32.7 | 28.1 |

Regional Ventilation and Perfusion in Patients in Group I.

| Patient | Ventilation Index | | | | | | Perfusion Index | | | | | | Dynamic Distribution Index | | | | | |
|---------|-------------------|------|------|------|------|------|-----------------|------|------|------|------|------|----------------------------|------|------|-------|------|------|
| | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. |
| L.Car | 0.38 | 0.92 | 1.19 | 0.36 | 1.61 | 1.22 | 0.35 | 0.77 | 1.54 | 0.42 | 1.28 | 1.36 | 27.6 | 32.8 | 49.0 | 24.2 | 39.6 | 50.0 |
| A.S. a) | 1.47 | 0.83 | 0.74 | 1.45 | 1.10 | 0.45 | 1.14 | 0.99 | 0.46 | 1.34 | 1.09 | 0.76 | 26.0 | 28.4 | 25.6 | 132.0 | 54.7 | 34.8 |
| b) | 1.46 | 1.11 | 0.54 | 1.34 | 1.05 | 0.63 | 1.04 | 1.01 | 0.89 | 1.01 | 1.12 | 0.86 | 73.0 | 62.0 | 21.0 | 85.0 | 36.0 | 22.0 |
| G.P. | 1.39 | 0.79 | 0.97 | 1.44 | 1.09 | 0.89 | 1.62 | 0.51 | 1.20 | 0.76 | 0.90 | 1.21 | 37.0 | 42.0 | 43.3 | 74.0 | 46.8 | 34.4 |
| W.Sm. | 0.55 | 0.94 | 1.56 | 1.03 | 0.72 | 1.23 | 0.67 | 0.94 | 1.75 | 0.71 | 0.97 | 0.94 | 37.0 | 38.2 | 97.0 | 56.2 | 28.4 | 46.2 |
| J.K. | 0.84 | 1.66 | 1.11 | 0.44 | 0.97 | 0.50 | 0.67 | 0.86 | 1.14 | 0.44 | 0.77 | 1.49 | 50.0 | 134 | 25.4 | 24.3 | 30.4 | 30.0 |
| A.H. | 0.84 | 1.12 | 0.93 | 0.66 | 1.14 | 1.21 | 0.49 | 0.92 | 1.42 | 0.71 | 0.76 | 1.66 | 26.0 | 52.3 | 89.0 | 27.4 | 44.5 | 61.7 |
| J.Her. | 1.42 | 0.97 | 0.69 | 1.60 | 0.75 | 0.44 | 1.45 | 1.15 | 1.01 | 1.05 | 0.73 | 0.64 | 46.8 | 26.0 | 21.4 | 119 | 25.6 | 20.9 |
| N.S. | 0.50 | 2.10 | 1.06 | 1.08 | 0.66 | 0.43 | 0.57 | 1.11 | 1.80 | 0.63 | 0.35 | 0.37 | 23.2 | - | 105 | 24.7 | 22.6 | 21.0 |
| G.T. | 2.78 | 0.90 | 1.40 | 0.62 | 0.50 | 0.79 | 2.76 | 1.23 | 1.44 | 0.56 | 0.32 | 0.53 | 52.3 | 41.4 | 59.5 | 32.2 | 35.6 | 32.2 |

Regional Ventilation and Perfusion in Patients in Group I.

| Patient | Ventilation Index | | | | | | Perfusion Index | | | | | | Dynamic Distribution Index | | | | | | | | | | | |
|---------|-------------------|------|------|------|------|------|-----------------|------|------|------|------|------|----------------------------|------|------|------|------|------|-------|------|------|------|------|------|
| | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | R.M. | R.L. | L.U. | R.M. | R.L. | L.U. | L.M. | L.L. |
| A.V. | 0.60 | 0.35 | 1.00 | 0.51 | 0.77 | 2.04 | 0.77 | 0.58 | 0.99 | 0.74 | 0.74 | 1.75 | 27.6 | 27.0 | 33.3 | 30.8 | 32.9 | 32.9 | 116.0 | | | | | |
| S.B. | 1.08 | 1.04 | 0.71 | 1.23 | 1.24 | 0.78 | 1.01 | 1.08 | 0.94 | 0.72 | 1.10 | 1.08 | 116 | 42.7 | 36.4 | 79.3 | 121 | 121 | 46.3 | | | | | |
| C.B. | 1.53 | 0.67 | 0.48 | 2.31 | 0.85 | 0.80 | 1.53 | 1.22 | 0.75 | 0.72 | 0.93 | 0.79 | 59.0 | 35.8 | 30.0 | 74.6 | 34.0 | 34.0 | 26.3 | | | | | |
| T.C. | 0.77 | 1.40 | 1.03 | 0.88 | 1.18 | 0.46 | 0.47 | 1.22 | 1.26 | 0.60 | 1.39 | 0.84 | 28.4 | 104 | 25.2 | 39.1 | 71.5 | 71.5 | 21.0 | | | | | |
| R.F. | 1.15 | 0.85 | 0.91 | 1.67 | 1.44 | 0.38 | 1.13 | 1.18 | 1.17 | 1.08 | 1.10 | 0.58 | 105 | 43.3 | 50.0 | 158 | 57.5 | 57.5 | 16.6 | | | | | |
| J.Hud. | 1.30 | 1.02 | 0.59 | 1.23 | 1.14 | 0.75 | 1.22 | 1.19 | 0.45 | 1.30 | 1.17 | 0.64 | 64.2 | 66.0 | 36.5 | 44.0 | 44.0 | 44.0 | 35.5 | | | | | |
| L.Cun. | 1.02 | 0.95 | 1.37 | 1.03 | 1.00 | 0.65 | 1.02 | 1.30 | 1.32 | 0.83 | 0.81 | 0.65 | 71.0 | 50.3 | 42.0 | 120 | 57.5 | 57.5 | 24.6 | | | | | |
| D.C. | 1.40 | 1.09 | 1.04 | 1.18 | 0.76 | 0.37 | 1.00 | 1.19 | 1.41 | 0.74 | 0.67 | 0.35 | 90.5 | 73.5 | 49.1 | 40.0 | 30.8 | 30.8 | 30.3 | | | | | |
| A.M. | 0.89 | 0.93 | 1.02 | 1.02 | 0.89 | 1.34 | 0.40 | 0.70 | 1.54 | 0.51 | 0.99 | 1.82 | 44.2 | 44.2 | 47.2 | 99.5 | 80.0 | 80.0 | 76.7 | | | | | |
| J.C. | 1.34 | 1.40 | 1.12 | 0.48 | 0.85 | 1.25 | 0.51 | 1.10 | 1.42 | 0.63 | 0.69 | 1.28 | 25.4 | 14.1 | 50.5 | 16.7 | 16.5 | 16.5 | 29.5 | | | | | |
| G.P. | 0.70 | 0.87 | 1.29 | 0.61 | 1.01 | 1.15 | 0.42 | 0.52 | 1.61 | 0.49 | 0.87 | 1.46 | 18.2 | 63.4 | 103 | 24.3 | 38.4 | 38.4 | 77.0 | | | | | |

Regional Ventilation and Perfusion in Patients in Group I.

| Patient | Ventilation Index | | | | Perfusion Index | | | | Dynamic Distribution Index | | | | | | | | | |
|----------|-------------------|------|------|------|-----------------|------|------|------|----------------------------|------|------|------|------|------|------|------|-------|-------|
| | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | F.U. | F.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. |
| G1.H. a) | 0.27 | 0.36 | 1.33 | 0.29 | 0.48 | 1.31 | 0.67 | 0.52 | 1.49 | 0.44 | 0.63 | 2.06 | 20.2 | 27.4 | 72.0 | 19.9 | 20.4 | 41.7 |
| b) | 0.95 | 0.63 | 1.08 | 0.90 | 1.08 | 1.17 | 0.64 | 0.70 | 1.22 | 0.85 | 1.03 | 1.20 | 38.5 | 41.0 | 51.3 | 49.3 | 77.0 | 92.0 |
| c) | 0.91 | 0.87 | 0.97 | 0.71 | 0.07 | 1.32 | 0.58 | 0.71 | 1.03 | 0.89 | 1.17 | 1.32 | 43.7 | 43.5 | 69.0 | 60.5 | 60.0 | 65.3 |
| A.V.M.a) | 0.80 | 1.15 | 0.86 | 1.18 | 1.15 | 0.79 | 0.72 | 1.15 | 1.07 | 1.02 | 1.16 | 0.76 | 32.7 | 75.3 | 22.6 | 56.5 | 158 | 21.0 |
| b) | 1.18 | 1.36 | 0.61 | 0.93 | 1.11 | 0.48 | 0.51 | 0.69 | 0.54 | 1.56 | 1.73 | 0.93 | 35.0 | 32.7 | 19.8 | 40.3 | 123.0 | 24.9 |
| Gee. H. | | | | | | | | | | | | | | | | | | |
| a) | 0.82 | 1.19 | 1.19 | 0.98 | 0.98 | 0.51 | 0.71 | 1.49 | 1.49 | 0.60 | 0.70 | 0.51 | 100 | 112 | 53.2 | 41.7 | 125 | 38.4 |
| b) | 0.80 | 1.06 | 0.71 | 1.66 | 0.89 | 1.08 | 0.60 | 1.11 | 1.34 | 0.89 | 0.89 | 1.01 | 53.2 | 42.5 | 26.1 | 54.6 | 53.2 | 24.7 |
| E.P. a) | 0.51 | 0.40 | 1.76 | 0.67 | 0.48 | 2.46 | 0.60 | 0.50 | 1.81 | 0.32 | 0.54 | 2.02 | 26.8 | 31.1 | 28.6 | 31.0 | 38.2 | 52.2 |
| b) | 0.80 | 1.05 | 1.33 | 0.59 | 0.45 | 1.65 | 0.90 | 0.82 | 1.48 | 0.40 | 0.46 | 1.80 | 24.6 | 46.4 | 50.8 | 23.3 | 25.0 | 42.3 |
| A.P. | 0.57 | 0.38 | 1.75 | 0.80 | 0.76 | 1.44 | 0.36 | 0.41 | 1.70 | 0.54 | 0.60 | 1.90 | 26.6 | 30.7 | 85.0 | 23.0 | 37.8 | 146.0 |
| W.Sh a) | 1.33 | 1.08 | 0.49 | 1.15 | 1.25 | 0.70 | 1.47 | 1.51 | 0.73 | 0.84 | 1.00 | 0.57 | 48.6 | 40.4 | 10.3 | 78.7 | 85.0 | 20.6 |
| b) | 1.63 | 0.98 | 0.57 | 1.46 | 1.08 | 0.46 | 1.55 | 1.19 | 0.46 | 1.25 | 1.04 | 0.60 | 87.7 | 69.0 | 40.0 | 68.0 | 49.5 | 28.8 |

Regional Ventilation and Perfusion in Patients in Group I.

| Patient | Ventilation Index | | | | | | Perfusion Index | | | | | | Dynamic Distribution Index | | | | | |
|---------|-------------------|------|------|------|------|------|-----------------|------|------|------|------|------|----------------------------|------|------|------|------|------|
| | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. |
| J.W. a) | 2.20 | 0.70 | 0.38 | 1.71 | 0.82 | 0.45 | 1.31 | 0.77 | 0.73 | 1.54 | 1.00 | 0.75 | 34.0 | 21.6 | 17.7 | 63.8 | 42.0 | 26.7 |
| b) | 2.28 | 0.90 | 0.29 | 2.03 | 1.19 | 0.79 | 2.15 | 1.09 | 0.72 | 1.59 | 0.95 | 0.69 | 65.4 | 36.5 | 19.3 | 96.0 | 51.6 | 23.5 |
| W.P. a) | 1.26 | 1.10 | 0.42 | 1.26 | 1.40 | 0.34 | 0.81 | 1.23 | 0.91 | 0.88 | 1.16 | 0.83 | 65.8 | 47.2 | 12.8 | 50.0 | 21.4 | 14.5 |
| b) | 1.20 | 1.21 | 0.57 | 1.29 | 1.10 | 0.50 | 1.87 | 1.15 | 0.48 | 1.26 | 0.78 | 0.59 | 11.8 | 11.8 | 29.4 | 11.6 | 30.8 | 14.7 |
| R.P. | 0.50 | 1.28 | 1.08 | 0.48 | 1.21 | 1.18 | 0.49 | 0.97 | 1.58 | 0.25 | 0.95 | 1.42 | 20.2 | 94.5 | 82.7 | 16.9 | 13.6 | 11.2 |
| P.J. | 0.86 | 1.10 | 0.60 | 0.97 | 1.37 | 0.71 | 0.52 | 0.76 | 0.74 | 1.79 | 1.31 | 0.96 | 38.2 | 105 | 36.6 | 13.9 | 11.5 | 28.8 |
| H.K. | 0.81 | 1.36 | 1.08 | 0.66 | 0.81 | 1.04 | 0.54 | 0.87 | 1.12 | 1.23 | 0.79 | 1.34 | 42.0 | 43.7 | 47.4 | 47.4 | 29.2 | 31.3 |
| E.Sh.a) | 0.63 | 0.76 | 0.77 | 0.69 | 1.80 | 0.83 | 1.00 | 0.82 | 0.59 | 0.95 | 1.33 | 0.67 | 62.5 | 46.5 | 18.0 | 50.0 | 11.2 | 16.4 |
| b) | 1.03 | 1.25 | 0.93 | 0.79 | 0.85 | 0.75 | 0.81 | 1.39 | 1.19 | 0.52 | 0.92 | 0.91 | 63.7 | 58.5 | 28.6 | 25.9 | 71.0 | 33.8 |
| J.S. | 0.55 | 1.29 | 1.32 | 0.38 | 0.95 | 1.40 | 0.97 | 0.95 | 1.52 | 0.21 | 0.56 | 1.36 | 31.4 | 60.2 | 14.6 | 28.1 | 37.6 | 97.0 |
| C.Mah | 1.95 | 1.21 | 0.47 | 1.77 | 0.55 | 0.35 | 1.53 | 1.83 | 1.18 | 0.70 | 0.42 | 0.52 | 89.0 | 28.0 | 18.6 | 33.8 | 16.7 | 15.8 |
| N.H. | 1.02 | 0.69 | 0.69 | 1.03 | 1.15 | 0.55 | 1.48 | 0.74 | 0.49 | 1.09 | 1.51 | 0.31 | 24.2 | 20.0 | 19.4 | 47.0 | 60.5 | 22.8 |

Regional Ventilation and Perfusion in Patients in Group III.

| Patient | Ventilation Index | | | | | | Perfusion Index | | | | | | Dynamic Distribution Index | | | | | |
|---------|-------------------|------|------|------|------|------|-----------------|------|------|------|------|------|----------------------------|------|-------|------|------|------|
| | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. | R.U. | R.M. | R.L. | L.U. | L.M. | L.L. |
| E.N. | 1.28 | 0.85 | 0.84 | 1.69 | 0.96 | 0.67 | 1.01 | 0.93 | 0.67 | 1.29 | 1.15 | 0.99 | 60.0 | 46.2 | 20.8 | 53.2 | 70.3 | 23.6 |
| J.M. | 1.05 | 0.80 | 0.57 | 1.28 | 1.18 | 0.99 | 0.82 | 0.89 | 0.93 | 1.01 | 1.18 | 1.06 | 47.6 | 26.7 | 15.0 | 73.7 | 64.0 | 22.0 |
| E.G. | 1.09 | 0.88 | 1.00 | 1.19 | 0.82 | 1.04 | 0.65 | 1.00 | 1.22 | 0.84 | 0.81 | 1.44 | 77.5 | 59.2 | 52.2 | 66.8 | 39.6 | 40.4 |
| F.K. | 2.10 | 0.98 | 1.17 | 0.88 | 0.85 | 0.94 | 2.54 | 1.27 | 1.35 | 0.72 | 0.55 | 0.75 | 66.0 | 72.0 | 94.5 | 33.7 | 67.0 | 71.0 |
| R.T. | 1.81 | 1.07 | 0.43 | 1.73 | 1.58 | 0.40 | 1.97 | 1.09 | 0.69 | 1.14 | 1.51 | 0.75 | 95.2 | 28.1 | 12.5 | 95.2 | 27.0 | 15.0 |
| E.Sc. | 1.10 | 0.89 | 0.83 | 1.00 | 1.22 | 0.85 | 1.00 | 0.91 | 0.81 | 1.03 | 1.06 | 1.03 | 24.4 | 25.7 | 24.3 | 29.8 | 25.4 | 26.7 |
| E.Pat. | 0.87 | 0.94 | 1.30 | 0.93 | 0.99 | 0.97 | 0.72 | 0.94 | 1.43 | 0.63 | 1.06 | 1.12 | 69.5 | 66.6 | 73.6 | 78.8 | 81.3 | 103 |
| F.R. | 0.86 | 1.05 | 0.97 | 1.35 | 0.88 | 0.98 | 0.96 | 1.21 | 1.07 | 0.80 | 0.76 | 1.28 | 58.0 | 49.5 | 115.0 | 33.0 | 32.0 | 36.0 |

BIBLIOGRAPHY

Actuarius (c. 1300) from Badham (1814)

Aetius (c. 450 A.D.) from Badham (1814)

American Thoracic Society, Committee on Diagnostic Standards
for Non-Tuberculous Respiratory Diseases (1962).

Definitions and classifications of chronic bronchitis,
asthma, and pulmonary emphysema.

Am. Rev. resp. Dis., 85, 762.

Anderson, J.A., and Dunnill, M.S. (1965)

Observations on the estimation of the quantity of emphysema
in the lungs by the point-sampling method.

Thorax, 20, 462.

Armitage, G.H., and Arnott, W.M. (1949)

Effect of voluntary hyperpnoea on pulmonary blood flow.

J. Physiol. (Lond), 109, 64.

Badham, C. (1814)

An Essay on Bronchitis, etc.

Callow, London.

Baillie, M.

The morbid anatomy of some of the most important parts
of the human body.

Johnston and Nicol, London.

Baldwin, E. de F., Cournand, A., and Richards, D.W., Jnr. (1949)

Pulmonary insufficiency, III. A study of 122 cases of
chronic pulmonary emphysema.

Medicine (Baltimore), 28, 201.

Baldwin, E. de F., Harden, K., Greene, D.G., Cournand, A., and Richards, D.W. (1950)

Pulmonary insufficiency, IV - A study of 16 cases with large pulmonary air cysts or bullae.
Medicine (Baltimore), 29, 169.

Ball, W.C., Jr., Stewart, P.B., Newsham, L.G.S., and Bates D.V. (1962)

Regional pulmonary function studied with xenon¹³³.
J. clin. Invest., 41, 519.

Barnhard, H.J., Pierce, J.A., Joyce, J.W., and Bates, J.H. (1960)

Roentgenographic determination of total lung capacity.
A new method evaluated in health, emphysema and congestive cardiac failure.

Am. J. Med., 28, 51.

Bates, D.V., (1959)

Unusual forms of emphysema.

Am. Rev. resp. Dis. (Suppl.), 80, 172.

Bates, D.V., (1962) Indiscussion.

In: Ciba Foundation Symposium on Pulmonary Structure and Function. Churchill, London.

Edited by A.V.S. de Reuck and M. O'Connor. Churchill, London.

Bates, D.V., (1965) a. In discussion.

In: Progress in Research in Emphysema and Chronic Bronchitis.

Edited by R.S. Mitchell. Karger, Basel and New York.

Bates, D.V., (1965) b.

Measurement of regional ventilation and blood flow distribution.

In: Handbook of Physiology Respiration II. Chapter 57.

Edited by W.O. Fenn and H. Rahn.

Waverly Press. Baltimore, U.S.A.

Bates, D.V. and Christie, R.V. (1964)

Respiratory Function in Disease.

Saunders, Philadelphia and London.

Bates, D.V., Boucet, N.G., and Dormer, A.E. (1955).

The pulmonary diffusing capacity in normal subjects.

J. Physiol. (Lond.), 129, 237.

Bates, D.V., Knott, J.M.S., and Christie, R.V. (1956)

Respiratory function in emphysema in relation to prognosis.

Q. Jl. Med., 25, 137.

Bedell, G.N., and Marshall, R., DuBois, A.B., and Comroe, J.H., Jr.
(1956).

Plethysmographic determination of the volume of gas trapped
in the lungs.

J. clin. Invest., 35, 664.

Bedell, G.N., and Ostiguy, G. (1966)

The steady state and single breath diffusing capacity in
patients with types A and B chronic airways obstruction.

Paper to the Medical Research Society (London). (Unpublished)

Belcher, J.R., Capel, L., Pattinson, J.N., and Smart, J. (1959)

Hypoplasia of the pulmonary arteries.

Brit. J. Dis. Chest, 53, 253.

Bentivoglio, L.G., Beerel, F., Stewart, P.B., Bryan, A.C.,
Ball, W.C., Jr., and Bates, D.V. (1963).

Studies of regional ventilation and perfusion in pulmonary
emphysema using xenon¹³³.

Am. Rev. resp. Dis., 88, 315.

Bishop, P.J., (1959).

Samuel Johnsons lung.

Tubercle, Lond., 40, 478.

Bonet, T. (1679).

Sepulchretum, sive anatomia practica etc.

Chouët, Geneva.

Briscoe, W.A., and DuBois, A.B. (1958).

The relationship between airways resistance, airway
conductance and lung volume in subjects of different
age and body size.

J. clin. Invest., 37, 1279.

Briscoe, W.A., and Cournand, A. (1962).

The degree of variation of blood perfusion and of
ventilation within the emphysematous lung, and some
related considerations.

In: Ciba Foundation Symposium on Pulmonary Structure
and Function. p.304.

Edited by A.V.S. de Reuck, and M. O'Connor.

Churchill, London.

Briscoe, W.A., Hardewig, A., Emmanuel, G., Gurtner, P.H., Rochester, D.F., and Cournand, A. (1959).

The distribution of blood perfusion to the lung.

Fed. Proc., 18, 16.

Briscoe, W.A., and Nash, E.S. (1965).

The slow space in chronic obstructive pulmonary disease.

Respiratory Failure.

Ann. N.Y. Acad. Sci., 121, 706.

Burrows, B., Kasik, J.E., Niden, A.H., and Barclay, W.R. (1961).

Clinical usefulness of the single-breath pulmonary diffusing capacity test.

Am. Rev. resp. Dis., 84, 789.

Burrows, B., Niden, A.H., Fletcher, C.M., and Jones, N.L. (1964).

Clinical types of chronic obstructive lung disease in London and Chicago.

Am. Rev. resp. Dis., 90, 14.

Burrows, B., Strauss, R.H., and Niden, A.H., (1965).

Chronic obstructive lung disease, III. Inter-relationships of pulmonary function data.

Am. Rev. resp. Dis., 91, 861.

Burrows, B., Niden, A.H., Barclay, W.R., and Kasik, J.E. (1965).

Chronic obstructive lung disease, II. Relationship of clinical and physiologic findings to the severity of airways obstruction.

Am. Rev. resp. Dis., 91, 665.

Burrows, B., Fletcher, C.M., Heard, B.E., Jones, N.L. and Wootliff, J.S. (1966).

The emphysematous and bronchial types of chronic airways obstruction.

Lancet, 1, 830.

Cabot, R.C. (1927).

Physical diagnosis. 9th edition.
Wood, New York.

Cadigan, J.B., Marks, A., Ellicott, M.F., Jones, R.H., and Gaensler, E.A. (1961).

An analysis of factors affecting the measurement of pulmonary diffusing capacity by the single breath method.
J. clin. Invest., 40, 1495.

Campbell, E.J.M., (1958)

The Respiratory Muscles.

Lloyd-Luke, London.

Campbell, E.J.M., and Howell, J.B.L. (1963).

Sensation of breathlessness.

Br. med. Bull., 19, 36.

Carlens, E. (1949).

A new flexible double-lumen catheter for bronchospirometry.

J. thorac. Surg., 18, 742.

Christie, R.V. (1944).

Emphysema of the lungs.

Br. med. J., 1, 105.

Ciba Guest Symposium Report (1959), Edited by C.M. Fletcher.

Terminology, definitions and classification of chronic pulmonary emphysema and related conditions.

Thorax, 14, 286.

Comroe, J.H. Jnr., (1966).

In: Breathlessness. p. 236, 237.

Edited by J.B.L. Howell and E.J.M. Campbell.

Blackwell Scientific Publications. Oxford.

Comroe, J.H. Jnr., and Botelho, S. (1947)

The unreliability of cyanosis in the recognition of arterial anoxemia.

Am. J. Med. Sci., 214, 1.

Comroe, J.H., Jnr., Forster, R.E., II, DuBois, A.B., Briscoe, W.A., and Carlsen, E. (1962).

The Lung.

2nd. edition, Year Book Medical Publishers, Chicago.

Cotes, J.E. (1965).

Lung Function.

Blackwell Scientific Publications, Oxford.

Cournand, A., (1957)

Pulmonary circulation. Its control in man, with some remarks on methodology.

Science, (N.Y.), 125, 1231.

Cournand, A., (1965)

Chairman's introductory remarks.

Respiratory failure.

Ann. N.Y. Acad. Sci., 121, 885.

- Darke, C.S., Chrispin, A.R., and Snowden, B.S. (1960).
Unilateral lung transradiancy: A physiological study.
Thorax, 15, 74.
- Davidson, M. (1936).
Discussion on Emphysema.
Proc. R. Soc. Med., 29, 1315.
- Davidson, L. (1966)
Repeatability of respiratory data in the individual subject.
Thorax, 21, 47.
- Davies, L.G., Rosser, T.H.L., and West, L.R. (1965)
Autotransplantation of the lung in sheep.
Thorax, 20, 481.
- Davies, G.M., Simon, G., and Reid, L. (1966).
Pre and post operative assessment of emphysematous bullae.
Br. J. Dis. Chest., 60, 120.
- Dill, D.B., (1944).
Handbook of Respiratory Data in Aviation.
Committee on Medical Research, Washington.
- Dollery, C.T., and Gillam, P.M.S., (1963).
The distribution of blood and gas within the lungs measured
by scanning after administration of ^{133}Xe .
Thorax, 18, 316.
- Dollery, C.T., and Hugh-Jones, P. (1963).
Distribution of gas and blood in the lungs in disease.
Br. med. Bull., 19, 59.

Dornhorst, A.C. (1955).

Respiratory insufficiency.

Lancet, 1, 1185.

Dornhorst, A.C., Heaf, P.J.D., and Semple, S.J.G. (1957)

Unilateral "emphysema".

Lancet, 2, 873.

Douglas, C.G., and Priestley, J. (1948)

Human Physiology, 3rd. edition.

Clarendon Press, Oxford.

DuBois, A.B., Botelho, S.Y., and Comroe, J.H.Jnr., (1956)

A new method for measuring airway resistance in man using
a body plethysmograph: Values in normal subjects and in
patients with respiratory disease.

J. clin. Invest., 35, 327.

Ebert, R.V., and Pierce, J.A. (1963).

Pathogenesis of pulmonary emphysema.

Arch. Int. Med., 111, 34.

Eiseman, B., Petty, T., and Siden, W. (1959).

Experimental emphysema:- a review.

Am. Rev. resp. Dis., 80, 147.

Elmes, P.C., Fletcher, C.M., and Dutton, A.A.C. (1957)

Prophylactic use of oxytetracycline for exacerbations
of chronic bronchitis.

Br. med. J., 2, 1272.

Erikson, S. (1964).

Pulmonary emphysema and alpha₁ - antitrypsin deficiency.

Acta. med. scand., 175, 197.

Field, W.E.H., Davey, E.N., Reid, L., and Roe, F.J.C. (1966)

Bronchial mucus gland hypertrophy; its relation to symptoms and environment.

Br. J. Dis. Chest, 60, 66.

Fishman, A.P., (1965).

The roads to respiratory insufficiency.

Respiratory Failure.

Ann. N.Y. Acad. Sci., 121, 657.

Fletcher, C.M. (1952).

In: Discussion on the diagnosis of pulmonary emphysema.

Proc. R. Soc. Med., 45, 577.

Fletcher, C.M., Elmes, P.C., Fairbairn, A.S., and Wood, C.H. (1959)

The significance of respiratory symptoms and the diagnosis of chronic bronchitis in a working population.

Br. Med. J., 2, 257.

Fletcher, C.M., Hugh-Jones, P., McNicol, M.W., and Pride, N.B. (1963)

The diagnosis of pulmonary emphysema in the presence of chronic bronchitis.

Q. Jl. Med., 32, 33.

Fletcher, C.M., Jones, N.L., Burrows, B., and Niden, A.H. (1964)

American emphysema and British bronchitis.

Am. Rev. resp. Dis., 90, 1.

Floyer, Sir John. (1698)

A Treatise of the Asthma.

Wilkins, London.

Forster, R.E. (1957).

Exchange of gases between alveolar air and pulmonary capillary blood: pulmonary diffusing capacity.

Physiol. Rev., 37, 39.

Forster, R.E., (1965)

Interpretation of measurements of pulmonary diffusing capacity.

IN: Handbook of Physiology, Respiration II. Chapter, 59.

Edited by W.O. Fenn and H. Rahn. Waverly Press, Baltimore, U.S.A.

Fouche, R.F., Spears, J.R., and Ogilvie, C. (1960).

Unilateral emphysema.

Br. Med. J., 1, 1312.

Gairdner, W.T. (1850).

On the pathological anatomy of bronchitis and the diseases of the lung connected with bronchial obstruction.

Sutherland and Knox, Edinburgh.

Glazier, J.B., Hughes, J.M.B., Maloney, J.E., Pain, M.C.F., and West, J.B. (1966).

Decreasing alveolar size from apex to base in the upright lung.

Lancet, 2, 203.

Gee, S.J. (1915).

Medical Lectures and Aphorisms. 4th edition.

Oxford Univ. Press, London.

Greenhow, E.H. (1878).

On Bronchitis, and the Morbid Conditions Associated with it.

Longmans, Green. London.

Guz, A. (1966a)

Effects of Vagal Block in Normal Subjects.

In: Breathlessness. p. 65.

Edited by J.B.L. Howell and E.J.M. Campbell.

Blackwell Scientific Publications. Oxford.

Guz, A. (1966b).

Paper to the Medical Research Society. London. (Unpublished).

Haldane, J.S. (1922).

Respiration.

Yale Univ. Press, New Haven.

Haldane, J.S., and Graham, J.I. (1935).

Methods of Air Analysis. 4th edition.

Griffin. London.

Hammond, J.D.S., (1957).

The physical properties of the lungs in chronic cor pulmonale.

Clin. Sci., 16, 481.

Hentel, W., Longfield, A.N., Vincent, T.N., Filley, G.F., and

Mitchell, R.S. (1963).

Fatal chronic bronchitis.

Am. Rev. resp. Dis., 87, 216.

Higgins, I.T.T. (1957).

Respiratory symptoms, bronchitis, and ventilatory capacity
in random sample of an agricultural community.

Br. med. J., 2, 1198.

Himmelstein, A., Harris, P., Fritts, H.W.,Jnr., and Cournand, A. (1958)

Effect of severe unilateral hypoxia on the partition of
pulmonary blood flow in man.

J. thorac Surg., 36, 369.

Howell, J.B.L., (1965)

Breathlessness in pulmonary disease.

In: Breathlessness. p. 165.

Edited by J.B.L. Howell and E.J.M. Campbell.

Blackwell Scientific Publications. Oxford.

Hugh-Jones, P., (1952)

A simple standard exercise test and its use for measuring
exertion dyspnoea.

Br. med. J., 1, 65.

Hugh-Jones, P., (1963)

In: Symposium Chronic Respiratory Disorders.

R.C.P. Edinburgh.

Hugh-Jones, P., Ritchie, B.C., and Dollery, C.T. (1966)

Surgical treatment of emphysema.

Br. med. J., 1, 1133.

Hurst, A. (1959).

Familial emphysema - a review.

Am. Rev. resp. Dis. (Suppl.), 80, 179.

Hutchinson, J. (1846).

On the capacity of the lungs, and on respiratory functions,
with a view of establishing a precise and easy method of detecting
disease by the spirometer.

Med. Chir. Trans. (Lond.), 29, 137.

Jenner, Sir William Bt. (1871).

A System of Medicine. Vol. III, 475.

Edited by J.R. Reynolds. MacMillan. London.

Jensen, K.M. Miscall, L., and Steinberg, I. (1961).

Angiocardiography in bullous emphysema: its role in selection
of the case suitable for surgery.

Amer. J. Roentgenol., 85, 229.

Jones, N.L. (1966).

Pulmonary gas exchange during exercise in patients with
chronic airways obstruction.

Clin. Sci., 31, 38.

Knipping, H.W., Bolt, W., Venrath, H., Valentin, H., Ludes, H.,
and Endler, P. (1955).

Eine neue methode zur prüfung der herz-und lungenfunktion.

Dtsch. Med. Wchnschr., 80, 1146.

Knott, J.M.S., and Christie, R.V. (1951).

Radiological diagnosis of emphysema.

Lancet, 1, 881.

Knudson, R.J. and Gaensler, E.A. (1965)

Surgery for emphysema - a review.

Ann. Thor. Surg., 1, 332.

Krogh, A. and Lindhard, J.

The volume of the dead space in breathing and the mixing
of gases in the lungs of man.

J. Physiol. (Lond.), 51, 59.

Laennec, R.T.H., (1819).

De L'Auscultation Mediate, etc. 1, 211.

Brosson et Chaude, Paris.

Laennec, R.T.H., (1821)

A Treatise on the Diseases of the Chest, etc.

Translated by John Forbes.

Underwood, London.

Larson, R.K., (1965)

Historical note on emphysema.

Am. Rev. resp. Dis., 91, 277.

Laurenzi, G.A., Turino, G.M. and Fishman, A.P. (1962)

Bullous disease of the lung.

Am. J. Med., 32, 361.

Laws, J.W., and Heard, B.E. (1962)

Emphysema and the chest film: a retrospective radiological
and pathological study.

Br. J. Radiol., 35, 750.

Liebow, A.A., (1962).

In discussion. Symposium on Pulmonary Structure and Function.

Ciba Foundation. p. 331.

Edited by A.V. de Reuck and M. O'Connor. Churchill. London.

Macklem, P.T., and Becklake, M.R. (1963)

The relationship between the mechanical and diffusing

properties of the lung in health and disease.

Am. Rev. resp. Dis., 87, 47.

Macleod, W.M., (1954).

Abnormal transradiancy of one lung.

Thorax, 9, 147.

McMichael, J. (1939).

A rapid method of determining lung capacity.

Clin. Sci., 4, 167.

MacNamara, J., Prime, F.J., and Sinclair, J.D. (1959).

An assessment of the steady-state carbon monoxide method
of estimating pulmonary diffusing capacity.

Thorax, 14, 166.

Marshall, R. (1958).

Methods of measuring pulmonary diffusing capacity and
their significance.

Proc. R. Soc. Med., 51, 101.

Martin, W.J. (1949).

The physique of young adult males.

M.R.C. Memorandum No. 20, H.M.S.O., London.

Martin, C.J., Cline, F., and Marshall, H. (1953).

Lebar alveolar gas concentrations: effect on body position.

J. clin. Invest., 32, 617.

Martin, C.J., Marshall H., and Clive, F. (1953).

Lebar alveolar gas concentrations: effect of reduced lung
volumes.

J. clin. Invest., 6, 209.

Martin, C.J., and Young, A.C. (1957).

Ventilation-perfusion variations within the lung.

J. appl. Physiol., 11, 371.

Mattson, S.B., and Carlens, E. (1955).

Lebar ventilation and oxygen uptake in man: influence
of body position.

J. thoracic. Surg., 30, 676.

Mead, J. (1966)

Mechanical Factors in the Control of Breathing

In: Breathlessness. p.139.

Edited by J.B.L. Howell and E.J.M. Campbell.

Blackwell Scient. Publ., Oxford.

Meakins, J.C., and Davies, H.W. (1925).

Respiratory Function in Disease.

Oliver and Boyd, Edinburgh and London.

Medical Research Council, (1960).

Statement by committee on the aetiology of chronic bronchitis.

Br. med. J., 2, 1665.

Medical Research Council (1965)

Statement by committee on the aetiology of chronic bronchitis.

Lancet, 1, 775.

Millard, F.J.C., (1965)

The development and the electrocardiographic diagnosis of
right ventricular hypertrophy.

M.D. Thesis, London.

Mithoefer, J.C. (1965)

Breathholding

IN: *Handbook of Physiology, Respiration II, Chapter 38.*

Edited by W.O. Fenn and H. Rahn.

Waverly Press. Baltimore, U.S.A.

Morgagni, G.B., (1769)

De sedibus, et causis morborum, etc. 2 vols.
Ramondiniana, Venice.

English translation B. Alexander. London.

Nadel, J.A., and Tierney, D.F. (1961)

Effect of a previous deep inspiration on airway resistance
in man.

J. appl. Physiol. 16, 717.

Nash, E.S., Briscoe, W.A., and Cournand, A. (1965).

The relationship between clinical and physiological findings
in chronic obstructive disease of the lungs.

In: *Progress in Research in Emphysema and Chronic Bronchitis*, 2. 305,
Edited by R.S. Mitchell. Karger. Basel and New York.

Nicklaus, T.M., Stowell, D.W., Christiansen, W.R., Renzetti, A.D., Jnr.
(1966).

The accuracy of the roentgenologic diagnosis of chronic
pulmonary emphysema.

Am. Rev. resp. Dis., 93, 889.

Ogilvie, C. (1959).

Patterns of disturbed lung function in patients with
chronic obstructive vesicular emphysema.

Thorax, 14, 113.

Ogilvie, C., and Caterall, M. (1959)

Patterns of disturbed lung function in patients with emphysematous bullae.

Thorax, 14, 216.

Olsen, H.C., and Gilson, J.C. (1960).

Respiratory symptoms, bronchitis and ventilatory capacity in man: An Anglo-Danish comparison, with special reference to differences in smoking habits.

Br. med. J., 1, 450.

Prime, F.J., and Westlake, E.K. (1954).

The respiratory response to carbon dioxide in emphysema
Clin. Sci., 13, 321.

Rahn, H., and Fahri, L.E. (1965)

Ventilation, perfusion and gas exchange - the $\dot{V}A/Q$ concept
In: Handbook of Physiology - Respiration I. Chapter 30.
Edited by W.O. Fenn and H. Rahn.
Waverly Press, Baltimore, U.S.A.

Rankin, J., McNeill, R.S., and Forster, R.E. (1960)

Influence of increased alveolar CO tension on pulmonary diffusing capacity for CO in man.

J. appl. Physiol., 15, 543.

Reid, L. Mc.A. , (1954)

Pathology of chronic bronchitis.

Lancet, 1, 275.

Reid, L.Mc.A. (1960)

Measurement of the bronchial mucous gland layer: Diagnostic yardstick in chronic bronchitis
Thorax, 15, 132.

Reid, L.Mc.A. (1965)

The role of chronic bronchitis in the production of chronic obstructive pulmonary emphysema.

J. Am. Med. Wom. Ass., 20, 633.

Reid, L.Mc.A. and Millard, F.J.C. (1964)

Correlation between radiological diagnosis and structural lung changes in emphysema.

Clin. Radiol., 15, 307.

Reid, L.Mc.A. and Simon, G. (1964)

Unilateral lung transradiancy.

Thorax, 17, 230.

Richards, D.W. (1960).

Pulmonary emphysema: Etiologic factors and clinical forms.

Ann. int. Med., 53, 1105.

Riley, R.L., and Cournand, A. (1949)

Ideal alveolar air and the analysis of ventilation-perfusion relationships in the lungs.

J. appl. Physiol., 1, 825.

Riley, R.L., and Cournand, A. (1951)

Analysis of factors affecting partial pressures of oxygen and carbon dioxide in gas and blood of the lungs.

J. appl. Physiol., 4, 77.

Ruysch, F. (1701-16).

Thesaurus Anatomicus.

Walters, Amsterdam.

Sackner, M.A., Will, D.H., and Dubois, A.B. (1966)

The site of pulmonary vasomotor activity during hypoxia or serotonin administration.

J. clin. Invest., 45, 112.

Scadding, J.G. (1963).

Meaning of diagnostic terms in bronchopulmonary disease

Br. med. J., 2, 1425.

Scadding, J.G. (1965)

A case of chronic bronchitis with Macleod's Syndrome.

Symposium at the Institute of Diseases of the Chest, London.

Br. med. J., 1, 708.

Scadding, J.G. (1966)

Patterns of respiratory insufficiency.

Lancet, 1, 701.

Seebohm, P.M. and Bedell, G.N. (1963)

Primary pulmonary emphysema in young adults.

Am. Rev. resp. Dis., 87, 41.

Severinghaus, J.W. (1960)

Methods of measurement of blood and gas carbon dioxide during anaesthesia.

Anesthesiology, 21, 717.

Siebens, A.A., Grant, A.R., Kent, D.C., Klopstock, R., and Cincotti, J.J. (1957).

Pulmonary cystic disease: physiologic studies and results of resection.

J. thorac Surg., 33, 185.

Simon, G., (1964)

Radiology and emphysema

Clin. Radiol. 15, 293.

Simon, G., (1966)

Radiology in epidemiological studies and some therapeutic trials.

Br. med. J., 2, 491.

Simon, G., and Galbraith, H.J.B. (1953)

Radiology of chronic bronchitis.

Lancet, 2, 850.

Simon, G., and Medvei, V.C. (1962)

Chronic bronchitis: Radiological aspects of a five year follow up.

Thorax, 17, 5.

Simpson, T. (1958).

Chronic bronchitis and emphysema.

Tubercle, Lond., 39, 307.

Simpson, T., Heard, B., and Laws, J.W. (1963)

Severe irreversible airways obstruction without emphysema

Thorax, 18, 361.

Smart, J., Naimi, S., and Capel, L.H., (1961)

The relationship between effort intolerance, spirometry and blood gas analysis in patients with chronic obstructive airway disease.

Br. J. Dis. Chest, 55, 6.

Smyllie, H.C. Blendis, L.M., and Armitage, P. (1965)

Observer disagreement in physical signs of the respiratory system.

Lancet, 2, 412.

Society of Actuaries (1959)

Build and Blood Pressure Study, 1, 16.
Chicago.

Sutinen, S., Christoforedis, A.J., Klugh, G.A., and Pratt, P.C. (1965)

Roentgenologic criteria for the recognition of non
symptomatic pulmonary emphysema: etc.

Am. Rev. resp. Dis., 91, 69.

Sweet, H.C., Wyatt, J.P., Fritsch, A.J., and Kinsella, P.W. (1961)

Panlobular and centrilobular emphysema. Correlation of clinical
findings with pathologic patterns.

Ann. int. Med., 55, 565.

Svanberg, L. (1957)

Influence of posture on the lung volumes, ventilation and
circulation in normals. A spirometric-bronchspirometric
investigation .

Scand. J. clin. Lab. Invest., 9, Suppl. 25.

Tabakin, B.S., Adhikari, P.K., and Miller, D.B. (1959)

Objective long-term evaluation of the surgical treatment
of diffuse obstructive emphysema.

Am. Rev. resp. Dis., 80, 825.

Thurlbeck, W.M., (1963)

A clinico-pathological study of emphysema in an American hospital.

Thorax, 18, 59.

Thurlbeck, W.M., and Angus, G.E. (1963).

The relationship between emphysema and chronic bronchitis
as assessed morphologically.

Am. Rev. resp. Dis., 87, 815.

Thurlbeck, W.M., Fraser, R.G., and Bates, D.V., (1965)

The correlation between pulmonary structure and function
in chronic bronchitis, emphysema and asthma.

In: Progress in Research in Emphysema and Chronic Bronchitis,
2, 295.

Edited by R.S. Mitchell. Karger, Basel and New York.

Towers, M.K. (1966)

Chronic cor pulmonale.

Postgrad. med. J., 42, 506.

Wade, O.L., and Bishop, J.M. (1962).

Cardiac output and regional blood flow.

Blackwell Scientific Publications, Oxford.

Wade, O.L., and Gilson, J.C. (1951).

The effect of posture on diaphragmatic movement and vital
capacity in normal subjects with a note on spirometry as
an aid to determining radiological lung volumes.

Thorax, 6, 103.

Watson, W. (1794)

An account of what happened on opening the body of an
asthmatic person.

Phil. Trans. R. Soc. Med., 54, 239.

West, J.B. (1962).

Regional differences in gas exchange in the lung of the
erect man.

J. appl. Physiol., 17, 893.

West, J.B., (1965)

Topographical distribution of blood flow in the lung.

In: Handbook of Physiology, Respiration II, Chapter 58.

Edited by W.O. Fenn and H. Rahn. Waverly Press, Baltimore.

West, J.B., (1965)

Ventilation/blood flow and gas exchange.

Blackwell Scientific Publications, Oxford.

West, J.B., and Dollery, C.T. (1960).

Distribution of blood flow and ventilation-perfusion ratio

in the lung measured with radioactive CO₂.

J. appl. Physiol., 15, 405.

West, J.B., Holland, R.A.B., Dollery, C.T., and Matthews, C.M.E. (1962)

Interpretation of radioactive gas clearance rates in the lung.

J. appl. Physiol., 17, 14.

West, J.B., and Hugh-Jones, P. (1959).

Effect of bronchial and arterial circulation on alveolar

gas concentrations in the lobe of a dog's lung.

J. appl. Physiol., 14, 743.

West, J.B., and Hugh-Jones, P. (1961).

Pulsatile gas flow in bronchi caused by the heart beat.

J. appl. Physiol., 16, 697.

West, J.B. and Jones, N.L. (1965).

Effect of changes in topographical distribution of lung

blood flow on gas exchange.

J. appl. Physiol., 20, 825.

Whitfield, A.G.W., Smith, O.E., Richards, D.G.B., Waterhouse, J.A.H.,
and Arnott, W.M. (1951).

The correlation between radiological appearances and the
clinical and spirometric state in emphysema.

Q. Med. Jl., 20, 247.

World Health Organization, (1961).

Chronic Cor Pulmonale - Report of an Expert Committee.

Wld. Hlt. Org. techn. Rep. Ser. No. 213.

Williams, C.J.B. (1835)

The Pathology and Diagnosis of Diseases of the Chest, etc.

3rd. edition, Churchill, London.

Williams, H. (1965)

In discussion.

Progress in Research in Emphysema and Chronic Bronchitis.

Vol. 2. p. 304.

Edited by R.S. Mitchell, Karger. Basel and New York.

Williams, H., and Seriff, N.S. (1963).

Chronic obstructive pulmonary disease: an analysis
of clinical, physiologic and roentgenologic features.

Am. J. Med., 35, 20.

Williams, M.H. and Zohman, L.R., (1959)

Cardio-pulmonary function in chronic obstructive emphysema.

Am. Rev. resp. Dis., 80, 689.