

https://theses.gla.ac.uk/

Theses Digitisation:

https://www.gla.ac.uk/myglasgow/research/enlighten/theses/digitisation/

This is a digitised version of the original print thesis.

Copyright and moral rights for this work are retained by the author

A copy can be downloaded for personal non-commercial research or study, without prior permission or charge

This work cannot be reproduced or quoted extensively from without first obtaining permission in writing from the author

The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the author

When referring to this work, full bibliographic details including the author, title, awarding institution and date of the thesis must be given

Enlighten: Theses <u>https://theses.gla.ac.uk/</u> research-enlighten@glasgow.ac.uk

#### THE IDENTIFICATION OF MALE-SPECIFIC TRANSCRIPTS FROM

#### D.MELANOGASTER

A thesis submitted for the degree of Doctor of Philosophy at the University of Glasgow.

Steven R.H. Russell

Institute of Genetics, University of Glasgow.

November 1989

ProQuest Number: 11003325

All rights reserved

INFORMATION TO ALL USERS The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



ProQuest 11003325

Published by ProQuest LLC (2018). Copyright of the Dissertation is held by the Author.

All rights reserved. This work is protected against unauthorized copying under Title 17, United States Code Microform Edition © ProQuest LLC.

> ProQuest LLC. 789 East Eisenhower Parkway P.O. Box 1346 Ann Arbor, MI 48106 – 1346

The research reported within this thesis is my own work, except if otherwise indicated. This work has not been submitted for any other degree.

(一) 他们的"合当

# Steven R.H. Russell

he been started as a service to be a started as y -

TORNY LARCET TO YOR YREY LOOSON

· 11 ·

# **CONTENTS**

Contents						
Abbreviations						
Acknowledgements						
Summary				-	vii	
Chapter	1	_	INTRODUCTION	-	1	
	•				20	
Chapter	2	-	MATERIALS & METHODS	-	39	
	2.1	-	Materials	-	40	
	2.2	-	Methods	-	41	
	2.21	-	Nucleic Acid Isolation	-	41	
	2.211	-	In-vitro DNA Manipulation	-	43	
	2.2111	-	Electrophoresis And Blotting		43	
	2.2iv		Radiolabelled Nucleic Acids	-	44	
	2.2v	-	Construction Of Drosophila	-	45	
			cDNA Libraries			
	2.2vi	-	Screening Recombinant Lambda	-	47	
			Libraries			
	2.2vii	-	<u>In-situ</u> Hybridisation To	-	47	
			Polytene Chromosomes			
	2.3	-	Drosophila	-	48	
Chapter	3		ISOLATION OF SEX-SPECIFIC GENES	-	50	
			FROM DROSOPHILA			
	3.1		Introduction	-	51	
	3.2	-	Results	-	56	
	3.2i	-	Construction Of cDNA Libraries	-	56	
	3.2ii	_	Differential Screening Of The		59	
			Male cDNA Library			
	3.2iii	_	Differential Screening Of A	-	60	
			Drosophila Genomic Library			
	3.2iv		Reverse Northern Analysis	-	62	
	3.3	_	Discussion	-	64	
Chapter	4	_	IDENTIFICATION OF PREVIOUSLY	-	67	
-			ISOLATED MALE-SPECIFIC GENES			
	4.1	_	Introduction	-	68	
	4.2	_	Results	-	70	

	4.2i	_	Clone gK14	-	70
	4.2ii	-	Clone gK21	-	71
	4.3	-	Discussion	-	72
Chapter	5	_	NOVEL MALE-SPECIFIC GENES WITH	-	74
			PUTATIVE GERMLINE EXPRESSION		
	5.1	-	Introduction	-	75
	5.2	-	Results	-	78
	5. <b>2</b> i	-	Clone gS1	-	78
	5.2ii	-	Clone gS10	-	80
	5.2iii	-	Clone gK33	-	82
	5.3	-	Discussion	-	82
Chapter	6	-	MALE-SPECIFIC GENES WITH UNUSUAL	_	86
			GENOMIC ORGANISATION		
	6.1	-	Introduction	-	87
	6.2	-	Results	-	93
	6.2i	-	Clone gS4	-	93
	6.2ii	-	Clones gS8 And gS9	-	96
	6.3	-	Discussion	-	98
Chapter	7	_	A COMPLEX GENE ?	-	102
	7.1	_	Introduction	-	103
	7.2	-	Clones gS2 And gK15	-	103
	7.3	-	Discussion	-	106
Chapter	8	-	CONCLUDING REMARKS	-	112
		_	BIBLIOGRAPHY	_	116

A statistical and the second statis

1.1

iv

# ABBREVIATIONS

ATP	-	adenosine triphosphate					
bp		base pairs					
CDNA		complementary DNA					
Ci		Curies					
CM		centimetres					
cpm		counts per minute					
°c	-	degrees centigrade					
DNA		deoxyribonucleic acid					
dntp	-	deoxynucleotide triphosphate					
DTT		dithiothreitol					
EDTA	-	ethylenediamine tetra-acetic acid					
EtBr		ethidium bromide					
g		grammes					
g		centrifugal force equal to gravitational acceleration					
hr	-	hours					
kb	-	kilobases/kilobasepairs					
kd	-	kilodaltons					
1	_	litres					
L1,2,3	-	1st, 2nd, 3rd larval instars					
М	-	Molar					
min	-	minutes					
ml	-	millilitres					
mM	-	milliMolar					
mMol		milliMoles					
mRNA	-	messenger RNA					
NaPPi	-	sodium pyrophosphate					
ng	-	nanogrammes					
nt	-	nucleotides					
OD	-	optical density					
pfu	-	plaque forming units					
рН	-	acidity [-log <sub>10</sub> (Molar concentration of H <sup>+</sup> ions)]					
RNA	-	ribonucleic acid					
rpm	-	revolutions per minute					
rRNA	-	ribosomal RNA					
SDS	-	sodium dodecylsulphate					
Tris	-	tris(hydroxymethyl) amino ethane					
u	_	units					
uCi	-	microCuries					
ug	-	microgrammes					
ul	-	microlitres					
vol	-	volume					

I gratefully acknowledge the scientific and technical staff of the genetics department for support, encouragement and (for the most part) good natured tolerance. I am for making the last four indebted to my fellow students years an enjoyable and interesting experience, especially Dee Special thanks to everyone who has kept me and Mark. copiously supplied with the wares of T.C. Reids (probably most of the department). Kim has my gratitude for a rigorous scientific training and for coming top of the "buying Steve a drink" league table. Last but not least special thanks to Fiona not only for her editing skills which have helped make this thesis far more readable than it would have been otherwise but also for putting up with me for so long.



#### SUMMARY

A primary Lambda genomic library was screened with CDNA probes derived from male and female 3rd instar larval mRNA. Twenty clones which show male-specific or male-elevated expression were identified. Ten of these clones have been characterised in some detail. Two of the clones, qK14 and aK21. appear to be re-isolates of the previously identified male-specific genes mst349 and mst336 (DiBenedetto et al., 1987). These, along with five new isolates (gS4, qS8, qS9, gS10, gK33) show expression patterns on northern blots consistent with germline-specific expression. The transcripts are expressed in male larvae, pupae and adults but not in agametic males or XX individuals transformed to somatic males by the tra2 mutation (pseudomales).

Two of these clones, gS8 and gS9, are non-overlapping cross hybridise at high stringency indicating that but they contain the same or very similar sequences. Partial CDNA identify a single band in female DNA but multiple clones in male DNA, suggesting the transcribed sequence is Y bands associated. Many of these male-specific bands are conserved at least 3 different laboratory strains. Northern blots in using RNA from XO and XY males show similar levels of the transcript suggesting that the non sex-specific locus can be transcribed, at least in the absence of a Y chromosome. Insitu hybridisation to polytene chromosomes with either of the genomic clones give no signal suggesting that the non sexunder replicated specific location is genomic or heterochromatic.

Another of these germline-specific clones, gS4, appears to be heterochromatic. Southern blots of genomic DNA indicate the presence of several different repeats. In addition <u>in-</u> <u>situ</u> hybridisation to polytene chromosomes indicate that these repeats reside at a single genomic location: at the base of the left arm of chromosome 2.

Of the three remaining clones, gS1 identifies a weakly expressed 3.0kb male-specific transcript. Although this transcript is not detected in agametic males low levels of expression during the pupal stage suggests that it may not be germline specific.

The other two clones gS2 and gK15, are overlapping and

map uniquely to region 61F on the polytene chromosome map. These clones identify at least 3 male-specific transcripts on northern blots as well as a non sex-specific transcript which is elevated during the embryonic and pupal stages. The pattern of expression appears to be complex, all of the transcripts are developmentally regulated. One of the malespecific transcripts is expressed in agametic males and possibly in pseudomales. This transcript is also present in the embryo although it is not known at present if it is sexspecific at this stage.

## CHAPTER 1

#### INTRODUCTION

and standard with bring

a second design and a grade the second

n an trainn an an trainn 1995 - Maria Sangara, an trainn an trainn an trainn an trainn an trainn 1997 - Maria Sangara, an trainn an trainn

e e la companya de la

•

The end result of the pathways which regulate the different aspects of sexual dimorphism in the fly is the expression of the structural genes responsible for sexspecific phenotypes. It is here that we can see the similarity between the control of the sexual phenotype and way in which homeotic selector genes specify structural the features of the body segments. As with the homeotic genes the sex determination genes make binary decisions which result in a cell or tissue choosing a particular pathway of development. Thus by studying the way in which the sex determination hierarchy co-ordinately regulates sets of genes we may gain some insight into the way in which developmental processes are regulated. Furthermore sex-specific qene expression is not regulated in isolation from the rest of the organism: temporal and positional information must be coordinated along with the specification of sex to ensure that genes are expressed in the correct place at the correct time. Thus the study of sexual differentiation may give some us insight into the way in which different regulatory hierarchies interact to specify the appropriate expression of a particular gene.

In this chapter I shall review our current understanding of the regulation of sexual differentiation as a prelude to a discussion of a set of male-specific genes which I have isolated. As we shall see the extensive knowledge of the control of sex-specific gene expression offers a unique opportunity to investigate the molecular mechanisms which underlie complex problems in development.

Before discussing the genetic and molecular aspects of sex-specific gene expression I will briefly discuss the biology of some of the aspects of sexual differentiation.

# Development Of The Sexes

Pole cell nuclei, the precursors of the germline, separate from the somatic nuclei prior to the blastoderm stage. They proliferate at the posterior end of the embryo until there are approximately 50 cells. These cells are then internalised and migrate, during gastrulation, to the level of the 5th abdominal segment. Genetic evidence, which will be

discussed later, suggests that the pole cells have already determined their sex by this time. At approximately the 10th. hour of embryogenesis a few of the pole cells begin to associate with mesodermal cells and during the next hour the pole cells become completely encapsulated by somatic cells and the gonad is formed (Campos-Ortega and Hartenstein, 1985).

Sonnenblick (1950) reports that there is a bimodal distribution in the number of pole cells encapsulated. In one population an average of 5-7 pole cells are incorporated into the gonad, in the other a larger gonad contains an average of 9-13 pole cells. He proposed that this difference reflects the first manifestation of sexual dimorphism in the fly; the larger gonad being male and the smaller female. Although definitive confirmation of this hypothesis awaits, it is supported by studies of gonad development during the larval 1931). These show that upon hatching a stage (Keris, size difference in the gonads is apparent and can be reliably used to sex 1st instar larvae. The gonads of 1st instar larvae are also sexually dimorphic in terms of the position they occupy relative to the fat body. While the smaller female gonads are embedded in the fat body the larger male gonads are merely bordered by this tissue.

Ablation of the pole cells by UV irradiation prior to migration does not appear to affect the sexual their dimorphism of the somatic component of the gonad (Aboim, 1945). Similarly the effect of grandchildless mutations (eg.gS(1)N26 and gS(1)N441, Niki and Okada, 1981; agametic, Engstrom et al., 1982; grandchildless of D.subobscura, Fielding, 1967), which result in the loss of pole cells either prior to or after they reach the gonad, appear not to affect the sexual phenotype of the somatic component of the gonad. However in the absence of germ cells the gonads do not develop to full maturity, though they are recognisable as female contain all the somatic male or and appear to these Taken together components of the wild type gonad. observations suggest that the somatic component of the gonad sexually determined independently of the germline. This is is further supported by observations on the suggestion development of the gonad in individuals mutant in somatic sex determination genes. As will be discussed later the somatic determination hierarchy controls the sexual sex

differentiation of the somatic component of the gonad but plays no role in the sexual differentiation of the germline. Thus it is probable that in at least one tissue, the gonad, the sexually determined state is set early in development.

Apart from the gonads, which increase in size, overt sexual dimorphism is not apparent through most of the larval stages. Lauge (1982) however has reported that sexually dimorphic differentiation of the genital imaginal disk becomes apparent during the second larval instar. Major sexual differentiation occurs during metamorphosis at the genital stage. During this stage the gonads and the pupal disk differentiate to form the adult reproductive tract. The following outline, which is by no means complete, has been drawn from Lauge (1982), Mahawold and Kambysellis (1980) and Tokuyasu (1980). A diagram of the male and Lindsley and reproductive tracts is presented female internal in Figure 1.1.

Early investigators believed that the single genital disk contained indifferent cells capable of forming either male or female derivatives depending on the chromosomal constitution these cells. However clonal analysis (Nothiger of et al., 1977; Schupbach et al., 1978) demonstrates that the single genital disc contains three distinct primordia. One specifies structures, one specifies female structures and the male third is an indifferent primordium of the analia. The genital primordia specify the formation of the internal disk genitalia and the external structure of the 8th and successive abdominal segments.

The male reproductive system consists of a pair of coiled testes joined to the seminal vesicle by a short testicular duct or vas deferens. From the seminal vesicles the sperm exit, via the anterior ejaculatory duct, to the sperm pump then via the posterior ejaculatory duct to the and penis apparatus. The accessory glands, which produce secretions such as pheremonal peptides, join the anterior ejaculatory duct near the seminal vesicles. During the early stages of metamorphosis the testes develop apart from the disc derivatives, joining the vas deferens mid way through the pupal stage. Attachment to the vas deferens induces the testes to elongate and coil. Later in development the testes assume their characteristic yellow colour which spreads to the vas deferens.





Figure 1.1a: Diagram of the male internal reproductive tract from <u>D.melanogaster</u>. After Lindsley and Tokuyasu (1980).

b: Diagram of the female internal reproductive tract from<u>D.melanogaster</u>. After Mahowald and Kambysellis (1980). External male characteristics include dark pigmentation over all of the posterior tergites, fusion of the 6th and 7th segments and the genital arch which forms the claspers and supports for the penis apparatus.

The female portion of the genital disk produces a pair of laterel oviducts which extend anteriorly to meet the developing ovaries. Posteriorly the oviducts fuse to form a common oviduct leading to the uterus. A pair of spermatheaca and а single sperm receptacle open into the uterus. These function as sperm storage organs. A pair of accessory glands parovaria, which produce secretion for coating the or egg, also open into the uterus. Externally the female differs from male in abdominal pigmentation, which is and the reduced posterior margin of the tergites. restricted to the In addition the 6th and 7th segments remain separate the and genital arch forms the vaginal plate or ovipositor.

analysis and gynandromorph studies indicate Clonal that only one of the genital disk primordia develops in males or females and that the choice is determined by the somatic sex determination hierarchy. In true intersex flies, such as those produced by mutations in the gene doublesex, both primordia develop and structures derived from both are visible in the adult (Hildreth, 1965).

The other major site of external sexual dimorphism is the basitarsus of the fore-leq. In males there are generally 6 rows of transverse bristles and 1 row of longditudinally arranged thick heavily pigmented bristles known as the sex Females have on average 8 rows of transverse bristles comb. but no longditudinal bristles (Tokunaga, 1962). Intersexual individuals tend to have 6 rows of transverse bristles and a of bristles lying in a position intermediate between row transverse and longditudinal rows. These bristles are also intermediate in size and pigmentation when compared with the The third obvious normal bristles. sexual sex comb and of body size, females are larger than dimorpism is that It is not clear how this dimorphism is controled but males. sex transforming mutations in <u>Sex-lethal</u> can transform body somatic 1984). Mutations in the sex size (Cline, hierarchy do not affect this particular determination dimorphism.

As well as the above dimorphisms internal differences are also apparent in biochemistry (eg yolk protein synthesis) and

behaviour. These observations suggest that the sexually determined state may extend to most, if not all, somatic tissues. This hypothesis is supported by the finding that embryonically derived cell lines (KC Schnieder-2) and maintain either male or female expression of sex determination genes but not both (M.McKeown, pers.comm.).

# Spermatogenesis in Drosophila

This brief account of spermatogenesis is drawn from the descriptions of several authors ( Lindsley and Tokuyasu, 1980; Hennig, 1985; Lifschytz, 1987), due to constraints of space oogenesis will not be discussed and interested readers referred to the review by Mahawold and Kambysellis are (1980).

During spermatogenesis in <u>Drosophila</u> a group of 64 haploid sperm are produced from a single gonial cell. The production of mature spermatozoa takes approximately 10 days. cells are generated at the apex of the testis by the Gonial unequal division of a stem cell. The stem cells, which are in fact the pole cells which populated the embryonic qonad, anchored at the apex. The gonial cell is invested by remain apical somatic cells to form a cyst. The cyst will two contain the descendants of the gonial cell during the entire period of sperm development. The gonial cell within the cyst undergoes four rapid mitotic divisions to produce 16 primary Cytoplasmic bridges continue to link the spermatocytes. daughter cells. Thus the gametes within a cyst develop as а syncytium. Primary spermatocytes are transcriptionaly active the mRNA required for subsequent sperm produce all and Autoradiographic studied on tritiated uridine development. uptake have failed to detect transcription in post-meiotic stages of spermatogenesis (Olivieri and Olivieri, 1965). The spermatocytes grow for approximately four days, increasing in During this stage the nuclei are fold. volume some 25 at meiotic prophase. It is also during this stage arrested chromosome becomes active. At the end of the that the Y meiotic divisions produce the 64 spermatocyte stage spermatids which again remain interconnected by cytoplasmic bridges. As noted above it is generally accepted that there

is little or no postmeiotic transcription, spermatids are however translationally active.

The spermatids develop synchronously within the cyst, in can only be described as a remarkable morphological what differentiation, to produce the 1.8mm long sperm (the sperm of <u>D.hydei</u> are even longer, attaining a length of over 10mm !). The major features of this differentiation are: 1) The condensation and organisation of the mitochondria into the nebenkern structure. 2) The elongation of the centriole to form the axoneme and its subsequent association with the nebenkern. 3) The elongation of the axoneme and surrounding nebenkern derivatives to form the tail of the sperm. 4) The condensation and elongation of the spermatid nucleus. 5) Separation of each sperm into a membrane bound individual by the passage of a "cystic bulge", which also rids the sperm of excess cytoplasm. 6) The coiling and final release of the from the cyst into the seminal vesicle. To put this sperm morphological change into some perspective, if we consider the size of a pre-elongation spermatid to be equivalent to the size of a standard postage stamp, then the resulting sperm would be equivalent to a piece of sewing thread remarkable approximately 6 feet long! This is all the more when we consider that this differentiation occurs the in absence of any on-going transcription. In a sense it can be viewed as a gigantic piece of self assembly.

testis of the newly hatched larva contains The approximately 38 gonial cells (Sonnenblick, 1950; Keris, 1931) but no primary spermatocytes. According to Garcia-Bellido (1964) the first spermatocytes become apparent during larval instar. Unfortunately there the 1st are no descriptions of spermatocyte numbers during the larval stages. During the pupal stage there may be a maximum of approximately 100 spermatocyte cysts per testis (Garcia-Bellido, 1964). This author reports a reduction to around 50 cysts per testis post-eclosion. A genetic analysis estimates 30 cysts per testis during adult life (Lindsley and Tokuyasu, 1980), it is not clear if these differences are significant. spermatids are detected during the larval stages, the No first meiotic divisions occur around the time of pupariation.

As one may suspect such a complex process is susceptible to disruption by a large number of mutations (see Lifschytz 1987 for an extensive review). Apart from autosomal and X-

linked point mutations various chromosomal aberrations, including X to autosome translocations and certain proximal X deficiencies, also disrupt spermatogenesis. The Y chromosome, dispensable for the rest of development, also has a vital role in spermatogenesis. some of these unusual chromosomal mutations will be considered in chapter 6.

# The Genetics of Sex Determination

Sex in Drosophila melanogaster is determined by the ratio X chromosomes to sets of autosomes (X:A ratio). Bridges of (1921) demonstrated, by studying individuals with aneuploid and polyploid chromosomal constitutions, that an X:A ratio of > 1 results in female differentiation while a ratio of < 0.5 results in male differentiation. Individuals with an intermediate X:A ratio of 0.67, as with the genotype XX:AAA (triploid intersexes), develop as mosaics composed of both male and female tissue. The observation that haplo-X tissues still differentiate as phenotypically male when lacking a Y XX/XO mosaics or in XO individuals, chromosome, in as Y chromosome plays no part in demonstrates that the sex sex determining also suggests that the determination. It mechanism operates at the level of the individual cell. This latter suggestion is strongly supported by the results of somatic recombination experiments which generate clones of cells homozygous for sex transforming mutations (Baker and Ridge, 1980; Sanches and Nothiger, 1982).

The choice of a particular pathway of sexual development has three major consequences for the fly: 1) Somatic tissues differentiate in a sexually dimorphic way. 2) The cells of the germline are directed to the appropriate developmental 3) The level of transcription of X-linked genes is pathway. regulated in order to compensate for the imbalance in Хlinked gene dosage between males and females. This is by transcribing most genes on the single Х achieved chromosome of males at twice the rate of those same genes in the diplo-X female (Lucchesi and Manning, 1987). This is in contrast to the mammalian X inactivation mechanism of dosage compensation (Cattanach, 1975).

The use of genetic, and latterly molecular, techniques to

study mutations which perturb these functions has led to an enormous increase in our understanding of the control of sexual differentiation. This has been particularly fruitful in the case of those genes regulating somatic sex determination, with the result that the somatic sex determination hierarchy is one of the best understood examples of the molecular control of gene expression in any complex eukaryote. Rapid advances have been possible for а number of reasons: 1) The wealth of genetic and molecular techniques available to the <u>Drosophila</u> investigator. 2) The non-lethal nature of many of the mutations which perturb differentiation, allowing the examination of mutant **sex**ual phenotypes in adult individuals where sexually dimorphic features are most easily observed. 3) The ability to study lethal in mosaic individuals those mutations which are without experiencing interference from neighbouring nonmutant tissue.

I shall overveiw our current understanding of the control of sexual differentiation and follow this with a more detailed description of the individual genes involved. The somatic sex determination hierarchy, being the easiest to is understood in the greatest detail. Dosage study, compensation and germline sex are more poorly understood. Dosage compensation has recently been the subject of an extensive review (Lucchesi and Manning, 1987) and will only dealt with briefly in this discussion. A diagrammatic be representation of the genetic interactions is presented in Figure 1.2.

As mentioned above the primary determinant of sex is the X:A ratio. Early in development individual cells, in response to this ratio, set the state of activity of the key gene Sexlethal(Sxl) such that it is "on" in XX zygotes and "off" in zygotes (Cline, 1988). Sxl requires the activity of both XY maternal (eg <u>daughterless</u>) and zygotic (eg <u>sisterless-A</u>) genes to initiate or maintain the appropriate expression state. After assessment of the X:A ratio Sxl maintains the sexually differentiated state independently of the initiating signal. Somatic sex, germline sex and dosage compensation all appear to be regulated by <u>Sxl</u>. Downstream of this gene their for the most part mediated by independent control is pathways.

Sex-specific gene expression in the soma is regulated by



hierarchy of four known genes: transformer а (tra), <u>transformer2</u> (<u>tra2</u>), <u>doublesex</u> (<u>dsx</u>) and <u>intersex</u> (ix). No other regulatory genes have been identified in this pathway, although their existence cannot be discounted. Baker and (1980) proposed a model for the interactions Ridge of the above loci based on genetic evidence. Subsequent molecular studies (Nagoshi et al., 1988; Goralski et al., 1989) strongly support and refine this model. The female-specific activity of Sxl is required for the production of а functional <u>tra</u> mRNA. The <u>tra</u> gene product is in turn required, in conjunction with the product of the tra2 gene, to direct the production of female-specific dsx mRNA. The female-specific dsx gene product acts, in conjunction with the <u>ix</u> gene product, to repress male terminal differentiation functions. Female terminal differentiation functions are not repressed and hence female somatic development ensues. In the absence of Sxl function, as in XY individuals, tra femalespecific mRNA is not produced. tra2 alone is not sufficient to direct the production of the female-specific dsx mRNA and male-specific <u>dsx</u> mRNA is produced by default. The <u>dsx</u> male product has the opposite effect to the female product, female terminal differentiation functions are repressed and male functions are expressed. The molecular mechanisms by which <u>dsx</u> acts to control terminal differentiation are as yet unknown.

necessary for functions the Dosage compensation hyperexpression of X-linked genes in males are negatively regulated by <u>Sxl</u>. In males, where <u>Sxl</u> is absent, the products male-specific-lethal 1 and 2 and male less of the (collectively referred to here as <u>msl</u>s) genes are able to direct X chromosome hypertranscription, at least in the later stages of development. In the presence of <u>Sxl</u> these functions repressed. There is substantial evidence that other, as are unidentified, functions are required for correct dosage yet compensation in the early embryo (Cline, 1984; Gergen, 1987; Gergen and Wieschaus, 1988).

known about the correct development of the Less is germline. It has been established that the correct X:A ratio and the appropriate activity of <u>Sxl</u> are required (Schupbach, Steinman-Zwicky et al., 1989). The regulatory qenes 1985; in somatic sex and dosage compensation are not involved in the germline for sexual differentiation (Marsh required

and Wieschaus, 1978; Schupbach, 1982), however tra2 and mle gene products are required for the successful completion of spermatogenesis (Belote and Baker, 1983; Bachiller and Sanchez, 1986). The product of the Ovo locus is necessary for the establishment of the female germline (Oliver et al., 1987). Undoubtedly more genes remain to be uncovered which are necessary for germline sex. Interestingly recent experiments (Steinmann-Zwicky et al., 1989) have shown that in addition to the cell autonomous control of germline sex an inductive interaction between germline and soma can be demonstrated for XX germ cells. Thus, in this case at least, sex determination is not cell autonomous.

#### <u>Sex-lethal(Sxl)</u>

<u>Sxl</u> is a central gene which regulates all aspects of sex determination in Drosophila. The gene maps to salivary gland chromosome region 6F on the X chromosome. Much of our understanding of the role Sxl plays in sexual differentiation from the study of various mutations which perturb the comes function of this gene in a sex-specific fashion. Simply stated loss of function mutations are female-specific lethals whereas gain of function mutations are male-specific lethals. study of more subtle mutations in <u>Sxl</u> and its positive The regulators, principally in the laboratory of T.W. Cline, has allowed the separation of different <u>Sxl</u> functions and the elucidation of a genetic model for its action. Subsequent molecular cloning of the gene has provided some insight into how sexual differentiation is controlled at the molecular level.

#### Genetics of Sxl

Both null alleles (eg  $\underline{Sxl}^{f1}$ ) and deletions are homozygous lethal in diplo-X (female) individuals but have no discernible effect on the viability or development of haplo-X (male) individuals. Gain of function mutations (eg  $\underline{Sxl}^{M1}$ ) are dominant male lethal. They appear to have no effect on

otherwise wild type females, rather these alleles are able to females from mutations in positive regulators of <u>Sxl</u> rescue (eg <u>daughterless</u>, <u>da</u>) which would otherwise be female lethal (Cline, 1978, 1979). Due to the lethal nature of these mutations much of the early work focused on clonal analysis either in gynandromorphs or in homozygous clones generated by somatic recombination. These experiments indicated that: a) mutations are sex-transforming.  $Sxl^{f1}/Sxl^{f1}$ Sxl clones undergo male somatic differentiation in a diplo-X background Sx1<sup>M1</sup> (Cline, 1983) and somatic clones undergo female differentiation in a haplo-X background. b) The lethality associated with these mutations may be due to perturbations in dosage compensation as the clones were both underrepresented and poorly developed, indicating disruption of a vital function. These conclusions are supported by studies which examined the effect of these alleles in triploid intersexes (Cline, 1983). Here gain of function alleles are feminising and result in reduced expression of the X-linked genes Bar and Hairy wing. On the other hand loss of function alleles are masculinising and appear to some extent to Lucchesi increase the expression of X-linked genes. and provided evidence (1981)have also from Skirpsky transcriptional autoradiography studies which implicates Sxl in the regulation of dosage compensation.

By using a variety of selection procedures various hypomorphic <u>Sxl</u> alleles have been isolated. The study of these alleles in a variety of genetic backgrounds has allowed the dissection of <u>Sxl</u> function. The most informative of these will be discussed.

is a derivative of <u>Sxl<sup>M1</sup> which</u> is Sxl<sup>fm7,M1</sup> hemizygous males and homozygous viable in females (Cline, viable in 1984). It retains the capability of the parent allele to chromosomal females from maternal da mutations. rescue Females carrying this allele are somatically sex-transformed, indicating that female somatic sex determining functions have been lost. Somewhat surprisingly these females have male body size. Sxl is the only sex determining gene which affects this pronounced sexual dimorphism.

The viability in both sexes observed with this allele suggest that if <u>Sxl</u> controls dosage compensation then appropriate functions are being provided in both sexes. Two copies of this allele allow the repression of (female-lethal)

X chromosome hyperactivation genes whereas one copy is not sufficient for this purpose (ie males survive). Evidence for the involvement of  $\underline{Sxl}$  in establishing the correct level of Xchromosome transcription comes from an examination of this allele in  $Sxl^{fm7}$ , M1/Df(Sxl) hemizygous females. (XX) individuals to survive beyond the late fail larval stage, however when these individuals are simultaneously homozygous for any of the male-specific-lethal (msl) loci the lethality is partially rescued. This indicates that it is the inappropriate expression of <u>msl</u>s which is ,at least in part, responsible for the lethal effect. However this rescue is only observed with this particular <u>Sxl</u> allele. Other femalelethal alleles, used both in the above study and others (Cline, 1983; Skripsky and Lucchesi, 1982), are not rescued by <u>msl</u> mutations. This observation can be explained if we assume that there are two dosage compensation pathways, one operating early in development and the other, regulated by <u>msl</u>s, operating later. If this is correct we can reconcile Sxl<sup>fm7,M1</sup> the <u>msl</u> rescue of hemizygous individuals by assuming that this allele is defective primarily in its interaction with late dosage compensation functions. Consistent with this view is the late lethal period observed hemizygous females. That this allele is hemizygous male in retains the constitutive male-lethal viable but still mutation reflects the selection for a male viable derivative. This must have leaky interactions with dosage compensation functions to allow males to survive (Cline, 1984).

Other lines of evidence support the contention that there is two dosage compensation pathways: 1) There а are discrepancy between the lethal phases of Sxland msl mutations, the latter cause male lethality during the late larval or early pupal stages while the lethal former are considerably earlier (early larval for  $\underline{Sxl}^{M1}$  and late embryo Experiments on hypomorphic <u>runt</u> alleles  $Sxl^{\pm 1}$ ). 2) for indicate that this gene is dosage compensated in the early embryo during the blastoderm stage (Gergen, 1987; Gergen and Wieschaus, 1987). When individuals carry mutations in any, or the <u>msl</u>s there is little or no effect on runt all, of In contrast the mutation  $\underline{Sxl}^{f1}$ appears to expression. individuals, XX this is in expression increase runt early dosage to repress failure with а consistent compensation functions which do not involve <u>msl</u>s. The effect

of the <u>Sxl<sup>M1</sup></u> allele was also examined in these experiments, it did not show the expected repression of <u>runt</u>. Subsequent experiments (Steinmann-Zwicky, 1989) suggest that this allele is not constitutive for all <u>Sxl</u> functions when the X:A ratio is inappropriate for normal <u>Sxl</u> expression.

Genes involved in an early pathway for dosage compensation would be expected to show an embryonic male lethal phenotype when mutated. Two such loci have been reported, killer of male A and killer of male B (Pierre, 1972). Disappointingly further reports on these mutations have not been published nor are they listed in any of the stock centres.

The studies described above allow us to separate genetically two distinct <u>Sxl</u> functions; that regulating **so**matic sex and that which regulates dosage compensation. In a further series of experiments Cline was able to establish a third function for <u>Sxl</u>, that of a positive autoregulator. If fly of the genotype  $Sxl^{fm7,M1}/Sxl^+$  (XX) is produced from a da/da mothers these individuals develop with large patches of female soma. Since the wild type allele in this stock cannot be activated in the absence of maternal <u>da</u> product and the **mu**tant allele specifies male soma, the appearance of female somatic tissue must result from activation of the wild type copy by the partially constitutive  $Sxl^{fm7,M1}$  allele. Thus an autoregulatory function for <u>Sxl</u> is uncovered (Cline, 1984).

Studies involving other partial loss-of-function alleles (Sxl<sup>f9</sup>, Sxl<sup>fLS</sup>, Maine et al., 1985a; Salz et al., 1987) have further dissection of <u>Sxl</u> function. Somatic allowed recombination experiments show that homozygous <u>Sxl<sup>fLS</sup>(XX)</u> clones develop as male tissue. In similar experiments homozygous  $Sxl^{f9}(XX)$  clones have a female phenotype. However when these alleles are examined in <u>da</u> backgrounds <u>Sxl</u>fLS **ap**pears more wild type than <u>Sxl<sup>f9</sup></u>: ie. it is more able to tolerate reductions in the amount of <u>da</u> gene product. The above observations can be reconciled if we assume that <u>Sxl<sup>f9</sup></u> is defective in initiating stable expression (da sensitive) started it can maintain appropriate expression but once (female soma). The opposite would hold for <u>Sxl<sup>fLS</sup></u>: it can initiate expression but cannot maintain it.

A requirement for correct <u>Sxl</u> activity in the female germline has been demonstrated by Schupbach (1985). Pole cell transplantation experiments show that  $\frac{Sxl^{f1}}{Sxl^{f1}}$ (XX) germ

in wild type soma develop as undifferentiated cells multicellular cysts. The allele <u>Sxl</u>fs1 is homozygous female viable and wild type with respect to somatic function but it shows the same germline phenotype as the <u>Sxl<sup>f1</sup></u> null allele. Interestingly а heteroallelic combination  $Sxl^{fm7,M1}/Sxl^{M1,fm3}$ , which in somatic tissue produces male development, permits normal oogenesis with wild type eggs produced. Again different <u>Sxl</u> functions can being be separated by genetic means.

The picture which emerges from these studies is of a complex gene involved in assessing the primary sex determining signal, establishing the appropriate response to the signal, retaining a memory of the signal via an autoregulatory mechanism and finally transmitting that assessment to the regulatory genes which control somatic sex, germline sex and dosage compensation (Cline, 1984).

# Molecular Analysis of Sxl

The <u>Sxl</u> locus was cloned via a P-element insertion allele <u>Sxl<sup>fPb</sup></u> (Maine et al., 1985a; 1985b; Salz et al., 1987). Subsequently molecular studies have been used to investigate the organisation of the gene (Salz et al., 1987,1989; Bell et al., 1988). The results of these investigations can be summarised as follows: The Sxl transcription unit spans over 23kb of DNA and encodes multiple overlapping transcripts which are regulated in a sex-specific, tissue-specific and temporal way. Ten transcripts have so far been distinguished on northern blots.

Three of these are embryo specific and are transcribed only during the earliest stages of development (2-5 hr). They may be transcribed from an embryo-specific promoter located downstream from the start site used during the rest of development. The presence of <u>Sxl</u> alleles believed to be defective in pathway initiation and evidence from P-element mobilisation experiments are consistent with the presence of a regulatory element involved in establishing <u>Sxl</u> expression (Salz et al., 1987).

The embryo-specific transcripts are replaced by a set of seven transcripts which persist throughout the remainder of

the life cycle. Three of these transcripts are male specific four are female-specific. Two of the female-specific and transcripts depend on the presence of a germline. Sequence analysis of cDNA clones corresponding to some of the <u>Sxl</u> transcripts indicates that all of the male-specific mRNA species contain a 190bp exon which is not present in female embryonic mRNA's. It is not clear at present or if the embryo-specific primary transcripts contain this exon. The contains an in-frame termination codon which severely exon truncates the conceptual <u>Sxl</u> polypeptide. As male and female transcripts are believed to share the same start site it would appear, therefore, that sex-specific regulation of <u>Sxl</u> throughout the majority of the life cycle is not at the level of transcription: rather it is at the mRNA processing stage.

four female transcripts appear to differ The at their 3'ends by the use of alternative exons. At present it is not known if these sequence differences reflect functional differences. However given that two of these transcripts are germline specific it is highly probable that the transcripts encode different functions. All of the female-specific transcripts are found in unfertilised eggs. They appear to be rapidly degraded when the zygote-specific transcripts first is unclear what role these maternally derived appear. It transcripts play in early development but since Sxlthese the simplest view would be that autoregulates transcripts have no function. Rather they are present as a consequence of their transcription in the female germline. In of this it should be noted that Sxldependent support maternal effects cannot be detected (Cline, 1980; Oliver et al., 1988).

alleles all contain dominant male-lethal Five transposable element insertions within 1kb of the malespecific exon (Maine et al., 1985b). These all appear to be downstream of the 3' acceptor site for this exon. of Two insertions are <u>B104</u> elements which are transcribed in these the opposite direction to <u>Sxl</u>, two are <u>297</u> elements which are transcribed in the same direction, the fifth is of unknown The dominant gain of function phenotype associated origin. with these insertions could, in principal, be explained in several ways. The three dimensional structure of the primary transcript could be disrupted by these insertions such that splicing at the terminating exon is impaired (Baker, 1989).

Another possibility is that antisense transcripts from the B104 elements could base pair with the Sxl primary transcript and again disrupt splicing. These two suggestions would mean that the terminating exon cannot be included in the Sxl mRNA hence an active (male-lethal) <u>Sxl</u> polypeptide is always and produced. These mechanisms however do not readily explain the ability of these alleles to rescue mutations in positive In light of the finding that both the  $\underline{da}$ regulators. and <u>sis-b</u> products may be transcriptional activators (see below), an alternative explanation is needed. Perhaps the transposable elements provide alternative transcriptional start sites which allow for the production of primary transcripts devoid of the male-specific exon (Maine et al., 1985b). In this context it should be noted that the B104 element was isolated by virtue of its high level of expression in the early embryo, at the same time (2-5 hr) as zygotic <u>Sxl</u> expression begins (Sherer et al., 1982). Schupbach (1985) has suggested that <u>da</u> may be required to allow <u>Sxl</u> expression at this early stage in development when the there is little transcriptional activity in the rest of This suggests the possibility that these mutations genome. are male lethal not because of any disruption of splicing but because of the inappropriate expression of <u>Sxl</u> in the male embryo. In other words differential splicing is the mechanism <u>Sxl</u> uses to maintain the sexually determined state but sexspecific transcription may be the mechanism used to establish of preliminary evidence support this it. Two lines hybridisation experiments to 1) <u>in-situ</u> conjecture: blastoderm embryos with <u>Sxl</u> probes identify two populations embryo, one expressing high levels of <u>Sxl</u> and one not of (D.Coulter cited in Salz et al., 1989). 2) Maine al. et (1985a) report that two of the three embryo-specific Sxl transcripts are absent or severely reduced in the progeny of homozygous da mothers. Further experiments on the embryonic transcripts will be required to confirm this hypothesis. Particularly enlightening would be the identification of the 5' ends of these transcripts.

Sequence analysis of female-specific cDNA clones reveals a long open reading frame with the potential to produce a 39kd polypeptide. This protein would contain two domains which have sequence similarity to the RNA binding domains of several Ribonucleoproteins (RNP) from yeast, rat and man (see

Bell et al., 1988 for details). Many of the female-lethal loss function mutations are DNA rearrangements which disrupt of the coding sequence of this polypeptide. The finding that the Sxl gene product may be an RNA binding protein provides an attractive way to explain the autoregulatory function of <u>Sxl</u>. In this model a <u>Sxl</u> gene product can prevent the inclusion of translational terminating exon in the processed mRNA by the binding to the 3' acceptor site of this exon. When the Sxl polypeptide is absent the acceptor is not blocked and the exon included in the mRNA. Indirect evidence in support of this model for <u>Sxl</u> autoregulation has been obtained from studies on the tra gene and will be discussed below. (Bell et al., 1988; Sosnowski et al., 1989).

The molecular cloning of <u>Sxl</u> has given some working hypotheses as to how the gene functions. The presence of а male-specific translation terminating exon is consistent with the genetic evidence that no <u>Sxl</u> function is required in males. The presence of zygotic transcripts at the earliest development is consistent with clonal analysis stages of (Cline, 1984) and dosage compensation data (Gergen, 1987) which suggest that <u>Sxl</u> activity is required at or before the blastoderm stage. The presence of multiple female-specific transcripts is consistent with the different roles <u>Sxl</u> has to play in regulating sexual dimorphism in the fly. One major question remains unanswered; how is <u>Sxl</u> able to respond to the primary sex determining signal? Progress in this area is being made by isolating and characterising genes which regulate <u>Sxl</u>, some of which will be briefly considered below.

#### Regulators of Sex-lethal

Several loci have been reported which have effects suggesting an interaction with  $\underline{Sxl}$ . The most thoroughly characterised of these is <u>daughterless</u> (<u>da</u>).

<u>da</u>, which maps to salivary gland chromosome region 31E (2L), was first isolated as a recessive, temperature sensitive, maternal effect mutation resulting in the production of all male progeny (Bell, 1954). Cline (1978) demonstrated that this gene interacts with <u>Sxl</u> by isolating a <u>da</u> suppressor, which turned out to be the constitutive  $\underline{Sxl}^{M1}$ 

allele described above. Subsequent studies confirmed that da is required in XX zygotes to activate <u>Sxl</u> (Cline, 1979,1980). initial experiments were carried out using These the <u>da<sup>1</sup> allele. This allele is a zygotic</u> hypomorphic lethal in both sexes either when hemizygous or at 29<sup>0</sup>C when homozygous (Mange and Sandler, 1973; Cline, 1976). Several null alleles have now been isolated (Cronmiller and Cline, 1987; Caudy et al., 1988) and these show that <u>da</u> is required zygotically in for the formation of the peripheral both sexes nervous system. The phenotype of <u>da</u> nulls is similar to that observed with Achaete-scute (AS-C) mutations with regard to their effect on neural cell development. Caudy et al. (1988)suggest that the products of both genes may interact in neural development. The maternal effect of <u>da</u> appears to be strictly sex-specific.

The gene has been cloned (Cronmiller et al., 1988; Caudy 1988), and shown to produce two transcripts present et al., throughout the life cycle. These encode а conceptual polypeptide of 74kd, regions of which show striking sequence similarity with the myc oncogene, products of the AS-C (a startling finding in light of the similar complex phenotypes mentioned above) and more recently a 75% sequence identity with the DNA binding domain of Immunoglobulin Enhancer Binding protein. (Murre et al., 1989a). In addition the <u>da</u> polypeptide also contains a helix-loop-helix motif which has been implicated in protein dimerisation. This finding suggests a mechanism by which <u>da</u> may interact with Sx1. The da gene product, either alone or in conjunction with other polypeptides, could act as a transcriptional activator regulatory sequences for embryo-specific the the at Several lines of evidence transcripts discussed above. support this hypothesis: 1) temperature shift experiments with embryos from <u>da/da</u> mothers indicate that the maternal <u>da</u> required within the hours post first three is product oviposition (Cline, 1978). 2) <u>Sxl</u> activity is required in XX at approximately this time. 3) Sxl embryo-specific zygotes transcripts appear at approximately this time. 4) It has been reported that at least two of the embryo-specific transcripts are absent or severely reduced in embryos which lack <u>da</u> gene product (Maine et al., 1985a). An alternative explanation for consistent with these is also function, which <u>da</u> observations, is that it acts indirectly on <u>Sxl</u> by activating

another gene(s) which is required for <u>Sxl</u> expression.

Since the <u>da</u> gene product is maternal and appears to be in all eggs irrespective of their chromosomal active constitution it is unlikely to transmit information regarding ratio. It is more likely to be a the X:A factor without discriminatory ability required to permit the activation of Loci which assess and respond to the X:A ratio are Sxl. expected to be zygotic. Some are expected to be X-linked, so called numerator elements, and others autosomal denominator They are expected to behave in a genetically elements. predictable way. For numerator elements an increase in dosage should be detrimental to males as the apparent X:A ratio is increased. Decreasing the dosage should have the opposite effect and be detrimental to females. At least two loci have been identified which fulfil these and other genetic requirements for such factors: namely <u>sisterless-a</u> (<u>sis-a</u>) and <u>sisterless-b</u> (sis-b).

sis-a is an X-linked recessive female-specific lethal, it shows no maternal effect (Cline, 1986). Homozygous females can be rescued by  $\underline{Sxl}^{M1}$ . The lethal period is similar to that observed for <u>Sxl</u> nulls and clonal analysis indicates that <u>sis-a</u> activity is only required early in development. <u>sis-b</u> is also X-linked and has similar properties to sis-a (Cline, 1988; Torres and Sanchez, 1989). Several observations support the view that these genes are part of the X:A counting mechanism which activates <u>Sxl</u>. Increasing the dosage of either gene can be male-lethal and is feminising in triploids; females are rescued from reductions in Sxl or maternal <u>da</u> levels. Conversely reductions in dosage are masculinising in triploids and decrease the viability of females with reduced <u>Sxl</u> or <u>da</u> levels. The interaction of the <u>sis</u> genes with the <u>Sxl</u> alleles  $\underline{Sxl}^{f9}$  and  $\underline{Sxl}^{fLS}$ are especially informative. As previously discussed these alleles are believed to be defective in pathway initiation  $(Sxl^{\pm 9})$ and pathway maintenance  $(\underline{Sxl}^{fLS})$ , both <u>da</u> and <u>sis-a</u> show very similar phenotypes in combination with these Sxl alleles when tested for dominant synergistic effects; <u>sis-a</u> <u>Sxl<sup>f9</sup></u> (XX) double heterozygotes from <u>da</u>/+ mothers are not viable whereas <u>sis-a</u> <u>Sxl</u>fLS females from the same mothers are as viable as individuals wild type for <u>Sxl</u>. This seems to indicate that reductions in sis-a levels are only detrimental when coupled with <u>Sxl</u> alleles defective in pathway initiation. No effect

is detected when  $\underline{sis}-\underline{a}$  is coupled with  $\underline{Sxl}$  alleles which can initiate  $\underline{Sxl}$  expression normally but which are defective in maintaining expression. This result was previously shown for  $\underline{da}$  (Maine et al., 1986a).

The molecular nature of <u>sis-a</u> is not known but <u>sis-b</u> has recently been shown to be a product of the Achaete-scute complex, at least in part a product of the <u>T4</u> transcription unit of this complex (Torres and Sanchez, 1989). The product this locus appears to be a DNA binding protein with of the same DNA binding and helix-loop-helix motifs as the putative <u>da</u> polypeptide. Murre et al. (1989b) have shown that <u>da</u> and AS-C T3 gene products form heterodimers in-vitro. The AS-C T3 gene product appears to have a similar structure to the т4 These heterodimers can bind specifically product. DNA to fragments containing promoter elements, whereas the individual polypeptides cannot. This remarkable finding suggests that <u>da</u> and <u>AS-C</u> products interact by forming heterodimers required for transcriptional activation of both <u>Sxl</u> and genes involved in neurogenesis. This is compatible with the previous suggestion that da acts as a transcription factor directly at the putative <u>Sxl</u> embryo specific regulatory elements. An alternative explanation is that da AS-C is required to activate the expression of product transcription units, which are then required early in development to activate <u>Sxl</u> (<u>sis-b</u> function) and later in development for neurogenesis (<u>scute</u> function). It will be particularly informative to examine the effects of maternal da mutations on zygotic sis-b expression.

Whether the <u>sis-a</u> gene is a conventional protein coding gene remains to be determined. It is possible that it could be a binding site for a regulator which counts the number of X chromosomes by a titration effect. This suggestion echoes, respects, a model proposed by Chandra (1985)to in some for the assessment of the X:A ratio. He proposes а account maternally activated zygotic repressor which is titrated by binding sites on the X chromosome. While diplo-X individuals have enough sites to mop up the repressor and hence would allow <u>Sxl</u> expression, in Haplo-X individuals there would be insufficient number of sites and hence <u>Sxl</u> would be an repressed. The proposed repressor need not act on <u>Sxl</u> directly but instead on an X-linked positive regulator of <u>Sxl</u> such as the <u>sis-b</u> gene. Such a model is appealing in its

simplicity but is rather unsatisfactory when one considers that the level of repressor would have to be exquisitely balanced. Minor fluctuations either way would be lethal to one sex or the other. A similar model with a number of repressor-site interactions could perhaps buffer the system against changes in the level or activity of a single component. As to the existence of other numerator elements Cline suggests that more are present but that their number is limited. He proposes selecting for the dominant synergism discussed above to isolate such genes (Cline, 1988).

Several other genes which appear to interact with <u>Sxl</u> have been reported but are at present rather poorly characterised.

<u>Female sterile(1)</u><sup>1621</sup> also known as <u>liz</u> and <u>snf</u> (Steinmann-Zwicky, 1988; Oliver et al., 1988) is an X linked gene which has both maternal and zygotic functions required for <u>Sxl</u> activity. It is required in both the germ line and the soma. Some lethal effect is apparent which can be enhanced by reductions in <u>Sxl</u> or <u>da</u> levels and rescued by <u>Sxl<sup>M1</sup></u>.

<u>Daughterkiller</u> (<u>Dk</u>) is a dominant maternal effect female lethal, it can be rescued by  $\underline{Sxl}^{M1}$ . (Steinmann-Zwicky at al., cited by Lucchesi and Manning, 1987).

<u>female lethal(2)d</u> a temperature sensitive homozygous female lethal. It is reported that this allele can rescue males from the lethal effects of <u>Sxl<sup>M1</sup></u>. (San Juan and Sanchez, cited by Lucchesi and Manning, 1987). This suggests that it either acts downstream of <u>Sxl</u> or is an activation function which <u>Sxl<sup>M1</sup></u> is not constitutive for.

In summary genetic and molecular evidence suggest a model for the regulation of the sex determination gene <u>sex-lethal</u>. Zygotic functions (eg <u>sis-a</u>, <u>sis-b</u>) cell autonomously assess the X:A ratio early in development. When the X:A ratio is >1 these functions, in conjunction with maternal and zygotic functions (eg <u>da</u>, <u>liz</u>), activate <u>Sxl</u>. This initial activation may take the form of sex-specific transcription. Thereafter <u>Sxl</u>, by virtue of its autoregulatory capacity, is able to maintain the sexually differentiated state independent of the X:A ratio. This maintenance function probably operates at the

level of sex-specific RNA processing mediated by the Sxl gene product(s). The product(s) of the <u>Sx1</u> gene control dosage compensation, germline sex and somatic sex. This latter is probably regulated by the same pathway sex-specific splicing mechanism which maintains Sxl. Baker (1989) suggests that germline sex and dosage compensation may also be regulated by differential splicing. Functional studies of the different <u>Sxl</u> transcripts will be required to address this question.

## Somatic Sex Determination Hierarchy

discussed previously the genes which control somatic As sex have been well characterised at the genetic and molecular level. As with <u>Sxl</u> a detailed genetic analysis of <u>tra</u>, tra2, dsx and ix has provided a framework for subsequent molecular studies. Baker and Ridge (1980) carried out a clonal analysis of these genes which demonstrated cell autonomy of gene simultaneously activity. They also constructed flies homozygous for mutations in more than one of these genes. In way they were able to demonstrate a clear epistatic this relationship with dsx at the end of the pathway and tra and tra2 regulating dsx. Due to the similarity in phenotype the interaction between tra and tra2 could not be investigated. It was also unclear where <u>ix</u> acted in view of its similarity with dsx. Molecular cloning of three of these genes has helped to resolve these ambiguities and has provided strong support for Baker and Ridge's model. Each of the genes involved in the hierarchy will be discussed below.

# transformer(tra)

tra resides at 73A on the polytene chromosome map. The gene was first discovered by Sturtevant (1945) as a recessive mutation which transforms chromosomally female individuals into phenotypic males, there is no discernible effect on chromosomal males. In homozygous females all somatic tissue is phenotypically male but these individuals have a female

body size. addition these In individuals exhibit male courtship behaviour (McRobert and Tompkins, 1985). The gonads homozygous tra females develop as rudimentary testes of and King, 1960) however the status of the germ (Brown cells within these underdeveloped gonads is unclear. The appearance, at a low frequency, of oogenic and spermatogenic cells aerm is reported but in the majority of testes examined only a mass of degenerating tissue is found (Brown and King, 1960; Seidel, 1963; Steinmann-Zwicky et al., 1989). Pole cell transplantation experiments with tra/tra germ cells and wild type hosts indicate that tra is not required in the female germline for normal oogenesis (Marsh and Wieschaus, 1978; Schupbach, 1982). There is no apparent effect on dosage compensation as X-linked genes appear to be expressed at wild type levels (Brown and King, 1960). Thus it would appear that is required solely for female somatic differentiation. <u>tra</u> Somatic recombination experiments indicate that tra activity is cell autonomous and that it is required at least until pupariation in some tissues (Baker and Ridge, 1980).

The gene has been cloned (Butler et al., 1986; McKeown et al., 1987) and localised to a 3.5kb <u>EcoRI</u> fragment which is able to rescue <u>tra/tra</u> females in P-element transformation experiments. This fragment hybridises to four transcripts on northern blots, two of these correspond to unrelated genes which lie very close to the <u>tra</u> gene (the 3' gene actually overlaps with <u>tra</u>). The other two transcripts, which are expressed throughout development, are from the <u>tra</u> gene. Only the smaller of these two transcripts is female specific.

Sequence analysis of cDNA clones led to the surprising finding that <u>tra</u> was regulated by differential splicing (Boggs et al., 1987). Both tra transcripts use the same start site but differ by the use of alternative 3' splice acceptor sites within the first intron of the gene. The non sexsuch a spliced in way that is specific transcript translational terminating codons are included in all reading frames. The female specific transcript on the other hand uses acceptor approximately 200bp downstream of that used by an sex-specific transcript. The female-specific the non transcript encodes a long open reading frame capable of producing a 22kd polypeptide. This conceptual polypeptide shows no homology with known sequences but it is extremely basic suggesting that it could possibly bind to nucleic
acids. Sex-specific splicing of the wild-type gene has been used to monitor gene interactions within the sex determination hierarchy (Nagoshi et al., 1988). As expected only the activity of <u>Sxl</u> was required to produce the femalespecific transcript.

P-element transformation experiments with cDNA clones under the control of the heat shock promoter show that the non sex-specific transcript is without detectable function. Only the female-specific cDNA can rescue <u>tra/tra</u> females, moreover this cDNA alone is sufficient to transform XY individuals into somatic females. This surprising finding indicates that the tra gene is sufficient to direct all other genes downstream to female somatic development (McKeown et al., 1988). Expression of this construct in various mutant backgrounds has been used to verify the genetic model of somatic sex determination. In agreement with the model ectopic expression of tra is able to rescue sex-transforming <u>Sxl</u> mutations but not mutations in <u>tra2</u>, <u>dsx</u> or <u>ix</u>. The observation that <u>tra2</u> mutations are epistatic to tra indicates that tra2 does not act upstream of tra.

A more detailed examination of the splicing reaction has provided considerable insight into the regulation of <u>tra</u> (Sosnowski et al., 1989; Belote et al., 1989). When the whole intron is inserted into another gene, sex-specific splicing be detected suggesting that the specificity resides can entirely within the intron. Support for this comes also from the analysis of deletions within the intron when in its al., 1989). These (Sosnowski et normal environment experiments suggest that the non sex-specific splice site is the target for regulation such that in the presence of Sxl this splice site is only used 50% of the time. A comparison of the sequences at the regulated <u>tra</u> splice-site and the 3' splice site which proceeds the male-specific exon in the Sxl gene reveals a striking homology. The homology occurs at the polypyrimidine tract upstream of the 3' acceptor. A region of this type is believed to be necessary for successful splicing in all eukaryotic introns (Krainer and Maniatis, 1988). In both of the regulated introns this tract is composed almost exclusively of T residues. This is highly unusual and not а feature found in other introns. It has been suggested that а <u>Sxl</u> gene product binds at these splice sites to prevent their This would result, in the case of tra, the in use.

preferential use of the downstream female specific acceptor. A similar proposal has been put forward to explain the autoregulatory capacity of <u>Sxl</u> (Bell et al., 1988).

# Transformer 2 (tra2)

This gene maps cytologically to region 51B of the salivary gland chromosome map. Phenotypically it is similar to <u>tra</u>, homozygous females being transformed into somatic males (Watanabe, 1975). However in contrast to tra2 tra, homozygous males are sterile. This appears to reflect a requirement for tra2 in the male germline to allow the completion of spermatogenesis. The defect in spermatogenesis, a failure to elongate sperm heads, suggests that tra2 may be involved in X chromosome condensation (Belote and Baker, 1983). tra2 is not required in the female germline (Schupbach, 1982) or for dosage compensation (Fujihara et al., 1978).

The availability of hypomorphic and temperature sensitive tra2 alleles has been an invaluable aid in the investigation of somatic sex determination, particularly in the study of terminal differentiation genes. The temperature sensitive alleles tra2<sup>ts1</sup> and tra2<sup>ts2</sup> (Belote and Baker, 1982) have been used to estimate the temporal requirements for the action of the sex determination hierarchy in different somatic tissues. Temperature shift experiments indicate that is required from at least the mid 2nd larval instar tra2 through to the pupal stage to permit female differentiation. Different tissues however require tra2 activity at different times; for example female genital disk derivatives require the product throughout the entire period. However expression only required from just before to just after pupariation is prevent the formation of the male sex comb. The <u>tra2</u>ts1 to also been used to provide evidence for the allele has involvement of the sex determining hierarchy in regulating male courtship behaviour. Hemizygous <u>tra2</u>ts1 chromosomal females develop as female like intersexes at the permissive temperature. Under these conditions they do not appear to exhibit male courtship behaviour. If these individuals are shifted to the restrictive temperature post eclosion a

substantial proportion show male courtship behaviour after a delay of several days (Belote and Baker, 1987). The authors suggest that this behavioural change is indicative of a certain amount of plasticity in the regulation of sexual behaviour. It is not clear however if this plasticity is primarily a reflection of the intersexual nature of the females when they are raised at the permissive temperature.

The hypomorphic allele  $\underline{tra2}^{OFT}$  (Fujihara et al., 1978) is homozygous male fertile and homozygous females are partially transformed. Interestingly some of these individuals have apparently normal ovaries containing well developed eqqs. Hemizygous females have more testis like rudimentary gonads which are frequently associated with ovarian-like tissue. This allele could prove to be very useful in studying the proposed interaction between the germline and somatic component of the gonad uncovered by Steinmann-Zwicky et al. (1989).

tra2 has recently been cloned (Amrein et al., 1988; Goralski et al., 1989). Transcripts from the gene (it is not yet clear how many) are detected at all stages of development in both germline and soma. It appears that tra2 is not expressed sex-specifically or regulated at the RNA level, no difference having been detected between males and females respect to tra2 transcript levels. This is compatible with with the observation that expression of female-specific tra cDNA is sufficient to promote female somatic differentiation individuals. A conceptual <u>tra2</u> polypeptide of in XY approximately 250 amino acids has extensive similarity to RNA binding proteins found in hn-RNPs and sn-RNPs, both of these particles have been implicated in mRNA splicing. This finding suggests that tra2 may act directly to regulate the splicing of <u>dsx</u> mRNA. <u>tra</u> may act in conjunction with <u>tra2</u> in this role or tra may be required for the translation of the tra2 mRNA. These models are being actively investigated (Goralski et al., 1989; Burtis and Baker, 1989). The expression of tra2 in the female germline and male soma is somewhat puzzling given that no function for <u>tra2</u> has been detected in these tissues. The expression of <u>tra2</u> in the female germline is at very high levels, much higher than in the soma where the function is apparent. High levels of tra2 transcript are also detected in the 0-3hr embryo (Amrein et al., 1988; Goralski et al., 1989). Taken together these observations suggest that

there may be a maternal contribution of tra2 to the zygote. If this is the case any function for maternally derived tra2 must be redundant: recall that pole transcripts cell transplantation experiments indicate that <u>tra2</u> is not in the female germline. It is possible, then, required that expression in the female germline merely reflects tra2 the presence of a regulatory element which is needed for high expression in the male germline but which is also active in the female germline. This could also be an explanation for tra2 expression in the male soma.

#### <u>doublesex</u> (<u>dsx</u>)

First discovered by Hildreth (1965) as a recessive affecting both males and females dsx maps mutation to Both XX salivary gland chromosome region 84E. and XY individuals homozygous for null <u>dsx</u> alleles develop as morphologically identical intersexes apart from the sexual dimorphism in body size which is maintained. In contrast to triploid intersexes which are mosaics of male and female tissue, <u>dsx</u> intersexuallity is evident at the level of the cells are individual cell. Here it would appear that characteristics male and female expressing both simultaneously. This is particularly striking in the genital disc derivatives where both male and female structures are apparent in adults. The gonads of <u>dsx</u> individuals are, in most cases, irregular spherical masses resembling neither the male or female gonad, more than two gonads have never been observed. <u>dsx</u> is not required in the female germline (Schupbach, 1982).

Two alleles of the <u>dsx</u> gene have been particularly informative in the dissection of the role this gene plays in sexual differentiation.

 $\underline{dsx}^{D}$  (dominant): this allele has no effect on chromosomally male individuals but transforms females to intersexes when they carry a wild type copy of the gene. Hemizygous females or  $\underline{dsx}^{D}$ /+ females simultaneously homozygous for <u>tra</u> or <u>tra2</u> develop as somatic males.

 $dsx^{136}$ : when homozygous or heterozygous with a null allele chromosomal males are transformed to intersexes. There

is no effect on chromosomal females.  $\underline{dsx}^{D}$  is dominant to  $\underline{dsx}^{136}$ .  $\underline{dsx}^{D}/\underline{dsx}^{136}$  chromosomal males are normal whereas chromosomal females are intersexual. Finally, when examined in a homozygous <u>tra2</u> background,  $\underline{dsx}^{136}$  females are transformed to intersexes.

Baker and Ridge (1980) were able to reconcile these observations by proposing that <u>dsx</u> is a bifunctional gene one of two different states for the correct required in differentiation of sexual either males or females. Furthermore they proposed that the products of the tra, tra2 genes are required only for expression of ix and the dsx female state. Their model predicts that <u>dsx<sup>D</sup></u> represents constitutive expression of the male state.  $dsx^{136}$ , by their model, has lost the ability to express the male state but still requires the activity of tra, tra2 and ix for expression of the female state. They go on to suggest that dsx acts as a repressor of terminal differentiation functions <u>dsx<sup>M</sup> represses female differentiation genes</u> that such and  $\underline{dsx}^F$  represses male functions. In the absence of the  $\underline{dsx}$  gene both sets of terminal differentiation genes are expressed hence the observed intersexual phenotype. Surprisingly it would appear that simultaneous expression of  $\underline{\mathtt{dsx}}^M$  and  $\underline{\mathtt{dsx}}^F$ has the same phenotype (eg (XX)  $dsx^D/+$ ), suggesting that the products are mutually antagonistic. The model is consistent with dsx<sup>M</sup> being a default state of expression and with the tra, tra2 and ix gene products being required to override the default state to bring about female differentiation.

The molecular cloning of <u>dsx</u> sheds some light on the bifunctional nature of this gene (Baker and Wolfner, 1987; and Baker, 1989). It appears that the gene produces Burtis two major transcripts, one male-specific the other femalespecific. These result from the sex-specific splicing and alternative polyadenylation of a primary transcript. These transcripts initiate at the same start site. Other non sexspecific transcripts are also present, it is not yet clear if sex-specific functional role. The have any they are detected in 3rd instar larvae, pupae (where transcripts their expression is highest) and adults. Genetic evidence earlier for be present much predicts that they differentiation of the gonad during the embryonic stage (Hildreth, 1965) and the genital disk during the 2nd larval instar (Wieschaus and Nothiger, 1982). Whether failure to

detect the adult sex-specific transcripts during these earlier stages is due to a lack of sensitivity in the experiments or a requirement for other, as yet undetected, transcripts dsx early in development remains to be determined. Strong evidence that the sex-specific transcripts represent the functional <u>dsx</u> product comes from an examination four <u>dsx<sup>D</sup> alleles.</u> of They all contain disruptions of the female specific exon. Burtis and Baker (1989) propose that tra and tra2 regulate the sex-specific splicing of the <u>dsx</u> transcript, they suggest that regulation mediated at the 3' acceptor site of the female-specific is by the action of tra and tra products. In exon support of this they note that the sequence at this particular splice junction is a poor match to consensus Drosophila acceptor sequences. Further support comes from an examination of the same splice site in <u>D.virilis</u>. While the intron sequences upstream of the female specific exon are highly diverged the poor acceptor is well conserved. The model proposed by Burtis and Baker is a splice site activation mechanism where tra and tra2 act at this splice site to allow the use of the otherwise poor acceptor. This is in contrast to the splice blockage model proposed for <u>Sxl</u> and <u>tra</u> site genes. Α molecular analysis of the sex-specific splicing shows that <u>Sxl</u>, <u>tra</u> and <u>tra2</u> but not <u>ix</u> are required for the production the female-specific transcript (Nagoshi et al., 1988). of This is consistent with the genetic model and the idea of a male default state.

Although these data are encouraging other experiments been less successful. P-element transformation studies have been carried out using sex-specific clones CDNA have controlled by an actin promoter, these constructs show very little effect on homozygous dsx individuals. Similarly there are only minor effects on wild type individuals carrying а cDNA from the opposite sex. In both of these experiments it is mainly tergite pigmentation which is noticeably affected. There is however some elevation of yolk protein mRNA in males carrying dsx<sup>F</sup>. It is not yet clear whether or not the weak associated with these constructs is due to phenotype inappropriate expression from the actin promoter or whether reflects the lack of another essential dsx function. it Unfortunately the sequence of the <u>dsx</u> polypeptides has been less informative than one would have hoped. Male dsx encodes

a putative 57kd polypeptide and the female transcript one of 45kd. These polypeptides differ at their carboxy termini. No significant homology between the dsx polypeptides and known protein sequences has been detected. The polypeptides do, contain strings of repeated amino acids found however, in <u>Drosophila</u> regulatory proteins, although it other is not known if these regions are significant (see Burtis and Baker, for a full discussion). Clearly further studies of 1989 the dsx gene and the targets which it regulates are required before we know how the gene is acting.

#### <u>intersex</u> (<u>ix</u>)

Comparatively little is known about <u>ix</u> compared with other genes in the somatic sex determination hierarchy. Homozygous ix females are transformed to intersexes but there is no effect on males (Morgan et al., 1943; Baker and Belote, 1983). There is no information regarding the autonomy of <u>ix</u> but it seems likely, by analogy with the other sex determining genes, to act cell autonomously. In the double analyses of Baker and Ridge (1980) dsx, tra and tra2 mutant were all epistatic to <u>ix</u>. This would suggest that it is at the beginning of the pathway. The analysis was hampered, however, by the similarity of <u>dsx</u> and <u>ix</u> phenotypes. Molecular analysis of <u>tra</u> and <u>dsx</u> sex-specific splicing show that is not required for the production of female-specific transcripts from either of these genes (Nagoshi et al., 1988). This would place <u>ix</u> above <u>dsx</u> in the hierarchy. These observations are compatible if <u>ix</u> is required to act in conjunction with the  $dsx^{F}$  product(s) for its activity. Thus and tra2 would appear to be epistatic because they fail tra to produce <u>dsx<sup>F</sup></u> (Baker and Ridge, 1980, Baker et al., 1987). A molecular analysis of this gene may help to clarify its role in sexual differentiation. One attractive idea is that the <u>ix</u> product provides an opposite function to that encoded in the carboxy terminal of the  $\underline{dsx}^{M}$  polypeptide (B.Baker pers.com).

# Regulation Of Germline Sex

As we have seen correct activity of <u>Sxl</u> is required for germline sex determination. Several other genes affecting the development of the germline have been identified (eg. <u>agametic</u>, <u>tudor</u>, <u>grandchildless</u>) however these genes are not sex-specific, rather they disrupt pole cell development in both sexes. In addition most of these genes have effects on somatic development.

One gene, ovo (Busson et al., 1983), has been isolated required only for female germline development. which is Mutant males are not affected and develop as normal and individuals. Ovo, which maps to region 4E1 on the fertile Х chromosome, was initially identified by dominant alleles, the most severe of which, Ovo<sup>D1</sup>, results in female sterility. The of  $\underline{Ovo}^{D1}/+$  females are atrophied and ovaries almost completely devoid of germ cells. Clonal analysis indicates that <u>ovo</u> may be required continuously in the female germline (Perrimon, 1984). Loss of function alleles have been isolated and used to determine ovo requirement during development (Oliver et al., 1987). Here pole cell death is observed from late blastoderm onwards. This finding suggests that germ cells determine their sex at around the same time as somatic this is the case it is before they come into tissue. If to qonadal soma. <u>Sxl</u> has been shown be contact with the in the soma at this time but it is not yet known if active the germline this early. The germline it is active in mutations in <u>Sxl</u> and <u>ovo</u> are radically of phenotype different. Sxl mutations result in undifferentiated growth of germ cells whereas ovo mutations cause germ cell death. It is not clear as yet if, or how, ovo and Sxl interact but as these genes show a different phenotype an mutations in epistatic analysis may be possible. In this respect the allele <u>Sxl</u><sup>fs1</sup> may prove to be of some use as it allows normal somatic development. It would also be informative to female effect of <u>ovo</u> mutations on the pattern of Sxl assess the is important to establish any transcription. Clearly it germline sex understand are to if we interaction The region containing the ovo qene has determination. recently been cloned (Mevel-Ninio et al., 1989) and is under investigation, it would be of some interest to investigate ectopic expression of the gene in males, perhaps by P the

element transformation experiments.

recent study of The Steinmann-Zwicky et al. (1989) indicates that control of germline sex is not entirely cell autonomous: an interaction between germline and gonadal soma apparent. In these studies pole cells were transplanted is into agametic host embryos and their development assessed at the adult stage. In ovaries germ cells develop according to their X:A ratio; XX germ cells undergo oogenesis with fertile eggs being produced. XY germ cells initiate spermatogenesis, however they do not progress beyond the spermatocyte stage. In testes both XY and XX germ cells enter spermatogenesis, XY cells produce functional sperm while XX cells degenerate after the spermatocyte stage. These observations led the authors to conclude that germ cells have an inherent ability determine their sex but that compatible soma is required to to support full development. However in the case of the XX germ cells an inductive interaction with the gonadal soma in the determination of germline sex also seems to be apparent. This can be interpreted as either a signal from ovarian necessary to initiate the female pathway of tissue development or a testicular signal capable of imposing spermatogenic development irrespective of chromosomal sex. against the latter hypothesis; XY cells enter Arguing the the sex of soma, spermatogenesis irrespective of а signal from the male gonad imposing spermatogenesis somatic would appear to be redundant.

Steinmann-Zwicky et al. also studied pole cells carrying Sxl mutations, again transplanted into agametic various for <u>Sxl<sup>f1</sup></u> XX pole cells homozygous enter the hosts. spermatogenic pathway in both ovaries This and testes. observation supports the previous studies of Schupbach (1985) requirement in the female germline. cells XX pole Sxl on carrying a copy of the semi-constitutive  $\underline{Sxl}^{M1}$ allele oogenesis irrespective of somatic sex. Thus this initiate allele is able to over-ride the somatic induction previously suggest that the somatic This would seem to observed. induction involves an interaction between the soma and <u>Sxl</u> or one of its positive regulators. Female soma may be required to activate and/or maintain <u>Sxl</u> expression or male soma may repress <u>Sxl</u>. Again the latter explanation would appear less likely; if <u>Sxl</u> is never expressed in the male germline why have a mechanism to repress it? The  $\underline{Sxl}^{M1}$  allele has no

effect on haplo-X germ cells. This supports previous suggestions that this allele is not constitutive for all functions in situations where the X:A ratio is inappropriate for <u>Sxl</u> activation (Cline, 1988; Steinmann-Zwicky, 1989).

Identification of other genes involved in germline sex determination is clearly a high priority if we are to unravel the regulation of this process. Two routes are available at present, one is to examine sex-specific sterile mutations in the hope of identifying those in which germ cell development arrested at an early stage. This is how the ovo gene was is Such an undertaking is however fairly daunting identified. given the large number of sterile mutants so far isolated (for example over 600 genes are estimated to be mutable to male sterility, Lindsley and Lifschyts 1972). An alternative approach may be to isolate genes expressed sex-specifically early in development either by +/- screening or by using subtracted cDNA probes. Single sex embryo populations can be obtained easily using fly stocks which contain the Responder element of <u>Segregation Distorter</u> translocated on to the sex (Lyttle, 1989; Walker et al., 1989). These chromosomes strains allow the production of embryo populations which are greater than 95% single sex. By examining the expression of, and making mutants in, genes isolated by the above method one would hope to identify candidates for genes involved in germline sex determination.

#### Sex-specific Gene Expression In The Soma

Several genes have been isolated which show sexually dimorphic expression in somatic tissues. Two basic types of regulation are apparent; direct regulation where the somatic sex determining hierarchy is required continuously to ensure the appropriate expression of a gene, and indirect regulation where the hierarchy specifies the formation of a sexually dimorphic tissue which in turn determines the expression of a gene.

#### Direct Regulation

The yolk protein (YP's) genes are the only known examples

genes subject to this form of regulation. These three Хof genes are expressed in the female fat body and linked the follicle cells associated with vitellogenic oocytes. It is only in the fat body that direct regulation is apparent, expression in the follicle cells is by virtue of tissue specificity. YP1 and YP2 are closely linked separated by around 1.2kb. They are transcribed divergently. YP3 is located about 1000kb away from the other two genes (Shepherd et al., 1985)

Several investigators have studied the expression of studies these genes. Of particular relevance are on the effects of mutations in the sex determination hierarchy (Bownes and Nothiger, 1981; Ota et al., 1981; Brennan et al., 1982). The results of these investigations may be summarised by stating that YP expression in the female fat body requires the expression of the female specific dsx product. Belote et (1985) conducted temperature shift experiments with the al. tra2<sup>ts1</sup> allele to further demonstrate that tra2, and by inference dsx, is required continuously for YP expression in for tra2<sup>ts1</sup> the adult. If an XX individual homozygous is raised at the restrictive temperature it develops as а somatic male and does not synthesis YP's. If it is then shifted down to the permissive temperature YP synthesis is initiated. Similarly YP synthesis declines dramatically if а similar individual is raised at the permissive temperature and then shifted up to the restrictive temperature. in contrast YP synthesis in the follicle cells of the ovary is not directly affected by these temperature shifts (M. Bownes pers. comm.). Molecular cloning of the YP1 gene and P-element transformation experiments with upstream sequences have identified a 125bp enhancer-like element which is sufficient to direct fat body specific and sex-specific expression to a heterologous gene (Garabedian et al., 1986). An additional element which confers follicle cell specific expression has also been identified (Shepherd et al., 1985). At present it is unclear how the enhancer element is regulated by the sex determining hierarchy. The dsx product may act directly at element, or indirectly by regulating the expression this of The some other factor required for YP expression. direct regulation of these genes is probably a special feature in the control of sex-specific gene expression as the fat body is not a sex-specific tissue. The majority of sex-specific

genes are expressed in a tissue specific manner. Nevertheless the YP genes offer a unique opportunity to investigate directly the molecular mechanisms by which the sex determination hierarchy operates.

# Indirect Regulation

All of the genes so far isolated which are indirectly regulated by the somatic sex determination hierarchy are expressed in the reproductive apparatus. In the female the chorion genes are expressed in the somatic follicle cells of the ovary. Expression is dependent on the presence of an ovary and a functional germline (DiBenedetto et al., 1987). males, genes expressed specifically in the accessory In glands and the ejaculatory duct have been identified. These include esterase-6 (Oakshott et al., 1987) and glucose dehydrogenase (Cavener et al., 1985), pheremone peptides 70A (Chen et al., 1988) and mst355 (Monsma and Wolfner, 1988) and several accessory gland transcripts of unknown function: mst-1, 2, 3, 35 (Schafer, 1986a) and mst316 (DiBenedetto et al., 1987). Expression of these accessory gland transcripts has been examined in chromosomal females carrying mutations in sex determining genes. As expected their expression various dependent upon the presence of an accessory qland. was mst316 Chapman and Wolfner (1988) examined the expression of mst355 in <u>tra2</u><sup>ts1</sup> temperature shift experiments. and They irreversibly found that the expression of these genes was determined at approximately the same time as accessory gland formation is irreversibly determined. By further examination these genes it should be possible to identify the of regulatory elements which confer tissue specificity. The key questions will then be what factors act at these elements and how are these factors regulated.

interesting example of sex-specific gene expression One glucose dehydrogenase gene (<u>gld</u>). is the This qene is in both sexes during the pupal stage where it expressed appears to be required for normal eclosion. However in adults expression is restricted to the male ejaculatory duct where gene product is believed to have a role in reproductive the behaviour. It would be of some interest to examine the regulatory elements of this gene to try and separate the different modes of expression.

Before leaving sex-specific genes in the soma it is worth noting that many genes may not be regulated by simple all or nothing expression. For example many genes may show sexual dimorphism merely in terms of the level or the site of expression (eg pigment distribution on the adult cuticle). The study of these genes may permit a more subtle understanding of how gene expression is regulated. However methods to isolate this class of genes are at present limited.

All of the somatic sex-specific genes so far isolated are during the adult stage. expressed However phenotypic differences between the sexes are apparent at earlier stages, most notably in the gonad. Identification of genes expressed specifically in the soma of the developing qonad, particularly those genes expressed early in development, would be most useful in attempting to elucidate regulatory mechanisms (see below).

#### Germline-Specific Transcripts

Rather more progress has been made in the study the of regulatory sequences controlling genes expressed in the germline. however we are hampered because Here our understanding of germline sex determination is rather poor. Of the genes that have been studied mst(3)gl9 and B2-tubulin have been the subject of particularly extensive examination. Both genes are only expressed in the male germline. Kuhn et al. (1988) have shown that the 102bp upstream of the mst(3)gl9 start site is sufficient to confer spermatocytespecific expression on a heterologous gene. In addition they demonstrate that 110nt of 5' untranslated sequence and 10ntof coding sequence are sufficient to allow appropriate germline translational control of a <u>lacZ</u> fusion construct. The B2-tubulin gene has been dissected by Michiels et al. (1989), they find that the 52bp upstream of the start site is sufficient to direct testis specific expression. Within this region they identified a 14bp element (BUE1) which is in sequence and position in the homologous gene conserved If this sequence is mutated expression from <u>D.hydei</u>. is However in contrast to the element found at the abolished.

YP1 gene, BUE1 does not act as an enhancer but as a conventional promoter element. The BUE1 sequence is not found upstream of the mst(3)g19 gene. Whether this reflects a requirement for these genes at different times in spermatogenesis or that a variety of sequences confer sperm specific transcription remains to be determined.

The isolation and characterisation of more genes showing sexually dimorphic expression, both in the soma and the germline, will be important in our attempts to define regulatory elements. The identification of regulatory factors acting at these elements is dependent upon the ease with which reliable nuclear extracts can be isolated for DNA binding studies. In this respect sex-specific transcripts expressed early in development will be particularly useful as nuclear extracts can be prepared with ease from the In chapter 3 I will outline the Drosophila embryo. experimental approach I have chosen to try and identify genes in this class.

# CHAPTER 2

 $\phi_{i}$  ,  $\phi_{i$ 

MATERIALS AND METHODS

Solid and line line and the solution of the so

#### 2.1 MATERIALS

The reagents used during the course of this work were supplied by the following manufacturers.

a)<u>Enzymes</u>: Restriction endonucleases, T4 Polynucleotide kinase, T4 DNA ligase, <u>Ecoli</u> DNA polymeraseI, <u>E.coli</u> RNaseH, (BRL). Klenow fragment <u>E.coli</u> DNA polymeraseI, calf intestinal phosphatase, (Boeringer Mannheim). AMV reverse transcriptase, (Pharmacia). T4 DNA polymerase, (Anglian Biotechnology). <u>EcoRI</u> methylase, (New England Biolabs).

**b**)<u>Specific reagents</u>: Oligo(dT)-cellulose type 7, oligo(dT)<sub>12-18</sub>, dNTP's, RNAGUARD [human placental ribonuclease inhibitor],  $pd(N_6)$ , Sephadex G50, (Pharmacia). Biotin-16-dUTP, (Boeringer Mannheim). Streptavidin-biotinylated horseradish peroxidase complex, Hybond-N membrane, Lambda invitro packaging kit (Amersham). <u>EcoRI</u> linkers, (BRL). Ultragel AcA34, (LKB).

c)Others: All other chemicals and reagents were purchased from generally available sources with no preference for any one manufacturer. Wherever possible AnAlar or Biochemical grade chemicals were used.

<u>Buffers and Media</u>: Most common buffers and media recipes were obtained from the Cold Spring Harbour Molecular Cloning manual (Maniatis et al., 1982). Buffers for restriction and modifying enzymes were prepared according to the manufacturers recommendations.

#### 2.2 METHODS

#### 2.2i Nucleic Acid Isolation:

a)<u>Plasmids</u>: Large scale plasmid isolation was carried out by the alkaline lysis method (Birnboim and Doly, 1979) as described in Maniatis et al.(1982) or on a small scale by the boiling method (Holmes and Quigley, 1981).

**b)**Bacteriophage lambda: Large scale isolation of lambda DNA was performed by a modification to the protocol of R.Buckland Heidelberg). A high titre stock was prepared from a (EMBL plate lysate using a fresh single plaque as starting This stock was used to infect appropriate material. host (OD<sub>660</sub> approx. 0.3) in a 21 flask containing 200ml cells of 10mM MgSO4. The culture was incubated at  $37^{\circ}C$  with L-broth, shaking until lysis was apparent (4-5hrs.), 1ml vigorous of CHCl<sub>3</sub> and 8g of NaCl was added and shaking continued for а further 10-15min. The debris was pelleted (15,000g, 15min) and 20g of PEG 8000 was added to the recovered supernatant. PEG was dissolved at room temperature and the The samples left at 4<sup>O</sup>C overnight to precipitate the phage. The were was pelleted (15,000g, 15min) all precipitate of the supernatant carefully removed and the pellets resuspended in 11ml of phage buffer. The phage suspension was vortexed with an equal volume of CHCl<sub>3</sub>, the phases separated (27,000g, 10min) and 10ml of the aqueous phase added to 7.2g of CsCl. Once the CsCl had dissolved the samples were decanted into quickseal tubes and spun for 12hr at 49,000rpm in a Beckman Ti70 rotor. The phage band was removed from the side with an 18G syringe needle and dialysed against TE (3 changes 11 each). DNA was purified by sequential extractions with phenol, 1:1 phenol/CHCl<sub>3</sub> and CHCl<sub>3</sub>. The DNA was then dialysed against TE as above and stored at 4<sup>O</sup>C. Yields for EMBL3 and NM1149 recombinants were generally 100-200ug.

c)Lambda minipreps: Small scale isolation of lambda DNA for the analysis of recombinants was performed by a plate lysate method. (Maniatis et al., 1982).

d)Drosophila DNA: <u>Drosophila</u> high molecular weight modifications to the chromosomal DNA was prepared by procedure of Kaiser and Murray (1985). One gram of flies was in liquid N<sub>2</sub> with a precooled mortar and pestle and ground homogenised with 10ml of freshly prepared ice cold buffer (10mM Tris-HCl pH7.6, 60mM NaCl, 10mM EDTA, 0.15mM spermine, 0.15mM spermidine, 0.5%(v/v) Triton X-100) in a Wheaton 15ml 30ml homogeniser (A pestle). The solution was decanted into a corex tube on ice and the nuclei pelleted (6,000g, 10min) at 4<sup>0</sup>C. The supernatant was carefully removed and the nuclei resuspended in 10ml of homogenisation buffer by gentle pipetting. The nuclei were pelleted and resuspended a further times and then finally transferred to a 50ml Falcon tube 3 where they were lysed by the addition of 1ml of 20% (w/v)sarcosine followed by gentle inversion of the tube. RNaseA was added to a final concentration of 100ug/ml and the lysate incubated at 37°C for 30-60min. Proteinase K was added to a final concentration of 100ug/ml and incubation continued for 2-12hr. The lysate was extracted, once with phenol, three times with 1:1 phenol/CHCl<sub>3</sub>, twice with CHCl<sub>3</sub> and the aqueous phase was concentrated with iso-butanol to a volume of 1-2ml. solution was dialysed for 24-48hr against several This changes of TE and stored at  $4^{\circ}$ C. Yields of DNA were generally 200ug/g of starting material.

e)Drosophila RNA: RNA for Northern blot analysis was isolated by a modification of the method of Chomzynski and Sacchi (1987) that is described in the laboratory manual of K.Stanley (EMBL Heidelberg), followed by batch purification of poly(A)<sup>+</sup> mRNA using oligo(dT)-cellulose. One gram of in a tissue was homogenised at full speed Silverstone homogeniser for 2-3min in 10ml of denaturing solution (4M guanidinium isothiocyanate, 0.1M Tris-HCl pH8, в-0.1M mercaptoethanol, 0.1%(v/v) antifoam-A). To the homogenate the following were added sequentially, 1ml 2M NaOAc pH4, 10ml phenol and 2ml CHCl3. The mixture was vortexed, left to stand ice for 15-30min and centrifuged (12,000g, 10min). on The phase was removed to a fresh tube, 1vol of aqueous isopropanol added and the mixture left at -20<sup>O</sup>C for 1hr to allow the RNA to precipitate. The precipitate was pelleted 5ml of denaturing 10min) and resuspended in (12,000g, solution. The RNA was again precipitated as described above,

washed in 70% EtOH, and finally resuspended in 1ml of oligo(dT)-cellulose binding buffer (0.5M NaCl, 10mM Tris-HCl pH 7.5, 1mM EDTA, 0.1% (w/v) sarcosine). Oligo(dT)-cellulose (50mg/g tissue) was hydrated overnight at  $4^{\circ}$ C in H<sub>2</sub>O, treated for 15 min at room temperature with 0.1M NaOH and washed (5x 10ml sterile H<sub>2</sub>O, 5x 10ml binding buffer). The RNA was heated 68°C for 5min, chilled on ice, thoroughly mixed with to the oligo(dT)-cellulose in a 15ml Falcon tube and incubated at room temperature for 30min with gentle agitation. The oligo(dT)-cellulose was pelleted by centrifugation (1500g bench-top centrifuge) and washed 5 times using 10ml of binding buffer for each wash. The poly(A)<sup>+</sup> mRNA was eluted by resuspending the resin in 0.5ml of sterile H<sub>2</sub>O and incubating at 58°C for 15-30min with occasional shaking. The oligo(dT)cellulose was pelleted (3000g bench-top centrifuge) and the supernatant transferred to a 1.5ml microfuge tube. RNA The was precipitated by the addition of 50ul 3M NaOAc pH5, 1ml at -20<sup>O</sup>C for 1-12hr and collected EtOH, stored by centrifugation in a microfuge for 30min. The RNA was resuspended in 100ul of sterile  $H_2O$  and stored at -70<sup>O</sup>C. Yields were generally 50ug of RNA/g of tissue.

#### 2.2ii: In-vitro DNA manipulation

Restriction endonuclease digestion of DNA, subcloning of DNA into plasmid vectors and introduction of recombinant plasmids into <u>E.coli</u> were performed using minor modifications of standard procedures (Maniatis et al., 1982).

#### 2.2iii:Electrophoresis and blotting

Electrophoresis of DNA was carried out in TBE agarose gels. Fragment sizes were calculated according to the method of Helling et al. (1974) using restriction digested  $\lambda$ cI857 s7 DNA as a standard (Sanger et al., 1982). For hybridisation gels were blotted to Hybond-N membranes using recommended procedures (Maniatis et al., 1982). Hybridisations were carried out either in 6X SSC, 1% SDS, 10X Denhardts, 100ug/ml denatured herring testis DNA, 15ug/ml polyA, 0.005% NaPPi at 68°C or in 50% formamide, 5X SSPE, 1% SDS, 10X Denhardts,

100ug/ml denatured herring testis DNA, 15ug/ml polyA, 0.005% NaPPi at 42°C. Blots were washed in 2X SSC, 1% SDS, 0.1% 68<sup>0</sup>C where low stringency was required NaPPi at or with additional washes in 0.1X SSC, 1% SDS, 0.1% NaPPi at 68<sup>O</sup>C for stringency. Blots were exposed to either Kodak Xomat-S hiqh film at -70<sup>0</sup>C with intensifying or XAR-5 screens. RNA carried out by electrophoresis analysis was in agarose formaldehyde gels as described by Maniatis et al. (1982)conditions followed by transfer to Hybond-N using the recommended by the manufacturer. Blots were hybridised in the 50% formamide solution and washed as described above for DNA high stringency. Transcript sizes were estimated by at comparison with an RNA size ladder (BRL).

# 2.2iv:Radiolabeled nucleic acid

For blotting experiments <sup>32</sup>P labelled DNA was prepared either by nick translation of plasmid and phage DNA (Rigby et al., 1977) or by random priming of gel purified restriction fragments (Feinberg and Vogelstein, 1983). All probes were purified by Sephadex G50 chromatography in columns prepared 105from disposable 1ml syringes and were generally used at 10<sup>6</sup>cpm/ml of hybridisation solution. First strand CDNA was labelled to high specific activity in the following way: 100of a<sup>32</sup>PdCTP (800Ci/mMol) was dried down 150uCi in а siliconised microfuge tube, to this the following was added order, 4ul 5X 1st strand buffer (250mM Tris-HCl pH8.3, in NaPPi, 50mM MgCl<sub>2</sub>, 50mM DTT), 1ul 80mM 30u 700mM KCl, RNAGUARD, 2ul dNTP mix (10mM dAGT), 2ug oligo(dT)<sub>12-18</sub>, 1ug poly(A)<sup>+</sup> mRNA, 20u AMV reverse transcriptase and sterile  $H_20$ to a final volume of 20ul . The reaction was incubated at 42<sup>0</sup>C for 30min. 1ul of Then 10mM dCTP was added and incubation continued for a further 30min. The RNA was hydrolysed by the addition of 1 volume of 0.6M NaOH, 20mM EDTA and incubation at 65<sup>O</sup>C for 30min. Radiolabelled cDNA was purified on Sephadex G50 columns and incorporation estimated by Cherenkov counting a small aliquot (Maniatis et al, 1982). Specific activities of >1x10<sup>8</sup> cpm/ug mRNA were obtained for best probes, lower incorporation appeared be the to associated with impure mRNA template.

Single sex 3rd instar cDNA libraries were constructed as follows:

a) RNA preparation. RNA was prepared by modifications to the procedure of Chirgwin et al. (1979). Hand sexed 3rd late instar larvae were homogenised, 30 at a time, in 1ml 4M of guanidinium isothiocyanate, 50mM Tris-HCl pH7.6, 10mM EDTA, 2% (v/v) sarcosine, 140mM B-mercaptoethanol, 25mM NaCitrate, 0.1% Antifoam-A. 30ml of pooled homogenate was layered onto a 15ml cushion of 5.7M CsCl, 0.1M EDTA in a Beckman polyallomer SW28 centrifuge tube which was spun at 25,000rpm for 40hr. After centrifugation the supernatant and cushion were removed by aspiration and the pellet resuspended in 3ml of the above denaturing solution. This solution was in turn layered onto a 1.5ml CsCl cushion in a polyallomer SW50 tube and spun for 20hr at 35,000rpm. The supernatant and cushion were removed with care, the pellets allowed to drain and the RNA resuspended in 1ml of sterile H<sub>2</sub>O. Poly(A)<sup>+</sup>mRNA was isolated chromatography on three sequential oligo(dT)-cellulose by columns (Aviv and Leder, 1972) following the manufacturers instructions except that sarcosine replaced SDS in all buffers (sarcosine is less susceptible to precipitation in high salt than SDS) and the  $poly(A)^+$  was eluted with sterile water.

b)First and second strand cDNA synthesis: This was carried out by a modification to the procedure of Gubler and Hoffman (1983) as described in the Amersham cDNA synthesis kit but manufacturers. reagents from different Reaction using products were sized on 1.5% alkaline agarose gels (Maniatis 1982) and yields estimated determining al., by et radioactivity incorporated into acid precipitable material The cDNA was extracted twice with (Berger, 1987). а 1:1 of phenol/CHCl<sub>3</sub> (in this and all subsequent mixture extractions organic phases were combined and reextracted with TE), and the combined aqueous phases were extracted 3 times with ether. The cDNA was precipitated (in siliconised microfuge tubes) by the addition of 1vol 4M  $NH_AOAc$  and 2vol incubated on ice for 30min and pelleted (12,000g, EtOH, 30min). The pellet was resuspended in 50ul of  $H_2O_1$ , subjected

to a second precipitation as described above, rinsed with 70% EtOH, air dried briefly and resuspended in 10ul of  $H_2O$ . То the cDNA 4ul 5X R1 methylase buffer (0.5M NaCl, 0.5M Tris-HCl 5mM EDTA), 2ul 1.3mM S-Adenosyl Methionine, 20u EcoRI рН8, 37<sup>0</sup>C methylase was added and the reaction incubated at for 60min. The cDNA was purified by two phenol/CHCl<sub>3</sub> and two ether extractions, precipitated with 0.1vol 3M NaOAc pH5, 2vol EtOH and resuspended in 10ul H<sub>2</sub>O.

c)Addition of Linkers and Ligation To Vector: EcoRI linkers were kinased and added to the cDNA as described by Huynh et al, (1985). It is imperative that the free linker molecules are all removed prior to ligation with the vector. This was achieved in the following way: the linkerd cDNA was extracted, once with phenol/CHCl<sub>3</sub> and once with ether, precipitated with 0.1vol 3M NaOAc pH5, 0.6vol Isopropanol at room temperature for 30min and recovered by centrifugation (30min, 12,000g). This is a selective precipitation in which small molecules are left in the supernatant. The pellet was resuspended in 20ul of STE (0.4M NaCl, 10mM Tris-HCl pH8, 1mM and applied to a column of Ultragel AcA34. The EDTA) matrix had previously been extensively washed with STE and poured into a siliconised disposable 1ml pipette. The column was developed with STE and 50ul fractions were collected. The latter were assayed for radioactivity and the peak fractions containing cDNA were pooled and precipitated with 2vol EtOH. The cDNA was resuspended in 20ul H<sub>2</sub>O and the concentration determined by spotting 1ul onto an ethidum bromide plate along with standards. The cDNA was ligated into the  $\lambda$ insertion vector, NM1149 (Murray, 1983) using a vector:cDNA molar ratio of 2:1. In practice 10ug of EcoR1 cut NM1149 was precipitated with 100ng of cDNA, resuspended in 7ul 10mM MgCl<sub>2</sub> and incubated at 42<sup>o</sup>C for 15min to allow the cohesive ends of lambda to anneal. To this tube 1ul 10X ligase buffer (600mM Tris-HCl pH7.5, 100mM MgCl<sub>2</sub>, 100mM B-mercaptoethanol, 10mM EDTA), 1ul 10mM ATP, 1u T4 DNA ligase was added and the reaction was incubated overnight at 15<sup>0</sup>C. The ligation mix packaged into infectious phage particles using а was commercial kit and the resultant phage suspension titered on L87 and NM514(hfl) E.coli strains.

Plaque hybridisations were carried out by a modification to the method of Benton and Davis (1977). Five to ten thousand recombinant phage were plated per 10cm x 10cm L-agar plate with  $10^8$  host cells and 0.6% top L-agarose, 10mM MgSO<sub>4</sub>. The plates were incubated at 37°C for 6-8hr (or until the plaques just visible) and cooled at 4<sup>0</sup>C for at least were 1hr. Hybond-N filter replicas were taken from plates (up to 6 filters/plate), denatured (0.5M NaOH, 1.5M NaCl) for 3-5min, neutralised (0.5M Tris-HCl pH7.5, 1.5M NaCl) for 5min and washed in 2X SSC for 15min. The filters were air dried and then baked at 80°C for 1hr. They were then immersed in 18 Triton X-100, blotted briefly on 3MM paper and soaked in 2X SSC for 15min. For optimal results filters were prehybridised individual plastic bags for 2-12hr in 10ml of aqueous in hybridisation solution (see section on DNA blotting above) at 68<sup>0</sup>C in a shaking water bath. The prehybridisation solution removed and the probe added with 3ml of fresh was hybridisation solution. Levels of background were found to be greatly reduced when hybridisation solution and probes were filtered through 0.22um filters. The filters were incubated at 68<sup>0</sup>C 8hr (recombinant probes) or 48hr (cDNA probes) for with shaking. After hybridisation the filters were treated as described above for DNA blots.

#### 2.2Vii: In-Situ Hybridisation to Polytene Chromosomes

Isolation and treatment of <u>Drosophila</u> salivary glands was performed as described by Pardue (1986). Probes were EMBL3 or NM1149 recombinants labelled with Bio-16-UTP by nick translation (Rigby et al., 1977). Hybrids were detected with Streptavidin-Biotinylated Horseradish Peroxidase complex and diaminobenzidine/H<sub>2</sub>O<sub>2</sub> as described by Pardue (opp.cit). The slides were stained with Giemsa and the coverslips were mounted with DPX. Chromosomes were photographed using a Leitz Vario-orthomat camera system and Kodak Ectachrome 160 Tungsten film.

Plasmid and Lambda vectors

DS941	STRAIN			. ••	YP-1	Actin A2	aDH2023 aDH1523	рИС19 рЕНВL18 NM1149	PLASHID
( <u>rec</u> F143 <u>pro</u> A7 <u>str</u> 31 <u>thr</u> 1 <u>leu6</u> <u>tsx33 mlt</u> 12 <u>his4 arg</u> E3 <u>lac</u> Iq <u>lac</u> Z <b>A</b> m15)	GENOTYPE	<u>E.coli strains</u>	TABLE 2.1		BamHI-BglII fragment of yolk protein 1 in pGEM-1	EcoRI fragment of Actin 5C in pBR322	Sall fragment of sgs3 in pBR322 HindIII fragment of sgs4 in pBR322	amp <sup>r</sup> amp <sup>r</sup> f1(G) $\lambda b538 \frac{sr1}{3}^{0} \frac{imm^{4}}{3}^{4}$ $\frac{sr1}{6}^{0} \frac{shnd}{11}^{11}$	GENOTYPE
D. Sherratt	SOURCE/REFERENCE			t	Hung and Wensink	Fyrberg et al	Kaiser et al Muskavitch and Hogness	Yanisch-Perron et al Dent <u>e et</u> al Hurray	REFERENCE
				• •	(1981)	(1981)	(1986) (1980)	(1985) (1985) (1983)	

TABLE 2.2

L87 NH514 NH621

(<u>supE supF hsdR</u> (<u>hsdR argH galE</u> (<u>hsdR mcrA mcrB</u>

<u>trpE</u> metD tonA) <u>galX</u> <u>strA lyc</u>B7) <u>supE44</u> <u>rec</u>D1009)

Amersham Murray (1983) Whittaker et al (1988)

#### 2.3 DROSOPHILA

A description of the drosophila stocks used in this work can be found in table 2.3. All designations are as described by Lindsley and Grell (1968) or Lindsley and Zimm (1987). Flies were kept at 18°C or 25°C depending on genotype and reared on Glasgow fly food (10g agar, 15g sucrose, 30g glucose, 35g yeast, 15g maize meal, 10g wheatgerm, 30g treacle, 10g soya flour, 0.1% Nipagen, 0.5% propionic acid per litre of water).

Flies carrying homozygous mutations in sex transforming loci were generated as follows.

<u>tra2</u>: <u>tra2<sup>ts1</sup> bw</u>/CyO;BsY males were crossed to virgin Df(2L)trix/CyO females, mated for 3 days at 18<sup>o</sup>C and transferred to 29<sup>o</sup>C. Straight winged progeny were homozygotes <sup>7</sup> and XY flies were identified by their Bar eye phenotype.

Germline deficient flies were produced in two ways, both of which utilise maternal effect mutations.

<u>Tudor</u> ( $wc^8$ ): Virgin F1 homozygotes were collected from a  $wc^8$ <u>bw</u> <u>sp</u>/CyO stock and self crossed, the F2 generation fail to form pole cells. To assess the penetrance of the phenoype 150 F2 males were mated to virgin wild type females. No larvae were detected in a two week period.

<u>Oskar</u>: A similar crossing scheme was used for  $osk^{301}$  flies but due to the temperature sensitive nature of this mutation the F2 generation was produced at  $18^{\circ}C$ 

XXY females were produced from a C(1)M3/FM6/Y stock.

XO males were produced from a C(1)RM/In(1)E Y/O stock.

In the later stages of this work single sex non-mutant flies for the preparation of mRNA for northern blots were generated with the segregation distorter stocks constructed by T.W. Lyttle and kindly provided by J. Lucchesi (Lyttle, 1989; Walker et al., 1989), Table 2.4 lists the genotypes of the strains used. The following is a description of the crossing scheme used.

TABLE 2.4

SD No4 +/+/Y ; SD-ARM 28 / In(2L)Cy In(2R)Cy, <u>S<sup>2</sup> Cy cn bw</u>

SD No1 C(1)DX, Y £ / T(X;Y)22-3, Y Y f. Rsp<sup>S</sup> B<sup>S</sup> / Y ; In(2L)Cy, CY E(SD) Rsp<sup>1</sup> bw / cn bw SD No3 +/+/Dp(2;Y)CB25-4, Y<sup>+</sup> Y<sup>S</sup>.Y<sup>L</sup> Rsp<sup>S</sup> B<sup>S</sup>; In(2L)Cy, CY E(SD) Rsp<sup>1</sup> bw / cn bw C(1)DX, <u>x</u> <u>f</u> /+/ Y ; SD-ARM 28 / In(2L)Cy In(2R)Cy, <u>S</u><sup>2</sup> <u>Cy cn bw</u> SD No2

SD STRAINS

(1968) Belote and Lucchesi (1980) (1987) (1986) (1968) (1973) (1985) (1968) (1968) Boswell and Mahawold Schalet and Lefevre Lindsley and Grell Lindsley and Grell Lindsley and Grell Lindsley and Grell Belote and Baker Lehmann et al REFERENCE 0 TABLE 2.3 C(1)RM, Y M / In(1)E Y<sup>L</sup>.Y<sup>S</sup>, Y M <u>f</u> tra2<sup>ts1</sup> bw / CyO ; BsY C(1)M3 / FM6 / Y ; <u>bw st</u> STRAIN GENOTYPE p<sup>p</sup> osk<sup>301</sup> e / TM3 Df(2R)trix / CyO <mark>ыс<sup>8</sup> bw зр / </mark>СуО (MT) Oregon R (WT) Canton S (WT) m56 i.

Drosophila stocks

<u>XY flies</u>: Curly-winged, brown-eyed males from SD No1 were mated to curly winged, star-eyed virgin females from SD No2. Curly winged vermilion-eyed male progeny were mated with virgin OregonR females. The resultant progeny were >96% male.

XX flies: Curly-winged, brown-eyed males from SD No3 were crossed with curly-winged star-eyed virgin females from SD No4. Curly-winged male progeny with wild type eyes were mated with OregonR virgin females. The progeny from this cross were >97% female.

TABLE 2.4

SD No4 +/+/Y ; SD-ARM 28 / In(2L)Cy In(2R)Cy, <u>S<sup>2</sup> Cy cn bw</u>

SD No1 C(1)DX, Y f / T(X;Y)22-3, Y Y f. Rsp<sup>S</sup> B<sup>S</sup> / Y ; In(2L)Cy, CY E(SD) Rsp<sup>1</sup> bw / cn bw SD No3 +/+/Dp(2;Y)CB25-4, Y<sup>+</sup> Y<sup>S</sup>.Y<sup>L</sup> Rsp<sup>S</sup> B<sup>S</sup>; In(2L)Cy, CY E(SD) Rsp<sup>1</sup> bw / cn bw C(1)DX, <u>Y</u> <u>F</u> /+/ Y ; SD-ARM 28 / In(2L)Cy In(2R)Cy, <u>S</u><sup>2</sup> <u>Cy cn bw</u> SD STRAINS SD No2

(1968) (1987) (1968) (1973) Belote and Lucchesi (1980) (1968) (1968) (1985) (1986) Boswell and Mahawold Schalet and Lefevre Lindsley and Grell Lindsley and Grell Lindsley and Grell Lindsley and Grell Belote and Baker Lehmann et al REFERENCE 0 **TABLE 2.3** C(1)RM, Y M / In(1)E Y<sup>L</sup>.Y<sup>S</sup>, Y M f C(1)M3 / FM6 / Y ; <u>bw st</u> tra2<sup>ts1</sup> bw / CyO ; BsY STRAIN GENOTYPE p<sup>p</sup> osk<sup>301</sup> e / TM3 Df(2R)trix / CyO wc<sup>8</sup> bw sp / CyO Canton S (WT) (HT) Oregon R (WT) m56 i.

Drosophila stocks

#### CHAPTER 3

### ISOLATION OF SEX-SPECIFIC GENES FROM D.MELANOGASTER

To help further our understanding of the sex specific control of gene expression I set out to isolate Drosophila genes expressed only in one sex. By studying such genes it is to be hoped that some insight can be gained into the molecular mechanisms that act to regulate genes in a tissue specific and temporal way. Sexual dimorphic gene expression in <u>D.melanogaster</u> is a convenient system in which to study this problem for several reasons. Firstly this is because the phenotype, either male or female, is specific to the whole organism (with the caveat mentioned at the end of chapter 1 regarding spatial differences in gene expression). This facilitates isolation of large quantities of RNA populations which differ only in the transcripts of interest. Secondly the regulatory hierarchy which leads to somatic sexual differentiation, and to a lesser extent dosage compensation and germline sex determination, is well characterised at the genetic and molecular level. This allows the initial characterisation of sex-specific genes by straightforward genetic tests. Thirdly the wealth of genetic and molecular techniques available to the <u>Drosophila</u> investigator allows identification of DNA sequences that confer specific This can be routinely accomplished regulation <u>in-vivo</u>. by reintroducing cloned regulatory sequences, together with appropriate reporter genes, into the fly and studying the expression of the reporter (Spradling, 1986).

There are several ways in which one could envisage isolating genes expressed sex-specifically. The classical way screen for a sex-specific be to mutagenise and would approach has been very successful in phenotype. Such an identifying new alleles of the genes involved in the control somatic sex determination as well as the male-specificof lethal genes. Such screens are relatively straightforward to out when the phenotype is sex-specific lethal or sex carry transforming. However the majority of genes expressed sexspecifically may be expected to have a less dramatic phenotype (e.g. sterility) when mutated . In such cases it is far more difficult to determine the molecular nature of the phenotype or indeed the tissue in which the gene is normally mutant line generated would require а expressed. Each laborious examination of individuals in order to establish а

morphological defect associated with the mutation. Since it has been estimated that approximately 10% of the mutable loci in the Drosophila genome can be mutated to give a male sterile phenotype (Lindsley and Lifschytz, 1972; Schafer, 1986a) this approach would involve screening a prohibitively large number of individuals (>600). A molecular analysis of any mutation isolated in this way would still require localisation and isolation of the gene which in itself can be a difficult and time consuming process.

More recently an elegant approach has been developed by O'Kane and Gehring (1987) utilising a P-element vector containing the <u>E.coli</u> B-galactosidase gene (LacZ). The lacZ gene is under the control of the weakly constitutive Pelement promoter which permits only low levels of <u>lacZ</u> transcription. If this vector is induced to transpose to a new chromosomal location where it comes under the influence of a genomic enhancer element then elevated <u>lacZ</u> expression can readily be detected by staining for B-galactosidase activity. The pattern of staining, to a first approximation, reflects the developmental specificity of the particular enhancer element. By using a vector containing an antibiotic resistance gene and an <u>E.coli</u> origin of replication the genomic DNA flanking the insertion site can be easily This approach is already being widely used to isolated. identify early markers of neural cell development (for example Bellen et al., 1989; Ethan et al., 1989; Wilson et al., 1989) and will undoubtedly be more widely used in the study of sexual differentiation. Indeed some insertions which appear to have staining localised to the embryonic germ cells already been identified (Bellen et al., 1989). It must have be born in mind however that insertions may be influenced by more than one regulatory element and hence give misleading patterns of expression. Also the enhancer elements detected by this technique may be located some distance away from the genes they control making it rather more difficult to isolate these genes. A third difficulty may also arise in detecting enhancers" which are not associated with any "naked particular gene.

An alternative approach to the isolation of sex-specific genes is differential screening of a cDNA or genomic DNA library; so called +/- screening. With this method duplicate filter replicas of plaques are taken from a library and

screened with probes derived from two different mRNA populations, one population containing the transcripts of interest and the other deficient or containing reduced levels of these sequences. Clones which hybridise with one probe but show no or reduced hybridisation with the other may then be examined further as candidates for differentially expressed genes. Such an approach has been successful in the isolation of stage-specific transcripts from developing Xenopus embryos (Dworkin and Dawid, 1980) and transcripts regulated during Dicyostelium development (Williams and Lloyd, 1978). The major advantage of the differential screening approach is that clones identified in this way correspond directly to the transcript of the differentially expressed gene and relatively little subsequent manipulation is required to obtain the purified sequence. On the other hand a severe disadvantage is that such a sequence is isolated with no knowledge of it's biological function other than the fact that it is differentially expressed.

This technique would appear to be well suited to the isolation of sex-specific transcripts from Drosophila since the main requirement is two RNA populations which differ in the abundance of the transcripts significantly of interest. The method has been successfully employed by other workers in this field (Schafer, 1986a; Di Benedetto et al., 1987). However these groups did not identify an important class of genes, namely sex-specific transcripts expressed in early development. In one case the screen was deliberately biased against detecting such transcripts; Schafer, being primarily interested in spermatogenesis, prepared probes from selected adult mRNA of low molecular weight (6-9S) and size screened out any clones which also hybridised with embryonic mRNA. In the other Di Benedetto et al screened with sexed adult and pupal probes and identified 7 new sex-specific genes. However none of these showed detectable expression in embryo. As the somatic and germline components of the the gonad are almost certainly sexually differentiated during the embryonic stage, isolation of transcripts from these tissues would be a considerable asset to molecular studies of sexspecific gene expression. With this in mind I set out to isolate sex-specific transcripts using mRNA from male and larvae, this being the earliest third instar female developmental stage which could be reliably sexed in large

enough quantities.

Before embarking upon such a screening procedure there are several important considerations.

1) As discussed above the primary requirement is the identification of suitably different mRNA populations to use in the preparation of cDNA probes. In this case male and female 3rd. instar larvae.

2) The relative abundance of the transcripts one wishes identify is important when considering this experimental to approach. Generally speaking +/- screening will allow the isolation of sequences present as 0.1% of the mRNA population (Sargent, 1987). If careful attention is paid when preparing the probes and hybridisation conditions are optimised this limit can be reduced by an order of magnitude. If one is interested in isolating genes expressed at a lower level than 0.01% then subtracted probes must be employed (Sargent, 1987). As the production, of subtracted probes is more technically demanding than +/- screening I decided to opt initially for the simpler approach.

3) Finally the type of library to be screened, either cDNA or genomic, must be decided upon and several considerations are relevant to this matter.

 a) Number of clones to be screened: for a genomic library this parameter (N) is determined by the insert size and may be estimated from the equation,

 $N = \frac{\ln(1-p)}{\ln(1-x/y)}$ 

(Kaiser and Murray, 1985) where p is the probability of finding a particular clone in a library with inserts of x kb, from an organism with a haploid genome size of y kb. Using a <u>Drosophila</u> library containing 20kb inserts one would need to screen approximately 30,000 primary recombinants to have a 99% probability of finding a particular sequence. The use of an amplified library would require screening of a larger number of clones in order to overcome misrepresentation bias.

For a cDNA library the number of clones to be screened (N) depends on the abundance of the required transcript in the mRNA population and is estimated from the equation

#### ln(1-z)

(Sargent, 1987) where p is again the probability of finding the sequence and z is the fractional abundance the of sequence in the mRNA population. Using the limit of 0.01% of the mRNA population detectable in a +/- screening then we would need to screen 50,000 cDNA clones to have 998 а probability of finding a transcript in this abundance class; almost twice as many as for a <u>Drosophila</u> genomic library. If however the subject of investigation was an organism with а significantly larger genome than <u>Drosophila</u> then screening a cDNA library would require fewer clones than a genomic library.

b) Insert characteristics; each individual clone in a cDNA library should contain sequences representing only а single gene (or gene family). The insert of a clone from а sequences genomic library however can potentially contain which represent more than one gene. In practice this means that if a differentially expressed gene is linked to another non-differential gene on the same clone the hybridisation signal of the latter may obscure the differential signal of the former. Although this problem may be avoided by using a this probe to screen а genomic library subtracted consideration would favour the use of a cDNA library in +/screens.

Screening considerations: When a complex C) mRNA population is used to generate a probe then the intensity of hybridisation signal generated by a given transcript is the generally directly proportional to the abundance of that sequence in the mRNA population (Williams, 1981). This means that clones which show strong hybridisation code for abundant transcripts. With a genomic library, clones which code for abundant mRNAs are generally present at the same frequency as clones which code for transcripts in the low abundance class. is not the case for a cDNA library where the frequency This of clones containing a particular sequence is dependent on the abundance of that sequence in the mRNA population. In practise this means that the clones which give the strongest hybridisation signals may be present many times on а particular filter. This consideration effectively limits the density at which a cDNA library can be screened using the

+/- technique as the signals from the abundant sequences will mask hybridisation to other clones. Again this problem can be overcome by the use of subtracted probes. With these considerations in mind I decided to construct and screen larval instar cDNA libraries and also screen, third in conjunction with K.Kaiser, a primary genomic library to try identify sex-specific transcripts. and We felt that by screening both types of library we would maximise our chances of identifying new sex-specific genes.

#### 3.2 RESULTS

#### 3.2i Construction of cDNA Libraries

Drosophila male and female 3rd larval instar CDNA libraries were constructed as described in the methods section. Table 3.1 shows the yields of the cDNA products obtained. For both templates the first strand synthesis products are within the range expected (30-50%: Berger et al., 1983). The yield of globin first strand cDNA, which approaches the maximum expected, may reflect more efficient synthesis due to the small size of the template. Second strand products are again approximately as expected (90-100%), the yield of male second strand at 109% probably reflects pipetting error when dispensing aliquots for acid precipitable counting. The low yield of globin second strand was consistently observed in several experiments. No attempt for was made to determine the the reason this. under alkaline conditions was used Electrophoresis to determine the size and structure of the cDNA. Figure 3.1a shows an autoradiogram of synthesis products analysed on а 1.5% alkaline agarose gel. The size distribution (0.4 to is approximately equivalent for first and second 2.5kb) strand products indicating a low level of hairpin structure formed during first strand synthesis. The discrete bands visible at approximately 750 and 1100nt. are consistently observed when I synthesise cDNA from larval mRNA. I do not if these represent copies of abundant mRNAs (which are know not visible in neutral RNA gels) or are derived from self priming rRNA molecules. Since the mRNA was purified by three

TABLE 3.1

	FEMALE	MALE	GLOBIN
% INCORPORATION	2.7	2.4	3.0
YEILD 1st STRAND	380ng	340ng	420ng
%mRNA TRANSCRIBED	38	34	42
% INCORPORATION	2.7	2.6	1.1
YEILD 2nd STRAND	380ng	370ng	150ng
% 1st STRAND TRANSCRIBED	100	109	36

TABLE 3.2

	TITER 		<u>187</u> .nm514 RATIO	ADJUSTED BACKGROUND pfu/m1	TOTAL RECOMBINANTS pfu/m1
RELIGATED VECTOR	6 4x10	4 4x10	100		
VECTOR +	6	6	1.5	4	6
FEMALE cDNA	3x10	2x10		3x10	1.9x10
VECTOR +	6	5	5.8	4	<b>5</b>
MALE cDNA	<b>3</b> .5x10	6x10		3.5x10	5.7x10
rounds of oligo(dT)-cellulose chromatography the latter possibility is considered less likely, although an mRNA population is never completely depleted of rRNA.

The double stranded cDNA was then methylated, ECORI linkers added and the resulting molecules ligated into the lambda insertion vector NM1149. The recombinants were packaged into phage particles using a commercial packaging kit. A control with self-ligated vector was included to asses the effectiveness of the biological selection. The packaged phage were titered on non-selective (L87) and selective (NM514) E.coli hosts. The titers are shown in Table 3.2. The ratio of titers on L87 to NM514 should be high in the absence of insert and much lower when insert is present indicating that the biological selection is operating as expected, this is observed for both libraries. The titers also allow the number of recombinants within the library to be estimated. An adjusted background is calculated by subtracting the number plaques expected on NM514 in the absence of inserts of from the number actually obtained in the presence of the inserts (the value expected on NM514 is estimated by dividing the total number of plaques on the L87 host by the L87:NM514 ratio obtained from the self-ligated vector.). This figure however is a minimal estimate and does not take into account damage to the ends of the vector arms which may be caused by impurities present in the cDNA. Damage to the vector arms can "fool" the biological selection resulting in false positives. Both of the libraries appear to contain approximately 95% recombinants, this figure is higher than expected (70 - 80%)and may reflect damage to the vector arms. A more realistic figure for the level of recombinants in the library can be obtained by physical analysis. To this end 48 plaques were selected at random from each library, miniprep DNA prepared, <sup>32</sup>P datp digested with <u>EcoRI</u> and end labelled with using "Klenow" polymerase. After electrophoresis in 1% agarose gels the DNA was visualised by autoradiography (not shown). In the male library 37 phage contained inserts (78%) and of these 4 (8%) contained more than one EcoRI fragment, the average size was 850bp. The female library sample contained 38 phage with inserts (79%), 9 (23%) of which contained more than one EcoRI fragment, the average size in this case was 890bp. There are possible explanations for the presence of more than two one fragment in a particular phage, one is that the cDNA contains

Figure 3.1a: Autoradiogram of cDNA synthesis products electrophoresed in a 1.5% alkaline-agarose gel. Lanes 1) Male 1st strand, 2) Male 2nd strand, 3) Female 1st strand, 4) Female 2nd strand, 5) Globin 1st strand, 6) Globin 2nd strand. Size markers indicated are in kilobases. Overnight exposure.

<u>Figure 3.1b</u>: Autoradiogram of <u>sqs3</u> and <u>sqs4</u> positive cDNA clones end labelled and electrophoresed in a 1.0% agarose gel. Lanes 1-3 female library <u>sqs3</u>; 4 & 5 female library <u>sqs4</u>; 6-8 male library <u>sqs3</u>; 9 & 10 male library <u>sqs4</u>; 11 NM1149. Marker sizes are in kilobasepairs, exposure time 2 hours.

Figure 3.1c: Autoradiogram of northern blots probed with various DNAs. Even numbered lanes contain male mRNA (1ug), odd numbered lanes contain female mRNA (1ug). Each pair of lanes were probed with the following; 1,2 aDM1523; 3,4 male <u>sgs4</u> clone 9; 5,6 aDM2023; 7,8 male <u>sgs3</u> clone 7; 9,10 male <u>sgs3</u> clone 8. Sizes are in kilobases. The blots were washed at high stringency and exposed overnight.

<u>Figure 3.1d</u>: Autoradiogram of duplicate filters from male cDNA library probed with cDNA derived from male and female 3rd instar larvae as indicated. The blots were washed at high stringency and exposed overnight.









3·1d

internal EcoRI site the other is that two different an CDNA have become inserted into the same phage. molecules These possibilities may be distinguished by using the inserts as probes to northern blots. This was not carried out as it was felt that the low level of "double inserts" present in the male library would not interfere significantly with the screening. The relatively high level of double inserts in the female library may present some screening problems.

intention was to use these As the libraries for differential screening experiments I considered it prudent to examine for the presence of known sequences them at the expected abundance. То this end 7000 plaques from each library were screened with Sqs3 and Sqs4 probes (aDM2023 and aDM1523: Kaiser et al., 1986). The male library yielded 43 <u>Sgs3</u> and 12 <u>Sqs4</u> positives, the female 50 and 15. Although there are no direct abundance measurements of these published ( the abundance of the transcripts transcripts changes through larval development), the ratio of Sqs4 to Sgs3 transcripts has been documented during studies on dosage compensation to be approximately 0.25 (Kaiser et al., 1986). figure is in reasonable agreement with the ratios Ι This observe with the libraries (0.28 and 0.3). Three Sgs3 and two library. DNA <u>Sqs4</u> positives were purified from each was extracted by the miniprep method, digested with EcoRI and end labelled with <sup>32</sup>PdATP and "Klenow" polymerase. The insert size was determined by agarose gel electrophoresis: an autoradiogram of this gel is shown in Figure 3.1b. The sizes approximately 1100nt of the <u>Sqs3</u> and <u>Sqs4</u> transcripts are Meyerowitz, (Hofmann and Korg, 1987) and 1300nt (Crosby and the male respectively. As the autoradiogram shows 1986) library clones examined contain some inserts of approximately correct size (lanes 6-10). The clone in lane 8 has more the than one insert, it was not determined which one was the Sgs3 1-5) contained positive. The female library clones (lanes smaller smaller inserts, the clone in lane 2 has an insert than 250bp which is barely visible on this exposure. DNA from three of the positives from the male library (lanes 7,8 and and plasmids containing Sgs3 and Sgs4 inserts were nick 9) translated and used as hybridisation probes to northern blots male and female mRNA. The mRNA samples were run on the of for and the filter divided same blotted together gel, hybridisation with separate probes. Figure 3.1c shows an

autoradiogram from this experiment. All the clones tested appear to hybridise to the correct transcript, the clone from lane 8 also hybridises to a weaker transcript of higher molecular weight which is consistent with the additional inserts in this phage being derived from another transcript.

То summarise, male and female third larval instar CDNA libraries were successfully constructed, the libraries contain inserts of known transcripts at approximately the expected frequency. In the case of the male library at least some of the inserts are near full length and recognise the correct transcript on Northern blots. I concluded that the for differential libraries were suitable screening experiments. Half of each library was amplified (Maniatis et al., 1982) for pilot experiments. When screened with the Sqs3 probe the amplified library gave a similar number of positives as the primary library (data not shown).

# 3.211 Differential Screening Of The Male cDNA Library

Approximately 15,000 plaques from the primary male cDNA library were plated at a density of 1,500 per 10  $\,\mathrm{cm}^2$ plate and two filter replicas taken from each plate. One set was screened with <sup>32</sup>P cDNA derived from male 3rd instar larval mRNA, the other set with <sup>32</sup>P cDNA from female 3rd instar The probes were prepared as described in the larval mRNA. methods section and were used at specific activities of >10<sup>8</sup>cpm/ug of cDNA. The hybridisations were carried out in aqueous solution and washed at high stringency as described in the methods section. An example of a pair of filters from far these experiments is shown in Figure 3.1d. Α one of greater number of plaques than expected show a signal after only overnight exposure. Furthermore the signals were of even intensity, again an unexpected result. The relatively number of plaques hybridising is approximately 75% of the total plated which eliminates the possibility that the probe was contaminated with Lambda sequences. This template conclusion was corroborated when screening genomic libraries with probes synthesised from the same template. The number of positives detected is in very close agreement with the number plaques judged to contain inserts by physical analysis, of

this suggests that the hybridisation was due to the presence of an insert. It is possible that the hybridisation was between the oligo(dT) tail of the cDNA probe and the dA-dT tract present in the cDNA inserts, although polyA was present in the hybridisation solution to prevent such hybridisation. To try and test this possibility I synthesised cDNA probes primed from random hexamers rather than  $oligo(dT)_{12-18}$ . However these experiments were hampered by high levels of background consistently observed with these probes. Similar background was also observed when the genomic library and dot of isolated phage DNA were screened. It is not clear blots why such high background was observed but it may be specific the random hexamers and reverse transcriptase. to No background was observed when plasmid probes were random primed with hexamers and Klenow polymerase, however plasmid probes were not synthesised with hexamer primers and reverse transcriptase. Alternatively the background may have been due impurities in the cDNA which oligo-dT priming was to not sensitive to. Numerous experiments were performed to try and eliminate this problem. Templates and probes were phenol extracted or filtered through nitro-cellulose membranes prior Different concentrations of hexamers ranging over to use. several orders of magnitude were tried. Unfortunately none of these measures were successful and the cause of these phenomena remain obscure. In view of the success achieved in screening the genomic library and pressure of time I decided discontinue attempts at screening the cDNA library. The to libraries however will still be useful for isolation of cDNA copies of genes identified in the +/- screens of the genomic library.

# 3.2111 Differential Screening Of A Drosophila Genomic Library

The <u>Drosophila</u> genomic library used was constructed in the lambda vector EMBL3 by K.Kaiser. It contains 6-18kb inserts derived from Oregon-R DNA partially digested with <u>Sau3A</u>. Two differential screens were carried out.

<u>Screen 1</u>: approximately 20,000 primary recombinants were screened at a density of 5,000 plaques per  $10 \text{ cm}^2$  plate. Duplicate filters were taken from each plate and screened

Figure 3.2a: Autoradiogram of duplicate filters from a <u>Drosophila</u> genomic library probed with cDNA derived from male and female L3 mRNA. See text under Screen 2 for details. Holes represent putative positives. Exposure was for two weeks.

<u>Figure 3.2b</u>: Autoradiogram from secondary screen of positives from screen 2. See text for details. Exposure was for two weeks.



<sup>32</sup>P cDNA from male and female third instar larvae. with The probes were labelled to a specific activity of >10<sup>8</sup>cpm/ug of cDNA. Hybridisations in aqueous solution and high stringency washes were carried out as described in the methods section. After exposing the filters for two weeks approximately 10% of the plaques gave signals on the autoradiograms. Of these 15 were judged to be male specific or differential. Α region corresponding to the positive signal was picked and the phage streaked out to obtain single plaques. Ten plaques from each of the isolates were picked out into an array and screened again. Five of the primary positives failed to give differential signals. The remaining phage designated S1-S10 screened again and a further 2 lost. These 8 positives were screened for a fourth time along with the positives were isolated from the screen described below.

Screen 2: in this experiment 30,000 primary recombinants screened at a density of 8,000 plaques per plate using were the same conditions as above. In this case 54 plaques were identified putative male specific/differential, the as majority showing differential signals. A pair of filters from is shown in Figure 3.2a. Positives were aqain this screen streaked out and screened in an array. A row of Lambda clones containing an <u>Sgs4</u> insert were included to asses the degree 54 of hybridisation to each filter (Figure 3.2b). Of the screened 34 positives remained these were designated K1-K34. A tertiary screen reduced this number to 20.

Final screening: The 28 plaques from both screens were into an array (3 plaques per picked out isolate) and 4 32<sub>P</sub> taken from the plate. These were screened with filters cDNA from new preparations of sexed larval mRNA and also with derived from sexed adult mRNA, the screens are shown probes in Figure 3.3. The upper panel details the signals for each individually while the lower panel shows the whole clone similar The majority of the clones show autoradiogram. expression with larval and adult probes which confirms that larvae were correctly sexed. It is also that apparent the of the clones are still not single phage populations. some is probably a result of the previous screenings having This been performed in close packed arrays where it is difficult to avoid some cross contamination. This is especially so when period while the the plates are left for a two week the hybridisation and autoradiography are carried out. On

Figure 3.3a and b: Autoradiograms of quaternary screening of "male-specific" positives from screens 1 and 2. The lower panels (3.3b) show the complete filters screened with cDNA derived from male and female larvae and adults as indicated. The upper panel (3.3a) has these same filters organised by individual clone for convenience. In this case the rows for each clone represent, from top to bottom: Male L3, Female L3, Male adult, female adult. The exposure time was 1 week.









Q



gS1	gS2	gS4	gS5	gS7	
g S8	gS9	g\$10	gK6	gK8	
g K 10	gK11	gK12	gK14	gK15	
gK16	gK17	gK18	gK25	gK 24	
gK22	g <b>K 2</b> 1	gK27	gK 29	gK 30	
gK 31	gK 33	gK34			

basis of this experiment the clones were divided into 3 groups.

<u>Group 1</u>: Male specific, the 13 clones in this group (gS2gK33) show no or very little hybridisation with female probes. These clones probably contain genuine male specific genes.

<u>Group 2</u>: Male differential, the 7 clones in this class (gS1-gK31) show reduced hybridisation with female probes as compared to male probes. Two possibilities could account for this pattern of hybridisation; One is that the transcript is expressed at a higher level in males than in females. The other is that more than one transcribed sequence is located the cloned fragment and only one of on these is male specific. This question can be resolved by dissecting the clone into smaller fragments and analysing them individually.

Group 3: ambiguous, the remaining 8 clones were placed in category. Three of them showed no hybridisation, this qS7, gK11 and gK29. Two appear to be larval specific, qK24 and gK34. Of the remaining three gS5 gives a weak signal and is difficult to asses, gK12 shows adult male specific expression but is not sex-specific in larvae. gK25 appears to contain 3 different phage, one is not sex-specific, one larval is specific and the third is male differential.

After screening the clones four times at least 20 remain which show male specific/differential expression. For further analysis I selected 8 clones from group 1: gS2, gS4, gS9, gS10, gK14, gK15, gK21 and gK33, two clones from group 2: gS1 and gS8, and two from group 3: gS5 as I was unsure of it's status and gS7. This clone appeared negative in this screen isolate in previous screens so I went back to the but not third screen and used that in subsequent from the experiments. The remaining clones are being analysed by another worker.

# 3.2iv Reverse Northern Analysis

To permit a more detailed analysis of the putative malespecific genes a "reverse northern" analysis was performed. In these experiments recombinant phage DNAs (purified on a large scale) were digested with restriction enzymes,

<u>Figure 3.4a-c</u>: EtBr stained 0.8% agarose gel containing <u>BamH1</u> digested DNA from the set of male-specific clones selected for further analysis (0.5ug/lane). The clone numbers are given along the top. Size markers are in kilobasepairs. B is an autoradiogram of a southern blot of this gel probed with cDNA derived from male adult mRNA, C is an autoradiogram of a blot from a duplicate gel probed with cDNA from female mRNA. Both blots were washed at high stringency and exposed for three days.



В

С





electrophoresed in duplicate 0.8% agarose and qels transferred to nylon filters. The filters were probed in parallel, one with cDNA prepared from male adult mRNA, the other with cDNA prepared from female adult mRNA. Probes were prepared to specific activities of >10<sup>8</sup>cpm/ug of cDNA. After hybridisation, washing at high stringency aqueous and the filters were identify autoradiography compared to restriction fragments with male-specific hybridisation. Two controls were included to asses the sex-specificity of the probes. YP1-1 is a fragment of the yolk protein 1 gene in the plasmid vector pGem1 (Kindly provided by M.Bownes). cS9 is а 650bp cDNA which I isolated during pilot experiments, it is described fully in chapter 6. At the time of these adult non sex-specific control was experiments an not available in our laboratory. Somewhat fortuitously clone qS7 appears to fulfil this function.

The reverse northern analysis was performed on phage DNA digested with SalI, BamHI and HindIII. The qel and autoradiograms for the BamHI digests are shown in Figure 3.4. upper panel shows the ethidium bromide stained gel, the The middle panel shows hybridisation with male cDNA and the lower with female cDNA. Several bands hybridise strongly with male show little or no hybridisation with female. The CDNA and CDNA male specific control cS9 only hybridises with male (band at 650bp). Weak hybridisation between the vector and female cDNA is also apparent. A band of 5kb, which represents undigested vector DNA not visible on the stained qel, is detected by hybridisation female cDNA. In addition weak between female cDNA and the arms of the Lambda vector is spurious hybridisation apparent. The reason for this is female-specific control, YP1, unknown. With the strong female cDNA is observed. However weak hybridisation with hybridisation with male cDNA is also detected. I estimate by eye that hybridisation to YP-1 with the male probe is around lower than with the female probe. This probably 100 fold reflects slight contamination of male with female, mRNA however the analysis is not unduly affected by this. The combined results of these experiments are given in table 3.3.

The majority of the clones show male specific hybridisation to a small restriction fragment. Several of the male-specific fragments were subcloned into plasmid vectors for further analysis and will be described in the following

TABLE 3.3

	<u>Sall</u>	<u>BamHI</u>	<u>HindIII</u>
g\$1	<u>1.6Kb</u> (Y)	(¥)	¥ (3.7Kb)
<b>g</b> \$2	10.5Kb (6.5Kb)	<u>5.8,1.6,0.8Kb</u> (4.2Kb)	(¥)
g\$4	v	٧	Ŷ
g\$5	(8.5,5.4Kb)	(¥)	(8.0,5.0Kb)
g\$7	(2.1,1.3Kb)	(1.8Kb)	(¥)
g\$8	(7.0Kb)	(12.0Kb)	( <u>5.5Kb)</u>
gS9	11.0Kb	4.1Kb	<u>3.2Kb</u>
g\$10	7.5, <u>1.6Kb</u>	<u>(</u> ¥)	(V), <u>2.6Kb</u>
gK14	11.0Kb (3.4Kb)	6.2Kb (Y)	9.0Kb (V)
gK15	1.1Kb (6.5Kb)	1.6,0.8Kb (4.2Kb)	(¥) (5.0Kb)
gK21	(¥)	(¥)	5.5Kb (4.0Kb)
gK33	۷	<b>V</b>	<u>0.6Kb</u>

Table 3.3: Restriction fragments identified by reverse northern blots. The sizes are in Kilobasepains. Underlined fragments were sub-cloned into plasmid vectors for further analysis. Those fragments enclosed in brackets hybridised with cDNA from both sexes. V denotes hybridisation to cloned sequences which remain attached to the Lambda vector arms. chapters. In addition a number of restriction fragments hybridise weakly with cDNA from both sexes. This may be due to the presence of adjacent non sex-specific genes.

As is apparent clone gS7 is not sex-specific and was omitted from subsequent analyses. Clone gS5 shows very weak hybridisation which appears if anything to be female elevated. Northern blot analysis with this clone failed to detect any sex-specific transcripts (not shown). Consequently clone will not be discussed further. Ten clones which this show male-specific hybridisation remain. From the restriction it appears that gS2 and gK15 overlap as they digests share fragments. common <u>BamHI</u> and SalI То determine the of the other clones blots of <u>SalI</u> relatedness digested DNA were probed with restriction fragments from each clone (not In this way the overlap between gS2 and gK15 shown). was confirmed and cross hybridisation between gS8 and gS9 was detected although these two clones do not appear to overlap. A more detailed analysis of each clone will be presented in the following chapters.

#### 3.3 DISCUSSION

Screening of the genomic library appears to have successfully identified cloned genomic fragments which contain genes for male specific transcripts. The isolation of clones which are male differential is comparable 20 to the screens carried out by other workers. Schafer (1986a) 35,000 clones and isolated 12 which defined 5 new screened genes, DiBenedetto et al. (1987) screened 15,000 clones and 13 which defined 7 new genes and 3 previously **iso**lated isolated chorion and yolk protein genes. In their experiments these groups screened amplified libraries which may be both underrepresented for many sequences. By using a primary library we may have increased our chances of finding previously unidentified genes. The choice of larval templates for the preparation of the probes may have directed our screens to a different class of genes than those previously isolated. When we screen the genomic library with probes derived from sexed adults many more positives are apparent

than with the larval probes (K.Kaiser & S.Russell, unpublished data). These adult sex-specific clones probably represent transcripts derived from the highly differentiated structures which develop from the genital disc. As the gonad is the only major sexually dimorphic structure apparent during the larval stages of development, it is possible that the clones identified in our screens are transcribed specifically in this tissue. It is also noteworthy that no female-specific clones were identified in our screen. This reinforces the belief that the clones represent gonad specific transcripts as a) the female gonad is considerably larval stage smaller than the male during the and b) gametogenesis does not commence in the female until late in the pupal stage whereas primary spermatocytes can be detected the testis as early as the 1st larval instar in (Garcia-Bellido, 1964)

With regards to the screening there are steps which could Identification of positives the be improved upon. on autoradiograms of the primary screen was quite time consuming. As the plaques were distributed at random it was difficult to identify the differential signal amongst the majority which were not sex-specific. On the other hand the secondary screens where all of the plaques were ordered in an array was considerably easier. Purifying the primary also problematic. I do not know how positives was many genuine positives were lost during the purification but it would be surprising if none were. Poorly growing clones may have been missed when plaques were streaked out although every effort was made to select plaques of differing size. These problems could be overcome in a number of ways; the use of subtracted probes would make identification of positives much easier. However the purification of the positives would still be as problematic. If one were to screen the whole library in arrays identification and purification of clones would be greatly simplified. Thirty thousand plaques may seem like a prohibitively large number to plate in arrays but а device for picking out and replicating 1000 plaques at a time has recently been developed in our laboratory (MacKenzie et This may make such an approach feasible. The al., 1989). library other possibility is to construct a subtracted cDNA enriched for then relatively few clones, which are the sequences of interest, need be screened.

Disappointingly my attempts at screening cDNA libraries were unsuccessful. As discussed earlier the data suggests that the observed hybridisation was due to the presence of an insert, possibly mediated by the polyA tail. It is possible that the polyA added to the hybridisation solution was degraded and hence not effective as a blocking agent. The use of random primed cDNA should have overcome this problem but the levels of background obtained with these probes prevented this avenue from being exploited. I have little idea as to why this problem was encountered. Similar background has been observed by other workers in our laboratory, specifically the pD(N)<sub>6</sub> and source of reverse transcriptase used in with an alternative source this work. Perhaps of reverse primers or perhaps slightly transcriptase and/or longer primers may be more productive.

#### CHAPTER 4

## IDENTIFICATION OF PREVIOUSLY ISOLATED MALE-SPECIFIC GENES

#### 4.1 INTRODUCTION

The belief that we have isolated genuine male-specific would be strengthened by the identification of genes previously characterised genes, hence it is worth considering briefly male-specific genes isolated by other workers. Schafer (1986a; 1986b) screened 35,000 phage from the Maniatis library (Maniatis et al., 1978) with cDNA derived selected adult male mRNA. size He identified 195 from positives of which he discarded 183 as these hybridised with CDNA from embryos or adult females. The remaining 12 clones identified 5 genomic regions. Four; mst(2)Ag1 at 57D, mst(3)Ag3 at 95F, mst(2)Ag35 at 51F and a small gene family 75C, all encode small (around mst(3)Aq2 at 500nt) mRNAs expressed in the paragonia. These transcripts appear to be indirectly regulated by the somatic sex determination hierarchy. The fifth, mst(3)Gl9, at 87F encodes а 750nt transcript specific to the male germline.

DiBenedetto et al. (1987) screened 15,000 plaques from the same library with cDNA derived from male and female adult mRNA. In all they identified 13 sex-specific clones, 6 malespecific and 7 female-specific. Three of the female-specific clones were re-isolates of chorion or yolk protein genes and were type 1 ribosomal spacer sequences. The remaining 2 2 identified a novel gene fst(3)230. Of the male-specific genes mst(3)316 at 95E encodes a 350nt transcript expressed in the paragonia. The remaining 5; mst(2)323 at 25F, mst(2)325 at 47A, mst(3)349 at 66D, mst(3)345 at 95E and mst(3)336 at 98C all encode male-specific transcripts expressed in the germline. A few other male-specific genes have been cloned and these have been discussed in chapter 1.

an attempt to characterise our male-specific clones In several experimental approaches were used. In the first instance crude restriction maps were deduced by a series of restriction enzyme double digests. These maps merely provide working reference and it should be emphasised that they а have not been confirmed by sub-cloning or partial digestion. Genomic organisation of the cloned sequences was determined Southern blotting and in-situ hybridisation to polytene by chromosome spreads. In this way genomic location and copy number could be determined.

Preliminary characterisation of the expression of the male-specific transcripts was achieved via northern blotting

experiments. Information on the developmental regulation of the transcripts was obtained from northern blots of mRNA from different developmental stages. The use of sexed material from some stages provided confirmation of the sex-specificity observed in cDNA hybridisation experiments. The <u>SD</u> stocks discussed in the materials section were used to generate single-sex fly populations from which mRNA was isolated.

Hybridisation to mRNA from flies carrying different mutations or defects allows preliminary separation into those sequences transcribed in the germline and those transcribed the soma. The  $\underline{tra2}$  mutation, which transforms XX in somatic to phenotypic maleness, allows one to determine tissue whether or not a particular gene is under the control of the somatic sex-determination hierarchy. If a male-specific gene expressed in XX; tra2/tra2 pseudomales the most likely is conclusion is that it is expressed in somatic tissues and It should be noted however that few regulated by <u>dsx</u>. а in the spermatocytes are detected rudimentary testes of al., 1989) the pseudomales (Steinmann-Zwicky et so possibility of germline expression cannot be eliminated if weak expression is detected. If a male-specific transcript is not expressed in pseudomales then it is either not expressed somatically or it is not regulated by the somatic sex determination hierarchy. One caveat to this is with regard to tissues which are regulated by the hierarchy but which fail to develop fully in pseudomales such as the somatic component of the gonad. Here a low level of expression from the reduced tissue in pseudomales may not be detected on northern blots.

expression can be investigated using agametic Germline individuals. In these experiments offspring were generated from mothers homozygous for tudor (tud) (Boswell and Mahawold, 1985) and <u>oskar<sup>301</sup> (osk</u>)(Lehmann and Nusslein-Volhard, 1986) pole-cells mutations. The offspring develop without and consequently their gonads are underdeveloped and devoid of germ-cells. Failure to detect male-specific transcripts in indicative of germline specific these individuals is or germline dependent expression. Germline dependent expression is exemplified by chorion gene expression which occurs in the somatic follicle cells of the ovary but which requires a in order to normally developing oocyte be expressed (DiBenedetto et al., 1987). Expression of a male-specific gene in both pseudomales and agametic males argues strongly

for somatic expression. Failure to detect the transcript in either suggests germline expression (with the caveat discussed regarding the underdeveloped gonad). Expression in agametic males but not in pseudomales suggests somatic expression not regulated by the somatic sex-determination hierarchy, for example genes involved in dosage compensation. Expression of a male-specific gene in a pseudomale but not in an agametic male would indicate an early germline function but is not expected as very few XX germ cells enter spermatogenesis in pseudomales.

In the case of germline specific transcripts it would also be of some interest to investigate the effect of mutations affecting spermatogenesis. There are a large number of mutations which disrupt spermatogenesis and an investigation of them all would be somewhat time consuming. obtain some preliminary information expression of male-То specific genes was examined in males lacking a Y chromosome and in males (XY) homozygous for the tra2 mutation (XO) as these genotypes are associated with aberrant both of spermatogenesis. mRNA derived from females carrying Y а chromosome was also included on some blots (genotype XXY). For the most part hybridisation to this mRNA was identical to normal XX females. Lanes containing this mRNA are shown in some figures but will not be discussed further.

### 4.2 RESULTS

Two of the clones isolated in our screen (gK14 and gK21) appear to correspond to previously identified male-specific genes.

4.2i qK14: this clone maps to salivary gland chromosome region 66D/E as shown in Figure 4.1a. A crude restriction map the clone is shown in Figure 4.1c. The map was of derived from the double digests shown in Figure 4.2a and also by probing blots of the digests with the 3.2kb Sall restriction fragment marked on the restriction map (not shown). The restriction fragments identified on reverse northern blots are shown on the restriction map (black bar). In addition to male-specific fragments the hatched box indicates weak the from both sexes. The diagram also hybridisation to cDNA

- Figure 4.1a: Photomicrographs of <u>in-situ</u> hybridisation to polytene chromosomes using biotinylated lambda gK14. The large panel shows hybridisation to the left arm of chromosome 3. The insert shows hybridisation at region 66DE.
- Figure 4.1b: Photomicrographs of polytene chromosome spreads hybridised with biotinylated lambda gK21. The upper panel shows hybridisation to the right arm of chromosome 3. The lower panel shows hybridisation to an asynapsed chromosome at region 98CD. The signal is stronger on one of the homologues.
- <u>Figure 4.1c</u>: Crude restriction maps of the lambda clones gK14 and gK21. Enzymes; B=<u>BamHI</u>, H=<u>HindIII</u>, E=<u>EcoRI</u>, S=<u>SalI</u>. The black bar represents hybridisation with cDNA derived from males. The hatched bar represents hybridisation with cDNA derived from either sex. The dotted line indicates the 3.2Kb <u>SalI</u> fragment used as a probe. The arrows depict the overlap with the mst(3)349 and mst(3)336 clones isolated by DiBenedetto et al. (1987).



indicates the extent of the apparent overlap with the clone mst(3)349 identified by DiBenedetto et al. (1987).

When the whole phage is used as a probe to Northern blots transcripts are detected (Figure 4.2b). 3 Two of these, approximately 1.6 and 1.8kb, are not sex-specific and are weakly expressed in pupae, adults, agametic individuals and tra2 homozygotes. Both of these transcripts may be encoded on restriction fragments defined as non sex-specific the on reverse northern blots. The third transcript, of approximately 2.5kb, is male-specific. It is expressed in 3rd larvae (L3) (the lane is underloaded and not visible instar this exposure), pupae and adults. It is not detected in in (lanes & 11) or pseudomales agametic males 9 (lane 8) suggesting that the transcript is germline specific. However it is expressed normally in XO males and tra2/tra2 males.

pattern of expression, the chromosomal location and The the restriction map of gK14 are very similar to those reported by DiBenedetto et al. (1987) for mst(3)349. Their genomic clone also hybridises at 66D and identifies 1 malespecific and 2 non-specific transcripts. The male-specific transcript is expressed in the same stages and shows the same profile in mutants as the male-specific transcript from gK14. Expression of the non sex-specific transcripts of gK14 also appear to be similar to those reported for mst(3)349.

4.2ii qK21: This clone was localised by <u>in-situ</u> to polytene chromosome region hybridisation 98C/D on the right arm of chromosome 3 (Figure 4.1b). Restriction enzyme and a crude restriction map are presented in digests Figures 4.2c and 4.1c respectively. The reverse northern analysis identified a male-specific 5kb HindIII fragment and non-specific HindIII fragment of 4kb, these are marked on a genomic the restriction map. The apparent overlap with the clone mst(3)336 identified by DiBenedetto et al. (1987) is indicated on the diagram. Probing genomic DNA southern also blots with the whole <u>SalI</u> insert indicates that the sequences are single copy in the Drosophila genome (not shown).

Using the whole clone as a probe to northern blots identifies 2 male-specific transcripts of approximately 1.3 and 1.6kb (Figure 4.2d). The 1.3kb transcript is detected in

<u>Figure 4.2a</u>: EtBr stained 0.8% agarose gel containing restriction digests of Lambda gK14 DNA (lug/lane). Enzymes; R=<u>EcoRI</u>, B=<u>BamHI</u>, H=<u>HindIII</u>, s=<u>SalI</u>. Size markers are in kilobase pairs.

Northern blot analysis with various poly-A mRNAs Figure 4.2b: (5ug/lane). The probe used was lambda gK14, the blot was washed at high stringency and exposed for 3 days. The stripped blot was reprobed with actin probe Act-A2, washed at high stringency and overnight. Lanes: 1) Male L3, 2) Female L3, exposed 3) Male pupal, 4) Female pupal, 5) Male adult, 6) Female adult, 7) XY;<u>tra2/tra2</u>, 8) XX;<u>tra2/tra2</u>, 9) Agametic male (tud), 10) Agametic female (tud), 11) Agametic male (osk), 12) XO male, 13) XXY female. The size is in kilobases.

<u>Figure 4.2c</u>: EtBr stained 0.8% agarose gel containing restriction digested lambda gK21 DNA. Enzyme designations as for figure 4.2a. Sizes in kilobasepairs.

<u>Figure 4.2d</u>: Northern blot analysis using Lambda gK21 DNA as a probe. Lane designations and conditions as for figure 4.2b.



L3, pupae and adults, it is not expressed in agametic males or pseudomales. An identical pattern of expression is found by DiBenedetto et al. for a similar sized transcript from mst(3)336. The 1.6kb transcript is expressed strongly in male male pupae and male adults however weak expression L3, in females agametic males and pseudomales is also detected. DiBenedetto et al. also report a 1.6kb transcript but do not report weak expression in females or mutants. The reason for this difference is at present unclear, it may be due to different sensitivity of the northern blots. Alternatively it may be a transcript encoded on the restriction fragments identified as non sex-specific on reverse northerns blots. al. do not report the probe used for their DiBenedetto et northern blots so a comparison is not possible at present. In addition to the two transcripts described above a small approximately 0.5kb is also detected. It transcript of is found in male pupae and male adults but longer exposures of the autoradiogram indicate that it is also present in females although at lower abundance. Dibenedetto et al. report a non sex-specific 0.5kb transcript but they do not indicate elevated expression in males. Given the strength of the signal for the transcript on my northern blots it is difficult to believe that this is a transfer or hybridisation artefact. Although considering the odd shape of the band this in mind. Alternatively possibility should be borne the difference between my data and that of DiBenedetto et al. may reflect strain differences. As these authors do not show an autoradiogram it is difficult to assess whether or not these are meaningful. Subcloning to identify differences transcript-specific probes will be required to confirm these observations.

A higher molecular weight transcript of approximately 2kb is also detected on this blot but not on others, this may represent a processing intermediate or a transcript from the non-specific area of the clone.

## 4.3 DISCUSSION

The majority of the data presented in this chapter suggests that two of the male-specific genomic clones

identified in our screen correspond to genes previously characterised. By <u>in-situ</u> hybridisation the clones map to the same chromosome region. The restriction maps of the clones correspond fairly well. Northern blot analysis identify very similar, although not identical, transcripts and patterns of expression. While formal proof would require cross hybridisation between our clones and those of DiBenedetto et al. the data presented are sufficient to conclude that it is highly probable that they are the same genes

DiBenedetto et al. examined the expression of their genes by in-situ hybridisation to tissue sections. In this way they determined that expression of the male specific transcripts was restricted to the lumen of the testis. These experiments complement the conclusions of my northern blot experiments which indicated that expression was germ-line specific. I extend the data of DiBenedetto et al. by finding no effect on expression of the male-specific transcripts from removal of the Y chromosome. The isolation of previously characterised genes provides powerful internal controls for the behaviour of the mutant fly lines and for the reliability of the insitu hybridisation localisations. In this respect it should be noted that the similarity between our sequences and those of DiBenedetto et al. was not uncovered until I had completed northern blots, restriction maps and genomic localisation.

In the succeeding chapters I shall describe the characterisation of 6 novel male-specific genes.

#### CHAPTER 5

"你就是我们的你,你们们的你们的你,你们还能是你能能吗?""你们的你?"

# NOVEL GENES WITH PUTATIVE GERMLINE EXPRESSION

A statistical descent of the second second

#### 5.1 INTRODUCTION

Many genes appear to be involved in the process of spermatogenesis in Drosophila. This is evident from the large number of loci which can be mutated to give a male sterile phenotype. Lifschytz (1987) has suggested that these genes can be divided into two broad categories; Those with products specific to the male germline and those whose products are required in both the germline and in the soma. The latter class are believed to be male sterile because the process of is likely to be more sensitive than spermatogenesis general somatic metabolism to reductions or perturbations in the products of these genes. The rational behind this suggestion is that as the germline is the ultimate carrier of genetic information to the next generation strong selection for is fitness imposed. This hypothesis is supported by the finding that at least some male sterile mutations are allelic to organismal lethal mutations (Lifschytz, 1987).

a cellular differentiation process Spermatogenesis is which involves a complex series of morphological changes. Furthermore the majority of sperm differentiation occurs in the absence of any substantial ongoing transcription; the process relies on mRNA transcribed during the premeiotic stage (see chapter 1). This is not to say spermatocyte that during sperm there is no regulation of gene expression differentiation. On the contrary extensive translational operate to provide functional gene controls appear to products at the appropriate times in development. the Thus male isolation and investigation of genes expressed in the germline may facilitate an investigation of the regulation of tissue specific gene transcription and of the regulation of translation. The results of such studies on testismRNA specific B-tubulin and mst(3)gl9 have been discussed in chapter 1.

As to the function of molecules specifically involved in spermatogenesis we can envisage a requirement for at least two classes of genes.

Class 1: In the first instance structural proteins necessary formation of sperm-specific organelles (eq the for the This class of molecules is axoneme). nebenkern and specific B-Tubulin, major а testis exemplified by the of the axoneme. In addition this tubulin subunit component

been shown, by mutational analysis, to be required has for meiotic spindle formation (but not the mitotic spindle of gonial cells) and to form post-meiotic cytoplasmic microtubules (Kemphues et al., 1982). Whether or not the testis specificity of this tubulin subunit is due to structural differences with respect to other B-Tubulins or to unique sperm-specific regulatory sequences is at present unclear. Other structural molecules include the product of the testis-specific transcript mst(3)ql9 examined by Kuhn et al (1988). This transcript encodes a small 56 amino acid polypeptide composed almost exclusively of cys-gly-pro Mammalian sperm also contain cys-pro rich proteins repeats. which are believed to have elastic properties (Guraya, 1987). Nucleic acid probes from this gene detect a sperm-specific transcript in mice. The authors propose, by analogy with mammalian sperm, that the mst(3)gl9 polypeptide may be а structural component involved in sperm tail elasticity.

Hennig and co-workers (cited in Hennig, 1985) identified immunologically a 155kd sperm-specific polypeptide in <u>D.hydei</u> which appears to be a structural component of the nebenkern. Interestingly they find that this polypeptide is specific Y chromosome loop until it is localised to a incorporated into the nebenkern. In the absence of a Y chromosome the level of the polypeptide is considerably reduced. As the polypeptide is not believed to be Y encoded Y chromosome may be playing a regulatory role in the the translation or storage of this polypeptide. More recently Bonaccorsi et al (1988) have demonstrated similar properties a <u>D.melanogaster</u> sperm-specific antigen using the for same antisera.

for the presence in <u>Drosophila</u> of protamine Evidence as those arginine rich, sperm-specific histones such like, found in mammalian sperm is at present contradictory. Henniq (1985)has failed to detect such molecules in the sperm of D.hydei. On the other hand Das et al (1964) and Hauschteck-Jungen and Hartl (1982) have convincing evidence for these proteins being present at the time of chromosome condensation <u>D.melanogaster</u>. This apparent contradiction may reflect in technical difficulties in detecting these proteins in <u>D.hydei</u> That these differences reflect species variation sperm. appears less likely considering the similarity between these species in most other aspects of spermatogenesis.

Class 2: Another class of sperm-specific molecules that one may expect to find are those involved in the regulation sperm-specific processes. We can of imagine molecules involved in regulating tissue-specific gene expression at both the transcriptional and translational level. We may also find genes whose products direct the series of morphological changes necessary to produce a functional sperm. Although there are no firm examples of these types of molecule two interesting genes are worth mentioning.

The ref(2)P gene product is involved in sigma virus replication and is also necessary for male fertility (Contamine et al., 1989). The sterile phenotype in homozygous males, a failure to differentiate the nebenkern, results from The mitochondria a failure in mitochondrial aggregation. appear to degenerate instead of coalescing into the nebenkern (Dezelee et al., 1989). The gene has recently been cloned and shown to encode a polypeptide of 76kd which contains а mitochondrial pre-sequence and a zinc finger region (Dezelee allow et al., opp.cit.). The mitochondrial pre-sequence may polypeptide to associate with the outer membrane of the the mitochondrion and promote aggregation. The zinc finger motif could perform a regulatory role or may be involved in the storage of the polypeptide until it is required.

Leroy et al (1989) have recently isolated a murine spermspecific transcript (PL10) which encodes a polypeptide with similarity to eIF-4A. In mammals, as in Drosophila, there is considerable translational control of stored transcripts spermatogenesis (protamine mRNA is stored for more during than a week prior to translation. Peschon et al., 1987). The related to translation initiation finding that PL10is suggests that it may have a role in regulating factors the translation of stored transcripts. In cross-hybridisation experiments using a PL10 gene probe related sequences are detected in <u>Drosophila</u>. In addition there is extensive similarity with the <u>Drosophila</u> <u>vasa</u> gene product which is believed to regulate translation in the early embryo (Lasko and Ashburner, 1988). Whether or not related molecules are to found in the male germline of <u>Drosophila</u> remains be to be determined.

In the following chapter I will present the initial characterisation of 3 novel genes putatively expressed in the male germline. It is hoped that a structural and mutational

analysis of these genes may provide further insight into the molecular events underlying the complex process of spermatogenesis in <u>Drosophila</u>.

#### 5.2 RESULTS

5.2i Clone qS1: This clone shows rather weak hybridisation on reverse northern blots. Long exposures of these blots identify a 1.6kb SalI fragment that hybridises cDNA. Male-specific hybridisation to other only to male restriction digests identifies fragments attached to the lambda vector arms. A crude restriction map of the right hand end of the clone is shown in Figure 5.2d. The region of malespecific cDNA hybridisation is indicated. The 1.6kb SalI fragment was used to isolate a 1.0kb cDNA clone from the amplified male L3 cDNA library described in chapter 3. When the cDNA library was screened only 0.03% of the plaques gave a positive signal. This low frequency is consistent with the level of hybridisation on reverse northern low blots. The cDNA was subcloned as an EcoRI fragment into the plasmid vector pEMBL18 to give plasmid pCS1. To ensure that the CDNA related to the "male-specific" sequences within is the DNA was digested with a variety of genomic clone, qS1 restriction enzymes, electrophoresed, blotted and probed with pCS1 (Figure 5.2a). The 1.6kb <u>SalI</u> fragment and fragments attached to the right hand arm of the lambda vector are detected. Hybridisation to undigested DNA is also apparent in and <u>BamHI</u> digests. This pattern of hybridisation is EcoRI identical to the hybridisation observed on reverse northern blots. These data confirm the partial restriction map for the right hand end of the genomic clone but other sites toward the left hand end of the clone have not been ordered.

genomic clone was used as a probe to salivary gland The polytene chromosome spreads and hybridises uniquely to region chromosome 3 82D at the base of the right arm of (Figure 5.1a,b). When pCS1 is used to probe southern blots of male and female Or DNA and DNA from other Drosophila strains (Cs and m56i) single bands are detected (Figure 5.2b). The sizes of these bands are consistent with the restriction map of the genomic clone. In addition the blot confirms that the

<u>Figure 5.1</u>: Photomicrographs of <u>in-situ</u> hybridisation to polytene chromosome spreads with biotinylated lambda clones.

- a): Probe gS1 hybridising to the base of the right arm of chromosome 3.
- b): Close up view of the above spread showing hybridisation at region 82D.
- c): Probe gK33, the arrow points to the weak signal on the right arm of chromosome 3.
- d): Close up of a different spread hybridised with gK33 showing hybridisation to region 89B.


B

D

A





gene is present as a single copy in the <u>Drosophila</u> genome and demonstrates that the sequence is conserved in other strains.

A northern blot analysis (Figure 5.2c) using the pCS1 probe detects 2 transcripts of low abundance. A major malespecific transcript of approximately 3.0kb confirms the malespecific hybridisation detected on reverse northern blots. transcript is detected in male L3 (lane 4: barely This visible on this exposure), pupae, adults, tra2/tra2 males and males. The transcript is not detected in agametic males, XO pseudomales or developmental stages prior to L3. Although pattern of expression is consistent with expression this in the male germline the low abundance of the transcript may preclude detection in cases where its level is substantially reduced. This would be the case if it were expressed in the somatic portion of the gonad. In addition low expression during the pupal stage would not be expected from а qene sensitive nuclease spermatocytes. A more transcribed in protection assay will be required to assess expression in individuals. In-situ hybridisation to tissue these mutant sections will be required to definitively localise the tissue of expression. Despite these ambiguities Ι can safely required for conclude that the presence of a germline is wild-type levels of expression.

A minor 0.9kb transcript of lower abundance is detected in most, if not all, developmental stages. It is not sexspecific in adults (lanes 15 and 16) but may be elevated in males. The transcript is particularly pronounced in both embryos and male pupae (lanes 1 and 6). It appears to be reduced in agametic males and females (lanes 12-14) as well as pseudomales (lane 11). It is of particular interest to the elevated expression during embryogenesis and note metamorphosis, both stages where major differentiation events individuals occurring. Reduced expression in with are rudimentary gonads is suggestive of germline associated Again it will be of considerable interest expression. to determine the site of expression by <u>in-situ</u> hybridisation to tissue sections.

It is possible that these two transcripts are related by differential RNA processing or by the use of different transcription start-sites within the same gene. In some mRNA preparations a high molecular weight transcript of approximately 4kb is detected (arrow in Figure 5.2c lane 10),

<u>Figure 5.2a</u>: EtBr stained 0.8% agarose gel of restriction enzyme digested lambda gS1 DNA (lug/lane) and southern blot probed with pCS1. The blot was washed at high stringency and exposed for 3 hours. Enzymes: R=EcoRI, H=HindIII, B=BamHI, S=SaII. Size markers in kilobasepairs.

<u>Figure 5.2b</u>: Southern blot of restriction enzyme digested <u>Drosophila</u> DNA (2ug/lane) electrophoresed on 0.8% agarose gel and probed with pCS1. The blot was washed at high stringency and exposed for 2 days. Size markers in kilobasepairs. Lanes: 1) m56i;<u>EcoRI</u>, 2) Cs;<u>EcoRI</u>, 3) Female Or;<u>EcoRI</u>, 4) Female Or;<u>BamHI</u>, 5) Female Or;<u>HindIII</u>, 6) Male Or;<u>EcoRI</u>, 7) Male Or;<u>BamHI</u>, 8) Male Or;<u>HindIII</u>.

Figure 5.2c: Northern blot analysis with various poly-A mRNAs (5ug/lane). The probe was pCS1, the blots were washed at high stringency and exposed for 2 weeks. The stripped blots were reprobed with actin probe Act-A2, washed at high stringency and exposed overnight. Lanes: 1) Embryo, 2) L1, 3) L2, 4) Male L3, 5) Female L3, 6) Male pupal, 7) Female pupal, 8) Male adult, 9) Female adult, 10) XY;tra2/tra2, 11) XX;tra2/tra2, 12) Agametic male (tud), 13) Agametic female (tud), 14) Agametic male (osk), 15) Male adult, 16) Female adult, 17) XO male. The sizes are in kilobases.

<u>Figure 5.2d</u>: Restriction map of the right hand end of Lambda clone gS1. Enzymes: B=<u>BamHI</u>, H=<u>HindIII</u>, R=<u>EcoRI</u>, S=<u>SalI</u>. Sites in brackets have not been ordered. Black bar indicates male-specific hybridisation identified on reverse northern blots.



this may represent a processing intermediate.

5.211 Clone qS10: The reverse northern analysis of this clone detects two <u>SalI</u> fragments (1.8 and 6.5kb) and а HindIII fragment (2.6 kb). The 2.6kb HindIII fragment was used to isolate a 0.6kb cDNA clone from the amplified male larval library. This was sub-cloned into the plasmid vector pEMBL18 to give plasmid pCS10. Only 0.05% of the plaques plated gave a positive signal, this is rather lower than would have been predicted from the strong signal on the reverse northern blots (and from the intensity of signal on northern blots, see below). This may reflect poor growth of containing this insert during amplification phage or а structural feature of the transcript which is refractory to reverse transcription. When the cDNA is used to probe blots of restriction digested gS10 DNA the fragments identified on reverse northerns are detected (Figure 5.3a). A deduced restriction map is shown in Figure 5.3d. Notice that a small (0.5kb) HindIII restriction fragment interrupts the continuity of the region showing male-specific hybridisation. This fragment is not detected by the cDNA clone (not shown) by the reverse northern analysis. It may represent or an intron or a duplication of the gene; current data do not differentiation between these two allow possibilities. Further sub-cloning and restriction mapping will be required confirm the restriction map shown and determine the to transcriptional organisation.

pCS10 is used to probe Southern blots of different When genomic DNAs restriction fragment polymorphisms are detected (Figure 5.3b). A 3.1kb EcoRI fragment is detected in Or and m56i (lanes 1,2,5 and 6). This fragment contained DNA is within the genomic clone. Cs DNA does not contain this fragment but in its place there is a 4.8kb fragment (lanes 3 4). This fragment is also visible in female Or DNA and and over-spill from the adjacent lane may represent or contamination with other DNA, It is not detected in male DNA. A 6.2kb EcoRI fragment is present in Or and Cs DNA, 1.4kb of this fragment is attached to the right hand end of the vector the genomic clone. This fragment is not present in m56i in DNA, it is replaced by a fragment of approximately 10kb.

<u>Figure 5.3a</u>: EtBr stained 0.8% agarose gel of restriction digested gS10 DNA (lug/lane) and Southern blot probed with pCS10. The blot was washed at high stringency and exposed overnight. Enzymes: R=EcoRI, H=HindIII, B=BamHI, S=SalI. Size markers in kilobasepairs.

Figure 5.3b: Sothern blots of restriction enzyme digested Drosophila DNA electrophoresed in 0.8% agarose gels (2ug/lane). The blots were probed with pCS10, washed at high stringency and exposed for 3 days. Lanes: 1) Male Or; <u>EcoRI</u>, 2) Female Or; <u>EcoRI</u>, 3) Male Cs; <u>EcoRI</u>, 4) Female Cs; <u>EcoRI</u>, 5) Male m56i; <u>EcoRI</u>, 6) Female m56i; <u>EcoRI</u>, 7) Female Or; <u>HindIII</u>, 8) Male Or; <u>HindIII</u>. Size markers are in kilobasepairs.

Figure 5.3c: Northern blot analysis with various poly-A mRNAs (5ug/lane). The probe used was pCS10, the blot was washed at high stringency and exposed for 3 days. The stripped blot was reprobed with actin probe Act-A2, washed at high stringency and exposed overnight. Lanes: 1) XXY, 2) XO, 3) Agametic male (osk), 4) Agametic female (tud), 5) Agametic male (tud), 6) XX;tra2/tra2, 7) XY;tra2/tra2, 8) Female adult, 9) Male adult, 10) Female pupal, 11) Male pupal, 12) Female L3, 13) Male L3. Size in kilobases.

<u>Figure 5.3d</u>: Restriction map of the left hand end of the Lambda clone gS10. Enzymes: B=<u>BamHI</u>, H=<u>HindIII</u>, R=<u>EcoRI</u>, S=<u>SalI</u>. The black bars represent male-specific hybridisation identified on reverse northern blots.



In addition to these strain differences a band of 7kb is reproducibly detected in both male and female Or DNA. This fragment is not accounted for in the genomic clone. When HindIII digested Or DNA is probed only the two fragments 7 predicted from the genomic clone are detected (lanes and 8). Contamination of the genomic DNA with plasmid sequences is considered unlikely as the band is not detected with other probes or by pCS10 in the <u>HindIII</u> digests. The presence of unrelated sequences on the cDNA clone also seems unlikely as these additional fragments are not detected in digests with addition only a single transcript other enzymes. In is detected on northern blots when pCS10 is used as а probe. Partial digestion by EcoRI is improbable as the extra bands are the same intensity in male and female DNA. A duplication sequences to another genomic location cannot of cS10 be eliminated. However this would have to alter the EcoRI sites not the <u>HindIII</u> sites. Since we know that the but gene is polymorphic between strains one possibility is that the Or DNA is contaminated with DNA from another strain. Alternatively the Or stock may contain a sub-population with one of sites. a rearrangement at the EcoRI Further fly blots will be required to experiments with single investigate these possibilities.

When either the genomic clone or the cDNA clone are used to probe polytene chromosome preparations no signal can be Incorporation of biotin into the probe detected. was achieved with control clones. equivalent to that Cohybridisation with control clones showed that hybridisation and detection were operating as expected. Two possibilities may account for this; either gS10 sequences are located at а constriction and I failed to identify the signal or the under-replicated sequences are located in an or heterochromatic region. In-situ hybridisation with tritiated probes and hybridisation to mitotic larval neuroblast spreads may allow localisation of these sequences.

northern blot analysis was carried out using the pCS10 Α (Figure 5.3c). A single transcript of approximately probe is present in male L3, pupae and adults. Ιt 0.9kb is not females, pseudomales or agametic males. This in detected germline specific expression. The suggests level of the transcript does not appear to be reduced by the lack of a Y chromosome or by the tra2 mutation. On the contrary the level

<u>Figure 5.4a</u>: Autoradiogram of a southern blot of restriction enzyme digested gK33 DNA electrophoresed in a 0.8% agarose gel (lug/lane). The blot was probed with the subclone pGK33H, washed at high stringency and exposed overnight. Enzymes: R=EcoRI, H=HindIII, B=BamHI, S=SalI. Size markers in kilobasepairs.

Figure 5.4b: Southern blot of EcoRI digested Drosophila genomic DNA electrophoresed on 0.8% agarose gel (2ug/lane). The blot was probed with pGK33H, washed at high stringency and exposed for 2 days. Lanes: 1) Male Or, 2) Female Or, 3) Male Cs, 4) Female Cs, 5) Male m56i, 6) Female m56i. Sizes are in kilobasepairs.

<u>Figure 5.4c</u>: Northern blot analysis with various poly-A mRNAs (5ug/lane). The probe used was pGK33H, the blot was washed at high stringency and exposed for 3 days. The stripped blot was reprobed with actin probe Act-A2, washed at high stringency and exposed overnight. Lanes: 1) Embryo, 2) L1, 3) L2, 4) Male L3, 5) Female L3, 6) Male pupal, 7) Female pupal, 8) Male adult, 9) Female adult, 10) XY;<u>tra2/tra2</u>, 11) XX;<u>tra2/tra2</u>, 12) Agametic male (<u>tud</u>), 13) Agametic female (<u>tud</u>), 14) Agametic male (<u>osk</u>), 15) XO male, 16) XXY female. The size is in kilobases.

<u>Figure 5.4d</u>: A restriction map of the left hand end of the Lambda clone gK33. Enzymes: B=<u>BamHI</u>, H=<u>HindIII</u>, R=<u>EcoRI</u>, S=<u>SalI</u>. The black bar symbolises the extent of male-specific hybridisation detected on reverse northern blots and also the subclone pGK33H.



of expression appears to be elevated in XO males. Confirmation of this observation will require a more quantitative analysis than the one presented.

5.2iii Clone qK33: in-situ hybridisation to polytene chromosomes with the whole genomic clone shows unique hybridisation on the right arm of chromosome at 89B 3 (Figure 5.1c,d). The reverse northern analysis identifies а small 0.6kb <u>HindIII</u> fragment that contains all of the hybridising to male cDNA. This fragment sequences was subcloned into the pEMBL vector to give plasmid pGK33h. Labelled plasmid DNA was hybridised to blots of restriction digested DNA (Figure 5.4a) and a crude restriction map of qK33 the left hand end deduced (Figure 5.4d). pGK33h detects a single 2.5kb EcoRI fragment in genomic DNA digests. The fragment is conserved in the other strains examined (Figure 5.4b).

Northern blot analysis (Figure 5.4c) with the pGK33h probe detects a smear from 0.5 to 0.7kb, this is consistent with the entire transcribed region being contained within the fragment when a polyA tail taken 0.6kb <u>HindIII</u> is into account. The transcript(s) are detected in male L3, pupae (where it is particularly prominent), adults and at similar levels in XO males and homozygous tra2 males. The transcript not detected in stages prior to L3, in females is or in agametic males. (Faint hybridisation to mRNA from pseudomales observed on this autoradiogram is considered to be an artefact probably due to over-spill or contamination, it is detected on other blots.) The pattern of expression is not suggestive of germline specific transcription.

## 5.3 DISCUSSION

features of the male-specific transcripts Several described in both this chapter and the previous chapter (gK14 are consistent with expression in the and gK21) male The failure to detect the transcripts in agametic germline. that the transcripts are dependent on the males shows the male germline for expression. The lack of of presence

expression in pseudomales strengthens this conclusion (with caveats discussed in the introduction to the chapter 4 regarding the underdeveloped gonad). The localisation of gK14 gK21 male-specific transcripts to the germline and by DiBenedetto et al (1987) confirm this conclusion, at least for these two clones. The developmental profile of expression also supports germline expression of these sequences. Low levels of the male-specific transcripts are found in the L3 stage where only a few transcriptionally active spermatocytes are found. The transcript levels increase dramatically during stage and are present at high levels the pupal during the adult stage. In addition the male-specific transcripts of gS10 and gK33 appear to be elevated during the pupal stage, paralleling the rise in spermatocyte numbers reported by Garcia-Bellido (1964). Although the evidence for germline is good the possibility of somatic, expression germline dependent, expression cannot be excluded without in-situ **hybrid**isation data. In the case of the male-specific transcript identified by qS1 this will be particularly important as this transcript is present at low abundance. It not likely that reduced expression of this transcript is in underdeveloped gonads would have been detected by the northern blot analysis presented here. The low level of expression in the pupal stage suggests that this transcript may not be expressed in spermatocytes.

If these transcripts are involved in spermatogenesis then neither the lack of a Y chromosome nor the presence of the appear substantially to affect their tra2 mutation transcription. This observation is similar to studies on the mst(3)gl9 transcript by Schafer (1986b) and studies on spermspecific tubulin expression (Hulsebos and Brand cited by Hennig 1985). These studies indicate that in XO males tubulin levels are normal but protein levels are transcript of the dramatically reduced. The apparent elevation qS10 males will require male-specific transcript in XO confirmation by quantitative studies.

The genomic locations of gS1 and gK33 give no clues as to their function, no male sterile genes or genes expected to have male-specific expression have been mapped in the vicinity of the clones described here. gK33 maps close to the <u>Stubble</u> region where there are a variety of chromosomal aberrations that could be used to generate mutations at the

gK33 locus. Alternatively a novel mutagenesis screen developed in our laboratory (Kaiser and Goodwin, in press) may be used to generate transposon insertions at or near the male-specific genes (see chapter 7).

present the genomic location of gS10 is unknown. In At the first instance the ambiguities surrounding the additional ECORI fragments detected on genomic blots will have to be resolved. As mentioned single fly DNA blots may be the best way to investigate this question. The inability to detect the this clone on polytene chromosome spreads location of suggests that it may be located in heterochromatic DNA. As we shall see in the next chapter several heterochromatic regions been detected which are involved in spermatogenic have Initially the localisation of gS10 sequences may processes. achieved by hybridisation to mitotic spreads, where be heterochromatin is available for hybridisation. If gS10 is located in heterochromatic DNA we would expect to find moderately repetitive DNA in the genomic clone, this has yet to be determined. An alternative explanation for the failure detecting gS10 sequences is that they are within in а constriction or locally underreplicated region. Hybridisation to mitotic spreads will help to identify if this is the case.

Little can be said at present concerning the nature of the products encoded by these male-specific transcripts. The low molecular weight transcripts encoded by gK33 and gS10 are other transcripts expressed reminiscent of in the male germline. For example the transcripts of mst(3)gl9 (450nt) and the murine protamine genes (800nt: Peschon et al., 1987). heterogeneous size of the gK33 transcript is similar The to that observed for murine sperm-specific protamine genes where size heterogeneity has been shown to reflect translational regulation by transcript deadenylation (Braun et al 1989). Whether the same is true for gK33 remains to be determined. small size of this gene makes it an ideal candidate for The experiments to identify transcriptional and fusion qene translational regulatory sequences, such an approach has been successful with the mst(3)gl9 gene (Kuhn et al 1988).

Of the three genes described here gS1 looks to be the interesting as the low transcript abundance suggests most that is does not encode an abundant structural protein. It some interest to elucidate the relationship will be of the male-specific and the non-specific transcripts. between

The elevated expression of the non-specific transcript during stages undergoing major differentiation tempts me to speculate on a role in gene regulation or nucleic acid metabolism. A sequence analysis of the cDNA clone is underway in our laboratory. Perhaps this may provide a clue as to the function encoded by this transcript.

Magnetical 2000 - Control Control Control Control
Michiel Activity - Control Control Control Control
Approximation active Control Control Control Control
Approximation Control Control Control Control Control
Approximation Control Control Control Control Control Control
Approximation Control Control Control Control Control Control Control
Approximation Control C

### CHAPTER 6

GENES WITH UNUSUAL GENOMIC STRUCTURE

and the second second

n kalenda in solar de la colori Esteste o egiptere de la colori d

NAL OF THE EAST OF STATES SHOULD AND THE TOTAL RELATES THE LASSING STATES STATES WE THERE ADDING STATES STATES STREETED AND THE CONTRACTOR STATES STATES STATES CONTRACTOR STATES STATES STATES CONTRACTOR STATES STATES STATES STATES FROM STATES STATES STATES TO STATES STATES STATES STATES

#### 6.1 INTRODUCTION

Most heterochromatic DNA appears to be composed of verv highly repetitive satellite DNA and contain few conventional genetic loci. However embedded within the simple sequence DNA mutable genetic loci are detected. Furthermore some of these loci have effects on mutations at normal It is some of these "genes" which I spermatogenesis. shall discuss below.

The Y Chromosome: The Y chromosome of D.melanogaster accounts for around 12% of the haploid genome. Cytologically appears to be entirely heterochromatic. It exhibits it intense staining with dyes specific to this class of DNA and underreplicated in polytene nuclei (Gatti remains et al., 1976). Bridges (1916) demonstrated that the Y chromosome is devoid of essential genes but that it is necessary for male fertility. Subsequent studies in D.melanogaster indicate that least 6 separable loci on the Y there are at chromosome necessary for male fertility. Two of these are on the short arm, ks-1 and ks-2, and four are on the long arm, kl-1, kl-2, kl-3 and kl-5 (Kennison, 1981). Deletions encompassing any of these fertility factors result in characteristic defects in the spermatogenesis, either early or late depending on particular factor deleted (Hardy et al., 1981). In D.hvdei appear to be 16 Y-linked complementation groups there necessary for male fertility (Hennig, 1985). For many years the fertility factors remained somewhat the nature of enigmatic. However genetic, cytological and molecular studies are beginning to shed some light on these unusual genes.

Much of the early work focused on the Y chromosome of structures visible <u>D.hydei</u> due to the prominent in the of this species ( Hennig, 1985, for spermatocyte nuclei а comprehensive review). During the spermatocyte stage the Y chromosome forms five characteristic structures reminiscent of the lampbrush loops of amphibian oocytes. The loops have Nooses, Clubs, Tubular Ribbons, each been named Pseudonucleoli and Threads on the basis of their distinctive morphology. A detailed genetic and cytological analysis has the indicated that the loops correspond to at least some of mutable fertility factors and suggests that they constitute

genetically discrete entities (Hennig, 1985). More recently Bonaccorsi et al. (1988) have demonstrated the presence of three similar loops in the spermatocyte nuclei of <u>D.melanogaster</u> and correlated them with 3 of the fertility factors, kl-3, kl-5 and ks-1.

A limited molecular analysis has been initiated and some Y chromosome DNA has been isolated (Hennig, 1985; Lifschytz, 1987). Broadly speaking the Y DNA so far characterised can be divided into 4 classes; 1) DNA specific to the Y chromosome but not to a specific region of the Y. This class is exemplified by the Y<sub>T</sub>I, Y<sub>T</sub>II and Y<sub>T</sub>III DNA clones from D.hvdei (Trapitz et al., 1988; Wlaschek et al., 1988). These sequences are found to be highly repetitive and transcribed in several of the lampbrush loops on the Y. 2) Y-specific and loop-specific DNA such as the Y20N, Y18Cl and Y22Tr clones isolated from <u>D.hydei</u> by Haraven et al. (1986). These sequences appear to be restricted to particular a loop Again they are repetitive and transcribed. 3 and structure. 4) Y associated DNA which is found on the Y chromosome as well as at other genomic locations. Class 3 DNA is highly repetitive satellite-like DNA exemplified by the <u>D.hydei</u> MY3 clone (Voqt and Hennig, 1986). Class 4 sequences are moderately repetitive and appear to resemble retroviral like transposable elements (Huijser et al., 1988; Brand and Hennig, 1989). These sequences are transcribed predominantly, though not exclusively, in the male germline. It is not known if they have any functional role. In addition it would appear that individual copies of these transposon-like sequences have diverged as multiple copies are only detected by а particular probe when low stringency hybridisation conditions are used.

The repetitive sequences transcribed from the lampbrush loops appear to have no protein coding potential. In addition no cytoplasmic transcripts have been detected. It is believed in long that these sequences are organised and transcribed tandem arrays hundreds of kilobases in length (Lifschytz et It has been proposed that the transcribed al., 1983). lampbrush loops perform a structural role in spermatogenesis or are required to store or compartmentalise proteins and RNA 1985; for post-meiotic development (Hennig, required Lifschytz, 1987). Several antibodies have been isolated which recognise antigens specifically associated with particular

lampbrush loops (Kloetzel et al., 1981; Tischendorf et al., 1989; Hennig, 1985). Some antigens are associated with specific structural components of the sperm at later stages in spermatogenesis, for example the 155kd antigen associated with the nebenkern discussed in chapter 5 (Hennig, 1985).

is not clear at present is if the lampbrush loops What are the only manifestation of the fertility factors. In this respect it should be noted that three of the D.melanogaster fertility factors have no known associated structures 1988). However this may reflect (Bonaccorsi et al., the destruction of such structures during the fixation process to cytological examination. These authors prior also demonstrate that some deletions of kl-3 which result in sperm dysfunction have no discernible effect on the lampbrush loop associated with this factor. At present it is still а possibility that regions of the fertility factors produce conventional protein products. In this respect we should also that some Y-linked sterile mutations can be induced note by same frequency as X-linked male steriles EMS at the (Kennison, 1983). These data suggest that some Y-linked point mutations although this has yet to be steriles are proved. Point mutations would not be expected to disrupt the function of tandemly repeated lampbrush loop structural transcripts. An argument against the occurrence of extensive point mutations is the failure of most experimenters to find complementing mutations within fertility factors. One author reported 25 non-allelic complementation groups on the Y has after EMS treatment (Williamson, 1972). Although these data are controversial and have not been confirmed recall that 16 fertility genes have been localised on the D.hydei Y chromosome. One explanation for the high level of EMS induced male steriles on the Y chromosome is that it may be more susceptible to EMS induced rearrangements than the other chromosomes. In this respect several workers have noted that induced mutations are twice as frequent on the Y than X-rav on the X (Kennison, 1983, and references therein). If indeed there are point mutations in fertility factors it will be of considerable interest to ascertain if the mutations are in coding or regulatory sequences.

<u>X-Chromosome Heterochromatin</u>: Approximately 50% of the length of the X chromosome is heterochromatic (Xh). mitotic In the salivary gland polytene chromosomes the Xh is not polyteneised and is associated with the chromocenter (Schalet Lefevere, 1976). As with most other heterochromatic and regions Xh appears to contain few conventional genes. However some DNA rearrangements within this region have detrimental effects on normal spermatogenesis. Unfortunately the nature of the rearrangements and how they disrupt spermatogenesis is at present poorly understood. Furthermore, if a conventional gene necessary for normal spermatogenesis were located on both the X and the Y it is unlikely that conventional genetic analysis would detect such a locus.

Cooper (1964) and Yamamoto and Miklos (1977) have identified at least two regions in Xh which are required for X-Y bivalent formation at meiotic prophase. These regions are apparently distinct from the <u>bobbed</u> (<u>bb</u>) loci containing the ribosomal RNA cistrons. Deletions of either of these regions substantially increases non-disjunction at male, but not female, meiosis (bivalent formation at female meiosis is mediated by euchromatic sequences). The precise location of the pairing sites on the Y chromosome has not been reported but seem to lie in the proximal ends of each arm (Cooper, By examining rearrangements which move these 1964). sites around the X chromosome Yamamoto and Miklos (1977) concluded that they represented specific sequences embedded within homogeneous satellite DNA. The precise nature of these pairing sequences remains to be determined.

Evidence that other Xh regions are required for proper spermatogenesis comes from the observations of several workers on sterility associated with certain Xh deficiencies. In a comprehensive study Rhaman and Lindsley (1980) examined such deletions and found that male sterility could be ascribed to some aberrations. However they found that the deletions induced sterility only when covered by duplications of Xh, carried either on the Y or on an autosome. In addition the behaviour of any particular deficiency was dependent upon the covering duplication used; a deficiency would induce sterility with one duplication but not with another. То account for their complex data they proposed the presence of three Xh sites or factors; one proximal to and two distal to At present it is not clear what is the nature of these <u>bb</u>.

sites nor if they are equivalent to the pairing sites previously identified (this is unlikely as the the deletions cause sterility and not non-disjunction). Equally it is also unclear whether the deficiencies uncover coding regions or sites necessary for correct chromosome behaviour during spermatogenesis. The authors favour the latter explanation and postulate that they have uncovered sites required for precocious X chromosome inactivation.

other region which is found on both the X and Y One chromosomes is the Stellate (Ste) locus (Hardy et al., 1984). The allelic state of the X-linked <u>Ste</u> locus determines the of which form in the morphology crystals primary spermatocytes of XO males. In <u>Ste<sup>+</sup>/O</u> males the crystals are needle shaped whereas in Ste/O males the crystals have a star difference shaped morphology. The region responsible for the morphology was mapped to 12C-13A on the Х in crystal chromosome. A specific region proximal to, but not including, for kl-2 was found to be the Y chromosome site responsible the appearance of the crystals. This was dubbed the stellate sequences on the X and Y have been control region. The analysed the molecular level by Livak (1984). A lambda at genomic DNA clone DM2L1 was isolated by virtue of its homology with a transcript elevated in the testes of XO males relative to XY males. The DM2L1 clone was found to have an unusual structure. In the first instance it was found to be devoid of recognition sites for many common hexanucleotide recognising restriction enzymes. Further examination with which have tetranucleotide recognition sequences enzymes uncovered a repeat structure; both CfoI and HincII released a tandem repeat of 1250bp. The repeats were found to be present and the Y chromosomes of <u>D.melanoqaster</u> both the Х on sequences were found mainly on the Y chromosome of (related D.mauritiana and <u>D.simulans</u>). In addition it was found that Ste phenotype was correlated with the number of repeats the chromosome: thus <u>Ste</u> alleles have many copies on the Х (approximately 200) whereas <u>Ste<sup>+</sup></u> alleles have few copies. The homologous regions on the Y chromosome map to the stellate control region and seem to be present at a stable level of around 80 copies.

Hybrid selection experiments identify a polypeptide of approximately 17kd present in both XO and XY testes. The

nature of the polypeptide and its role, if any, in normal spermatogenesis is at present unknown. The high level of associated with XO testes led Livak to propose expression the expression of the X-linked copies were in some that way negatively regulated by the Y sequences. In this model the absence of the Y sequences results in the production of excess Ste gene product which precipitates to form the It is not known if this regulation crystals. is transcriptional, translational or at the level of product stability. The morphology of the crystals is believed to be a reflection of X-linked Ste gene dosage.

One other system affecting spermatogenesis has elements localised within heterochromatic DNA. The <u>Segregation</u> <u>Distorter</u> (<u>SD</u>) system is a meiotic drive phenomena which results in the enhanced recovery of one member of a pair of homologous chromosome in the progeny of "distorting" fathers. There are three major components of the system; <u>Segregation</u> <u>Distorter</u> (<u>Sd</u>) a dominant allele which acts in trans, <u>Responder</u> (<u>Rsp</u>) the site of <u>Sd</u> action and <u>Enhancer of Sd</u> (<u>E(Sd)</u>) a modifier which increases the strength of drive.

The phenotypic effect of the <u>SD</u> system is to cause degeneration of sperm bearing the <u>Rsp</u><sup>S</sup> (sensitive) allele when an <u>Sd</u> allele is present in the genome. It would appear that chromatin condensation is prevented in spermatids carrying a sensitive responder. It has been suggested that the transition from somatic to sperm-specific histones is disrupted (Hauschteck-Jungen and Hartl, 1982).

Sd has been mapped to region 37D on the left arm of chromosome 2 (Brittnacher and Ganetsky, 1983). Deletions to be without effect on otherwise wild type appear individuals. It is therefore unclear what role, if any, Sd in normal spermatogenesis. The locus has been plays translocated to different chromosomal locations where it is still effective in causing drive.

The <u>Rsp</u> locus is located within the heterochromatin of the right arm of chromosome 2, very close to the centromere. Broadly speaking <u>Rsp</u> has two allelic states; insensitive  $(\underline{Rsp}^{i})$  and sensitive  $(\underline{Rsp}^{S})$ . All <u>Sd</u> bearing chromosomes isolated from the wild carry <u>Rsp</u><sup>i</sup> and are not subject to distortion. <u>Rsp</u><sup>S</sup> bearing chromosomes are recovered at a greatly reduced frequency when present in trans with an <u>Sd</u>

bearing chromosome (Brittnacher and Ganetsky, 1989; Lyttle et al., 1986). The <u>Rsp</u> locus has been cloned and appears to be composed of tandem repeats of a 240bp <u>XbaI</u> fragment (Wu et al., 1988). <u>Rsp<sup>i</sup></u> alleles appear to have substantially fewer copies of the repeat than <u>Rsp<sup>S</sup></u> alleles. No coding potential is detected within the repeat, nor can any transcripts be detected using the repeat as a probe to northern blots. Thus it is considered that the repeats act as a binding site for a trans-acting factor (Wu et al. opp.cit.).

E(Sd) is also located in a heterochromatic region; in the distal end of the heterochromatin of 2L (Brittnacher and Ganetsky, 1984; Sharp et al., 1985). E(Sd) has been defined by deletion and translocation analysis. It acts in trans to increase the strength of drive and has also been reported to exhibit weak drive on its own (ie. in the absence of Sd) (Sharp et al., 1985). The heterochromatin of chromosome 2 has extensively mutagenised by Hilliker (1976). In been EMS mutagenesis screens he recovered several lethals and the previously characterised eye colour mutation <u>light</u> within 2L heterochromatin. It would appear that none of these are allelic to <u>E(Sd)</u> (T.W. Lyttle, pers.com.). The apparent failure to find EMS induced mutations suggests that the locus may have an unconventional structure. It will be of interest to discover if <u>E(Sd)</u> is a conventional protein coding gene, if it produces a trans-acting RNA or if it has a trans-acting structural role.

In the following results section I shall present the preliminary characterisation of three genomic clones which encode male-specific transcripts. These clones have an unusual genomic structure, one appears to be heterochromatic and the other two are located primarily on the Y chromosome.

#### 6.2 RESULTS

<u>6.2i</u> Clone gS4. gS4 was localised, by <u>in-situ</u> hybridisation to polytene chromosomes, to the proximal end of section 40 on the left arm of chromosome 2 (Figure 6.1). As this region is associated with the chromocenter precise localisation has proved to be difficult. It appears however that this genomic clone contains DNA which only hybridises to

Α

restrictles, mets

В



figure 6.1: Photomicrographs of <u>in-situ</u> hybridisation to polytene chromosome spreads with biotinylated lambda gS4.

- a): Hybridisation to the base of 2L. There is a slight signal at the chromocenter, this is not observed in most preparations.
- b): Hybridisation to a different spread showing the signal restricted to 2L at region 40.

2L. Stretched chromosome preparations show hybridisation only to 2L and not to general heterochromatin or the chromocenter (not shown).

The molecular structure of the cloned DNA has some It was apparent that most unusual features. common enzymes which cut at hexanucleotide restriction recognition sites do not cut gS4. Thus there are no sites for BamHI, <u>HindIII, SalI, PstI</u> or <u>KpnI</u> within the qS4 insert. EcoRI, When digested with <u>Smal</u> or <u>Xhol</u> however a prominent band of 1.3kb is released, the intensity of the band suggests that several copies of a 1.3kb sequence are present in the insert of gS4 (Figure 6.2b). The simplest interpretation is that these sequences are arranged as a tandem array. In addition the two enzymes also release a fragment of 1.8kb. This may represent a diverged repeat (with the insertion of a 0.5kb fragment for example) or gS4 may contain sequences from the of the genomic repeat. The latter suggestion end is considered less likely because the 1.8kb fragment is produced by both enzymes. The size of the whole insert is estimated to be approximately 14kb which suggests there are approximately 8 copies of the 1.3kb fragment within gS4. At present the exact order of the repeats cannot be predicted as the position of the 1.8kb fragment is not known. A simple restriction map of the ordered site is shown in Figure 6.2d.

Additional repeat like structures are identified when the whole lambda clone is used as a probe to southern blots of These are shown in Figure 6.2a and genomic DNA. can be summarised as follows: With EcoRI digested DNA (lanes 1 - 4) most hybridisation is to DNA of greater than 20kb although lower molecular weight fragments are detected with long exposures. Some of these smaller fragments are conserved in other strains examined. XhoI (lanes 5-10) digestion the reveals the major 1.3kb band which is present in the genomic clone. In addition a ladder of bands increasing in molecular weight is also detected. Most of these bands are not multiples of 1.3kb and may represent diverged repeats. Α similar, if not identical, pattern is obtained by <u>Smal</u> digestion (not shown). Interestingly many of the bands are conserved in the 3 strains examined. Minor differences between male and female DNA are not believed to be significant and may reflect a degree of partial digestion (see below). Analysis of genomic DNA with enzymes which cut

<u>Figure 6.2a</u>: Southern blots of restriction enzyme digested <u>Drosophila</u> DNA (2ug/lane) probed with Lambda clone gS4. All the blots were washed at high stringency and exposed for 12 hours. <u>Lanes 1-4</u> : <u>EcoRI</u> digested DNA, 1) m56i, 2) Cs, 3) Male Or, 4) Female Or. 0.8% agarose gel, size is in kilobasepairs. <u>Lanes 5-10</u> : <u>XhoI</u> digested DNA, 5) Male Or, 6) Female Or, 7) Male Cs, 8) Female Cs, 9) Male m56i, 10) Female m56i. 0.8% agarose gel, sizes in kilobasepairs. <u>Lanes 11-14</u> : <u>CfoI</u> digested DNA, 11) m56i, 12) Cs, 13) Male Or, 14) Female Or, <u>Lanes 15-19</u> : Or DNA digested with; 15) <u>HpaII</u>, 16) <u>HaeII</u>, 17) <u>HincII</u>, 18) <u>ThaI</u>, 19) <u>Sau3A</u>. Lanes 11-19 1.2% agarose gel, size markers in basepairs.

<u>Figure 6.2b</u>: EtBr stained 0.8% agarose gel of restriction digested lambda clone gS4 (lug/lane). The enzymes used were, H=<u>HindIII</u>, Sa=<u>SalI</u>, Sm=<u>SmaI</u> X=<u>XhoI</u>. The sizes are in kilobasepairs.

Northern blot analysis with various poly-A mRNAs Figure 6.2c: (5ug/lane). The probe used was lambda gS4, the blot was washed at high stringency and exposed for the indicated times. The stripped blot was reprobed with actin probe Act-A2, washed at high stringency exposed overnight. and Lanes: 1) XXY, 2) XO, 3) Agametic male (<u>osk</u>), 4) Agametic female (tud), 5) Agametic male (<u>tud</u>), 6) XX;<u>tra2/tra2</u>, 7) XY;<u>tra2/tra2</u>, 8) Female adult, 9) Male adult, 10) Female pupal, 11) Male pupal, 7) XY:tra2/tra2, 12) Female L3, 13) Male L3, 14) L2, 15) L1, 16) Embryo. Sizes are in kilobases.

<u>Figure 6.2d</u>: A simple restriction map of the orderd sites in the Lambda clone gS4. The extent of the unmapped repeat is indicated below the diagram. Enzymes; Sa=SalI, Sm=SmaI, Xb=XbaI, Xh=XhoI.



at tetranucleotide recognition sequences reveals additional repeat-like structure (lanes 11-19). Hybridisation to <u>ThaI</u> digested Or DNA (lane 18) is almost exclusively to a 500bp fragment. Faint minor bands are detected at 1.0 and 1.1kb, again these may represent diverged repeats, the end of the repeats, or partial digestion. Hybridisation to HpaII and HincII digested Or DNA (lanes 15,17) reveals major fragments 1.1kb as well as additional bands of lower intensity. of Sau3A releases 3 strongly hybridising fragments of 380, 470 1000bp (lane 19). In contrast <u>CfoI</u> and <u>HaeII</u> do and not appear to recognise repeats. The <u>CfoI</u> digests (lanes 11 - 14) suggest that there is no difference between male and female Or DNA with this enzyme. The pattern of restriction fragments with this enzyme is different between the strains examined. The Cs and m56i strains have not yet been examined with the enzymes which recognise the repeats.

It would appear then that the lambda clone gS4 is composed of <u>SmaI</u>, <u>XhoI</u>, <u>HpaI</u>, <u>HincII</u> and <u>Sau3A</u> repeats with a major 500bp <u>ThaI</u> repeat. A more detailed analysis of the structure of this clone is under-way at present in our laboratory. The pattern of restriction enzyme digestion strongly suggests that gS4 represents heterochromatic DNA.

The transcripts recognised by the gS4 sequences also show some interesting properties. Male-specific RNAs are detected when the whole clone is used to probe northern blots (Figure 6.2c). The transcripts migrate as a smear from 400-700nt. If a polyA tail is taken into account this size range is compatible with the transcripts emanating from a single ThaI repeat. Alternatively the DNA homologous to the transcripts may be located at one end of the lambda clone and be represented as a unique sequence (ie one of the faint ThaI bands or the 1.8kb Sall band). The transcripts are detected embryos, male L3, male pupae and male adults. They are in also found in <u>tra2/tra2</u> males (lane 7) and XO males (lane 2) these individuals the level appears to but in be substantially reduced (in the 1 week exposure the XO lane is obscured by background however it appears reduced in the overnight exposure). No expression is detected in agametic males, pseudomales or females. This pattern of expression is consistent with male germline specific transcription.

6.211 Clones qS8 and qS9: As mentioned in chapter 3 these genomic clones share related sequences although they do not appear to overlap. The reverse northern analysis detects restriction fragments which hybridise predominantly with male within both of these clones (see Figure 6.4b). CDNA qS8 contains a 7.0kb SalI fragment and a 5.5kb HindIII fragment. These fragments hybridise weakly, but noticeably, with female cDNA. gS9 contains 11.0kb SalI, 4.1kb BamHI and 3.2kb HindIII fragments. In this case hybridisation with female CDNA is barely detectable. Experiments to identify the sequences within these clones which hybridise with female cDNA have yet to be performed. The data presented below suggest that they are independent of male-specific sequences.

The 3.2kb HindIII fragment from gS9 and the 5.5kb HindIII fragment from gS8 were used to isolate cDNA clones from the amplified male L3 cDNA library. The gS9 cDNA fragment isolated was 650bp long and when cloned into the pUC19 vector gave plasmid pCS9. 0.4% of the plated cDNA library plaques were positive in the screen. The gS8 cDNA isolated was 500bp and was cloned into the pEMBL vector to give pCS8. long In this case only 0.2% of the plated plaques were positive. Given the homology between these clones this difference is unexpected but may be a consequence of the library screens carried out at different times. To investigate the being extent of the sequences shared by qS8 and qS9 Southern blots from restriction enzyme digests of the genomic phage clones were probed with the cDNA clones. An example is shown in Figure 6.3a. Blots of qS8 and gS9 DNA were probed with pCS9 and washed at high stringency. For both of the genomic clones hybridising fragments are the ones identified by the the reverse northern experiments. An identical pattern of hybridisation was detected with the pCS8 probe (not shown). experiments allowed the construction of the crude Similar maps shown in Figure 6.4b. Thus I conclude that restriction these non-overlapping lambda clones contain the same or very similar "male-specific" sequences.

investigate the genomic organisation of these То sequences southern blots of male and female DNA were probed with the pCS9 probe (Figure 6.4a). The majority of the is restricted to male DNA indicating that hybridisation the on the Y chromosome. In addition it is reside sequences apparent that several copies of pCS9 sequences are present on

<u>Figure 6.3a</u>: EtBr stained gel restriction digested DNA from Lambda clones gS8 and gS9 (lug/lane) and a Southern blot probed with pCS9. The blot was washed at high stringency and exposed for 2 hours. Enzymes: E=EcoRI, B=BamHI, H=HindIII, S=SaII. The size markes are in kilobasepairs.

<u>Figure 6.3b</u>: Northern blot analysis with various poly-A mRNAs (5ug/lane). The probe used was pCS9, the blots were washed at high stringency and exposed for 3 days. The stripped blots were reprobed with actin probe Act-A2, washed at high stringency and exposed overnight. Lanes: 1) Embryo, 2) Male L3, 3) Female L3, 4) Male pupal, 5) Female pupal, 6) Male adult, 7) Female adult, 8) XO, 9) XX;tra2/tra2, 10) XY;tra2/tra2, 12) Agametic female (tud), 13) Agametic male (tud). The size is in kilobases.





the Y. Female DNA also hybridises with the pCS9 probe though this case the hybridisation is to a major band in in each some blots weaker bands are detected in digest. In female significance of this is at present unclear. The DNA, the bands in female DNA are unlikely to result from contamination not male DNA as the intensity of the hybridisation is with consistent with this explanation. From these digests it would appear that the qS9 clone originates from the Y chromosome. The presence of a 0.5kb EcoRI fragment in this clone which is present only in male DNA is consistent with this. The predicted restriction map of gS8 correlates with the major bands detected in female DNA. In particular a 5.5kb HindIII fragment, a 4.9kb EcoRI fragment and a high molecular weight BamHI fragment are detected. Experiments to prove that gS8 is not Y-linked are underway. Blots of male and female DNA from Cs and m56i (lanes 7-12) indicate that this genomic Or, organisation is very similar in the three strains.

The 3.2kb <u>HindIII</u> fragment from the gS9 genomic clone was also used to probe similar genomic southern blots (lanes 13-Hybridisation specific to the Y chromosome is still 16). detected however far greater similarity between male and female DNA is apparent. The number of bands detected with the sub-clone is consistent with the presence of genomic moderately repetitive DNA on this genomic fragment. When this experiment is extended using the whole gS9 genomic clone as a probe an even greater number of common sequences are detected (not shown). Again this is consistent with the presence of moderately repetitive DNA in the genomic clone.

Neither the genomic clones or the cDNA clones give а detectable signal when used as probes to polytene chromosome spreads. Co-hybridisations with control clones indicate that and detection conditions hybridisation were not the unfavourable. The lack of hybridisation to Y chromosome sites Y chromosome is surprising considering the is not underreplicated in salivary glands. In the case of the non Y sequences this suggests that they may be underreplicated or heterochromatic.

**Transcr**iptional analysis using the pCS9 clone a male-specific transcript of (Figure 6.3b) detects approximately 1.4kb. The transcript is present in male ьз, while the level appears unusually high on this blot, lower levels of transcript are detected at this stage on other

Figure 6.4a: Southern blot analysis of Drosophila genomic DNA (2ug/lane). Blots were probed with pCS9 (Lanes 1-12) or the 3.2Kb HindIII fragment from gS9 marked on the restriction map below (lanes 13-16). Electrophoresis was carried out in 0.8% agarose gels. Blots were washed at high stringency and exposed for 3 days. Lanes: 1) Female Or; <u>EcoRI</u>, 2) Female Or; <u>BamHI</u>, 3) Female Or;<u>HindIII</u>, 4) Male Or; EcoRI, 5) Male Or; BamHI, 6) Male Or; HindIII. 7-16 all <u>EcoRI</u> digests; 7) Male Or, Lanes 8) Female Or, 9) Male Cs, 10) Female Cs, 11) Male m56i, 12) Female m56i, 13) Male Or, 14) Female Or, 15) Male Cs, 16) Female Cs. Size markers are in kilobasepairs.

<u>Figure 6.4b</u>: Restriction maps of gS9 and the right hand end of gS8 Lambda clones. The black bar represents male-specific hybridisation identified by reverse northern experiments. The black bar under gS9 also identifies the 3.2Kb <u>HindIII</u> fragment used as a probe. Enzymes E=EcoRI, H=HindIII, B=BamHI, S=SaII.



<u>6.4</u>b



<u>,1Kb</u>,

 $--- \underbrace{ \begin{vmatrix} HS & B & B & H & H & S \\ \hline HS & I & I & I & I \\ \hline I & I & I & I & I \\ E & & E & E \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I & I & I \\ \hline HS & I \\ \hline$ 

blots. A possible explanation for this is that in previous experiments hand-sexed wild-type larvae were employed whereas in this case the material was generated from <u>SD</u> stocks. This will require further characterisation. The transcript is also male-specific in pupae, adults and tra2/tra2 males. There is expression in females, pseudomales or agametic males. no In XO males the level of the transcript appears to be reduced, however it is clearly not abolished (lane 8). This indicates that the non-Y location is capable of being transcribed, at least in the absence of the Y. It does not indicate whether or not the Y linked sequences are transcribed. The transcript is not detected in females carrying a Y chromosome (genotype XXY, not shown). This pattern of expression is again consistent with transcription specific to the male germline.

# 6.3 DISCUSSION

structure of the gS4 clone is reminiscent of The heterochromatic DNA in that several different repeats are apparent when genomic DNA is probed with this clone. At present we are cloning and mapping the <u>Smal</u> and <u>Thal</u> repeats try and develop a more detailed picture of the structure to of this region. We also need to establish whether or not the repeats are the site of transcription for the male-specific transcripts. At present this seems be the most likely possibility. However transcribed sequences at the ends of the clone, or within the 1.8kb <u>SmaI</u> fragment, cannot lambda be excluded. It will also be of considerable interest to determine the nature of the transcripts, in particular whether they perform a structural or a protein coding role. Experiments to address this question are in progress by cloning and sequencing cDNA copies of the transcript.

As one of the major criteria for isolating this sequence was oligo-dT primed reverse transcription it seems reasonable to assume that the transcript is polyadenylated and hence most likely to be protein coding. It should be noted however that heterochromatic DNA is renowned for its high A-T content so that the presence of an A-tract within the transcript cannot be excluded. However the restriction enzymes which cut the repeats detected by gS4 all recognise G-C rich sequences,

so the presence of long stretches of A residues is considered to be less likely.

gS4 male-specific The expression pattern of the transcript argues strongly for germline specific expression. Particularly compelling is the apparent reduction of levels in XO and tra2/tra2 individuals the transcript and absence of expression in agametic males. Interestingly the phenotypes of both XO and tra2 defects on spermatogenesis can considered as chromosomal phenomena (tra2 mutations have be resembling defects in X chromosome inactivation). effects This suggests that the gS4 transcripts are sensitive to perturbations in chromosome behaviour during spermatogenesis. This may be further tested by examining transcript levels in T(X:A) individuals.

some respects the structure of the gS4 clone is In reminiscent of the <u>Stellate</u> locus. It too is composed of tandem repeats and produces a low molecular weight germline specific transcript. However the effect of the Y chromosome on the transcription of gS4 and <u>Ste</u> appears to be completely opposite. Ste expression is elevated in XO males whereas qS4 transcripts appear to be reduced. The significance of the similarity between these two loci cannot be assessed until information on gS4 is obtained, but one is led to further wonder if a repetitive structure will be found to be a more general feature of genes expressed in the male germline.

localisation of qS4 sequences to the base of 2L The is intriguing and it will be important to characterise this further using mitotic metaphase spreads. In particular we need to determine if the sequences are within а heterochromatic or a euchromatic portion of the chromosome. fortuitously the euchromatic/heterochromatic Somewhat in 2L appears to be marked by a constriction in boundary mitotic spreads. In the absence of further data it is <u>E(Sd)</u>. This tempting to speculate that gS4 may represent locus maps at the distal end of 2L heterochromatin and appears to produce a trans-acting factor involved in chromosome behaviour during spermatogenesis. at However E(Sd) has only been associated with aberrant present spermatogenesis. It is not clear if <u>E(Sd)</u> has any role in spermatogenesis. Recall also that the Rsp locus is normal heterochromatic and repetitive. A repetitive nature for E(Sd)could explain the apparent lack of EMS induced mutations. Ι

intend to examine this possibility in collaboration with T.W. Lyttle, who is at present constructing appropriate E(Sd) deletions and rearrangements.

further aspect of gS4 is of interest, namely One the presence of transcripts in the embryo. I intend to investigate this expression in sexed embryos generated from discussed in chapter 2. I also wish the <u>SD</u> stocks to establish when by using staged collections during embryogenesis gS4 is transcribed. It is possible, although unprecedented, that the gene is not transcribed in the embryo but that the transcript is delivered by the sperm !).

genomic organisation of the gS8/gS9 sequences is of The considerable interest. As discussed in the introduction to this chapter other workers have identified Y associated DNA It would appear that the sequences recognised sequences. by the pCS9 cDNA clone are not related to any class of previously described Y associated sequences. The pCS9 sequences are not highly repetitive and furthermore they are as a discrete, apparently poly-adenylated, transcribed transcript. It is also unlikely that pCS9 sequences are related to the Y-associated retrotransposon elements isolated from <u>D.hydei</u>. The conservation of restriction pattern and Y chromosome location between the strains examined suggest that represent gS9 male-specific transcript sequences do not transposable elements, since one would expect to find transposons at a variety of chromosomal locations and at different sites in different strains. A sequence analysis of the cDNA clones is at present under-way to try and determine the nature of the gS9 gene product.

The genomic sequences flanking the gS9 transcription unit are moderately repetitive which appears to be typical of heterochromatic Y chromosome DNA. As to the location of the autosomal homologues I presume at present that they are also heterochromatic. Localisation on mitotic spreads will be required for two reasons; In the first instance to determine the autosomal location and in the second to examine the organisation on the Y chromosome. In particular I wish to determine if the sequences are clustered, perhaps at а be particular fertility factor. This question may also employing the set of Y chromosome deletions approached by generated by Kennison (1983).

The location of gS9 sequences on the Y and autosomal (or
X-linked) sites is again reminiscent of the <u>Ste</u> locus. Digestion of the gS9 genomic clone with restriction enzymes which release the <u>Ste</u> repeats fails to identify such repeats in gS9. Furthermore in contrast to the <u>Ste</u> transcript the gS9 transcript is not elevated in the absence of the Y chromosome.

Investigation of the transcription of gS9 sequences poses some difficult problems. We know that the autosomal locus can be transcribed in the absence of the Y, but whether or not it is expressed in the presence of the Y is rather more If the gS8 clone does indeed difficult to investigate. represent the autosomal location then we hope to find flanking sequences unique to this location. If this is the case we it may be possible to disrupt this location by transposon mutagenesis (see chapter 7). If gS8 does not represent the autosomal location this can be isolated by screening a female genomic library with the cDNA clones. This has already been carried out however the 12 positive clones isolated have not yet been analysed. Even if this is possible we may still learn little about the pattern of transcription. may be to employ duplications Another approach and deficiencies for both Y and autosomal sequences and analyse transcript levels in such individuals. Of course we must remember that it is still not clear whether or not we are dealing with a single autosomal site. Extensive southern blot analysis with calibrated blots will be required to address this question.

One final point of interest will be to examine other members of the <u>melanogaster</u> subgroup as well as other <u>Drosophila</u> species for the presence of Y linked or autosomal copies of gS9 sequences. This may allow evolutionary aspects of Y chromosome structure to be investigated.

### CHAPTER 7

<mark>propressionae</mark>ns en les anno el El <mark>major de la</mark> general coj nombre

distis spit Anotini oso sociosodo o Additisto o

anternatura Aurin (18-

e et de la galeria de la casa de l

Carl Are

### A COMPLEX GENE?

人名德国法尔 化过滤器 医放下过去 法保养 日本 建

what there have note a contract the writing of the

a solo o sende andre se andre se solo o de antre solo o se antre se antre solo de la solo de la solo de la solo Antre solo de antre se se solo de la solo de l Antre solo de antre se sonadore de la solo de

In this chapter I shall describe the analysis of the two remaining genomic clones, gS2 and gK15. These clones encode a male-specific transcripts which number of appear to be with a complex way. Several Drosophila expressed in genes complex transcriptional patterns have been characterised. The majority of these are involved in regulating developmental pathways. As was described in chapter 1 the <u>Sxl</u> and <u>dsx</u> genes produce multiple developmentally regulated transcripts. Other complex genes include Antennapedia, Ultrabithorax, Achaete-<u>Scute</u> and <u>decapentapleqic</u> (For a review of complex genes see Scott, 1987). It has been proposed that complex genes reflect informational economy in the Drosophila genome; by producing with different functions and complex multiple products regulation may be required to "build а fly" fewer qenes In this respect the number of genes (Scott, 1987). in the genome has been estimated at around 5,000 Drosophila (by saturation mutagenesis of small regions, see Spradling and for a review) to 15,000 (by mRNA complexity Rubin (1981) measurements, Hough-Evans et al., 1979). Both these estimates seem rather low when one considers the complexity of the fly. structures will be uncovered for Perhaps complex an molecular increasing number of Drosophila genes as the cloning of the genome progresses.

discussed earlier (Chapter 3) the lambda clones qS2 As and gK14 are overlapping, I will present evidence that these genomic clones encode at least three male-specific transcripts and at least one non sex-specific transcript. In the male-specific transcripts is a good addition one of gene expressed sex-specifically in the candidate for a embryo.

### 7.2 RESULTS

<u>Clones qS2 and qK15</u>: The restriction fragments of qK15 that encode sex-specific transcripts, as identified by northern analysis, are contained within gS2. Hence reverse my discussion will be restricted the the majority of to latter clone.

Α







<u>Figure 7.1</u>: Photomicrographs of polytene chromosome preparations hybridised with a biotinylated lambda gS2 probe.

- a): A whole spread showing hybridisation toward the end of the left arm of chromosome 3.
- b): Close up of a different chromosome showing hybridisation at region 61F. Identical signals are observed with the gK15 probe (not shown).

Restriction enzyme digests of both clones are presented in Figure 7.2a. Common EcoRI, BamHI and HindIII fragments are detected. The preliminary restriction map of gS2 shown in Figure 7.2d deduced by probing Southern blots of was restriction digests with various sub-clones (not shown). The reverse northern analysis identified three BamHI restriction fragments (5.8, 1.6 and 0.8kb) which hybridise to male, but not female, cDNA. Two additional <u>BamHI</u> fragments of 0.4 and separate the "male-specific" <u>BamHI</u> 0.7kb fragments. At present it is unclear whether or not the 0.4kb BamHI fragment hybridises with cDNA. This small fragment would have migrated out of the gel in the reverse northern experiments. The 0.7kb fragment from qS2 was present on the gel and does not appear hybridise with cDNA from either sex. This fragment to may represent an intron or an intergenic space. The gK15 qenomic begins within this 0.7kb fragment. No male-specific clone hybridisation to the left hand end of the vector was detected in the reverse northern experiments. However in this case the reverse northern experiments were hampered by spurious hybridisation between female cDNA and vector sequences. Α more extensive restriction analysis of this region is required to confirm the restriction map. In addition the 0.4 o.7kb BamHI fragments will be used to probe northern and blots to clarify their relationship to the transcripts discussed below.

A 4.2kb <u>BamHI</u> fragment hybridises weakly with cDNA derived from both sexes. At present this fragment has not been investigated but will be used to probe northern blots in future experiments.

The three "male-specific" <u>BamHI</u> restriction fragments were sub-cloned into the pEMBL18 vector and are designated pS2/5.8B, pS2/1.6B and pS2/0.8B respectively. The pS2/5.8B and pS2/1.6B plasmids have been restriction mapped and this is summarised in Figure 7.2d.

Southern blots of <u>EcoRI</u> digested male and female DNA from three <u>D.melanogaster</u> strains were probed with the pS2/5.8B plasmid (Figure 7.2b). Male and female <u>Or</u> and <u>m56i</u> DNAs show identical patterns of bands. This is consistent with the restriction map of the cloned DNA with the exception of a small 200bp <u>EcoRI</u> fragment, which would have migrated out of the gel. This small fragment is present in the pS2/5.8B subclone. <u>Cs</u> DNA contains a restriction fragment polymorphism.

<u>Figure 7.2a</u>: EtBr stained 0.8% agarose gel containing restriction enzyme digests of the overlapping lambda clones gS2 and gK15 (0.5ug/lane). Enzymes R=<u>EcoRI</u>, B=<u>BamHI</u>, H=<u>HindIII</u>, S=<u>SalI</u>. The size markers are in kilobasepairs.

Figure 7.2b: Southern blot of EcoRI digested Drosophila genomic DNA electrophoresed in a 0.8% agarose gel (2ug/lane). The probe was the plasmid pS2/5.8B. The blot was washed at high stringency and exposed for 2 days. Lanes: 1) Male Or, 2) Female Or, 3) Male Cs, 4) Female Cs, 5) Male m56i, 6) Female m56i. Sizes are in kilobasepairs.

Figure 7.2c: Northern blot analysis with various poly-A mRNAs (5ug/lane). The probe used was pS2/5.8B, the blot was washed at high stringency and exposed for 2 weeks. The stripped blot was reprobed with actin probe Act-A2, washed at high stringency and exposed overnight. Lanes: 1) Embryo, 2) (1, 3) L2, 4) Male L3, 5) Female L3, 6) Male pupal, 7) Female pupal, 8) Male adult, 9) Female adult, 10) XY;tra2/tra2, 11) XX;tra2/tra2, 12) Agametic male (tud), 13) Agametic female (tud), 14) Agametic male (osk), 15) XO male, 16) XXY female. The sizes are in kilobases.

<u>Figure 7.2d</u>: Restriction maps of the lambda clone gS2 and subclones derived from it. Enzymes: B=BamHI, E=EcoRI, H=HindIII, P=PstI, S=SaII, SS=SstI, Xh=XhoI, Xb=XbaI. The large arrow indicates the overlap with Lambda gK15. The black bars indicate male-specific, and the hatched bar indicates non sex-specific hybridisation detected on reverse northern blots.









In place of the 1.0kb <u>EcoRI</u> fragment it contains a fragment of 1.2kb. This is consistent the absence of the <u>EcoRI</u> site which produces the 200bp fragment in the genomic clone, but has yet to be confirmed on high resolution gels.

When either of the two genomic clones is used to probe salivary gland polytene chromosome preparations, unique hybridisation to region 61F on 3L is detected (Figure 7.1). Interestingly a <u>doublesex</u> cognate has also been mapped to this region (see discussion)

Α transcriptional analysis using probes from gS2 subclones been initiated and a complex pattern of has transcripts is detected. Northern blots probed with the pS2/5.8B subclone identify three transcripts of approximately 5.0, 2.0 and 1.7kb (Figure 7.2c). The 2.0kb transcript does not appear to be sex-specific. It is detected in embryos, Г3 is not yet clear if it is sex-specific at this stage (it as the female L3 RNA, lane 5, is slightly underloaded), pupae of sexes, where its level is elevated (lanes 6-7), and both adults of both sexes (lanes 8-9). It also appears to be present in agametic and tra2/tra2 individuals of both sexes (lanes 10-14).

The 5.0kb transcript is expressed in male L3, reduced in and elevated again in male adults. It is not male pupae detected in females, pseudomales or agametic males and is reduced in XO males (lane 15). These data, in particular the reduced expression in XO males, suggest that the 5.0kb transcript is germline-specific.

1.7kb transcript is expressed in embryos, male LЗ, The male pupae, male adults and weakly, but detectably in agametic males (lanes 12 and 14). It may also be present at low levels in pseudomales (lane 11). It does not appear to be expressed in females at any stage. This pattern is consistent germline associated, but not germline specific, with expression.

Preliminary data from the other subclones (not shown) indicate that the pS2/1.6B probe detects the 5.0kb transcript discussed above and an additional male-specific transcript of This new transcript appears to be expressed in male 1.2kb. L3, pupae and adults. It is not detected in agametic males or This pattern of expression suggests germline pseudomales. specific expression. The pS2/0.8B probe also detects the 5.0kb transcript and the 1.2kb transcript. At present I am

investigating these genomic clones further with a view to establishing the organisation of these transcribed sequences.

## 7.3 DISCUSSION

The region of the genome represented by these clones appears to be transcriptionally complex. Particular points of interest are: The reduced expression of the 5.0kb 1) pupae. 2) The elevated levels of transcript the 2.0kb in transcript during the pupal stage as well as its expression 3) The expression of the 1.7kb male-specific in embryos. The reduced but detectable transcript in embryos. 4) expression of this transcript in agametic males and possibly pseudomales.

pattern of expression of the 5.0kb transcript The superficially resembles a germline-specific transcript. However reduced expression during the pupal stage would not be expected from a gene transcribed in this tissue. number of Biological evidence indicates that the (transcriptionally active) spermatocytes is elevated in the pupal stage compared with the larval or adult stages (Garcia-Bellido, 1964). This is strongly supported by an examination of other germline-specific transcripts (see chapters 4 and 5). During the pupal stage the first meiotic divisions occur the male germline. However this would not be expected to in result in the reduced expression of a germline-specific transcript. Furthermore the level of the 5.0kb transcript rises again during the adult stage where there are many meiotically active cells. If this transcript is expressed in the male germline its developmental profile is highly unusual. If it is not expressed in the germline we are faced some puzzling data. The lack of expression in agametic with individuals be explained by germline associated can expression but the reduction in expression in XO males is accounted for less readily. Marsh and Wieschaus (1978) have shown, by pole cell transplantation, that the Y chromosome is not required in the male soma for normal spermatogenesis. Hence the absence of the Y chromosome may not be expected to alter the expression of a somatically expressed gene. In this should recall the observations of respect however we

Steinmann-Zwicky et al. (1989) on germline-soma interactions. Although their experiments indicate that haplo-X germ cells will initiate spermatogenesis irrespective of the constitution of the soma they will not successfully complete or development unless they are in male soma (XY XO). sperm This suggests that a germline-soma interaction required is for correct spermatogenesis. The nature of this interaction is obscure at present but probably involves the somatic cells of gonad. It is possible that a germline autonomous the disruption in spermatogenesis (eg. the lack of a Y chromosome or the absence of a germline) could affect the expression of a somatic gene involved in the germline-soma interaction.

is tempting to speculate that the reduced expression Ιt the 5.0kb transcript is connected with the elevated of expression of the non-sex-specific 2.0kb transcript. The expression of the 2.0kb transcript is of some interest. In particular its elevated levels in the embryo as well as in pupae. A closer inspection of the expression in sexed larvae will be required to determine the status of this transcript at this stage. The function of this transcript is obscure at present, it does not appear to be associated with the germline as the levels in agametic and wild-type individuals similar. If the expression of these two transcripts is are it could reflect the use of alternative linked an transcription start site, differential polyadenylation or differential mRNA splicing. Transcript specific probes will be required to investigate this phenomena in northern blot and tissue section <u>in-situ</u> hybridisation experiments. Alternatively the expression of these transcripts may not be connected. Their close proximity on the 5.8kb BamHI fragment may reflect two very close but unrelated genes. This situation has been observed previously: with the tra gene for example (see chapter 1). Another possibility is that the 2.0kb transcript may be expressed from within an intron of the gene for the 5.0kb transcript. This situation has also been observed before: the sqs4 and piq1 genes are located al.,1987). intron of the <u>dunce</u> gene (Chen et within an However the <u>dunce</u> intron in question is approximately 79kb long. I have no evidence for a giant intron in the gene which transcript. encodes the 5.0kb However until the start-site is defined this transcriptional remains а possibility.

The reduced expression of the 1.7kb transcript in agametic males suggests germline associated expression. However in this case the detectable expression in germline deficient individuals indicates that if the gene is expressed in the germline then this is not its only site of expression. A leaky phenotype of the grandchildless mutations employed to generate agametic individuals is not considered likelv as other, more abundant, germline transcripts described in this work are not detected in the agametic males (Chapter 4). At present the 1.7kb transcript is a candidate for a gene expressed in the somatic component of the gonad. If this is the case we may reasonably hope to find that the embrvonic expression of this transcript is also male-specific. Ι am investigating this possibility using two approaches: Bv isolating mRNA from single sex embryos generated from the SDstocks described in chapter 2 and by in-situ hybridisation to sections. An alternative explanation for the reduced tissue expression in agametic males is that the gene is expressed at high levels in the male germline but at lower levels in male somatic tissue. The expression of the 1.7kb transcript in pseudomales requires clarification. The very weak signal northern blot is insufficient to firmly detected on the conclude that the transcript is expressed in these individuals. A more sensitive nuclease protection assay will required to assess the expression in pseudomales. be Expression in pseudomales argues for male somatic expression regulated by the sex determination hierarchy. Reduced expression in these individuals compared with wild-type males expression in an underdeveloped tissue such as the suggests qonad.

the 1.7kb transcript turns out not to be expressed in If pseudomales one possibility is that it represents а dosage compensation function. Dosage compensation in the embryo and germline is not mediated by the msl genes the male in (Cline,1984; Gergen and Wieschaus, 1988; Schupbach, 1985; Sanchez, 1986). However it should noted Bachiller and be is no evidence at present for dosage compensation in there the male germline. If we presume that dosage compensation is active in the germline and that the same mechanism is used to provide appropriate X-linked gene expression in the embryo the pattern of expression of the 1.7kb transcript may then reflect such a function: high levels of expression the in

embryo, low levels during the larval stage where there are only a few germline cells requiring dosage compensation and the <u>msl</u> genes have taken over somatic dosage compensation functions. Finally increasing expression as the germ cell population rises during the pupal and adult stages. Removal of the germline results in greatly reduced transcript levels. Residual levels of transcription may be expected by analogy with the <u>msl</u> genes. Male-specific dosage compensation is not only function of at least one <u>msl</u> gene. the The mle qene product is required in both sexes in a neurological role where it is found to be allelic to the <u>nap</u> mutation (B.Baker M. Kernan, R. Kreber, B. Ganetsky; abstract pers.comm; from 29th Annual Drosophila Research Conference.). Thus we might expect that genes required for embryonic and/or germline dosage compensation have other functions which are not sexspecific. This may also explain why genes involved in isolated embryonic dosage compensation have not been in genetic screens.

At present the relationship between the 1.7kb transcript and the 5.0 and 2.0kb transcripts discussed above is not They may be transcribed divergently from within known. the 5.8kb <u>BamHI</u> fragment or be splicing products of the same primary transcript. In order to investigate the complex transcriptional pattern of this small DNA fragment Ι am initiating several lines of experimentation. In the first instance the whole region will be sub-cloned and used to generate strand-specific probes to identify the transcripts on northern blots. Transcript specific probes identified in this way will be used to isolate cDNA each copies of transcript from appropriate cDNA libraries (In addition to the cDNA libraries described in this work I have also constructed libraries from embryo and adult body mRNA). Sequence analysis of the cDNAs and the genomic region will help achieve a more detailed picture of the transcriptional organisation. The sequence analysis may also provide some clues as to the function of these transcripts.

The location of a <u>dsx</u> cognate to the same region of the genome is of considerable interest. The <u>dsx</u> cognate sequences were isolated by low stringency hybridisation with a <u>dsx</u> gene probe (Belote et al 1985). The authors report that no sex-specific transcripts were detected when <u>dsx</u> cognate clones were used to probe northern blots but pupal specific

transcripts were detected. I have tried probing Southern blots of <u>Drosophila</u> DNA with the pS2/5.8B clone under the same conditions of stringency used to isolate the dsx cognates. No additional sequences have been detected. However gS2 clone also contains a restriction the fragment which hybridises with cDNA from both sexes contains and qK15 sequences further 3' to this. It will be of some interest to; a) Probe blots of gS2 and qK15 with a dsx probe and b) Probe Southern blots of Drosophila DNA with other restriction fragments from these two genomic clones. It is possible that the <u>dsx</u> cognate clone isolated by Belote et al. from this region contains only a portion of the gene. The dsx gene itself spans some 30kb of DNA, contains large introns and shows a complex pattern of transcription. At present we know that the 5.0kb transcript from gS2 spans at least 8kb of DNA and that the transcriptional pattern within this is region complex containing at least three other transcripts.

well as a molecular approach I intend to initiate a As mutational analysis of this region. This will be achieved by mobilising transposable elements and detecting insertions at or near the gS2 sequences. By using a novel method developed in our laboratory P-element insertions in cloned genes can be readily detected (Kaiser and Goodwin, in press). Briefly, insertions are detected using the polymerase chain reaction (PCR) with a primer unique to cloned sequences and another specific to the terminal repeats of the P-element. If а Pelement inserts at or near the gS2 locus this can easily be detected by the appearance of a novel PCR amplification product. Such is the sensitivity of the PCR reaction that one individual carrying the insertion can be detected in a several thousand flies which not. population of do Furthermore if the crosses are arranged such that a female is carrying the insertion in a heterozygous form a sib-selection scheme can be employed. With this method DNA is prepared from the embryos produced by a large population of mutagenised females (one of which hopefully carries an insertion in the gene of interest). The PCR reaction will detect the presence insertions in this population. The females of desired may be subdivided into smaller populations and the process then repeated. After a few rounds of selection the population will reduced to one or a few females carrying the insertion. be Progeny from these females may then be thoroughly examined

for phenotypic and molecular disruptions. Such a method of generating mutations circumvents many of the uncertainties of conventional genetic analyses. In particular the sibselection procedure is carried out using heterozygous females. Thus insertions may be obtained without biasing for phenotype. In addition this scheme also allows the recovery of silent insertions and insertions with very weak phenotypic effects which may go undetected in conventional mutagenesis strategies. Undoubtedly this method will have wide applications in the mutagenesis of cloned genes and hopefully some will provide phenotypic information on the function of of the genes described in this work.

e electron an anna ber escart an anna projec namesen (blive techiname) tone her te her hand her so of sectors a ad a set 了我<mark>爱你你想不是我们没</mark>我,没想了我们就在一般就想到了。""你们这个人,这个人们的,你们就 化铁煤料橡胶 计形式容许算机的复数形式 美国教教会会 网络中国人名 法公司的 计标识 e transfer - estal julier state de la companya de l **通過**過過過過過過過一個的人<mark>過過還</mark>一個的人。但是這一個的人,也不是有一個的人的人,這一個人的人類。這 for a marting to fourther the formal of interaction of the elerer "The character all cales there a alterated is the state there was any the structure of the space of the description of the second statements o **主要**做你要想到这些你的,我们就是你们这些了,我们不能不能能是这些人,我们能能 "高粱的烧糕,你有我们一定我们没有我的算法快乐,我们认为你的难见,没有一个多少。"他们 and a set of the set of

### CONCLUDING REMARKS

Ι initiated this project in order to isolate genes expressed only in one sex. In particular I am interested in sexually dimorphic gene expression in the embryo. This is primarily because the Drosophila embryo is a good source of reliable nuclear extracts for transcription, DNA binding and splicing studies (Biggen and Tjian, 1988; Rio, 1988). The availability of nuclear extracts may the permit identification and characterisation of regulatory molecules which act at specific promoter elements. In addition regulatory proteins, once identified in this way, can be isolated from cDNA expression libraries by virtue of their ability to recognise specific DNA sequences (Vinson et al., 1988). Α second reason for wishing to identify genes expressed sex-specifically in the embryo is to investigate the mechanisms which regulate germline sex. As I described in very little is known about germline chapter 1 sex determination. The isolation and characterisation of genes expressed early in the development of the germline may allow investigation of this problem using the methods outlined an above.

Whether or not the aim of the project has been successfully realised remains to be determined. Twenty lambda clones were isolated which appear to encode male-specific or Ten of these have been male-elevated transcripts. investigated in some detail. Two genomic clones, gS2 and gS4, encode transcripts expressed in embryos. These transcripts male-specific in the larval, pupal and adult stages. At are sexed embryos has not present the expression in been determined. If they are sex-specific in the embryo then I am in a position to initiate the lines of investigation outlined The clone qS2 discussed in chapter 7 is particularly above. interesting in view of the complex transcriptional pattern of northern blots. In addition one the observed on transcripts is detected in agametic males, indicating somatic expression. The data suggests that the expression of this transcript is germline associated. It is this clone which Ι intend to concentrate my efforts on at present. The lines of experimentation I intend to follow were discussed in chapter 7.

Some of the other genes which I characterised have interesting features which merit further investigation. The Y associated sequences detected by gS8 and gS9 appear to be unusual. No other Y chromosome DNA with this particular organisation has been reported: Multiple copies on the Y chromosome and probably a single copy elsewhere in the genome. It should be noted that low stringency hybridisations with the pCS9 cDNA probe does not detect additional sequences in female DNA (not shown). This suggests that these sequences do genetic not represent diverged remnants of mobile elements. The expression of the male-specific transcript appears to be restricted to the male germline. No expression detected in agametic males or females (even when the is females have a Y chromosome). In addition the expression in males is not dependent on the Y chromosome, although the levels of the transcript appear reduced in XO males. Experiments are at present underway in our laboratory to map the structure of these two genomic clones more carefully. It will also be important to try and identify the qenomic location of these sequences by hybridisation to mitotic chromosomes.

The repetitive structure of the qS4 clone is also unusual. As discussed in chapter 6 it would appear that this genomic clone consists of heterochromatic-like DNA. This is suggested by a number of different repeat-like structures recognised by different enzymes and by the localisation of sequences near the chromocenter. At present it is these not clear how the repeats are organised. No simple structure which accommodates the 1.3kb <u>Smal</u> and 0.5kb ThaI repeats springs to mind. This clone is being dissected and mapped in our laboratory at present.

The isolation of the gS4 and gS9 clones, which appear to contain repetitive DNA, justifies the decision to use а primary genomic library for these screens. An amplified library may well have been depleted of repetitive DNA due to recombination during propagation in E.coli. In this respect a recD E.coli strain (nm621) was used throughout this work to propagate recombinant phage. The <u>recD</u> mutants appear to be more tolerant of repetitive DNA that standard <u>recA</u> strains (Whittaker at al., 1988).

Of the remaining genes gK14, gK21, gS10 and gK33 appear to be genes transcribed in the male germline. There is good,

although not conclusive, evidence that gK14 and qK21 represent previously characterised male germline-specific genes. All four of the above clones identify transcripts with very similar patterns of expression: weak in male L3, very strong in male pupae and lower in male adults. This is а pattern of expression which fits well with the of counts spermatocyte numbers during development (Garcia-Bellido, 1965). In addition these transcripts are not detected in agametic males or chromosomal females transformed to somatic by the <u>tra2</u> mutation. These data maleness strongly support the belief that the genes are only expressed in the male germline.

expression of the male-specific transcript detected The by the gS1 clone is unusual. Although I included this gene in the "putative male germline" section (chapter 5) there are suggestions that it may not be expressed in this tissue. In particular the weak expression during the pupal stage as compared to the adult stage. The increasing expression from L3 to adults and the apparent lack of expression in agametic males leads me to suspect that this gene may be expressed in the gonadal soma. If this is the case then this gene may be best candidate for a gene expressed sex-specifically the in the embryo. Indeed if we consider that the male gonad in the embryo contains fewer than a hundred cells (Sonnenblick, 1950) we may not expect to detect a gene expressed in this tissue by northern blotting. This suggests to me that the embryonic expression observed for the gS2 and gs4 transcripts may not be sex-specific, as they are germline associated during later stages in development.

In general the +/- screening technique used in this work been successful in isolating genes expressed has sexspecifically. Most, although not all, of the genes described work appear to be expressed at fairly low levels this in during the larval stage (the stage used to prepare the CDNA probes). In order to isolate genes expressed at a much lower may be expected of abundance, as the transcripts for regulatory proteins, I am considering initiating a new screen. I intend to use the SD strains (chapter 2) to produce sexed embryo populations and generate subtracted probes to genomic libraries. In this way I hope to identify screen regulatory molecules expressed male-specifically in the early The reason for screening for male-specific genes is embryo.

that it is far easier to transform a female into a male, using somatic sex determination mutations, than perform the converse male to female transformation. The hope is that by characterising sex-specific molecules, in particular the way in which they interact with the known sex determination genes, a more thorough understanding of the control of sexspecific gene expression may be achieved.

# **BIBLIOGRAPHY**

and a start of the second s The second sec

Aboim, A.N. (1945). Developpement Embryonaire Et Post-Embryonaire Des Gonades Normales Et Agametiques De <u>Drosophila</u> Melanogaster. Rev.Suisse Zool. 52: 53-154.

- Amrein, H., Gorman, M. and Nothiger, R. (1988). The Sex Determining Gene <u>transformer-2</u> Of <u>Drosophila</u> Encodes A Putative RNA Binding Protein. Cell 55: 1025-1035.
- Aviv, H. and Leder, P. (1972). Purification Of Biologically Active Globin mRNA By Chromatography On Oligothymidylic Acid-Cellulose. PNAS 69: 1408-1412.
- Bachiller, D. and Sanchez, L. (1986). Mutations Affecting Dosage Compensation In <u>Drosophila</u> <u>melanogaster</u>: Effect On The Germline. Dev.Biol. 118: 379-384.
- Baker, B.S. (1989). Sex In Flies: The Spice Of Life. Nature 340: 521-524.
- Baker, B.S. and Belote, J.M. (1983). Sex Determination And Dosage Compensation In <u>Drosophila</u> <u>melanogaster</u>. Ann.Rev.Genet. 17: 345-393.
- Baker, B.S., Nagoshi, R.N. and Burtis, K.C. (1987). Molecular Genetic Aspects Of Sex Determination Of <u>Drosophila</u>. BioEssays 6: 66-70.
- Baker, B.S. and Ridge, K.A. (1980). Sex And The Single Cell I: On The Action Of Major Loci Affecting Sex Determination In <u>Drosophila melanogaster</u>. Genetics 94: 383-423.

Baker, B.S. and Wolfner, M.F. (1987). A Molecular Analysis Of <u>doublesex</u> A Bifunctional Gene That Controls Both Male And Female Sexual Differentiation In <u>Drosophila melanogaster</u>. Genes and Dev. 2: 477-489.

Bell, A.E. (1954).
A Gene In Drosophila That Produces All Male Progeny.
Genetics 39: 958-959.

- Bellen, H.J., O'Kane, C.J., Wilson, C., Grossniklaus, U., Pearson, R.K. and Gehring, W.J. (1989). P-Element-Mediated Enhances Detection, A Versatile Method To Study Development In <u>Drosophila</u>. Genes and Dev. 3: 1288-1300.
- Bell, L.R., Maine, E.M., Schedl, P. and Cline, T.W. (1988). <u>Sex-lethal A Drosophila</u> Sex Determination Switch Gene Exhibits Sex-Specific RNA Splicing And Sequence Similarity To RNA Binding Proteins. Cell 55: 1037-1046.
- Belote, J.M. and Baker, B.S. (1982). Sex Determination In <u>Drosophila melanogaster</u>: Analysis Of <u>transformer-2</u> A Sex Transforming Locus. PNAS 79: 1568-1572.
- Belote, J.M. and Baker, B.S. (1983).
  The Dual Function Of A Sex Determination Gene In Drosophila
  melanogaster.
  Dev.Biol. 95: 512-517.
- Belote, J.M. and Baker, B.S. (1987). Sexual Behaviour: It's Control During Development And Adulthood In <u>Drosophila melanogaster</u>. PNAS 84: 8026-8030.

- Belote, J.M., Handler, A.M., Wolfner, M.F., Livak, K.J. and Baker, B.S. (1985). Sex-specific Regulation Of Yolk Protein Gene Expression In Drosophila melanogaster. Cell 40: 339-348.
- Belote, J.M. and Lucchesi, J.C. (1980). Male Specific Lethal Mutations Of <u>Drosophila melanogaster</u>. Genetics 96: 165-186.

Belote, J.M., McKeown, M., Boggs, R.T., Ohkawa, R. and Sosnowski, B.A. (1989). Molecular Genetics Of <u>transformer</u>, A Genetic Switch Controlling Sexual Differentiation In <u>Drosophila</u>. Dev.Genet. 10: 143-154.

- Benton, W.D. and Davis, R.W. (1977). Screening Lambda gt Recombinant Clones By Hybridisation To Single Plaques <u>in-situ</u>. Science 196: 180-182.
- Berger, S.L. (1987). Quantifying <sup>32</sup>P Labelled And Unlabelled Nucleic Acids. In Methods In Enzymology 152, Berger, S.L. and Kimmel, A.R. : Academic Press.
- Berger, S.L., Wallace, D.M., Puskas, R.S. and Eschenfeldt, W.H. (1983). Reverse Transcriptase And Its Associated Ribonuclease H : Interplay Of Two Enzyme Activities Controls The Yield Of Single-Stranded cDNA. Biochem. 22: 2365-2372.
- Biggen, M.D. and Tjian, R. (1988) Transcription Factors That Activate The <u>Ultrabithorax</u> Promoter In Developmentally Staged Extracts. Cell 53: 699-711
- Birnboim, H.C. and Doly, J. (1979). Rapid Alkaline Extraction Procedure For Screening Recombinant Plasmid DNA. NAR 7: 1513-1523.

Boggs, R.T., Gregor, P., Idriss, S., Belote, J.M. and McKeown, M. (1987). Regulation Of Sexual Differentiation In <u>Drosophila</u> <u>melanogaster</u> Via Alternative Splicing Of RNA From The <u>transformer</u> Gene. Cell 50: 739-747.

- Bonaccorsi, S., Pisano, C., Puoti, F. and Gatti, M. (1988).
  Y Chromosome Loops In <u>Drosophila melanogaster</u>.
  Genetics 120: 1015-1034.
- Boswell, R.E. and Mahawold, A.P. (1985). <u>tudor</u>, A Gene Required For Assembly Of The Germ Plasm In <u>Drosophila melanogaster</u>. Cell 43: 97-104.
- Bownes, M. and Nothiger, R. (1981). Sex Determining Genes And Vitellogenin Synthesis In <u>Drosophila melanogaster</u>. Mol.Gen.Genet. 182: 222-228.
- Brand, R.C. and Hennig, W. (1989). An Abundant Testis RNA Species Shows Sequence Similarity To Y Chromosomal And Other Genomic Sites In <u>Drosophila hydei</u>. MGG 215: 469-477.
- Braun, R.E., Peschon, J.J., Behringer, R.R., Brinster, R.L. and Palmiter, R.D. (1989). Protamine 3' Untranslated Sequences Regulate Temporal Translational Control And Subcellular Localisation Of Growth Hormone In Spermatids Of Transgenic Mice. Genes & Dev 3: 739-802.

Brennan, M.D., Weiner, A.J., Goralski, T.J. and Mahawold, A.P. (1982). The Follicle Cells Are Major Sites Of Vitellogenin Synthesis In <u>Drosophila melanogaster</u>. Dev.Biol. 89: 225-236. Bridges, C.B. (1916). Non-Disjunction As Proof Of The Chromosomal Theory Of Heredity. Genetics 1: 1-52, 107-163.

- Bridges, C.B. (1921).
  The Origin Of Variations In Sexual And Sex-Limited
  Characteristics.
  Am.Naturalist 56: 51-63.
- Brittnacher, J.G. and Ganetzky, B. (1983). On The Components Of Segregation Distortion In <u>Drosophila</u> <u>melanogaster</u>: II Deletion Mapping And Dosage Analysis Of The <u>Sd</u> Locus. Genetics 103: 659-673.
- Brittnacher, J.G. and Ganetzky, B. (1984). On The Components Of Segregation Distortion In Drosphila melanogaster: III Nature Of <u>Enhancer Of Sd</u>. Genetics 107: 423-434.
- Brittnacher, J.G. and Ganetzky, B. (1989). On The Components Of Segregation Distortion In <u>Drosophila</u> <u>melanogaster</u>: IV Construction Of Free Duplications For The <u>Rsp</u> Locus. Genetics 121: 739-750.
- Brown, E.H. and King, R.C. (1960). Studies On The Expression Of The transformer Gene Of <u>Drosophila melanogaster</u>. Genetics 46: 143-156.
- Burtis, K.C. and Baker, B.S. (1989). <u>Drosophila doublesex</u> Gene Controls Somatic Sexual Differentiation By Producing Alternatively Spliced mRNAs Encoding Related Sex-specific Polypeptides. Cell 56: 997-1010.
- Busson, B., Gans, M., Komitopoulou, K. and Masson, M. (1983). Genetic Analysis Of Three Dominant Female Sterile Mutations Located On The X-chromosome Of <u>Drosophila melanogaster</u>. Genetics 105: 309-325.

Butler, B., Pirrotta, V., Irminger-Finger, I. and Nothiger, R. (1986). The Sex Determination Gene <u>transformer</u> Of <u>Drosophila</u>, Molecular Cloning And Transformation Studies EMBO 5: 3607-3613.

Campos-Ortega, J. and Hartenstein, V. (1985).
The Embryonic Development Of <u>Drosophila</u> Melanogaster.
: Springer-Verlag.

- Cattanach, B.M. (1975). Control Of Chromosome Inactivation. Ann.Rev.Gen. 9: 1-18.
- Caudy, M., Vassin, H., Brand, M., Tuma, R. and Jan, Y.N. (1988). <u>daughterless</u>, A <u>Drosophila</u> Gene Essential For Both Neurogenesis And Sex Determination Has Sequence Similarities To myc And <u>Achaete-scute</u> Complex. Cell 55: 1061-1067.
- Cavener, D., Corbett, G., Cox, D. and Whetten, R. (1985). Isolation Of The Eclosion Gene Cluster And The Developmental Expression Of The gld Gene In <u>Drosophila melanogaster</u>. EMBO 5: 2939-2948.
- Chandra, S.H. (1985). Sex Determination A Hypothesis Based On Non-coding DNA. PNAS 82: 1165-1169.
- Chapman, K.B. and Wolfner, M.F. (1988). Determination Of Male-specific Gene Expression In Drosophila Accessory Glands. Dev.Biol. 126: 195-202.
- Chen, C., Malone, T., Beckendorf, S.K. and Davis, R.W. (1987). At Least Two Genes Reside Within A Large Intron Of The <u>dunce</u> Gene Of <u>Drosophila</u>. Nature 329: 721-724.

- Chen, P.S., Stumm-Zollinger, E., Aigaki, T., Balmer, J., Bienz, M. and Bohlen, P. (1988). A Male Accessory Gland Peptide That Regulates Reproductive Behaviour Of Female <u>Drosophila melanogaster</u>. Cell 54: 291-298.
- Chirgwin, J.M., Przybyla, A.E., MacDonald, R.J. and Rutter, W.J. (1979). Isolation Of Biologically Active RNA From Sources Enriched In Ribonuclease. Biochem. 24: 5294-5299.
- Chomczynski, P. and Sacchi, N. (1987). Single Step Method Of RNA Isolation By Acid Guanidinium Thiocyanate-Phenol-Chloroform Extraction. Anal. Biochem. 162: 156-159.

Cline, T.W. (1976).

A Sex Specific , Temperature Sensitive Maternal Effect Of The <u>daughterless</u> Mutation Of <u>Drosophila melanogaster</u>. Genetics 84: 723-742.

Cline, T.W. (1978).

Two Closely Linked Mutations In <u>Drosophila</u> That Are Lethal To Opposite Sexes And Interact With <u>daughterless</u>. Genetics 90: 683-698.

Cline, T.W. (1979).

A Male-Specific Lethal Mutation In <u>Drosophila</u> That Transforms Sex.

Dev.Biol. 72: 266-275.

Cline, T.W. (1980).

Maternal And Zygotic Sex Specific Gene Interactions In Drosophila melanogaster. Genetics 96: 903-926.

Cline, T.W. (1983).

The Interaction Between <u>daughterless</u> And <u>Sex-lethal</u> In Triploids: A Lethal Sex Transforming Maternal Effect Linking Sex Determination And Dosage Compensation In Droophila. Dev.Biol. 95: 260-274. Cline, T.W. (1984). Autoregulatory Function Of A Drosophila Gene Product That Establishes And Maintains The Sexually Determined State. Genetics 107: 231-277. Cline, T.W. (1986). A Female-specific Lethal Lesion In An X-Linked Positive Regulator Of The Drosophila Sex Determination Gene Sex-lethal. Genetics 113: 641-663. Cline, T.W. (1988). Evidence That sis-a And sis-b Are Two Of Several Discrete "Numerator Elements" Of The X:A Sex Determination Signal In Drosophila That Switch Sex-lethal Between two Alternate Stable Expression States. Genetics 119: 829-862. Contamine, D., Petijean, A.M. and Ashburner, A.M. (1989). In Press Genetics Cooper, K.W. (1964). Meiotic Conjunctive Elements Not Involving Chiasmata. PNAS 52: 1248-1255. Cronmiller, C. and Cline, T.W. (1987). The Drosophila Sex Determination Gene daughterless Has Different Functions In The Germline Versus Soma. Cell 48: 479-487. Cronmiller, C., Schedl, P. and Cline, T.W. (1988). Molecular Characterisation Of daughterless, A Drosophila Sex Determination Gene With Multiple Roles In Development.

Crosby, M.A. and Myerowitz, E.M. (1986). <u>Drosophila</u> Glue Gene <u>sgs3</u>: Sequences Required For Puffing And Transcriptional Regulation. Dev.Biol. 118: 593-607.

Genes & Dev. 2: 1666-1676.

- Das, C.C., Kaufmann, B.P. and Gay, H. (1964). Histone Protein Transition in <u>Drosophila melanogaster</u> I: Changes During Spermatogenesis. Exp.Cell.Res 35: 507-514.
- Dente, L., Sollazo, M., Baldari, C., Cesareni, G. and Cortese, R. (1985). The pEMBL Family Of Single-Stranded Vectors. In DNA Cloning I, Glover, D.M. : IRL Press.

Dezelee, S., Bras, F., Contamine, D., Lopez-Ferber, D.S. and Teninges, D. (1989). Molecular Analysis Of ref(2)P, A Drosophila Gene Implicated In Sigma Rhabdovirus Multiplication And Necessary For Male Fertility. EMBO 8: 3437-3446.

DiBenedetto, A.J., Lakich, D.M., Kruger, W.D., Belote, J.M., Baker, B.S. and Wolfner, M.F. (1987). Sequences Expressed Sex-specifically In <u>Drosophila</u> <u>melanogaster</u> Adults. Dev.Biol. 119: 242-251.

Dworkin, M.B. and Dawid, I.B. (1980). Use Of A Cloned Library For The Study Of Abundant PolyA+ mRNA During <u>Xenopus laevis</u> Development. Dev.Biol. 76: 449-464.

Engstrom, L., Caulton, J.H., Underwood, E.M. and Mahawold, A.P. (1982). Developmental Lesions In The agametic Mutant Of Drosophila melanogaster. Dev.Biol. 91: 163-170.

Ethan, B., Vaessin, H., Shepherds, S., Lee, K., McCall, K., Barbels, S., Ackerman, L., Carretto, R., Uemura, T., Grell, E., Jan, L.Y. and Jan, Y.N. (1989). Searching For Pattern And Mutation In The Drosophila Genome With P-LacZ Vectors. Genes and Dev. 3: 1273-1287.

- Feinberg, A.P. and Vogelstein, B. (1983).
  A Technique For Radiolabelling DNA Restriction Fragments To
  High Specific Activity.
  Anal.Biochem. 132: 6-13.
- Fielding, C.J. (1967).
  Developmental Genetics Of The Mutant grandchildless Of
  Drosophila subobscura.
  J.Embryol.Exp.Morphol. 17: 375-384.
- Fujihara, T., Kawabe, M. and Oishi, K. (1978).
  A Sex Transformation Gene In Drosophila melanogaster.
  J.Heredity 69: 229-236.
- Fung, S.T.C. and Gowen, J.W. (1957).
  The Developmental Genetics Of A Sex-Limited Gene In
  Drosophila melanogaster.
  J.Exptl.Zool. 134: 515-532.
- Fyrberg, E.A., Bond, B.J., Hershey, N.D., Mixter, K.S. and Davidson, N. (1981). The Actin Genes Of <u>Drosophila</u> Are Highly Conserved But Intron Positions Are Not. Cell 24: 107-116.
- Garabedian, M.J., Shepherd, B.M. and Wensink, P.C. (1986).
  A Tissue-specific Transcription Enhancer From The Drosophila
  Yolk Protein 1 Gene.
  Cell 45: 859-867.
- Garcia-Bellido, A. (1964). Beziehungen Zwischen Vermerungshachstum Und Differenzierung Von Mannlichen Keimzellen Von Drosophila melanogaster. Wilhelm Roux Arch. 155: 594-610.
- Gatti, M., Pimpinelli, S. and Santini, G. (1976). Characterisation Of <u>Drosophila</u> Heterochromatin I: Staining And Decondensation With Hoechest 33258 And Quinacrine. Chromasoma 57: 351-375.

- Gergen, P.J. (1987). Dosage Compensation In <u>Drosophila</u>: Evidence That <u>daughterless</u> And <u>Sex-lethal</u> Control X Chromosome Activity At The Blastoderm Stage Of Embryogenesis. Genetics 117: 477-485.
- Gergen, P.J. and Wieschaus, E. (1987). Dosage Requirements For <u>Runt</u> Expression In The Segmentation Of <u>Drosophila</u> Embryos. Cell 45: 289-299.
- Goralski, T., Edstrom, J.E. and Baker, B.S. (1989). The Sex Determination Locus <u>transformer-2</u> Of <u>Drosophila</u> Encodes A Polypeptide With Similarity To RNA Binding Proteins. Cell 56: 1011-1018.
- Gubler, U. and Hoffman, B.J. (1983).
  A Simple And Very Efficient Method For Generating cDNA
  Libraries.
  Gene 25: 263-269.
- Guraya, S.S. (1987).
  Biology Of Spermatogenesis And Spermatozoa In Mammals.
  : Springer-Verlag.
- Haraven, D., Zuckerman, M. and Lifschytz, E. (1986). Origin And Evolution Of The Transcribed Repeated Sequences Of The Y Chromosome Lampbrush Loops Of <u>Drosophila hydei</u>. PNAS 83: 125-129.
- Hardy, R.W., Lindsley, D.L., Livak, K.J., Lewis, B., Siversten, A., Joslyn, G., Edwards, J. and Bonaccorsi, S. (1984). Cytogenetic Analysis Of A Segment Of The Y Chromosome Of <u>Drosophila melanogaster</u>. Genetics 107: 591-610.
- Hardy, R.W., Tokuyasu, K.T. and Lindsley, D.L. (1981). Analysis Of Spermatogenesis In <u>Drosophila melanogaster</u> Bearing Deletions For Y Chromosome Fertility Genes. Chromosoma 83: 593-617.

Hauschteck-Jungen, E. and Hartl, D. (1982). Defective Histone Transition During Spermiogenesis In Heterozygous <u>Segregation Distorter</u> Males Of <u>Drosophila</u> <u>melanogaster</u>. Genetics 101: 57-69.

Helling, R.B., Goodman, H.M. and Boyer, H.W. (1974). Analysis Of The <u>R.Ecoli</u> Fragments Of DNA From The Lambdoid Bacteriophage And Other Viruses By Agarose Gel Electrophoresis. J.Virol. 14: 1235-1244.

Hennig, W. (1985).

Y Chromosome Function And Spermatogenesis In <u>Drosophila hydei</u>. Adv.In Genet. 23: 179-234.

Hildreth, P.E. (1965).

<u>doublesex</u> A Recessive Gene That Transforms Males And Females Of <u>Drosophila</u> Into Intersexes. Genetics 51: 659-678.

Hilliker, A.J. (1976).

Genetic Analysis Of The Centromeric Heterochromatin Of Chromosome 2 Of <u>Drosophila melanogaster</u>: Deficiency Mapping Of EMS Induced Lethal Complementation Groups. Genetics 83: 765-782.

Hofmann, A. and Korg, G. (1987).

Upstream Sequences Of Dosage Compensated And Non-compensated Alleles Of The Larval Secretion Protein <u>sgs4</u> In <u>Drosophila</u>. Chromosoma 96: 1-7.

Holmes, D.S. and Quigley, M. (1981).
A Rapid Boiling Method For The Preparation Of Bacterial
Plasmids.

Anal. Biochem. 114: 193-197.

Hough-Evans, B.R., Jacobs-Lorena, M., Cummings, M.R., Britten, R.J. and Davidson, E.H. (1980). Complexity Of RNA In Eggs Of <u>Drosophila melanogaster</u> and <u>Musca domestica</u>. Genetics 95: 81-94. Huijser, P., Kirchhoff, C., Lankenau, D.H. and Hennig, W. (1988). Retrotransposon-Like Sequences Are Expressed In Y Chromosomal Lampbrush Loops Of <u>Drosophila hydei</u>. J.Mol.Biol. 203: 689-697.

- Hung, M.C. and Wensink, P.C. (1981). The Sequence Of The <u>Drosophila melanogaster</u> Gene For Yolk Protein 1. NAR 9: 6407-6419.
- Huynh, T.V., Young, R.A. and Davis, R.W. (1985). Constructing And Screening cDNA Libraries In Lambda gt10 And gt11. In DNA Cloning I, Glover, D.M. : IRL Press.
- Kaiser, K., Furia, M. and Glover, D.M. (1986). Dosage Compensation At The sgs4 Locus Of Drosophila <u>melanogaster</u>. J.Mol.Biol. 187: 529-536.
- Kaiser, K. and Murray, N.E. (1985). The Use Of Lambda Replacment Vectors In The Construction Of Representative Genomic DNA Libraries. In DNA Cloning I, Glover, D.M. : IRL Press.
- Kemphues, K.J., Kaufman, T.C., Raff, R.A. and Raff, E.C. (1982). The Testis-Specific B-Tubulin Subunit Has Multiple Functions In Spermatogenesis. Cell 31: 655-670.

Kennison, J.A. (1981).
The Genetic And Cytological Organisation Of The Y Chromosome
Of Drosophila melanogaster.
Genetics 98: 529-548.

Kennison, J.A. (1983).

Analysis Of Y-Linked Mutations To Male Sterility in Drosophila melanogaster. Genetics 103: 219-234. Keris, J. (1931).
The Growth Of The Gonads In Drosophila melanogaster.
Genetics 16: 212-224.

- Kloetzel, P.M., Knust, E. and Schwochau, M. (1981). Analysis Of Nuclear Proteins In Primary Spermatocytes Of <u>Drosophila hydei</u> The Correlation Of Nuclear Proteins With The Function Of The Y Chromosome Loops. Chromosoma 84: 67-86.
- Krainer, A.R. and Maniatis, T. (1988). RNA Splicing. In Transcription And Splicing, Hames, D.B. and Glover, D.M. : IRL Press.
- Kuhn, R., Schafer, U. and Schafer, M. (1988). Cis-Acting Regions Sufficient For Spermatocyte-Specific Transcriptional And Spermatid-Specific Translational Control Of The <u>Drosophila melanogaster</u> Gene <u>mst(3)gl9</u>. EMBO 7: 447-454.
- Miklos, G.L.G. (1985) Localised Highly Repetitive DNA Sequences In Vertebrate And Invertebrate Genomes. In Molecular Evolutionary Genetics, MacIntyre, R.J. :Plenum Press
- Lasko, P.F. and Ashburner, M.A. (1988). The Product Of The <u>Drosophila</u> Gene <u>vasa</u> Is Very Similar To Eukaryotic Translation Initiation Factor 4A. Nature 335: 611-617.

Lauge, G. (1982).
Development Of The Genetalia And Analia.
In Handbook Of Drosophila Development, Ransom, R.
:Elsevier Press.

Lehmann, R.E. and Nusslein-Volhard, C. (1986).
Abdominal Segmentation, Pole Cell Formation And Embryonic
Polarity Require The Localised Activity Of A Maternal Gene In
Drosophila.
Cell 47: 141-152.

- Leroy, P., Alzari, P., Sasoon, D., Wolgemuth, D. and Fellous, M. (1989). The Protein Encoded By A Murine Male Germline-Specific Transcript Is A putative ATP-Dependant Helicase. Cell 57: 549-559.
- Lifschytz, E. (1987). The Genetic Control Of Spermatogenesis. Int.Rev.Cyt. 109: 211-258.
- Lifschytz, E., Haravan, D., Azriel, A. and Brodsly, H. (1983).
  DNA Clones And RNA Transcripts Of Four Lampbrush Loops From
  The Y Chromosome Of <u>D.Hydei</u>.
  Cell 32: 191-199.
- Lifschytz, E. and Haraven, D. (1977). Gene Expression And The Control Of Spermatid Morphogenesis In <u>Drosophila melanogaster</u>. Dev.Biol. 58: 276-294.
- Lindsley, D.L. and Grell, E.H. (1968). Genetic Variations Of <u>Drosophila melanogaster</u>. : Carnegie Institute Of Washington.
- Lindsley, D.L. and Lifschytz, E. (1972).
  The Genetic Control Of Spermatogenesis In Drosophila.
  In The Genetics Of Spermatozoon, Beatty, R.A. and GluecksohnWaelsch, S. : Boytrykkeriet Forum Copenhagen.
- Lindsley, D.L. and Tokuyasu, K.T. (1980).
  Spermatogenesis.
  In The Genetics And Biology Of Drosophila, Ashburner, M.A.
  and Wright, T.R.F. : Academic Press.
- Lindsley, D.L. and Zimm, G. (1985). The Genome Of <u>Drosophila melanogaster</u>. : Dros.Inf.Ser. 62.

- Livak, K.J. (1984). Organisation And Mapping Of A Sequence On The <u>Drosophila</u> <u>melanogaster</u> X And Y Chromosomes That Is Transcribed During Spermatogenesis. Genetics 107: 611-634.
- Lucchesi, J.C. and Manning, J.E. (1987). Gene Dosage Compensation In <u>Drosophila melanogaster</u>. Adv.Genet. 24: 371-430.
- Lucchesi, J.C. and Skripsky, T. (1981). The Link Between Dosage Compensation And Sex Determination In <u>Drosophila melanogaster</u>. Chromosoma 82: 217-227.

Lyttle, T.W. (1989). The Effect Of Novel Chromosome Position And Variable Dose On The Genetic Behavior Of The <u>Responder</u> Element Of The <u>Segregation Distorter</u> System Of <u>Drosophila melanogaster</u>. Genetics 121: 751-763.

- Lyttle, T.W., Brittnacher, J.G. and Ganetzky, B. (1986). Detection Of <u>Rsp</u> And Modifier Variation In The Meiotic Drive System <u>Segregation Distorter</u> (SD) Of Drosphila melanogaster. Genetics 114: 183-202.
- Mackenzie, C., Stewart, B. and Kaiser, K. (1989).
  A Device For Creating Orderd Arrays Of Plaques Or Colonies.
  Technique 1: 49-52.

Mahawold, A.P. and Kambysellis, M.P. (1980). Oogenesis. In The Genetics And Biology Of <u>Drosophila</u>, Ashburner, M.A. and Wright, T.F.R. : Academic Press.

Maine, E.M., Salz, H.K., Schedl, P. and Cline, T.W. (1985a). <u>Sex-lethal</u> A Link Between Sex Determination and Sexual Differentiation in <u>Drosophila melanogaster</u>. CSHSQB 50: 595-604.

- Maine, E.M., Salz, H.K., Cline, T.W. and Schedl, P. (1985b). The <u>Sex-lethal</u> Gene Of <u>Drosophila</u>: DNA Alterations Associated With Sex-Specific Lethal Mutations. Cell 43: 521-529.
- Mange, A.P. and Sandler, L. (1973).
  A Note On The Maternal Effect Mutants <u>daughterless</u> And <u>abnormal oocyte</u> In <u>Drosophila melanogaster</u>.
  Genetics 73: 73-86.
- Maniatis, T., Fritsch, E.F. and Sambrook, J. (1982). Molecular Cloning: A Laboratory Manual. : CSH.
- Maniatis, T., Hardison, R.C., Lacy, E., Laver, J., OConnell, C., Quon, D., Sim, G.K. and Efstradiatis, A. (1978). Isolation Of Structural Genes From Libraries Of Eukaryotic DNA. Cell 15: 687-701.
- Marsh, J.L. and Wieschaus, E. (1978). Is Sex Determination In The Germline And Soma Controlled By Different Mechanisms ? Nature 272: 249-251.
- McKeown, M., Belote, J.M. and Baker, B.S. (1987).
  A Molecular Analysis Of transformer A Gene In Drosophila
  melanogaster That Controls Female Sexual Differentiation.
  Cell 48: 489-499.
- McKeown, M., Belote, J.M. and Boggs, R.T. (1988). Ectopic Expression Of The Female transformer Gene Product Leads To Female Differentiation Of Chromosomally Male <u>Drosophila</u>. Cell 53: 887-895.
- McRobert, S.P. and Tompkins, L. (1985). The Effect Of <u>transformer</u>, <u>doublesex</u> And <u>intersex</u> Mutations On The Sexual Behaviour Of <u>Drosophila melanogaster</u>. Genetics 111: 89-96.
Mevel-Ninio, M., Mariol, M.C. and Gans, M. (1989). Mobilisation Of The gypsy And copia Retrotransposons In <u>Drosophila melanogaster</u> Induces Revertion Of The <u>ovoD</u> Dominant Female Sterile Mutation: Molecular Analysis Of Revertant Alleles. EMBO 8: 1549-1558.

Michiels, F., Gasch, A., Kaltschmidt, B. and Renkawitz-Pohl, R. A 146bp Promoter Element Directs The Testis Specificity Of The <u>Drosophila</u> B2-Tubulin Gene. 1989 EMBO 8: 1559-1565.

- Monsma, S.A. and Wolfner, M.F. (1988). Structure And Expression Of A <u>Drosophila</u> Male Accessory Gland Gene Whose Product Resembles A Peptide Pheromone Precursor. Genes and Devl. 2: 1063-1073.
- Morgan, T.H., Redfield, H. and Morgan, L.V. (1943). Maintainance Of A <u>Drosophila</u> Stock Center In Connection With Investigations On The Constitution Of Germinal Material In Relation To Heredity. Carnegie Inst.Wash.Yearbook 42: 171-174.

Murray, N.E. (1983).
Phage Lambda And Molecular Cloning.
In Lambda II, Hendrix, R.W., Roberts, J.W., Stahl, F.W. and
Weisberg, R.A. : CSH.

Murre, C., McCaw, P.S. and Baltimore, D. (1989a).
A New DNA Binding And Dimerisation Motif In Immunoglobulin
Enhancer Binding, <u>daughterless</u> and myc Proteins.
Cell 56: 777-783.

Murre, C., McCaw, P.S., Vaessin, H., Caudy, M., Yan, L.Y., Yan, Y.N., Cabrera, C.V., Buskin, J.N., Hauschka, S.D., Lasser, A.B., Weintraub, H. and Baltimore, D. (1989b). Interactions Between Heterologus Helix-Loop-Helix Proteins Generate Complexes That Bind Specifically To A Common DNA Sequence. Cell 58: 537-544.

- Muskavitch, M.A.T. and Hogness, D.S. (1980). Molecular Analysis Of A Gene In A Developmentaly Regulated Puff Of <u>Drosophila melanogaster</u>. PNAS 77: 7362-7366.
- Nagoshi, R.N., McKeown, M., Burtis, K.C., Belote, J.M. and Baker, B.S. (1988). The Control Of Alternative Splicing At Genes Regulating Sexual Differentiation In <u>Drosophila melanogaster</u>. Cell 53: 229-236.
- Niki, Y. and Okada, M. (1981). Isolation And Characterisation Of <u>grandchildless-Like</u> Mutants In <u>Drosophila melanogaster</u>. Wilhelm Rouxs Arch. 190: 1-10.
- Nothiger, R., Dudendorfer, A. and Epper, F. (1977). Gynandromorphs Reveal Two Separate Primordia For Male And Female Genetalia In <u>Drosophila melanogaster</u>. Wilhelm Rouxs Arch. 181: 367-373.
- Oakshott, J.G., Collet, C., Phillis, R.W., Nielson, K.M., Russell, R.T., Chambers, G.K., Ross, V. and Richmond, R.C. (1987). Molecular Cloning And Characterisation Of Esterase-6, A Serine Hydrolase Of <u>Drosophila</u>. PNAS 84: 3359-3363.
- O'Kane, C.J. and Gehring, W.J. (1987). Detection <u>in-situ</u> Of Genomic Regulatory Elements In <u>Drosophila</u>. PNAS 84: 9123-9127.
- Oliver, B., Perrimon, N. and Mahawold, A.P. (1987). The <u>ovo</u> Locus Is Required For Sex-specific Germline Maintainance In <u>Drosophila</u>. Genes & Devl. 1: 913-923.
- Oliver, B., Perrimon, N. and Mahowald, A.P. (1988). Genetic Evidence That The <u>san-fille</u> Locus Is Involved In <u>Drosophila</u> Sex Determination. Genetics 120: 159-171.

- Olivieri, G. and Olivieri, A. (196). Autoradiographic Study Of Nuclic Acid Synthesis During Speramtogenesis In <u>Drosophila</u> Melanogaster. Mutation Res. 2: 366-380.
- Ota, T., Fukunaga, A., Kawabe, M. and Oishi, K. (1981). Interactions Between Sex-transforming Mutants Of <u>Drosophila</u> <u>melanogaster</u> I: Hemolymph Vitellogenins And Gonad Morphology. Genetics 99: 429-441.
- Pardue, M.L. (1986). In-Situ Hybridisation To DNA Of Chromosomes And Nuclei. In <u>Drosophila</u>: A Practical Approach, Roberts, D.B. : IRL Press.
- Perrimon, N. (1984). Clonal Analysis Of Dominant Female Sterile Germline Dependant Mutations In <u>Drosophila melanogaster</u>. Genetics 108: 927-939.
- Peschon, J.J., Behringer, R.R., Brinster, R.L. and Palmiter, R.D. (1987). Spermatid Specific Expression Of Protamine 1 In Transgenic Mice. PNAS 84: 5316-5319.
- Pierre, A.M. (1972). New Mutants Report. Dros.Inf.Ser. 48: 16.

Rhaman, R. and Lindsley, D.L. (1980). Male Sterilising Interactions Between Duplications And Deficiencies For Proximal X Chromosome Material In <u>Drosophila</u> <u>melanogaster</u>. Genetics 90: 40-64.

Rigby, P.W.J., Dieckmann, M., Rhodes, C. and Berg, P. (1977). Labelling DNA To High Specific Activity In-Vitro By Nick Translation With DNA PolymeraseI. J.Mol.Biol. 113: 237-251. Rio, D. (1988)
Accurate And Efficient Pre-mRNA Splicing In Cell Free
Extracts.
PNAS. 85: 2904-2908

- Salz, H.K., Cline, T.W. and Schedl, P. (1987). Functional Changes Associated With Structural Alterations Induced By Mobilisation Of A P-Element Inserted In The <u>Sex-</u> <u>lethal</u> Gene Of Drosphila. Genetics 117: 221-231.
- Salz, H.K., Maine, E.M., Keyes, L.N., Samuels, M.E., Cline, T.W. and Schedl, P. (1989). The <u>Drosophila</u> Female-Specific Sex-Determination gene <u>Sex-</u> <u>lethal</u> Has Stage, Tissue and Sex Specific RNA's Suggesting Multiple Modes Of Regulation. Genes and Dev. 3: 708-719.
- Sanchez, L. and Nothiger, R. (1982). Clonal Analysis Of <u>Sex-lethal</u> A Gene Needed For Female Sexual Development In <u>Drosophila melanogaster</u>. Wilhelm Rouxs Arch. 191: 211-214.

Sanger, F., Coulson, A.R., Hong, G.F., Hill, D.F. and Petersen, G.B. (1982). Nucleotide Sequence Of Bacteriophage Lambda DNA. J.Mol.Biol. 162: 729-773.

Sargent, T.W. (1987).
Isolation Of Differentialy Expressed Genes.
In Methods In Enzymology 152, Berger, S.L. and Kimmel, A.R.
: Academic Press.

Schafer, U. (1986a).
Genes For Male-specific Transcripts In Drosophila
melanogaster.
Mol.Gen.Genet. 202: 219-225.

Schafer, U. (1986b).
The Regulation Of Male-Specific Transcripts By Sex
Determining Genes In Drosophila melanogaster.
EMBO 5: 3579-3582.

- Schalet, A. and Lefevre, G. (1973).
  The Localisation Of The Ordinary Sex Linked Genes In Section
  2D Of The Polytene X-Chromosome Of <u>Drosophila melanogaster</u>.
  Chromosoma 44: 183.
- Schalet, A. and Lefevre, G. (1976).
  The Proximal Region Of The X Chromosome.
  In The Genetics And Biology Of Drosophila, Ashburner, M.A.
  and Novitsky, E. : Academic Press.
- Scherer, G., Tschudi, C., Perara, J., Delius, H. and Pirrotta, V. (1982). <u>B104</u> A Newly Dispersed Gene Family In <u>Drosophila melanogaster</u> And Its Analogies With Retroviruses. J.Mol.Biol. 157: 435-451.
- Schulz, R.A. and Butler, B.A. (1989).
  Overlapping Genes Of <u>Drosophila melanogaster</u>: Organisation Of
  The <u>z600-gonadal-Eip28/29</u> Gene Cluster.
  Genes & Dev. 3: 232-242.

Schupbach, T. (1982).

Autosomal Mutations That Interfere With Sex Determination In Somatic Cells Have No Effect On The Germline. Dev.Biol. 89: 117-127.

**Schupbach**, **T**. (1985).

Normal Female Germ Cell Development Requires The Female X Chromosome To Autosome Ratio And Expression Of <u>Sex-lethal</u> In <u>Drosophila melanogaster</u>. Genetics 109: 529-548.

Schupbach, T., Wieschaus, E. and Nothiger, R. (1978).
The Embryonic Organisation Of The Genital Disk Studied In
Genetic Mosaics Of Drosophila melanogaster.
Wilhelm Rouxs Arch. 185: 249-270.

Scott, M.P. (1987). Complex Loci Of <u>Drosophila</u>. Ann.Rev.Biochem. 56: 195-227. Seidel, S. (1963).
Experimentelle Untersuchungen Uber Die Grundlagen Der
Sterilitat Von transformer Mannchen Bei Drosophila
melanogaster.
Wilhelm Rouxs Arch.Z.Vererbungsl. 94: 215-241.

- Sharp, C.B., Hilliker, A.J. and Holm, D.G. (1985).
  Further Characterisation Of Genetic Elements Associated With
  The Segregation Distorter Phenomenon Of Drosophila
  melanogaster.
  Genetics 110: 671-688.
- Shepherd, B., Garabedian, M.J., Hung, M.C. and Wensink, P.C. (1985). Developmental Control Of <u>Drosophila</u> Yolk Protein 1 Gene By Cis-acting DNA Elements. CSHSQB 50: 521-526.
- Skripsky, T. and Lucchesi, J.C. (1982). Intersexuality Resulting From The Interaction Of Sex-Specific Lethal Mutations In <u>Drosophila melanogaster</u>. Dev.Biol. 94: 153-162.

Sonnenblick, B.P. (1950).
The Early Embryogenesis Of Drosophila melanogaster.
In Biology Of Drosophila, Demerec, M. : Wiley & Sons.

- Sosnowski, B.A., Belote, J.M. and McKeown, M. (1989). Sex-Specific Alternative Splicing Of RNA From The <u>transformer</u> Gene Results From Sequence Dependant Splice Site Blockage. Cell 58: 449-459.
- Spradling, A.C. (1986).
  P-Element Mediated Transformation.
  In <u>Drosophila</u>: A Practical Approach, Roberts, D.B. : IRL
  Press.
- Spradling, A.C. and Rubin, G.M. (1981). <u>Drosophila</u> Genome Organisation: Conserved And Dynamic Aspects. Ann.Rev.Genet. 15: 219-264.

Steinmann-Zwicky, M. (1988).

Sex Determination In Drosophila: The X Chromosomal Gene liz
Is Required For Sxl Activity.
EMBO 7: 3889-3898.

Steinmann-Zwicky, M., Schmidt, H. and Nothiger, R. (1989). Cell Autonomous And Inductive Signals Can Determine The Sex Of The Germline Of <u>Drosophila melanogaster</u> By Regulating <u>Sexlethal</u>. Cell 57: 157-166.

Sturtevant, A.H. (1945).
A Gene In <u>Drosophila melanogaster</u> That Transforms Females
Into Males.
Genetics 30: 297-299.

Tischendorf, G., Liebrich, W., Trapitz, P., Wood, G. and Schwochau, M. (1989). A Group Of Non-Y Encoded <u>Drosophila hydei</u> Primary Spermatocyte Nuclear Glycoproteins Exhibit Epitopes Depending On Formation Of Y Chromosomal Giant Lampbrush Loops. Chromosoma 98: 144-152.

Tokunaga, C. (1962).

Cell Lineage And Differentiation Studied In The Male Fore-Leg Of <u>Drosophila melanogaster</u>. Dev.Biol. 4: 489-516.

Torres, M. and Sanchez, L. (1989).
The <u>scute</u> (T4) Gene Acts As A Numerator Element Of The X:A
Signal That Determines The State Of Activity Of <u>Sex-lethal</u> In
<u>Drosophila</u>.

EMBO 8: 3078-3086.

Trapitz, P., Wlaschek, M. and Bunemann, H. (1988). Structure And Function Of Y Chromosomal DNA II: Analysis Of Lambrush Loop Associated Transcripts In Nuclei Of Primary Spermatocytes Of <u>Drosophila hydei</u> By In-Situ Hybridisation Using Asymmetric RNA Probes From Different Families Of Repetitive DNA. Chromosoma 96: 159-170. Vinson, C.R., LaMarco, K.L., Johnson, P.F., Landschulz, W.H., and McKnight S.L. (1988) <u>In-situ</u> Detection Of Sequence-Specific DNA Binding Activity Specified By A Recombinant Bacteriophage. Genes & Dev. 2: 801-806

Vogt, P. and Hennig, W. (1986). Molecular Structure Of The Lambrush Loops Nooses Of The Y Chromosome Of <u>Drosophila hydei</u> II: DNA Sequences With Homologies To Multiple Genomic Locations Are Major Constituents Of The Loop. Chromosoma 94: 459-467.

Walker, E.S., Lyttle, T.W. and Lucchesi, J.C. (1989). Transposition Of The <u>Responder</u> Element Of The <u>Segregation</u> <u>Distorter</u> System To The X Chromosome In <u>Drosophila</u> <u>melanogaster</u>. Genetics 122: 81-86.

Watanabe, T.K. (1975).
A New Sex Transforming Gene On The Second Chromosome Of
Drosophila melanogaster.
Japn.J.Genet. 50: 269-271.

Whittaker, P.A., Campbell, A.J.B., Southern, E.M. and Murray, N.E. (1988). Enhanced Recovery And Restriction Mapping Of DNA Fragments Cloned In A New Lambda Vector. NAR 16: 6725-6728.

- Wieschaus, E. and Nothiger, R. (1982). The Role Of The <u>transformer</u> Gene In The Development Of The Genetalia And Analia of <u>Drosophila melanogaster</u>. Dev.Biol. 90: 320-334.
- Williams, J.G. (1981).
  The Preparation And Screening Of A cDNA Clone Bank.
  In Genetic Engineering I, Williamson, R. : Academic Press.

- Williams, J.G. and Lloyd, M.M. (1978). Changes In The Abundance Of Polyadenylated RNA During Slime Mould Development Measured Using Cloned Molecular Hybridisation Probes. J.Mol.Biol. 129: 19-35.
- Williamson, J.H. (1972). Allelic Complementation Between Mutants In The Fertility Factors Of The Y Chromosome In <u>Drosophila melanogaster</u>. Mol.Gen.Genet. 119: 43-47.
- Wilson, C., Pearson, R.K., Bellen, H.J., O'Kane, C.J., Grossniklaus, U. and Gehring, W.J. (1989). P-Element-Mediated Enhancer Detection: An Efficient Method For Isolating And Characterising Developmentally Regulated Genes In <u>Drosophila</u>. Genes and Dev. 3: 1301-1313.
- Wlaschek, M., Awgulewitsch, A. and Bunemann, H. (1988). Structure And Function Of Y Chromosomal DNA I: Sequence Organisation And Localisation Of Four Families Of Repetitive DNA On The Y Chromosome Of <u>Drosophila hydei</u>. Chromosoma 96: 145-158.
- Wu, C.I., Lyttle, T.W., Wu, M.L. and Lin, G.F. (1988). Association Between A Satellite DNA Sequence And The <u>Responder</u> Of <u>Segregation Distorter</u> in <u>Drosophila melanogaster</u>. Cell 54: 179-189.
- Yamamoto, M. and Miklos, G.L.G. (1977). Genetic Dissection Of Heterochromatin In <u>Drosophila</u>: The Role Of Basal X Heterochromatin In Meiotic Sex Chromosome Behaviour. Chromosome 60: 282-202

Chromosoma 60: 283-292.

Yanisch-Perron, C., Viera, J. and Messing, J. (1985). Improved M13 Phage Cloning Vectors And Host Strains. Gene 33: 103-119.



142